

## 8 Literaturverzeichnis

- [1] **Aaronson NK, Acquadro C, Alonso J et al. (1992)**  
International quality of life assessment (IQOLA) project.  
Qual Life Res 1 (5): 349-351
- [2] **Abholz HH, Pillau H**  
Anamnese, körperliche Untersuchung und Dokumentation.  
in: Kochen MM (Hrsg.): Allgemeinmedizin.  
Hippokrates Verlag, Stuttgart, 1992, S.110-116
- [3] **Achat H, Kawachi I, Levine S et al. (1998)**  
Social networks, stress and health-related quality of life.  
Qual Life Res 7 (8): 735-750
- [4] **Agréus L, Svärdsudd K, Nyrén O et al. (1995)**  
Irritable bowel syndrome and dyspepsia in the general population: overlap and lack of stability over time.  
Gastroenterol 109 (3): 671-680
- [5] **Agréus L, Svärdsudd K, Talley NJ (2001)**  
Natural history of gastroesophageal reflux disease and functional abdominal disorders: a population-based study.  
Am J Gastroenterol 96 (10): 2905-2914
- [6] **Allensbacher Jahrbuch der Demoskopie 1998 - 2002**  
Band 11, Noelle-Neumann E, Köcher R (Hrsg.)  
Verlag Sauer, München, 2002, S.233, 239, 257
- [7] **Allescher HD**  
Refluxkrankheit, Ösophagitis.  
In: Classen M, Diehl V, Kochsiek K: Innere Medizin.  
5. völlig überarbeitete Auflage, Verlag Urban & Fischer, München, 2004, S.1129-1136
- [8] **Arendt T, Liebe S**  
Chemische Ösophagusschäden.  
In: Classen M, Diehl V, Kochsiek K: Innere Medizin.  
5. völlig überarbeitete Auflage, Verlag Urban & Fischer, München, 2004, S.1137-1139
- [9] **Arendt T, Liebe S**  
Tumoren des Ösophagus.  
In: Classen M, Diehl V, Kochsiek K: Innere Medizin.  
5. völlig überarbeitete Auflage, Verlag Urban & Fischer, München, 2004, S.1139-1142
- [10] **Autorengruppe Psychosomatische Grundversorgung (1999)**  
Bado PSGV. Entwicklung und Erprobung einer Basisdokumentation für psychische, psychosoziale und psychosomatische Kontaktnlässe in der Primärversorgung.  
Psychotherapeut 44 (1): 46-50
- [11] **Balint M**  
Der Arzt, sein Patient und die Krankheit.  
10. Auflage, Verlag Klett-Cotta, Stuttgart, 2001
- [12] **Balint E, Norell JS**  
Fünf Minuten pro Patient. Eine Studie über die Interaktionen in der ärztlichen Allgemeinpraxis.  
10. Auflage, Suhrkamp Taschenbuch-Verlag, Frankfurt am Main, 1999
- [13] **Beach SR, Fincham FD, Katz J (1998)**  
Marital therapy in the treatment of depression: toward a third generation of therapy and research.  
Clin Psychol Rev, 18 (6): 635-661
- [14] **Becker S, Al Zaid K, Al Faris E (2002)**  
Screening for somatization and depression in Saudi Arabia: a validation study of the PHQ in primary care.  
Int J Psychiatry Med 32 (3): 271-283
- [15] **Bellach BM, Ellert U, Radoschewski M (2000)**  
Der SF-36 im Bundes-Gesundheitssurvey.  
Bundesgesundheitsbl Gesundheitsforsch Gesundheitsschutz 43 (3): 210-216

- [16] **Bellon JA, Delgado A, Luna JD et al. (1999)**  
Psychosocial and health belief variables associated with frequent attendance in primary care.  
Psychol Med 29 (6): 1347-1357
- [17] **Bennett EJ, Piesse C, Palmer K et al. (1998)**  
Functional gastrointestinal disorders: psychological, social, and somatic features.  
Gut 42 (3): 414-420
- [18] **Berardi D, Ceroni GB, Leggieri G et al. (1999)**  
Mental, physical and functional status in primary care attenders.  
Int J Psychiatry Med 29 (2): 133-148
- [19] **Bergmann E, Kamtsiuris P (1999)**  
Inanspruchnahme medizinischer Leistungen.  
Gesundheitswesen 61 (Sonderheft 2): 138-144
- [20] **Bergner M, Bobbitt RA, Carter WB et al. (1981)**  
The Sickness Impact Profile: development and final revision of a health status measure.  
Med Care 19 (8): 787-805
- [21] **Bertakis KD, Azari R, Helms LJ (2000)**  
Gender differences in the utilization of health care services.  
J Fam Pract 49 (2): 147-152
- [22] **Birchall H, Brandon S, Taub N (2000)**  
Panic in a general practice population: prevalence, psychiatric comorbidity and associated disability.  
Soc Psychiatry Psychiatr Epidemiol 35: 235-241
- [23] **Boyd EJS (1996)**  
The prevalence of esophagitis in patients with duodenal ulcer or ulcer-like dyspepsia.  
Am J Gastroenterol 91 (8): 1539-1543
- [24] **Brähler E**  
Körpererleben – ein vernachlässigter Aspekt der Medizin.  
In: Brähler E (Hrsg): Körpererleben. Ein subjektiver Ausdruck von Körper und Seele.  
Psychosozial-Verlag, Gießen, 1995, S. 3-18
- [25] **Brähler E, Felder H, Schumacher J (2003)**  
Wenn der Körper zur Last wird: Zum Einfluss von Alter, Geschlecht und Epoche auf körperliche Beschwerden.  
Psychosozial 26 (4): 105-120
- [26] **Brähler E, Gunzelmann T (1998)**  
30 Jahre Gießener Beschwerdebogen. Bibliographie zum GBB, GBB-KJ und GBB-24.  
Internet: <http://www.uni-leipzig.de/~medpsy>
- [27] **Brähler E, Möhring P**  
Der Körper im Beschwerdebild – Erfahrungen mit dem Gießener Beschwerdebogen.  
in: Brähler E (Hrsg): Körpererleben. Ein subjektiver Ausdruck von Körper und Seele.  
Psychosozial-Verlag, Gießen, 1995, S. 232-252
- [28] **Brähler E, Scheer JW**  
Der Gießener Beschwerdebogen. Handbuch.  
2. ergänzte und revidierte Auflage, Verlag Hans Huber, Bern, 1995
- [29] **Brähler E, Schumacher J, Brähler C (2000)**  
Erste gesamtdeutsche Normierung der Kurzform des Gießener Beschwerdeboogens GBB-24.  
Psychother Psychosom med Psychol 50 (1): 14-21
- [30] **Bühl A, Zöfel P**  
SPSS Version 10. Einführung in die moderne Datenanalyse unter Windows.  
Addison Wesley Verlag, München, 2000
- [31] **Bullinger M**  
Lebensqualität: Ein neues Bewertungskriterium für den Therapieerfolg.  
In: Pöppel E, Bullinger M, Härtel U: Kurzlehrbuch der medizinischen Psychologie.  
VCH-Verlag, Weinheim, 1994, S. 308-314

- [32] **Bullinger M**  
International validation and testing of quality of life scales in relation to Germany.  
In: Guggenmoos-Holzmann I, Bloomfield K et al. (Hrsg.):  
Quality of life and health: concepts, methods and applications.  
Blackwell-Wissenschafts-Verlag, Berlin 1995, S. 27-37
- [33] **Bullinger M (1997)**  
Gesundheitsbezogene Lebensqualität und subjektive Gesundheit.  
Psychother Psychosom Med Psychol 47 (3-4): 76-91
- [34] **Bullinger M, Kirchberger I**  
SF-36 Fragebogen zum Gesundheitszustand. Handanweisung.  
Hogrefe-Verlag, Göttingen, 1998
- [35] **Bullinger M, Kirchberger I, Ware J (1995)**  
Der deutsche SF-36 Health Survey.  
Z Gesundheitswiss 3 (1): 21-36
- [36] **Bundesverband der Arzneimittel-Hersteller e.V.**  
Der Arzneimittelmarkt in Deutschland in Zahlen. Unter besonderer Berücksichtigung der  
Selbstmedikation.  
15. Auflage, Krahe Druck GmbH, Unkel, 2002, Kapitel 11
- [37] **Candilis PJ, McLean RY, Otto MW et al. (1999)**  
Quality of life in patients with panic disorder.  
J Nerv Ment Dis 187 (7): 429-434
- [38] **Classen M**  
Erkrankungen der Magenschleimhaut.  
In: Classen M, Diehl V, Kochsiek K: Innere Medizin.  
5. völlig überarbeitete Auflage, Verlag Urban & Fischer, München, 2004, S.1143-1151
- [39] **Comberg HU**  
Definition der Allgemeinmedizin.  
In: Comberg HU, Klimm HD (Hrsg.): Allgemeinmedizin.  
3. überarbeitete Auflage, Georg Thieme Verlag, Stuttgart, 2001, S.1
- [40] **Coulehan JL, Schulberg HC, Block MR et al. (1997)**  
Treating depressed primary care patients improves their physical, mental and social functioning.  
Arch Intern Med 157 (10): 1113-1120
- [41] **Croudace T, Evans J, Harrison G et al. (2003)**  
Impact of the ICD-10 primary health care (PHC) diagnostic and management guidelines for mental  
disorders on detection and outcome in primary care.  
Br J Psychiatry 182 (1): 20-30
- [42] **De Waal MW, Arnold IA, Eekhof JA et al. (2004)**  
Somatoform disorders in general practice.  
Br J Psychiatry 184 (6): 470-476
- [43] **Deister A (a)**  
Angst- und Panikstörungen.  
In: Möller HJ, Laux G, Deister A: Psychiatrie und Psychotherapie.  
Thieme Verlag, Stuttgart, 2001, S. 106-133
- [44] **Deister A (b)**  
Essstörungen.  
In: Möller HJ, Laux G, Deister A: Psychiatrie und Psychotherapie.  
Thieme Verlag, Stuttgart, 2001, S. 268-278
- [45] **Deister A (c)**  
Somatoforme Störungen.  
In: Möller HJ, Laux G, Deister A: Psychiatrie und Psychotherapie.  
Thieme Verlag, Stuttgart, 2001, S. 254-267
- [46] **Deter HC (1998)**  
Psychosomatische Qualitätszirkel in der Arztpraxis sparen Geld.  
KV-Blatt Berlin 11: 31-32

- [47] **DeVault KR, Castell OD et al. (1999)**  
Updated guidelines for the diagnosis and treatment of gastroesophageal reflux disease.  
Am J Gastroenterol 94 (6): 1434-1442
- [48] **Dhillon PK, Farrow DC, Vaughan TL et al. (2001)**  
Family history of cancer and risk of esophageal and gastric cancers in the United States.  
Int J Cancer 93 (1): 148-152
- [49] **Diez-Quevedo C, Rangil T, Sanchez-Planell L et al. (2001)**  
Validation and utility of the Patient Health Questionnaire in diagnosing mental disorders  
in 1003 general hospital spanish inpatients.  
Psychosom Med 63 (4): 679-686
- [50] **Dimenäs E, Carlsson G, Glise H et al. (1996)**  
Relevance of norm values as part of documentation of quality of life instruments for use in upper gastro-  
intestinal disease.  
Scand J Gastroenterol 31 (Supplement 221): 8-13
- [51] **Dimenäs E, Glise H, Hallerbäck B et al. (1993)**  
Quality of life in patients with upper gastrointestinal symptoms.  
Scand J Gastroenterol 28 (8): 681-687
- [52] **Dimenäs E, Glise H, Hallerbäck B et al. (1995)**  
Well-being and gastrointestinal symptoms among patients referred to endoscopy  
owing to suspect duodenal ulcer.  
Scand J Gastroenterol 30 (11): 1046-1052
- [53] **Docherty JP (1997)**  
Barriers to the diagnosis of depression in primary care.  
J Clin Psychiatry 58 (Supplement 1): 5-10
- [54] **DSM-IV**  
Diagnostic and statistical manual of mental disorders.  
American Psychiatric Association, 1994  
Deutsche Fassung: Sass, H. et al. Hogrefe-Verlag, Göttingen, 1998
- [55] **Dupuy HJ**  
The Psychological General Well-Being (PGWB) Index.  
In: Wenger NK, Mattson ME, Furberg CF u.a. (Hrsg.): Assessment of quality of life  
in clinical trials of cardiovascular therapies.  
USA: Le Jacq Publishing Inc., 1984
- [56] **Eisen GM, Locke GR, Provenzale D (1999)**  
Health-related quality of life: a primer for gastroenterologists.  
Am J Gastroenterol 94 (8): 2017-2021
- [57] **Elkinton JR (1966)**  
Medicine and the quality of life.  
Ann Intern Med 64: 711-714
- [58] **Ellert U, Bellach BM (1999)**  
Der SF-36 im Bundes-Gesundheitssurvey: Beschreibung einer aktuellen Normstichprobe.  
Gesundheitswesen 61 (Sonderheft 2): 184-190
- [59] **Enck P, Dubois D, Marquis P (1999)**  
Quality of life in patients with upper gastrointestinal symptoms: results from the Domestic/International  
Gastroenterology Surveillance Study (DIGEST).  
Scand J Gastroenterol 34 (Supplement 231): 48-54
- [60] **Fischler B, Tack J, De Gucht V et al. (2003)**  
Heterogeneity of symptom pattern, psychosocial factors, and pathophysiological mechanisms  
in severe functional dyspepsia.  
Gastroenterology 124 (4): 903-910
- [61] **Frank L, Kleinman L, Rentz A et al. (2002)**  
Health-related quality of life associated with irritable bowel syndrome: comparison with other  
chronic disease.  
Clin Ther 24 (4): 675-689

- [62] **Franz, Matthias**  
Der Weg in die psychotherapeutische Beziehung.  
1. Auflage, Vandenhoeck & Ruprecht, Göttingen, 1997, S. 112-115
- [63] **Fritzsche K, Sandholzer H, Albota M et al. (1996)**  
Qualitätssicherung und Qualitätsmanagement in der Psychosomatischen Grundversorgung. Erste Ergebnisse eines Demonstrationsprojekts.  
Psychotherapeut 41: 326-330
- [64] **Fritzsche K, Sandholzer H, Brucks U et al. (1999)**  
Psychosocial care by general practitioners – where are the problems?  
Results of a demonstration project on quality management in psychosocial primary care.  
Int J Psychiatr Med 29 (4): 395-409
- [65] **Fritzsche K, Sandholzer H, Brucks U et al. (2000)**  
Psychosomatische Grundversorgung in der Hausarztpraxis.  
Z Ärztl Fortbild Qualitätssich 94 (2): 127-131
- [66] **Fritzsche K, Sandholzer H, Werner J et al. (2000)**  
Psychotherapeutische und psychosoziale Behandlungsmaßnahmen in der Hausarztpraxis.  
Psychother Psychosom Med Psychol 50 (6): 240-246
- [67] **Garratt AM, Ruta DA, Abdalla MI et al. (1993)**  
The SF-36 health survey questionnaire: an outcome measure suitable for routine use within the NHS?  
BMJ 306 (6890): 1440-1444
- [68] **Gater R, Tansella M, Korten A et al. (1998)**  
Sex differences in the prevalence and detection of depressive and anxiety disorders in general health care settings,  
Arch Gen Psychiatry 55: 405-413
- [69] **Gesellschaft für Rehabilitation bei Verdauungs- und Stoffwechselerkrankungen (1999)**  
Leitlinien für die Rehabilitation bei Verdauungs- und Stoffwechselerkrankungen: Peptische Krankheiten.  
Internet: [http://www.grvs.de/ll\\_grvs.htm](http://www.grvs.de/ll_grvs.htm)
- [70] **Glaeske G (1999)**  
Evidence-based medicine aus Sicht der Krankenkassen – ein Rahmen für qualifizierte Therapiefreiheit und verbesserte Wirtschaftlichkeit?  
Z Ärztl Fortbild Qualitätssich 93 (6): 421-426.
- [71] **Glaesmer H, Deter HC (2001)**  
Bedeutung der Arbeitsunfähigkeit durch psychische, psychiatrische und psychosomatische Erkrankungen im Rahmen der psychosomatischen Grundversorgung (PSGV) bei niedergelassenen Ärzten.  
Psychother Psychosom med Psychol 51 (3-4): A93
- [72] **Glise H, Wiklund I (2002)**  
Health-related quality of life and gastrointestinal disease.  
J Gastroenterol Hepatol 17 (Supplement): 72-84
- [73] **Goldberg D, Bridges K (1988)**  
Somatic presentations of psychiatric illness in primary care setting.  
J Psychosom Res 32 (2): 137-144
- [74] **Goodwin R, Olfson M (2001)**  
Treatment of panic attack and risk of major depressive disorder in the community.  
Am J Psychiatry 158 (7): 1146-1148
- [75] **Hach I, Rentsch A, Ruhl U et al. (2003)**  
Validität von Krankenscheindiagnosen psychischer Störungen.  
Gesundheitswesen 65 (6): 359-364
- [76] **Hahn SR, Kroenke K, Spitzer RL et al. (1996)**  
The difficult patient: prevalence, psychopathology, and functional impairment.  
J Gen Intern Med 11 (1): 1-8
- [77] **Halder SL, Locke GR III, Talley NJ et al. (2004)**  
Impact of functional gastrointestinal disorders on health-related quality of life: a population-based case-control study.  
Aliment Pharmacol Ther 19 (2): 233-242

- [78] **Haltenhof H, Oswald F, Bühler KE et al. (2000)**  
 Psychosoziale Kompetenz von Hausärzten und Versorgung psychisch Kranker: Erste Ergebnisse einer Fragebogenstudie.  
 Gesundheitswesen 62 (8-9): 462-467
- [79] **Hansen JM, Hallas J, Lauritsen JM et al. (1996)**  
 Non-steroidal anti-inflammatory drugs and ulcer complications: a risk factor analysis for clinical decision-making.  
 Scand J Gastroenterol 31 (2): 126-130
- [80] **Harms V**  
 Biomathematik, Statistik und Dokumentation.  
 HARMS Verlag, Kiel-Mönkeberg, 1998
- [81] **Haug TT, Svebak S, Wilhelmsen I et al. (1994)**  
 Psychological factors and somatic symptoms in functional dyspepsia. A comparison with duodenal ulcer and healthy controls.  
 J Psychosom Res 38 (4): 281-291
- [82] **Haug TT, Wilhelmsen I, Ursin H et al. (1995)**  
 What are the real problems for patients with functional dyspepsia?  
 Scand J Gastroenterol 30 (2): 97-100
- [83] **Hegerl U, Althaus D, Niklewski G et al. (2003)**  
 Optimierte Versorgung depressiver Patienten und Suizidprävention. Ergebnisse des „Nürnberger Bündnisses gegen Depression“.  
 Dtsch Ärztbl 100 (42): 2732-2737
- [84] **Henrich G, Herschbach P (2000)**  
 Questions on life satisfaction (FLZ<sup>M</sup>) – a short questionnaire for assessing subjective quality of life.  
 Eur J Psychol Ass 16: 150-159
- [85] **Herrmann H, Patrick DL, Diehr P et al. (2002)**  
 Longitudinal investigation of depression outcomes in primary care in six countries: the LIDO study. Functional status, health service use and treatment of people with depressive symptoms.  
 Psychol Med 32 (5): 889-902
- [86] **Herschbach P (2002)**  
 Das „Zufriedenheitsparadox“ in der Lebensqualitätsforschung.  
 Psychother Psychosom Med Psychol 52: 141-150
- [87] **Herschbach P, Henrich G, von Rad M (1999)**  
 Psychological factors in functional gastrointestinal disorders: characteristics of the disorder or of the illness behaviour?  
 Psychosom Med 61 (2): 148-153
- [88] **Higgins ES (1994)**  
 A review of unrecognized mental illness in primary care.  
 Arch Fam Med 3 (10): 908-917
- [89] **Holtmann G, Goebell H, Talley NJ (1994)**  
 Dyspepsia in consultants and non-consulters: prevalence, health-care seeking behaviour and risk factors.  
 Eur J Gastroenterol Hepatol 6: 917-924
- [90] **Howden CW, Hunt RH (1998)**  
 Guidelines for the management of Helicobacter pylori infection.  
 Am J Gastroenterol 93 (12): 2330-2338
- [91] **Howell S, Talley NJ (1999)**  
 Does fear of serious disease predict consulting behaviour amongst patients with dyspepsia in general practice?  
 Eur J Gastroenterol Hepatol 11 (8): 881-886
- [92] **Huber R, Koch D, Beiser I et al. (2004)**  
 Experience and attitudes towards CAM – a survey of internal and psychosomatic patients in a german university hospital.  
 Altern Ther Health Med 10 (1): 32-36
- [93] **ICD-10**  
 International statistical classification of diseases and related health problems.  
 World Health Organization, 1992, 10. Auflage. Deutsche Fassung: Weltgesundheitsorganisation, 1999

- [94] **Jackson JL, Houston JS, Hanling SR et al. (2001)**  
Clinical predictors of mental disorders among medical outpatients.  
Arch Intern Med 161 (6): 875-879
- [95] **Jhingran P, Cady RK, Rubino J et al. (1996)**  
Improvements in health-related quality of life with sumatriptan treatment for migraine.  
J Fam Pract 42 (1): 36-42
- [96] **Johnson JG, Spitzer RL, Williams JBW (2001)**  
Health problems, impairment and illnesses associated with bulimia nervosa and binge eating disorder among primary care and obstetric gynaecology patients.  
Psychol Med 31 (8): 1455-1466
- [97] **Jones R, Lydeard S (1989)**  
Prevalence of symptoms of dyspepsia in the community.  
BMJ 298 (6665): 30-32
- [98] **Junge B, Nagel M (1999)**  
Das Rauchverhalten in Deutschland.  
Gesundheitswesen 61 (Sonderheft 2): 121-125
- [99] **Kärger H, Obliers R, Körfer A et al. (1997)**  
Psychosomatische Grundversorgung.  
Z Allg Med 73: 211-217
- [100] **Karnowsky DA, Burchenal JH**  
The clinical evaluation of chemotherapeutic agents in cancer.  
In: MacLeod CM (Hrsg.): Evaluation of chemotherapeutic agents,  
Columbia University Press, New York, 1949, S. 191-205
- [101] **Kassenärztliche Vereinigung Berlin**  
Durchschnittliche Anzahl der Behandlungsfälle in Berliner Allgemeinarztpraxen in den Abrechnungsquartalen II/2001 bis I/2002.  
Abrechnungsstelle der KV Berlin, 2004
- [102] **Kiecolt-Glaser JK, Newton TL (2001)**  
Marriage and health: his and hers.  
Psychol Bull 127 (4): 472-503
- [103] **Kisley S, Goldberg D, Simon G (1997)**  
A comparison between somatic symptoms with and without clear organic cause: results of an international study.  
Psychol Med 27 (5): 1011-1019
- [104] **Klapow J, Kroenke K, Horton T et al. (2002)**  
Psychological disorders and distress in older primary care patients: a comparison of older and younger sample.  
Psychosom Med 64 (4): 635-643
- [105] **Koelz HR, Michel B (2004)**  
Nichtsteroidale Antirheumatika. Magenschutztherapie oder COX-2-Hemmer?  
Dtsch Arztebl 101 (45): 3041-3046
- [106] **Koloski NA, Talley NJ, Boyce PM (2000)**  
The impact of functional gastrointestinal disorders on quality of life.  
Am J Gastroenterol, 95 (1): 67-71
- [107] **Kroenke K (2003)**  
Patients presenting with somatic complaints: epidemiology, psychiatric comorbidity and management.  
Int J Methods Psychiatr Res 12 (1): 34-34
- [108] **Kroenke K, Jackson JL, Chamberlin J (1997)**  
Depressive and anxiety disorders in patients presenting with physical complaints: clinical predictors and outcome.  
Am J Med 103 (5): 339-347
- [109] **Kroenke K, Spitzer RL, Williams JB et al. (1994)**  
Physical symptoms in primary care. Predictors of psychiatric disorders and functional impairment.  
Arch Fam Med 3 (9): 774-779

- [110] **Kroenke K, Taylor-Vaisey A, Dietrich AJ et al. (2000)**  
Interventions to improve provider diagnosis and treatment of mental disorders in primary care.  
Psychosomatics 41 (1): 39-52
- [111] **Küchler T**  
Quality of life research in oncology.  
In: Guggenmoos-Holzmann I, Bloomfield K u.a. (Hrsg.): Quality of life and health: concepts, methods and applications.  
Blackwell-Wissenschafts-Verlag, Berlin, 1995, S. 89-96
- [112] **Kuykendall DH, Rabeneck L, Campell CJ et al. (1998)**  
Dyspepsia: how should we measure it?  
J Clin Epidemiol 51 (2): 99-106
- [113] **Kvien TK, Kassa S, Smedstad LM (1998)**  
Performance of the Norwegian SF-36 Health Survey in patients with rheumatoid arthritis.  
II. A Comparison of the SF-36 with disease-specific measures.  
J Clin Epidemiol 51 (11): 1077-1086
- [114] **Landolt-Theus P (1992)**  
Fälleverteilung in der Allgemeinmedizin.  
Der Allgemeinarzt 19 (3): 254-268
- [115] **Laubach W, Schumacher J, Mundt A et al. (2000)**  
Sozialschicht, Lebenszufriedenheit und Gesundheitseinschätzung. Ergebnisse einer repräsentativen Untersuchung der deutschen Bevölkerung.  
Soz Präventivmed 45 (1): 2-12
- [116] **Laux G (a)**  
Abhängigkeit und Sucht.  
In: Möller HJ, Laux G, Deister A: Psychiatrie und Psychotherapie.  
Thieme Verlag, Stuttgart, 2001, S. 306-330
- [117] **Laux G (b)**  
Affektive Störungen.  
In: Möller HJ, Laux G, Deister A: Psychiatrie und Psychotherapie.  
Thieme Verlag, Stuttgart, 2001, S. 73-105
- [118] **Lerner DJ, Levine S, Malspeis S et al. (1994)**  
Job strain and health-related quality of life in a national sample.  
Public Health 84 (10): 1580-1585
- [119] **Linden M, Gothe H, Ormel J (2004)**  
Ländervergleich Deutschland/Niederlande: Der Hausarzt als Gatekeeper.  
Dtsch Ärztebl 101 (39): 2600-2602
- [120] **Linden M, Maier W, Achberger M et al. (1996)**  
Psychische Erkrankungen und ihre Behandlung in Allgemeinarztpraxen in Deutschland.  
Nervenarzt 67 (3): 205-215
- [121] **Linzer M, Spitzer RL, Kroenke K et al. (1996)**  
Gender, quality of life, and mental disorders in primary care: results from the PRIME-MD 1000 Study.  
Am J Med 101 (5): 526-533
- [122] **Locke GR IIIrd (1998)**  
Prevalence, incidence and natural history of dyspepsia and functional dyspepsia.  
Baillieres Clin Gastroenterol 12 (3): 435-42
- [123] **Locke GR IIIrd, Weaver AL, Melton LJ 3rd et al. (2004)**  
Psychosocial factors are linked to functional gastrointestinal disorders: a population based nested case-control-study.  
Am J Gastroenterol 99 (2): 350-357
- [124] **Loerch B, Szegedi A, Kohnen R et al. (2000)**  
The primary care evaluation of mental disorders (PRIME-MD), german version: a comparison with the CIDI.  
J Psychiatr Res 34 (3): 211-220
- [125] **Löwe B, Gräfe K, Kroenke K et al. (2003)**  
Predictors of psychiatric comorbidity in medical outpatients.  
Psychosom Med 65 (5): 764-770



- [126] **Löwe B, Gräfe K, Zipfel S et al. (2003)**  
Detecting panic disorder in medical and psychosomatic outpatients: comparative validation of the Hospital Anxiety and Depression Scale, the Patient Health Questionnaire, a screening question, and physicians' diagnosis.  
J Psychosom Res 55 (6): 515-519
- [127] **Löwe B, Spitzer RL, Gräfe K et al. (2004)**  
Comparative validity of three screening questionnaires for DSM-IV depressive disorders and physicians' diagnosis.  
J Affect Disord 78 (2): 131-140
- [128] **Löwe B, Spitzer RL, Zipfel S et al.**  
PHQ-D – Gesundheitsfragebogen für Patienten. Komplettversion und Kurzform: Testmappe mit Manual.  
Pfizer GmbH, Karlsruhe, 2001
- [129] **Löwe B, Willand L, Eich W et al. (2004)**  
Psychiatric comorbidity and work disability in patients with inflammatory rheumatic disease.  
Psychosom Med 66 (3): 395-402
- [130] **Löwe B, Zipfel S, Herzog W**  
Gesundheitsfragebogen für Patienten (PHQ-D). Komplettversion und Kurzform mit vorläufigem Manual.  
Pfizer GmbH, Karlsruhe, 2001
- [131] **Mader FH, Weißgerber H**  
Allgemeinmedizin und Praxis.  
4. vollständig überarbeitete und erweiterte Auflage,  
Springer-Verlag, Berlin, Heidelberg, 2001, S.15-18
- [132] **Maier W, Linden M, Sartorius N (1996)**  
Psychische Erkrankungen in der Allgemeinpraxis. Ergebnisse und Schlussfolgerungen einer WHO-Studie.  
Dtsch Ärztebl 93 (18): 1202-1206
- [133] **Malfertheiner P, Holtmann G, Peitz U et al. (2001)**  
Leitlinien der Deutschen Gesellschaft für Verdauungs- und Stoffwechselkrankheiten zur Behandlung der Dyspepsie.  
Z Gastroenterol 39 (11): 937-956
- [134] **Malfertheiner P, Knippig C, Classen M**  
Peptisches Ulcus.  
In: Classen M, Diehl V, Kochsiek K: Innere Medizin.  
5. völlig überarbeitete Auflage, Verlag Urban & Fischer, München, 2004, S.1151-1164
- [135] **Manning WG, Wells KB (1992)**  
The effect of psychological distress and psychosocial well-being on use of medical services.  
Med Care 30 (6): 541-553
- [136] **Mansi C, Savarino V, Mela GS et al. (1993)**  
Are clinical patterns of dyspepsia a valid guideline for appropriate use of endoscopy? A report on 2253 dyspeptic patients.  
Am J Gastroenterol 88 (7): 1011-1015
- [137] **Mant JW, Jenkinson C, Murphy MF et al. (1998)**  
Use of the Short Form-36 to detect the influence of upper gastrointestinal disease on self-reported health status.  
Qual Life Res 7 (3): 221-226
- [138] **Mausner-Dorsch H, Eaton WW (2000)**  
Psychosocial work environment and depression: epidemiologic assessment of the demand-control model.  
Am J Public Health 90 (11): 1765-1770
- [139] **McHorney CA, Ware JE Jr, Lu JF et al. (1994)**  
The MOS 36-Item Short Form Health Survey (SF-36): III. Tests of data quality, scaling assumptions, and reliability across diverse patient groups.  
Med Care 32 (1): 40-66
- [140] **McHorney CA, Ware JE Jr, Raczek AE (1993)**  
The MOS 36-Item Short Form Health Survey (SF-36): II. Psychometric and clinical tests of validity in measuring physical and mental health constructs.  
Med Care 31 (3): 247-263

- [141] **Melchert HU, Knopf H, Pabel E et al. (2004)**  
Multi- und Komedikation von OTC-Präparaten. Eine Gegenüberstellung der alten und der neuen Bundesländer.  
Tägl Prax 45 (3): 605-619
- [142] **Moayyedi P, Forman D, Braunholtz D et al. (2000)**  
The proportion of upper gastrointestinal symptoms in the community associated with *Helicobacter pylori*, lifestyle factors, and nonsteroidal anti-inflammatory drugs.  
Am J Gastroenterol 95 (6): 1448-1455
- [143] **Murdaugh C (1997)**  
Health-related quality of life as an outcome in organizational research.  
Med Care 35 (Supp. 11): 41-48
- [144] **National Cancer Institute**  
NCI invitation conference on the economic impact of cancer clinical trials.  
Bethesda, United States of America, 1994
- [145] **Nease DE Jr, Volk RJ, Cass AR (1999)**  
Investigation of a severity-based classification of mood and anxiety symptoms in primary care patients.  
J Am Board Fam Pract 12 (1): 21-31
- [146] **Netter FH**  
Innere Medizin.  
Thieme-Verlag, Stuttgart, 2000, S. 814
- [147] **O'Malley PG, Wong PWK, Kroenke K et al. (1998)**  
The value of screening for psychiatric disorders prior to upper endoscopy.  
J Psychosom Res 44 (2): 279-287
- [148] **Olfson M, Shea S, Feder A et al. (2000)**  
Prevalence of anxiety, depression, and substance use disorders in an urban general medicine practice.  
Arch Fam Med 9 (9): 876-883
- [149] **Ormel J, Koeter MW, van den Brink W et al. (1991)**  
Recognition, management, and course of anxiety and depression in general practice.  
Arch Gen Psychiatry 48 (8): 700-706
- [150] **Ormel J, Von Korff M, Üstün TB et al. (1994)**  
Common mental disorders and disability across cultures: results from the WHO collaborative study on psychological problems in general health care.  
JAMA 272 (22): 1741-1748
- [151] **Osoba D (1994)**  
Lessons learned from measuring health-related quality of life in oncology.  
J Clin Oncol 12: 608-616
- [152] **Paterniti S, Niedhammer I, Lang T et al. (2002)**  
Psychosocial factors at work, personality traits and depressive symptoms.  
Br J Psychiatry 181 (2): 111-117
- [153] **Persoons P, Luyckx K, Desloovere C et al. (2003)**  
Anxiety and mood disorders in otorhinolaryngology outpatients presenting with dizziness: validation of the self-administered PRIME-MD Patient Health Questionnaire and epidemiology.  
Gen Hosp Psychiatry 25 (5): 316-323
- [154] **Pignone MP, Gaynes BN, Rusthon JL et al. (2002)**  
Screening for depression in adults: a summary of evidence for the U.S. Preventive Services Task Force.  
Ann Intern Med 136 (10): 765-776
- [155] **Pini S, Perkonnig A, Tansella M et al. (1999)**  
Prevalence and 12-month outcome of threshold and subthreshold mental disorders in primary care.  
J Affect Disord 56 (1): 37-48
- [156] **Radoschewski M, Bellach BM (1999)**  
Der SF-36 im Bundes-Gesundheitssurvey – Möglichkeiten und Anforderungen der Nutzung auf Bevölkerungsebene.  
Gesundheitswesen 61 (Sonderheft 2): 191-199
- [157] **Regan J, Yarnold J, Jones PW et al. (1991)**  
Palliation and life quality in lung cancer: how good are clinicians at judging treatment outcome?  
Br J Cancer 64 (2): 396-400

- [158] **Ren XS (1997)**  
Marital status and quality of relationships: the impact on health perception.  
Soc Sci Med 44 (2): 241-249
- [159] **Restellini A, Malè PJ, Nicolet T et al.**  
„Ich habe Magenschmerzen“.  
In: Raetzo MA, Restellini A (Hrsg.): Alltagsbeschwerden. Diagnostische und therapeutische Strategien in der allgemeinmedizinischen Praxis.  
Verlag Hans Huber, Bern, 1998, S. 253-262
- [160] **Revicki DA, Wood M, Maton PN et al. (1998)**  
The impact of gastroesophageal reflux disease on health-related quality of life.  
Am J Med 104 (3): 252-258
- [161] **Rieder A (2004)**  
Geschlechtsspezifische Unterschiede aus Sicht der Sozialmedizin.  
Tägl Prax 45 (2): 359-368
- [162] **Robinson JW, Roter DL (1999)**  
Counseling by primary care physicians of patients who disclose psychosocial problems.  
J Fam Pract 48 (9): 698-705
- [163] **Rojo-Moreno L, Livianos-Aldana L, Cervera-Martinez G et al. (2002)**  
The role of stress in the onset of depressive disorders. A controlled study in a spanish clinical sample.  
Soc Psychiatry Psychiatr Epidemiol 37 (12): 592-598
- [164] **Rothenbacher D, Peter R, Bode G et al. (1998)**  
Dyspepsia in relation to Helicobacter pylori infection and psychosocial work stress in white collar employees.  
Am J Gastroenterol 93 (9): 1443-1449
- [165] **Roy-Byrne PP, Stang P, Wittchen HU et al. (2000)**  
Lifetime panic-depression comorbidity in the National Comorbidity Study.  
Br J Psychiatry 176 (3): 229-235
- [166] **Sandholzer H, Brucks C, Cierpka M et al.**  
Qualitätssicherung in der Psychosomatik. Verbundstudie innerhalb des Demonstrationsprojektes zur Qualitätssicherung in der ambulanten Versorgung.  
Schriftenreihe des Bundesministeriums für Gesundheit, Band 118, Nomos-Verlag, Baden-Baden, 1999
- [167] **Sartorius N, Üstün B, Lecrubier Y et al. (1996)**  
Depression comorbid with anxiety: results from the WHO study on psychological disorders in primary health care.  
Br J Psychiatry 168 (Supplement 30): 38-43
- [168] **Schneider F, Harter M, Kratz S et al. (2003)**  
Unzureichender subjektiver Behandlungsverlauf bei depressiven Patienten in der haus- und nervenärztlichen Praxis.  
Z Ärztl Fortbild Qualitätssich 97 (Supplement 4): 57-66
- [169] **Schoepf D, Heun R, Weiffenbach O et al. (2003)**  
4-Wochen-Prävalenz somatoformer Störungen und assoziierte psychosoziale Beeinträchtigung.  
Nervenarzt 74 (3): 245-251
- [170] **Schulenburg JM v. d., Schöffski O (1994)**  
Überweisungs- und Einweisungsverhalten von Primärärzten nach dem GSG.  
Münch Med Wschr 136 (14): 202-205
- [171] **Schumacher J, Brähler E (1999)**  
Prävalenz von Schmerzen in der deutschen Bevölkerung.  
Der Schmerz 13 (6): 375-384
- [172] **Simon GE, Fleck M, Lucas R et al. (2004)**  
Prevalence and predictors of depression treatment in an international primary care study.  
Am J Psychiatry 161 (9): 1626-1634
- [173] **Simon GE, Von Korff M, Piccinelli M et al. (1999)**  
An international study of the relation between somatic symptoms and depression.  
N Engl J Med 341: 1329-1335

- [174] **Slevin ML, Plant H, Lynch D et al. (1988)**  
Who should measure quality of life, the doctor or the patient?  
Br J Cancer 57 (1): 109-112
- [175] **Spitzer RL, Kroenke K, Linzer M et al. (1995)**  
Health-related quality of life in primary care patients with mental disorders. Results from the PRIME-MD 1000 Study.  
JAMA 274 (19): 1511-1517
- [176] **Spitzer RL, Kroenke K, Williams JB (1999)**  
Validation and utility of a self-report version of PRIME-MD: the PHQ primary care Study.  
JAMA 282 (18): 1737-1744
- [177] **Spitzer RL, Williams JB, Kroenke K et al. (1994)**  
Utility of a new procedure for diagnosing mental disorders in primary care: the PRIME-MD 1000 Study.  
JAMA 272 (22): 1749-1756
- [178] **Spitzer RL, Williams JB, Kroenke K et al. (2000)**  
Validity and utility of the PRIME-MD patient health questionnaire assessment of 3000 obstetric-gynecologic patients: The PRIME-MD Health Questionnaire Obstetric-Gynecology Study.  
Am J Obstet Gynecol 183 (3): 759-769
- [179] **Stansfeld SA, Bosma H, Hemingway H et al. (1998)**  
Psychosocial work characteristics and social support as predictors of SF-36 health functioning: the Whitehall II Study.  
Psychosom Med 60 (3): 247-255
- [180] **Stansfeld SA, Fuhrer R, Shipley MJ et al. (1999)**  
Work characteristics predict psychiatric disorder: prospective results from the Whitehall II Study.  
Occup Environ Med 56 (5): 302-307
- [181] **Statistisches Bundesamt Deutschland**  
Gesundheitsberichterstattung des Bundes, Wiesbaden, 2002  
Internet: <http://www.gbe-bund.de>
- [182] **Statistisches Landesamt Berlin**  
Die zwölf Bezirke Berlins.  
Statistisches Taschenbuch 2001.  
Internet: <http://www.statistik-berlin.de>
- [183] **Stewart AL, Hays RD, Ware JE Jr (1988)**  
The MOS Short-form General Health Survey.  
Med Care 26 (7): 724-735
- [184] **Straus WL, Ofman JJ, MacLean C et al. (2002)**  
Do NSAIDs cause dyspepsia? A meta-analysis evaluating alternative dyspepsia definitions.  
Am J Gastroenterol 97 (8): 1951-1958
- [185] **Streich W, Braun B (2003)**  
Versorgungsqualität in der hausärztlichen Praxis – Ergebnisse einer bundesweiten Bevölkerungsbefragung in 2001/2002.  
Z Allg Med 79 (6): 270-277
- [186] **Südhoff T, Schmiegel W**  
Magentumoren.  
In: Classen M, Diehl V, Kochsiek K: Innere Medizin.  
5. völlig überarbeitete Auflage, Verlag Urban & Fischer, München, 2004, S.1166-1173
- [187] **Surtees PG, Wainwright NWJ, Khaw KT et al. (2003)**  
Functional health status, chronic medical conditions and disorders of mood.  
Br J Psychiatry 183 (4): 299-303
- [188] **Talley NJ, Phung N, Kalantar JS (2001)**  
ABC of the upper gastrointestinal tract: Indigestion: When is it functional?  
BMJ 323 (7324): 1294-1297
- [189] **Talley NJ, Weaver AL, Zinsmeister AR (1995)**  
Impact of functional dyspepsia on quality of life.  
Dig Dis Sci 40 (3): 584-589

- [190] **Tanum L, Malt UF (2001)**  
Personality and physical symptoms in nonpsychiatric patients with functional gastrointestinal disorder.  
J Psychosom Res 50 (3): 139-146
- [191] **Tress W, Kruse J, Heckrath C et al. (1997)**  
Psychogene Erkrankungen in hausärztlichen Praxen.  
Z psychosom Med 43: 211-232
- [192] **U.S. Preventive Services Task Force (2002)**  
Screening for depression: recommendations and rationale.  
Ann Intern Med 136 (10): 760-764
- [193] **Unützer J, Klap R, Sturm R et al. (2000)**  
Mental disorders and the use of alternative medicine: results from a national survey.  
Am J Psychiatry 157 (11): 1851-1857
- [194] **Üstün TB, Ayuso-Mateos JL, Chatterji S et al. (2004)**  
Global burden of depressive disorders in the year 2000.  
Br J Psychiatry 184 (5): 386-392
- [195] **Valenstein M, Kales H, Mellow A et al. (1998)**  
Psychiatric diagnosis and intervention in older and younger patients in a primary care clinic: effect of a screening and diagnostic instrument.  
J Am Geriatr Soc 46 (12): 1499-1505
- [196] **Wallace MB, Durkalski VL, Vaughan J et al. (2001)**  
Age and alarm symptoms do not predict endoscopic findings among patients with dyspepsia: a multi-centre database study.  
Gut 49 (1): 29-34
- [197] **Ware JE Jr, Gandek B (1998)**  
Overview of the SF-36 Health Survey and the International Quality of Life Assessment (IQOLA) Project.  
J Clin Epidemiol 51 (11): 903-912
- [198] **Ware JE Jr, Sherbourne CD (1992)**  
The MOS-36-Items Short Form Health Survey (SF-36): I. Conceptual Framework and Item Selection.  
Med Care 30 (6): 473-483
- [199] **Ware JE Jr, Snow KK, Kosinski M et al.**  
SF-36 Health Survey: Manual and Interpretation Guide.  
Boston, Massachusetts: The Health Institute, New England Medical Center, 1993
- [200] **Wensing M, Vingerhoets E, Grol R (2001)**  
Functional status, health problems, age and comorbidity in primary care patients.  
Qual Life Res 10 (2): 141-148
- [201] **Westbrook JI, McIntosh J, Talley NJ (2000)**  
Factors associated with consulting medical or non-medical practitioners for dyspepsia: an australian population-based study.  
Aliment Pharmacol Ther 14 (12): 1581-1588
- [202] **WHO**  
Preamble to the constitution of the World Health Organization as adopted by the international health conference, New York, 19-22 June, 1946: signed on 22 July 1946 by the representatives of 61 States and entered into force on 7 April 1948.  
Official Records of the WHO, 1948, No.2, S.100
- [203] **WHO World Mental Health Survey Consortium (2004)**  
Prevalence, Severity, and unmet need for treatment of mental disorders in the World Health Organization World Mental Health Surveys.  
JAMA 291 (21): 2581-2590
- [204] **Whooley MA, Simon GE (2000)**  
Managing depression in medical outpatients.  
N Engl J Med 343 (26): 1942-1950
- [205] **Wiklund I, Butler-Wheelhouse P (1996)**  
Psychological factors and their role in symptomatic gastroesophageal reflux disease and functional dyspepsia.  
Scand J Gastroenterol 31 (Supplement 220): 94-100

- [206] **Wilhelmsen I, Haug TT, Ursin H et al. (1995)**  
Discriminant analysis of factors distinguishing patients with functional dyspepsia from patients with duodenal ulcer.  
Dig Dis Sci 40 (5): 1105-1111
- [207] **Williams JW, Rost K, Dietrich AJ et al. (1999)**  
Primary care physicians' approach to depressive disorders.  
Arch Fam Med 8(1): 58-67
- [208] **Wilson D, Parsons J, Tucker G (2000)**  
The SF-36 summary scales: problems and solutions.  
Soz Präventivmed 45 (6): 239-246
- [209] **Wittchen HU, Hoyer J, Höfler M et al. (2001)**  
Hausärztliche Interventionen und Verschreibungsverhalten: Ergebnisse der GAD-P-Studie.  
MMW Fortschr Med 119 (Sonderheft 1): 36-41
- [210] **Wittchen HU, Jacobi F (2001)**  
Die Versorgungssituation psychischer Störungen in Deutschland. Eine klinisch-epidemiologische Abschätzung anhand des Bundes-Gesundheitssurveys 1998.  
Bundesgesundheitsbl Gesundheitsforsch Gesundheitsschutz 44: 993-1000
- [211] **Wittchen HU, Kessler RC, Beesdo K et al. (2002)**  
Generalized anxiety and depression in primary care: prevalence, recognition and management.  
J Clin Psychiatry 63 (Supplement 8): 24-34
- [212] **Wittchen HU, Krause P, Hoyer J et al. (2001)**  
Prävalenz und Korrelate generalisierter Angststörungen in der Allgemeinarztpraxis.  
MMW Fortschr Med 119 (Sonderheft 1): 17-25
- [213] **Wittchen HU, Müller N, Pfister H et al. (1999)**  
Affektive, somatoforme und Angststörungen in Deutschland – Erste Ergebnisse des bundesweiten Zusatzsurveys "Psychische Störungen".  
Gesundheitswesen 61 (Sonderheft 2): 216-222
- [214] **Wittchen HU, Winter S, Höfler M et al. (2000)**  
Häufigkeit und Erkennensrate von Depressionen in der hausärztlichen Praxis.  
MMW Fortschr Med 118 (Sonderheft 1): 22-30
- [215] **Wu LR, Parkerson GR, Doraiswamy PM (2002)**  
Health perception, pain, and disability as correlates of anxiety and depression symptoms in primary care patients.  
J Am Board Fam Pract 15: 183-190
- [216] **Yacavone RF, Locke GR, Provenzale D et al. (2001)**  
Quality of life measurements in gastroenterology: what is available?  
Am J Gastroenterol 96 (2): 285-297
- [217] **Zipfel S, Löwe B, Schneider A et al. (1999)**  
Lebensqualität, Depressivität und Krankheitsverarbeitung bei Patienten in der Wartezeit auf eine Herztransplantation.  
Psychother Psychosom Med Psychol 49 (6): 187-194