

5

BIBLIOGRAPHY

- Accola M.A., Huang B., Al Masri A., and McNiven M.A. (2002). The antiviral dynamin family member, MxA, tubulates lipids and localizes to the smooth endoplasmic reticulum. *J Biol Chem* **277**(24):21829-21835.
- Adayev T., Chen-Hwang M.C., Murakami N., Wang R., and Hwang Y.W. (2006). MNB/DYRK1A phosphorylation regulates the interactions of synaptojanin 1 with endocytic accessory proteins. *Biochem Biophys Res Commun* **351**(4):1060-1065.
- Agaton C., Galli J., Hoiden Guthenberg I., Janzon L., Hansson M., Asplund A., Brundell E., Lindberg S., Ruthberg I., Wester K., et al. (2003). Affinity proteomics for systematic protein profiling of chromosome 21 gene products in human tissues. *Mol Cell Proteomics* **2**(6):405-414.
- Al-Shahrour F., Minguez P., Vaquerizas J.M., Conde L., and Dopazo J. (2005). BABELOMICS: a suite of web tools for functional annotation and analysis of groups of genes in high-throughput experiments. *Nucleic Acids Res* **33**(Web Server issue):W460-464.
- Alexandrov A., Martzen M.R., and Phizicky E.M. (2002). Two proteins that form a complex are required for 7-methylguanosine modification of yeast tRNA. *Rna* **8**(10):1253-1266.
- Alvarez A.R., Sandoval P.C., Leal N.R., Castro P.U., and Kosik K.S. (2004). Activation of the neuronal c-Abl tyrosine kinase by amyloid-beta-peptide and reactive oxygen species. *Neurobiol Dis* **17**(2):326-336.
- Alvarez M., Estivill X., and de la Luna S. (2003). DYRK1A accumulates in splicing speckles through a novel targeting signal and induces speckle disassembly. *J Cell Sci* **116**(Pt 15):3099-3107.
- Antonarakis S.E., Adelsberger P.A., Petersen M.B., Binkert F., and Schinzel A.A. (1990). Analysis of DNA polymorphisms suggests that most de novo dup(21q) chromosomes in patients with Down syndrome are isochromosomes and not translocations. *Am J Hum Genet* **47**(6):968-972.
- Antonarakis S.E., Petersen M.B., McInnis M.G., Adelsberger P.A., Schinzel A.A., Binkert F., Pangalos C., Raoul O., Slaugenhaupt S.A., Hafez M., et al. (1992). The meiotic stage of nondisjunction in trisomy 21: determination by using DNA polymorphisms. *Am J Hum Genet* **50**(3):544-550.
- Antonarakis S.E., Warren A.C., McCormick M.K., Lewis J.G., Hieter P.A., and Chakravarti A. (1989). Molecular mapping of chromosome 21 and the region responsible for Down syndrome. *Prog Clin Biol Res* **311**:29-43.
- Auerbach D., Thaminy S., Hottiger M.O., and Stagljar I. (2002). The post-genomic era of interactive proteomics: facts and perspectives. *Proteomics* **2**(6):611-623.
- Augereau P., Badia E., Balaguer P., Carascossa S., Castet A., Jalaguier S., and Cavailles V. (2006). Negative regulation of hormone signaling by RIP140. *J Steroid Biochem Mol Biol* **102**(1-5):51-59.
- Bader G.D., Donaldson I., Wolting C., Ouellette B.F., Pawson T., and Hogue C.W. (2001). BIND--The Biomolecular Interaction Network Database. *Nucleic Acids Res* **29**(1):242-245.
- Bader G.D. and Hogue C.W. (2002). Analyzing yeast protein-protein interaction data obtained from different sources. *Nat Biotechnol* **20**(10):991-997.
- Baeuerle P.A. and Henkel T. (1994). Function and activation of NF-kappa B in the immune system. *Annu Rev Immunol* **12**:141-179.
- Bajo M., Fruehauf J., Kim S.H., Fountoulakis M., and Lubec G. (2002). Proteomic evaluation of intermediary metabolism enzyme proteins in fetal Down's syndrome cerebral cortex. *Proteomics* **2**(11):1539-1546.
- Balakirev M.Y., Tcherniuk S.O., Jaquinod M., and Chroboczek J. (2003). Otubains: a new family of cysteine proteases in the ubiquitin pathway. *EMBO Rep* **4**(5):517-522.
- Barabasi A.L. and Albert R. (1999). Emergence of scaling in random networks. *Science* **286**(5439):509-512.
- Barabasi A.L. and Oltvai Z.N. (2004). Network biology: understanding the cell's functional organization. *Nat Rev Genet* **5**(2):101-113.
- Barcena M., Radermacher M., Bar J., Kopperschlager G., and Ruiz T. (2007). The structure of the ATP-bound state of *S. cerevisiae* phosphofructokinase determined by cryo-electron microscopy. *J Struct Biol* **159**(1):135-143.
- Baxter L.L., Moran T.H., Richtsmeier J.T., Troncoso J., and Reeves R.H. (2000). Discovery and genetic localization of Down syndrome cerebellar phenotypes using the Ts65Dn mouse. *Hum Mol Genet* **9**(2):195-202.

5. Bibliography

- Beadle G.W. and Tatum E.L. (1941). Genetic Control of Biochemical Reactions in *Neurospora*. *Proc Natl Acad Sci U S A* **27**(11):499-506.
- Bendtsen J.D., Nielsen H., von Heijne G., and Brunak S. (2004). Improved prediction of signal peptides: SignalP 3.0. *J Mol Biol* **340**(4):783-795.
- Bertelsen V., Breen K., Sandvig K., Stang E., and Madshus I.H. (2007). The Cbl-interacting protein TULA inhibits dynamin-dependent endocytosis. *Exp Cell Res* **313**(8):1696-1709.
- Berthelsen J., Kilstrup-Nielsen C., Blasi F., Mavilio F., and Zappavigna V. (1999). The subcellular localization of PBX1 and EXD proteins depends on nuclear import and export signals and is modulated by association with PREP1 and HTH. *Genes Dev* **13**(8):946-953.
- Bertone P., Stole V., Royce T.E., Rozowsky J.S., Urban A.E., Zhu X., Rinn J.L., Tongprasit W., Samanta M., Weissman S., et al. (2004). Global identification of human transcribed sequences with genome tiling arrays. *Science* **306**(5705):2242-2246.
- Birney E., Stamatoyannopoulos J.A., Dutta A., Guigo R., Gingeras T.R., Margulies E.H., Weng Z., Snyder M., Dermitzakis E.T., Thurman R.E., et al. (2007). Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* **447**(7146):799-816.
- Blasko I., Marx F., Steiner E., Hartmann T., and Grubeck-Loebenstien B. (1999). TNFalpha plus IFNgamma induce the production of Alzheimer beta-amyloid peptides and decrease the secretion of APPs. *Faseb J* **13**(1):63-68.
- Bork P., Jensen L.J., von Mering C., Ramani A.K., Lee I., and Marcotte E.M. (2004). Protein interaction networks from yeast to human. *Curr Opin Struct Biol* **14**(3):292-299.
- Brandt B.R. and Rosen I. (1995). Impaired peripheral somatosensory function in children with Down syndrome. *Neuropediatrics* **26**(6):310-312.
- Brasch M.A., Hartley J.L., and Vidal M. (2004). ORFeome cloning and systems biology: standardized mass production of the parts from the parts-list. *Genome Res* **14**(10B):2001-2009.
- Brelje T.C., Wessendorf M.W., and Sorenson R.L. (1993). Multicolor laser scanning confocal immunofluorescence microscopy: practical application and limitations. *Methods Cell Biol* **38**:97-181.
- Brooker R.J. (1999). *Genetics : Analysis and Principles*. Menlo Park, California, USA, Addison-Wesley.
- Brosius U., Dehmel T., and Gartner J. (2002). Two different targeting signals direct human peroxisomal membrane protein 22 to peroxisomes. *J Biol Chem* **277**(1):774-784.
- Brown K.R. and Jurisica I. (2005). Online predicted human interaction database. *Bioinformatics* **21**(9):2076-2082.
- Brown W.T. (1987). Premature aging syndromes. *Curr Probl Dermatol* **17**:152-165.
- Buchin P.J., Levy J.S., and Schullinger J.N. (1986). Down's syndrome and the gastrointestinal tract. *J Clin Gastroenterol* **8**(2):111-114.
- Bussow K., Cahill D., Nietfeld W., Bancroft D., Scherzinger E., Lehrach H., and Walter G. (1998). A method for global protein expression and antibody screening on high-density filters of an arrayed cDNA library. *Nucleic Acids Res* **26**(21):5007-5008.
- Byrne J.L. and Ward K. (1994). Genetic factors in recurrent abortion. *Clin Obstet Gynecol* **37**(3):693-704.
- Cabin D.E., Gardiner K., and Reeves R.H. (1996). Molecular genetic characterization and comparative mapping of the human PCP4 gene. *Somat Cell Mol Genet* **22**(3):167-175.
- Canick J.A., Knight G.J., Palomaki G.E., Haddow J.E., Cuckle H.S., and Wald N.J. (1988). Low second trimester maternal serum unconjugated oestriol in pregnancies with Down's syndrome. *Br J Obstet Gynaecol* **95**(4):330-333.
- Carr J. (1994). Long-term-outcome for people with Down's syndrome. *J Child Psychol Psychiatry* **35**(3):425-439.
- Castet A., Herledan A., Bonnet S., Jalaguier S., Vanacker J.M., and Cavailles V. (2006). Receptor-interacting protein 140 differentially regulates estrogen receptor-related receptor transactivation depending on target genes. *Mol Endocrinol* **20**(5):1035-1047.
- Causier B. (2004). Studying the interactome with the yeast two-hybrid system and mass spectrometry. *Mass Spectrom Rev* **23**(5):350-367.
- Cavaillès V., Dauvois S., L'Horsset F., Lopez G., Hoare S., Kushner P.J., and Parker M.G. (1995). Nuclear factor RIP140 modulates transcriptional activation by the estrogen receptor. *Embo J* **14**(15):3741-3751.
- Chao M.V. (2003). Neurotrophins and their receptors: a convergence point for many signalling pathways. *Nat Rev Neurosci* **4**(4):299-309.
- Chapman R.S. and Hesketh L.J. (2000). Behavioral phenotype of individuals with Down syndrome. *Ment Retard Dev Disabil Res Rev* **6**(2):84-95.
- Chargaff E. (1950). Chemical specificity of nucleic acids and mechanism of their enzymatic degradation. *Experientia* **16**:201-209.

- Charrier E., Mosinger B., Meissirel C., Aguera M., Rogemond V., Reibel S., Salin P., Chounlamountri N., Perrot V., Belin M.F., *et al.* (2006). Transient alterations in granule cell proliferation, apoptosis and migration in postnatal developing cerebellum of CRMP1^{-/-} mice. *Genes Cells* **11**(12):1337-1352.
- Chaurasia G., Iqbal Y., Hanig C., Herzel H., Wanker E.E., and Futschik M.E. (2007). UniHI: an entry gate to the human protein interactome. *Nucleic Acids Res* **35**(Database issue):D590-594.
- Chettouh Z., Croquette M.F., Delobel B., Gilgenkrants S., Leonard C., Maunoury C., Prieur M., Rethore M.O., Sinet P.M., Chery M., *et al.* (1995). Molecular mapping of 21 features associated with partial monosomy 21: involvement of the APP-SOD1 region. *Am J Hum Genet* **57**(1):62-71.
- Chipuk J.E. and Green D.R. (2006). Dissecting p53-dependent apoptosis. *Cell Death Differ* **13**(6):994-1002.
- Chkheidze A.N. and Liebhaber S.A. (2003). A novel set of nuclear localization signals determine distributions of the alphaCP RNA-binding proteins. *Mol Cell Biol* **23**(23):8405-8415.
- Choi H.S., Kim C.S., Hwang C.K., Song K.Y., Law P.Y., Wei L.N., and Loh H.H. (2007). Novel function of the poly(C)-binding protein alpha CP3 as a transcriptional repressor of the mu opioid receptor gene. *Faseb J* **21**(14):3963-3973.
- Chou C.Y., Liu L.Y., Chen C.Y., Tsai C.H., Hwa H.L., Chang L.Y., Lin Y.S., and Hsieh F.J. (2008). Gene expression variation increase in trisomy 21 tissues. *Mamm Genome*.
- Coppus A., Evenhuis H., Verberne G.J., Visser F., van Gool P., Eikelenboom P., and van Duijn C. (2006). Dementia and mortality in persons with Down's syndrome. *J Intellect Disabil Res* **50**(Pt 10):768-777.
- Corder E.H., Saunders A.M., Strittmatter W.J., Schmechel D.E., Gaskell P.C., Small G.W., Roses A.D., Haines J.L., and Pericak-Vance M.A. (1993). Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* **261**(5123):921-923.
- Crick F.H. (1958). On protein synthesis. *Symp Soc Exp Biol* **12**:138-163.
- Cullen L.M. and Arndt G.M. (2005). Genome-wide screening for gene function using RNAi in mammalian cells. *Immunol Cell Biol* **83**(3):217-223.
- Cusick M.E., Klitgord N., Vidal M., and Hill D.E. (2005). Interactome: gateway into systems biology. *Hum Mol Genet* **14 Spec No. 2**:R171-181.
- D'Alton M. and Cleary-Goldman J. (2005). First and second trimester evaluation of risk for fetal aneuploidy: the secondary outcomes of the FASTER Trial. *Semin Perinatol* **29**(4):240-246.
- Dauphinot L., Lyle R., Rivals I., Dang M.T., Moldrich R.X., Golfier G., Ettwiller L., Toyama K., Rossier J., Personnaz L., *et al.* (2005). The cerebellar transcriptome during postnatal development of the Ts1Cje mouse, a segmental trisomy model for Down syndrome. *Hum Mol Genet* **14**(3):373-384.
- Davisson M.T., Schmidt C., Reeves R.H., Irving N.G., Akeson E.C., Harris B.S., and Bronson R.T. (1993). Segmental trisomy as a mouse model for Down syndrome. *Prog Clin Biol Res* **384**:117-133.
- Del Bo R., Comi G.P., Giorda R., Crimi M., Locatelli F., Martinelli-Boneschi F., Pozzoli U., Castelli E., Bresolin N., and Scarlato G. (2003). The 129 codon polymorphism of the prion protein gene influences earlier cognitive performance in Down syndrome subjects. *J Neurol* **250**(6):688-692.
- Delabar J.M., Chettouh Z., Rahmani Z., Theophile D., Blouin J.L., Bono R., Kraus J., Barton J., Patterson D., and Sinet P.M. (1992). Gene-dosage mapping of 30 DNA markers on chromosome 21. *Genomics* **13**(3):887-889.
- Demas G.E., Nelson R.J., Krueger B.K., and Yarowsky P.J. (1996). Spatial memory deficits in segmental trisomic Ts65Dn mice. *Behav Brain Res* **82**(1):85-92.
- Dennis G., Jr., Sherman B.T., Hosack D.A., Yang J., Gao W., Lane H.C., and Lempicki R.A. (2003). DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome Biol* **4**(5):P3.
- Dikic I. and Schmidt M.H. (2007). Malfunctions within the Cbl interactome uncouple receptor tyrosine kinases from destructive transport. *Eur J Cell Biol* **86**(9):505-512.
- Ding D.Q., Tomita Y., Yamamoto A., Chikashige Y., Haraguchi T., and Hiraoka Y. (2000). Large-scale screening of intracellular protein localization in living fission yeast cells by the use of a GFP-fusion genomic DNA library. *Genes Cells* **5**(3):169-190.
- Ding H.F., Bustin M., and Hansen U. (1997). Alleviation of histone H1-mediated transcriptional repression and chromatin compaction by the acidic activation region in chromosomal protein HMG-14. *Mol Cell Biol* **17**(10):5843-5855.
- Donev R., Newall A., Thome J., and Sheer D. (2007). A role for SC35 and hnRNP1 in the determination of amyloid precursor protein isoforms. *Mol Psychiatry* **12**(7):681-690.
- Down J.L. (1995). Observations on an ethnic classification of idiots [1866]. *Ment Retard* **33**(1):54-56.
- Emanuelsson O., Brunak S., von Heijne G., and Nielsen H. (2007). Locating proteins in the cell using TargetP, SignalP and related tools. *Nat Protoc* **2**(4):953-971.

5. Bibliography

- Engidawork E. and Lubec G. (2001). Protein expression in Down syndrome brain. *Amino Acids* **21**(4):331-361.
- Epstein C.J. (1990). The consequences of chromosome imbalance. *Am J Med Genet Suppl* **7**:31-37.
- Epstein C.J., Cox D.R., and Epstein L.B. (1985). Mouse trisomy 16: an animal model of human trisomy 21 (Down syndrome). *Ann N Y Acad Sci* **450**:157-168.
- Ermak G., Morgan T.E., and Davies K.J. (2001). Chronic overexpression of the calcineurin inhibitory gene DSCR1 (Adapt78) is associated with Alzheimer's disease. *J Biol Chem* **276**(42):38787-38794.
- Ewing R.M., Chu P., Elisma F., Li H., Taylor P., Climie S., McBroom-Cerajewski L., Robinson M.D., O'Connor L., Li M., *et al.* (2007). Large-scale mapping of human protein-protein interactions by mass spectrometry. *Mol Syst Biol* **3**:89.
- Faustino N.A. and Cooper T.A. (2003). Pre-mRNA splicing and human disease. *Genes Dev* **17**(4):419-437.
- Fernandez M.R., Porte S., Crosas E., Barbera N., Farres J., Biosca J.A., and Pares X. (2007). Human and yeast zeta-crystallins bind AU-rich elements in RNA. *Cell Mol Life Sci* **64**(11):1419-1427.
- Fields S. (2005). High-throughput two-hybrid analysis. The promise and the peril. *Febs J* **272**(21):5391-5399.
- Fields S. and Song O. (1989). A novel genetic system to detect protein-protein interactions. *Nature* **340**(6230):245-246.
- Fischer D.F., van Dijk R., Sluijs J.A., Nair S.M., Racchi M., Levelt C.N., van Leeuwen F.W., and Hol E.M. (2005). Activation of the Notch pathway in Down syndrome: cross-talk of Notch and APP. *Faseb J* **19**(11):1451-1458.
- FitzPatrick D.R., Ramsay J., McGill N.I., Shade M., Carothers A.D., and Hastie N.D. (2002). Transcriptome analysis of human autosomal trisomy. *Hum Mol Genet* **11**(26):3249-3256.
- Fok P.T., Huang K.C., Holland P.C., and Nalbantoglu J. (2007). The Coxsackie and adenovirus receptor binds microtubules and plays a role in cell migration. *J Biol Chem* **282**(10):7512-7521.
- Ford C.E. and Hamerton J.L. (1956). The chromosomes of man. *Nature* **178**(4541):1020-1023.
- Ford J., McEwan I.J., Wright A.P., and Gustafsson J.A. (1997). Involvement of the transcription factor IID protein complex in gene activation by the N-terminal transactivation domain of the glucocorticoid receptor in vitro. *Mol Endocrinol* **11**(10):1467-1475.
- Formosa T., Barry J., Alberts B.M., and Greenblatt J. (1991). Using protein affinity chromatography to probe structure of protein machines. *Methods Enzymol* **208**:24-45.
- Fortunato A. and Fraser A.G. (2005). Uncover genetic interactions in *Caenorhabditis elegans* by RNA interference. *Biosci Rep* **25**(5-6):299-307.
- Franklin R. and Gosling R.G. (1953). Molecular Configuration in Sodium Thymonucleate. *Nature* **171**:740-741.
- Friedel C.C. and Zimmer R. (2006). Inferring topology from clustering coefficients in protein-protein interaction networks. *BMC Bioinformatics* **7**:519.
- Fromm L. and Burden S.J. (2001). Neuregulin-1-stimulated phosphorylation of GABP in skeletal muscle cells. *Biochemistry* **40**(17):5306-5312.
- Fuenzalida K.M., Aguilera M.C., Piderit D.G., Ramos P.C., Contador D., Quinones V., Rigotti A., Bronfman F.C., and Bronfman M. (2005). Peroxisome proliferator-activated receptor gamma is a novel target of the nerve growth factor signaling pathway in PC12 cells. *J Biol Chem* **280**(10):9604-9609.
- Futschik M.E., Chaurasia G., and Herzog H. (2007). Comparison of human protein-protein interaction maps. *Bioinformatics* **23**(5):605-611.
- Gardiner K. (2006). Transcriptional dysregulation in Down syndrome: predictions for altered protein complex stoichiometries and post-translational modifications, and consequences for learning/behavior genes ELK, CREB, and the estrogen and glucocorticoid receptors. *Behav Genet* **36**(3):439-453.
- Gardiner K. and Costa A.C. (2006). The proteins of human chromosome 21. *Am J Med Genet C Semin Med Genet* **142**(3):196-205.
- Gardiner K., Davisson M.T., and Crnic L.S. (2004). Building protein interaction maps for Down's syndrome. *Brief Funct Genomic Proteomic* **3**(2):142-156.
- Gardiner K., Fortna A., Bechtel L., and Davisson M.T. (2003). Mouse models of Down syndrome: how useful can they be? Comparison of the gene content of human chromosome 21 with orthologous mouse genomic regions. *Gene* **318**:137-147.
- Gasteiger E., Gattiker A., Hoogland C., Ivanyi I., Appel R.D., and Bairoch A. (2003). ExpASY: The proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Res* **31**(13):3784-3788.
- Gaughan L., Logan I.R., Cook S., Neal D.E., and Robson C.N. (2002). Tip60 and histone deacetylase 1 regulate androgen receptor activity through changes to the acetylation status of the receptor. *J Biol Chem* **277**(29):25904-25913.
- Gavin A.C., Aloy P., Grandi P., Krause R., Boesche M., Marzioch M., Rau C., Jensen L.J., Bastuck S., Dumpelfeld B., *et al.* (2006). Proteome survey reveals modularity of the yeast cell machinery. *Nature* **440**(7084):631-636.

- Gavin A.C., Bosche M., Krause R., Grandi P., Marzioch M., Bauer A., Schultz J., Rick J.M., Michon A.M., Cruciat C.M., *et al.* (2002). Functional organization of the yeast proteome by systematic analysis of protein complexes. *Nature* **415**(6868):141-147.
- Georgieva S., Nabirochkina E., Dilworth F.J., Eickhoff H., Becker P., Tora L., Georgiev P., and Soldatov A. (2001). The novel transcription factor e(y)2 interacts with TAF(II)40 and potentiates transcription activation on chromatin templates. *Mol Cell Biol* **21**(15):5223-5231.
- Gerdes A.M., Horder M., and Bonnevie-Nielsen V. (1993). Increased IFN-alpha-induced sensitivity but reduced reactivity of 2',5'-oligoadenylate synthetase (2,5AS) in trisomy 21 blood lymphocytes. *Clin Exp Immunol* **93**(1):93-96.
- Giannone S., Strippoli P., Vitale L., Casadei R., Canaider S., Lenzi L., D'Addabbo P., Frabetti F., Facchin F., Farina A., *et al.* (2004). Gene expression profile analysis in human T lymphocytes from patients with Down Syndrome. *Ann Hum Genet* **68**(Pt 6):546-554.
- Giot L., Bader J.S., Brouwer C., Chaudhuri A., Kuang B., Li Y., Hao Y.L., Ooi C.E., Godwin B., Vitols E., *et al.* (2003). A protein interaction map of *Drosophila melanogaster*. *Science* **302**(5651):1727-1736.
- Gitton Y., Dahmane N., Baik S., Ruiz i Altaba A., Neidhardt L., Scholze M., Herrmann B.G., Kahlem P., Benkahla A., Schrinner S., *et al.* (2002). A gene expression map of human chromosome 21 orthologues in the mouse. *Nature* **420**(6915):586-590.
- Goehler H., Lalowski M., Stelzl U., Waelter S., Stroedicke M., Worm U., Droege A., Lindenberg K.S., Knoblich M., Haenig C., *et al.* (2004). A protein interaction network links GIT1, an enhancer of huntingtin aggregation, to Huntington's disease. *Mol Cell* **15**(6):853-865.
- Goffeau A., Barrell B.G., Bussey H., Davis R.W., Dujon B., Feldmann H., Galibert F., Hoheisel J.D., Jacq C., Johnston M., *et al.* (1996). Life with 6000 genes. *Science* **274**(5287):546, 563-547.
- Goldberg D.S. and Roth F.P. (2003). Assessing experimentally derived interactions in a small world. *Proc Natl Acad Sci U S A* **100**(8):4372-4376.
- Goshima Y., Nakamura F., Strittmatter P., and Strittmatter S.M. (1995). Collapsin-induced growth cone collapse mediated by an intracellular protein related to UNC-33. *Nature* **376**(6540):509-514.
- Gough N.R. (2002). Science's signal transduction knowledge environment: the connections maps database. *Ann N Y Acad Sci* **971**:585-587.
- Granholm A.C., Ford K.A., Hyde L.A., Bimonte H.A., Hunter C.L., Nelson M., Albeck D., Sanders L.A., Mufson E.J., and Crnic L.S. (2002). Estrogen restores cognition and cholinergic phenotype in an animal model of Down syndrome. *Physiol Behav* **77**(2-3):371-385.
- Greene M.E., Pitts J., McCarville M.A., Wang X.S., Newport J.A., Edelstein C., Lee F., Ghosh S., and Chu S. (2000). PPARgamma: observations in the hematopoietic system. *Prostaglandins Other Lipid Mediat* **62**(1):45-73.
- Gregory S.G., Barlow K.F., McLay K.E., Kaul R., Swarbreck D., Dunham A., Scott C.E., Howe K.L., Woodfine K., Spencer C.C., *et al.* (2006). The DNA sequence and biological annotation of human chromosome 1. *Nature* **441**(7091):315-321.
- Guipponi M., Vuagniaux G., Wattenhofer M., Shibuya K., Vazquez M., Dougherty L., Scamuffa N., Guida E., Okui M., Rossier C., *et al.* (2002). The transmembrane serine protease (TMPRSS3) mutated in deafness DFNB8/10 activates the epithelial sodium channel (ENaC) in vitro. *Hum Mol Genet* **11**(23):2829-2836.
- Haas S., Vingron M., Poustka A., and Wiemann S. (1998). Primer design for large scale sequencing. *Nucleic Acids Res* **26**(12):3006-3012.
- Haas S.A., Hild M., Wright A.P., Hain T., Talibi D., and Vingron M. (2003). Genome-scale design of PCR primers and long oligomers for DNA microarrays. *Nucleic Acids Res* **31**(19):5576-5581.
- Hager G.L., Lim C.S., Elbi C., and Baumann C.T. (2000). Trafficking of nuclear receptors in living cells. *J Steroid Biochem Mol Biol* **74**(5):249-254.
- Hahn J.S., Hu Z., Thiele D.J., and Iyer V.R. (2004). Genome-wide analysis of the biology of stress responses through heat shock transcription factor. *Mol Cell Biol* **24**(12):5249-5256.
- Hammerle B., Elizalde C., Galceran J., Becker W., and Tejedor F.J. (2003). The MNB/DYRK1A protein kinase: neurobiological functions and Down syndrome implications. *J Neural Transm Suppl*(67):129-137.
- Hamosh A., Scott A.F., Amberger J.S., Bocchini C.A., and McKusick V.A. (2005). Online Mendelian Inheritance in Man (OMIM), a knowledgebase of human genes and genetic disorders. *Nucleic Acids Res* **33**(Database issue):D514-517.
- Han D.K., Eng J., Zhou H., and Aebersold R. (2001). Quantitative profiling of differentiation-induced microsomal proteins using isotope-coded affinity tags and mass spectrometry. *Nat Biotechnol* **19**(10):946-951.
- Hart G.T., Ramani A.K., and Marcotte E.M. (2006). How complete are current yeast and human protein-interaction networks? *Genome Biol* **7**(11):120.
- Hattori M., Fujiyama A., Taylor T.D., Watanabe H., Yada T., Park H.S., Toyoda A., Ishii K., Totoki Y., Choi D.K., *et al.* (2000). The DNA sequence of human chromosome 21. *Nature* **405**(6784):311-319.

5. Bibliography

- Hennequin M., Morin C., and Feine J.S. (2000). Pain expression and stimulus localisation in individuals with Down's syndrome. *Lancet* **356**(9245):1882-1887.
- Herman B. (1989). Resonance energy transfer microscopy. *Methods Cell Biol* **30**:219-243.
- Hermjakob H., Montecchi-Palazzi L., Bader G., Wojcik J., Salwinski L., Ceol A., Moore S., Orchard S., Sarkans U., von Mering C., *et al.* (2004). The HUPO PSI's molecular interaction format--a community standard for the representation of protein interaction data. *Nat Biotechnol* **22**(2):177-183.
- Hershey A.D. and Chase M. (1952). Independent Functions of Viral Protein and Nucleic Acid in Growth of Bacteriophage *J Gen Physiol* **36**:39-56.
- Hester M., Thompson J.C., Mills J., Liu Y., El-Hodiri H.M., and Weinstein M. (2005). Smad1 and Smad8 function similarly in mammalian central nervous system development. *Mol Cell Biol* **25**(11):4683-4692.
- Heurgue-Hamard V., Champ S., Mora L., Merkulova-Rainon T., Kisselev L.L., and Buckingham R.H. (2005). The glutamine residue of the conserved GGQ motif in *Saccharomyces cerevisiae* release factor eRF1 is methylated by the product of the YDR140w gene. *J Biol Chem* **280**(4):2439-2445.
- Hieter P. and Boguski M. (1997). Functional genomics: it's all how you read it. *Science* **278**(5338):601-602.
- Hirschman L., Park J.C., Tsujii J., Wong L., and Wu C.H. (2002). Accomplishments and challenges in literature data mining for biology. *Bioinformatics* **18**(12):1553-1561.
- Holmstrom S., Van Antwerp M.E., and Iniguez-Lluhi J.A. (2003). Direct and distinguishable inhibitory roles for SUMO isoforms in the control of transcriptional synergy. *Proc Natl Acad Sci U S A* **100**(26):15758-15763.
- Hook E.B. (1981a). Rates of chromosome abnormalities at different maternal ages. *Obstet Gynecol* **58**(3):282-285.
- Hook E.B. (1981b). Unbalanced Robertsonian translocations associated with Down's syndrome or Patau's syndrome: chromosome subtype, proportion inherited, mutation rates, and sex ratio. *Hum Genet* **59**(3):235-239.
- Hoopengardner B., Bhalla T., Staber C., and Reenan R. (2003). Nervous system targets of RNA editing identified by comparative genomics. *Science* **301**(5634):832-836.
- Horton P., Park K.J., Obayashi T., Fujita N., Harada H., Adams-Collier C.J., and Nakai K. (2007). WoLF PSORT: protein localization predictor. *Nucleic Acids Res* **35**(Web Server issue):W585-587.
- Hu Y.H., Vanhecke D., Lehrach H., and Janitz M. (2005). High-throughput subcellular protein localization using cell arrays. *Biochem Soc Trans* **33**(Pt 6):1407-1408.
- Hu Y.H., Warnatz H.J., Vanhecke D., Wagner F., Fiebitz A., Thamm S., Kahlem P., Lehrach H., Yaspo M.L., and Janitz M. (2006). Cell array-based intracellular localization screening reveals novel functional features of human chromosome 21 proteins. *BMC Genomics* **7**:155.
- Hua S. and Sun Z. (2001). Support vector machine approach for protein subcellular localization prediction. *Bioinformatics* **17**(8):721-728.
- Huang E.J. and Reichardt L.F. (2001). Neurotrophins: roles in neuronal development and function. *Annu Rev Neurosci* **24**:677-736.
- Huang L., Kinnucan E., Wang G., Beaudenon S., Howley P.M., Huibregtse J.M., and Pavletich N.P. (1999). Structure of an E6AP-UbcH7 complex: insights into ubiquitination by the E2-E3 enzyme cascade. *Science* **286**(5443):1321-1326.
- Hubbard T.J., Aken B.L., Beal K., Ballester B., Caccamo M., Chen Y., Clarke L., Coates G., Cunningham F., Cutts T., *et al.* (2007). Ensembl 2007. *Nucleic Acids Res* **35**(Database issue):D610-617.
- Hublitz P., Kunowska N., Mayer U.P., Muller J.M., Heyne K., Yin N., Fritzsche C., Poli C., Miguet L., Schupp I.W., *et al.* (2005). NIR is a novel INHAT repressor that modulates the transcriptional activity of p53. *Genes Dev* **19**(23):2912-2924.
- Huh W.K., Falvo J.V., Gerke L.C., Carroll A.S., Howson R.W., Weissman J.S., and O'Shea E.K. (2003). Global analysis of protein localization in budding yeast. *Nature* **425**(6959):686-691.
- Huq M.D. and Wei L.N. (2005). Post-translational modification of nuclear co-repressor receptor-interacting protein 140 by acetylation. *Mol Cell Proteomics* **4**(7):975-983.
- Huret J.L., Delabar J.M., Marlhens F., Aurias A., Nicole A., Berthier M., Tanzer J., and Sinet P.M. (1987). Down syndrome with duplication of a region of chromosome 21 containing the CuZn superoxide dismutase gene without detectable karyotypic abnormality. *Hum Genet* **75**(3):251-257.
- Inoue H., Nojima H., and Okayama H. (1990). High efficiency transformation of *Escherichia coli* with plasmids. *Gene* **96**(1):23-28.
- Ishibashi T., Hayakawa H., and Sekiguchi M. (2003). A novel mechanism for preventing mutations caused by oxidation of guanine nucleotides. *EMBO Rep* **4**(5):479-483.
- Ito T., Chiba T., Ozawa R., Yoshida M., Hattori M., and Sakaki Y. (2001). A comprehensive two-hybrid analysis to explore the yeast protein interactome. *Proc Natl Acad Sci U S A* **98**(8):4569-4574.

- Ivakhno S. (2007). From functional genomics to systems biology. *Febs J* **274**(10):2439-2448.
- Jackson J.F., North E.R., 3rd, and Thomas J.G. (1976). Clinical diagnosis of Down's syndrome. *Clin Genet* **9**(5):483-487.
- Jacobs P.A., Baikie A.G., Court Brown W.M., and Strong J.A. (1959). The somatic chromosomes in mongolism. *Lancet* **1**(7075):710.
- James R.S., Ellis K., Pettay D., and Jacobs P.A. (1998). Cytogenetic and molecular study of four couples with multiple trisomy 21 pregnancies. *Eur J Hum Genet* **6**(3):207-212.
- Jeong H., Mason S.P., Barabasi A.L., and Oltvai Z.N. (2001). Lethality and centrality in protein networks. *Nature* **411**(6833):41-42.
- Johanson R.A., Sarau H.M., Foley J.J., and Slemmon J.R. (2000). Calmodulin-binding peptide PEP-19 modulates activation of calmodulin kinase II In situ. *J Neurosci* **20**(8):2860-2866.
- Jonsson U., Fagerstam L., Ivarsson B., Johnsson B., Karlsson R., Lundh K., Lofas S., Persson B., Roos H., Ronnberg I., et al. (1991). Real-time biospecific interaction analysis using surface plasmon resonance and a sensor chip technology. *Biotechniques* **11**(5):620-627.
- Joosten A.M., De Vos S., Van Opstal D., Brandenburg H., Gaillard J.L., and Vermeij-Keers C. (1997). Full monosomy 21, prenatally diagnosed by fluorescent in situ hybridization. *Prenat Diagn* **17**(3):271-275.
- Juarez J.C., Manuia M., Burnett M.E., Betancourt O., Boivin B., Shaw D.E., Tonks N.K., Mazar A.P., and Donate F. (2008). Superoxide dismutase 1 (SOD1) is essential for H₂O₂-mediated oxidation and inactivation of phosphatases in growth factor signaling. *Proc Natl Acad Sci U S A* **105**(20):7147-7152.
- Kadota M., Nishigaki R., Wang C.C., Toda T., Shirayoshi Y., Inoue T., Gojobori T., Ikeo K., Rogers M.S., and Oshimura M. (2004). Proteomic signatures and aberrations of mouse embryonic stem cells containing a single human chromosome 21 in neuronal differentiation: an in vitro model of Down syndrome. *Neuroscience* **129**(2):325-335.
- Kahlem P., Sultan M., Herwig R., Steinfath M., Balzereit D., Eppens B., Saran N.G., Pletcher M.T., South S.T., Stetten G., et al. (2004). Transcript level alterations reflect gene dosage effects across multiple tissues in a mouse model of down syndrome. *Genome Res* **14**(7):1258-1267.
- Kampa D., Cheng J., Kapranov P., Yamanaka M., Brubaker S., Cawley S., Drenkow J., Piccolboni A., Bekiranov S., Helt G., et al. (2004). Novel RNAs identified from an in-depth analysis of the transcriptome of human chromosomes 21 and 22. *Genome Res* **14**(3):331-342.
- Kanehisa M., Goto S., Hattori M., Aoki-Kinoshita K.F., Itoh M., Kawashima S., Katayama T., Araki M., and Hirakawa M. (2006). From genomics to chemical genomics: new developments in KEGG. *Nucleic Acids Res* **34**(Database issue):D354-357.
- Karlsson B., Gustafsson J., Hedov G., Ivarsson S.A., and Anneren G. (1998). Thyroid dysfunction in Down's syndrome: relation to age and thyroid autoimmunity. *Arch Dis Child* **79**(3):242-245.
- Kawahara Y., Ito K., Sun H., Kanazawa I., and Kwak S. (2003). Low editing efficiency of GluR2 mRNA is associated with a low relative abundance of ADAR2 mRNA in white matter of normal human brain. *Eur J Neurosci* **18**(1):23-33.
- Kedziora J. and Bartosz G. (1988). Down's syndrome: a pathology involving the lack of balance of reactive oxygen species. *Free Radic Biol Med* **4**(5):317-330.
- Kelly P.A. and Rahmani Z. (2005). DYRK1A enhances the mitogen-activated protein kinase cascade in PC12 cells by forming a complex with Ras, B-Raf, and MEK1. *Mol Biol Cell* **16**(8):3562-3573.
- Kennedy M.A., Venkateswaran A., Tarr P.T., Xenarios I., Kudoh J., Shimizu N., and Edwards P.A. (2001). Characterization of the human ABCG1 gene: liver X receptor activates an internal promoter that produces a novel transcript encoding an alternative form of the protein. *J Biol Chem* **276**(42):39438-39447.
- Kerley J.S., Olsen S.L., Freemantle S.J., and Spinella M.J. (2001). Transcriptional activation of the nuclear receptor corepressor RIP140 by retinoic acid: a potential negative-feedback regulatory mechanism. *Biochem Biophys Res Commun* **285**(4):969-975.
- Kikuchi T., Ichimiya S., Kojima T., Crisa L., Koshiba S., Tonooka A., Kondo N., Van Der Saag P.T., Yokoyama S., and Sato N. (2004). Expression profiles and functional implications of p53-like transcription factors in thymic epithelial cell subtypes. *Int Immunol* **16**(6):831-841.
- Kim Y.K., Furic L., Desgroseillers L., and Maquat L.E. (2005). Mammalian Staufen1 recruits Upf1 to specific mRNA 3'UTRs so as to elicit mRNA decay. *Cell* **120**(2):195-208.
- Kishino T., Lalonde M., and Wagstaff J. (1997). UBE3A/E6-AP mutations cause Angelman syndrome. *Nat Genet* **15**(1):70-73.
- Knipper M., Claussen C., Ruttiger L., Zimmermann U., Lullmann-Rauch R., Eskelinen E.L., Schroder J., Schwake M., and Saftig P. (2006). Deafness in LIMP2-deficient mice due to early loss of the potassium channel KCNQ1/KCNE1 in marginal cells of the stria vascularis. *J Physiol* **576**(Pt 1):73-86.
- Kochs G., Haener M., Aebi U., and Haller O. (2002). Self-assembly of human MxA GTPase into highly ordered dynamin-like oligomers. *J Biol Chem* **277**(16):14172-14176.

5. Bibliography

- Konecna A., Frischknecht R., Kinter J., Ludwig A., Steuble M., Meskenaitė V., Indermuhle M., Engel M., Cen C., Mateos J.M., *et al.* (2006). Calsyntenin-1 docks vesicular cargo to kinesin-1. *Mol Biol Cell* **17**(8):3651-3663.
- Korenberg J.R., Chen X.N., Mitchell S., Fannin S., Gerwehr S., Cohen D., and Chumakov I. (1995). A high-fidelity physical map of human chromosome 21q in yeast artificial chromosomes. *Genome Res* **5**(5):427-443.
- Korenberg J.R., Chen X.N., Schipper R., Sun Z., Gonsky R., Gerwehr S., Carpenter N., Daumer C., Dignan P., Distèche C., *et al.* (1994). Down syndrome phenotypes: the consequences of chromosomal imbalance. *Proc Natl Acad Sci U S A* **91**(11):4997-5001.
- Krause G., Winkler L., Mueller S.L., Haseloff R.F., Piontek J., and Blasig I.E. (2007). Structure and function of claudins. *Biochim Biophys Acta*.
- Kremer L., Dominguez J.E., and Avila J. (1988). Detection of tubulin-binding proteins by an overlay assay. *Anal Biochem* **175**(1):91-95.
- Krogan N.J., Cagney G., Yu H., Zhong G., Guo X., Ignatchenko A., Li J., Pu S., Datta N., Tikuisis A.P., *et al.* (2006). Global landscape of protein complexes in the yeast *Saccharomyces cerevisiae*. *Nature* **440**(7084):637-643.
- Krull M., Pistor S., Voss N., Kel A., Reuter I., Kronenberg D., Michael H., Schwarzer K., Potapov A., Choi C., *et al.* (2006). TRANSPATH: an information resource for storing and visualizing signaling pathways and their pathological aberrations. *Nucleic Acids Res* **34**(Database issue):D546-551.
- Kubota H., Hynes G., and Willison K. (1995). The eighth Cct gene, Cctq, encoding the theta subunit of the cytosolic chaperonin containing TCP-1. *Gene* **154**(2):231-236.
- Kumar A., Agarwal S., Heyman J.A., Matson S., Heidtman M., Piccirillo S., Umansky L., Drawid A., Jansen R., Liu Y., *et al.* (2002). Subcellular localization of the yeast proteome. *Genes Dev* **16**(6):707-719.
- Kuster B., Mortensen P., Andersen J.S., and Mann M. (2001). Mass spectrometry allows direct identification of proteins in large genomes. *Proteomics* **1**(5):641-650.
- Kwak Y.D., Brannen C.L., Qu T., Kim H.M., Dong X., Soba P., Majumdar A., Kaplan A., Beyreuther K., and Sugaya K. (2006). Amyloid precursor protein regulates differentiation of human neural stem cells. *Stem Cells Dev* **15**(3):381-389.
- Lamberti A., Caraglia M., Longo O., Marra M., Abbruzzese A., and Arcari P. (2004). The translation elongation factor 1A in tumorigenesis, signal transduction and apoptosis: review article. *Amino Acids* **26**(4):443-448.
- Lamesch P., Li N., Milstein S., Fan C., Hao T., Szabo G., Hu Z., Venkatesan K., Bethel G., Martin P., *et al.* (2007). hORFeome v3.1: a resource of human open reading frames representing over 10,000 human genes. *Genomics* **89**(3):307-315.
- Landgraf P., Rusu M., Sheridan R., Sewer A., Iovino N., Aravin A., Pfeffer S., Rice A., Kamphorst A.O., Landthaler M., *et al.* (2007). A mammalian microRNA expression atlas based on small RNA library sequencing. *Cell* **129**(7):1401-1414.
- Lange A.W., Rothermel B.A., and Yutzey K.E. (2005). Restoration of DSCR1 to disomy in the trisomy 16 mouse model of Down syndrome does not correct cardiac or craniofacial development anomalies. *Dev Dyn* **233**(3):954-963.
- Larsson M., Graslund S., Yuan L., Brundell E., Uhlen M., Hoog C., and Stahl S. (2000). High-throughput protein expression of cDNA products as a tool in functional genomics. *J Biotechnol* **80**(2):143-157.
- Lehner B. and Fraser A.G. (2004). A first-draft human protein-interaction map. *Genome Biol* **5**(9):R63.
- Lehner B. and Sanderson C.M. (2004). A protein interaction framework for human mRNA degradation. *Genome Res* **14**(7):1315-1323.
- Lehner B., Semple J.I., Brown S.E., Counsell D., Campbell R.D., and Sanderson C.M. (2004). Analysis of a high-throughput yeast two-hybrid system and its use to predict the function of intracellular proteins encoded within the human MHC class III region. *Genomics* **83**(1):153-167.
- Lejeune J., Turpin R., and Gautier M. (1959). Mongolism; a chromosomal disease (trisomy). *Bull Acad Natl Med* **143**(11-12):256-265.
- Levy J. (1991). The gastrointestinal tract in Down syndrome. *Prog Clin Biol Res* **373**:245-256.
- Li H.J., Haque Z.K., Chen A., and Mendelsohn M. (2007). RIF-1, a novel nuclear receptor corepressor that associates with the nuclear matrix. *J Cell Biochem*.
- Li J.F., Liu L.D., Ma S.H., Che Y.C., Wang L.C., Dong C.H., Zhao H.L., Liao Y., and Li Q.H. (2004a). HTRP--an immediate-early gene product induced by HSV1 infection in human embryo fibroblasts, is involved in cellular co-repressors. *J Biochem (Tokyo)* **136**(2):169-176.
- Li S., Armstrong C.M., Bertin N., Ge H., Milstein S., Boxem M., Vidalain P.O., Han J.D., Chesneau A., Hao T., *et al.* (2004b). A map of the interactome network of the metazoan *C. elegans*. *Science* **303**(5657):540-543.
- Li W. and Guan K.L. (2004). The Down syndrome cell adhesion molecule (DSCAM) interacts with and activates Pak. *J Biol Chem* **279**(31):32824-32831.
- Liebel U., Starkuviene V., Erfle H., Simpson J.C., Poustka A., Wiemann S., and Pepperkok R. (2003). A microscope-based screening platform for large-scale functional protein analysis in intact cells. *FEBS Lett* **554**(3):394-398.

- Lin H.Y., Michtalik H.J., Zhang S., Andersen T.T., Van Riper D.A., Davies K.K., Ermak G., Petti L.M., Nachod S., Narayan A.V., *et al.* (2003). Oxidative and calcium stress regulate DSCR1 (Adapt78/MCIP1) protein. *Free Radic Biol Med* **35**(5):528-539.
- Lin J., Blake M., Tang C., Zimmer D., Rustandi R.R., Weber D.J., and Carrier F. (2001). Inhibition of p53 transcriptional activity by the S100B calcium-binding protein. *J Biol Chem* **276**(37):35037-35041.
- Lowell B.B. (1999). PPARgamma: an essential regulator of adipogenesis and modulator of fat cell function. *Cell* **99**(3):239-242.
- Lu B., Jiang Y.J., Zhou Y., Xu F.Y., Hatch G.M., and Choy P.C. (2005). Cloning and characterization of murine 1-acyl-sn-glycerol 3-phosphate acyltransferases and their regulation by PPARalpha in murine heart. *Biochem J* **385**(Pt 2):469-477.
- Lu Z., Szafron D., Greiner R., Lu P., Wishart D.S., Poulin B., Anvik J., Macdonell C., and Eisner R. (2004). Predicting subcellular localization of proteins using machine-learned classifiers. *Bioinformatics* **20**(4):547-556.
- Lueking A., Horn M., Eickhoff H., Bussow K., Lehrach H., and Walter G. (1999). Protein microarrays for gene expression and antibody screening. *Anal Biochem* **270**(1):103-111.
- Lyle R., Gehrig C., Neergaard-Henrichsen C., Deutsch S., and Antonarakis S.E. (2004). Gene expression from the aneuploid chromosome in a trisomy mouse model of down syndrome. *Genome Res* **14**(7):1268-1274.
- Maglott D., Ostell J., Pruitt K.D., and Tatusova T. (2005). Entrez Gene: gene-centered information at NCBI. *Nucleic Acids Res* **33**(Database issue):D54-58.
- Makeyev A.V. and Liebhaber S.A. (2000). Identification of two novel mammalian genes establishes a subfamily of KH-domain RNA-binding proteins. *Genomics* **67**(3):301-316.
- Mann M. and Wilm M. (1994). Error-tolerant identification of peptides in sequence databases by peptide sequence tags. *Anal Chem* **66**(24):4390-4399.
- Mansilla F., Friis I., Jadidi M., Nielsen K.M., Clark B.F., and Knudsen C.R. (2002). Mapping the human translation elongation factor eEF1H complex using the yeast two-hybrid system. *Biochem J* **365**(Pt 3):669-676.
- Mao R., Zielke C.L., Zielke H.R., and Pevsner J. (2003). Global up-regulation of chromosome 21 gene expression in the developing Down syndrome brain. *Genomics* **81**(5):457-467.
- Marcell M.M. and Armstrong V. (1982). Auditory and visual sequential memory of Down syndrome and nonretarded children. *Am J Ment Defic* **87**(1):86-95.
- Marcell M.M. and Weeks S.L. (1988). Short-term memory difficulties and Down's syndrome. *J Ment Defic Res* **32** (Pt 2):153-162.
- Marheineke K. and Krude T. (1998). Nucleosome assembly activity and intracellular localization of human CAF-1 changes during the cell division cycle. *J Biol Chem* **273**(24):15279-15286.
- Maroun L.E. (1980). Interferon action and chromosome 21 trisomy. *J Theor Biol* **86**(3):603-606.
- Maroun L.E. (1995). Anti-interferon immunoglobulins can improve the trisomy 16 mouse phenotype. *Teratology* **51**(5):329-335.
- Maroun L.E., Heffernan T.N., and Hallam D.M. (2000). Partial IFN-alpha/beta and IFN-gamma receptor knockout trisomy 16 mouse fetuses show improved growth and cultured neuron viability. *J Interferon Cytokine Res* **20**(2):197-203.
- Marsischky G. and LaBaer J. (2004). Many paths to many clones: a comparative look at high-throughput cloning methods. *Genome Res* **14**(10B):2020-2028.
- Martinez-Frias M.L. (2005). The real earliest historical evidence of Down syndrome. *Am J Med Genet A* **132**(2):231.
- Maslen C.L., Babcock D., Robinson S.W., Bean L.J., Dooley K.J., Willour V.L., and Sherman S.L. (2006). CRELD1 mutations contribute to the occurrence of cardiac atrioventricular septal defects in Down syndrome. *Am J Med Genet A* **140**(22):2501-2505.
- Mathivanan S., Periaswamy B., Gandhi T.K., Kandasamy K., Suresh S., Mohmood R., Ramachandra Y.L., and Pandey A. (2006). An evaluation of human protein-protein interaction data in the public domain. *BMC Bioinformatics* **7 Suppl 5**:S19.
- McGinniss M.J., Kazazian H.H.J., Stetten G., Petersen M.B., Boman H., Engel E., Greenberg F., Hertz J.M., Johnson A., Laca Z., *et al.* (1992). Mechanisms of ring chromosome formation in 11 cases of human ring chromosome 21. *Am J Hum Genet* **50**(1):15-28.
- McInnis M.G., Chakravarti A., Blaschak J., Petersen M.B., Sharma V., Avramopoulos D., Blouin J.L., Konig U., Brahe C., Matisse T.C., *et al.* (1993). A linkage map of human chromosome 21:43 PCR markers at average intervals of 2.5 cM. *Genomics* **16**(3):562-571.
- McKusick V.A. (1997). Genomics: structural and functional studies of genomes. *Genomics* **45**(2):244-249.
- McNeil S., Zeng C., Harrington K.S., Hiebert S., Lian J.B., Stein J.L., van Wijnen A.J., and Stein G.S. (1999). The t(8;21) chromosomal translocation in acute myelogenous leukemia modifies intranuclear targeting of the AML1/CBFalpha2 transcription factor. *Proc Natl Acad Sci U S A* **96**(26):14882-14887.
- Mellgren G., Borud B., Hoang T., Yri O.E., Fladeby C., Lien E.A., and Lund J. (2003). Characterization of receptor-interacting protein RIP140 in the regulation of SF-1 responsive target genes. *Mol Cell Endocrinol* **203**(1-2):91-103.

5. Bibliography

- Merkatz I.R., Nitowsky H.M., Macri J.N., and Johnson W.E. (1984). An association between low maternal serum alpha-fetoprotein and fetal chromosomal abnormalities. *Am J Obstet Gynecol* **148**(7):886-894.
- Meyer G., Cabrera Socorro A., Perez Garcia C.G., Martinez Millan L., Walker N., and Caput D. (2004). Developmental roles of p73 in Cajal-Retzius cells and cortical patterning. *J Neurosci* **24**(44):9878-9887.
- Michiels F., Stam J.C., Hordijk P.L., van der Kammen R.A., Ruuls-Van Stalle L., Feltkamp C.A., and Collard J.G. (1997). Regulated membrane localization of Tiam1, mediated by the NH2-terminal pleckstrin homology domain, is required for Rac-dependent membrane ruffling and C-Jun NH2-terminal kinase activation. *J Cell Biol* **137**(2):387-398.
- Mikkelsen M. (1977). Down syndrome: cytogenetical epidemiology. *Hereditas* **86**(1):45-50.
- Min Jou W., Haegeman G., Ysebaert M., and Fiers W. (1972). Nucleotide sequence of the gene coding for the bacteriophage MS2 coat protein. *Nature* **237**(5350):82-88.
- Mohney R.P., Das M., Bivona T.G., Hanes R., Adams A.G., Philips M.R., and O'Bryan J.P. (2003). Intersectin activates Ras but stimulates transcription through an independent pathway involving JNK. *J Biol Chem* **278**(47):47038-47045.
- Moore C.S. (2006). Postnatal lethality and cardiac anomalies in the Ts65Dn Down syndrome mouse model. *Mamm Genome* **17**(10):1005-1012.
- Motonaga K., Itoh M., Becker L.E., Goto Y., and Takashima S. (2002). Elevated expression of beta-site amyloid precursor protein cleaving enzyme 2 in brains of patients with Down syndrome. *Neurosci Lett* **326**(1):64-66.
- Muresan Z. and Muresan V. (2004). A phosphorylated, carboxy-terminal fragment of beta-amyloid precursor protein localizes to the splicing factor compartment. *Hum Mol Genet* **13**(5):475-488.
- Murphy M., Insoft R.M., Pike-Nobile L., and Epstein L.B. (1995). A hypothesis to explain the immune defects in Down syndrome. *Prog Clin Biol Res* **393**:147-167.
- Nagy Z. (1999). Mechanisms of neuronal death in Down's syndrome. *J Neural Transm Suppl* **57**:233-245.
- Nakaya T. and Suzuki T. (2006). Role of APP phosphorylation in FE65-dependent gene transactivation mediated by AICD. *Genes Cells* **11**(6):633-645.
- Nimura K., Ishida C., Koriyama H., Hata K., Yamanaka S., Li E., Ura K., and Kaneda Y. (2006). Dnmt3a2 targets endogenous Dnmt3L to ES cell chromatin and induces regional DNA methylation. *Genes Cells* **11**(10):1225-1237.
- O'Doherty A., Ruf S., Mulligan C., Hildreth V., Errington M.L., Cooke S., Sesay A., Modino S., Vanes L., Hernandez D., et al. (2005). An aneuploid mouse strain carrying human chromosome 21 with Down syndrome phenotypes. *Science* **309**(5743):2033-2037.
- O'Rourke N.A., Meyer T., and Chandy G. (2005). Protein localization studies in the age of 'Omics'. *Curr Opin Chem Biol* **9**(1):82-87.
- Oitzl M.S., Reichardt H.M., Joels M., and de Kloet E.R. (2001). Point mutation in the mouse glucocorticoid receptor preventing DNA binding impairs spatial memory. *Proc Natl Acad Sci U S A* **98**(22):12790-12795.
- Okorokov A.L., Rubbi C.P., Metcalfe S., and Milner J. (2002). The interaction of p53 with the nuclear matrix is mediated by F-actin and modulated by DNA damage. *Oncogene* **21**(3):356-367.
- Olson L.E., Roper R.J., Baxter L.L., Carlson E.J., Epstein C.J., and Reeves R.H. (2004). Down syndrome mouse models Ts65Dn, Ts1Cje, and Ms1Cje/Ts65Dn exhibit variable severity of cerebellar phenotypes. *Dev Dyn* **230**(3):581-589.
- Olson L.E., Roper R.J., Sengstaken C.L., Peterson E.A., Aquino V., Galdzicki Z., Siarey R., Pletnikov M., Moran T.H., and Reeves R.H. (2007). Trisomy for the Down syndrome 'critical region' is necessary but not sufficient for brain phenotypes of trisomic mice. *Hum Mol Genet* **16**(7):774-782.
- Osman W., Laine S., and Zilliacus J. (2006). Functional interaction between the glucocorticoid receptor and GANP/MCM3AP. *Biochem Biophys Res Commun* **348**(4):1239-1244.
- Pacheco T.R., Coelho M.B., Desterro J.M., Mollet I., and Carmo-Fonseca M. (2006). In vivo requirement of the small subunit of U2AF for recognition of a weak 3' splice site. *Mol Cell Biol* **26**(21):8183-8190.
- Park K.S., Lee R.D., Kang S.K., Han S.Y., Park K.L., Yang K.H., Song Y.S., Park H.J., Lee Y.M., Yun Y.P., et al. (2004). Neuronal differentiation of embryonic midbrain cells by upregulation of peroxisome proliferator-activated receptor-gamma via the JNK-dependent pathway. *Exp Cell Res* **297**(2):424-433.
- Parker M.G., Christian M., and White R. (2006). The nuclear receptor co-repressor RIP140 controls the expression of metabolic gene networks. *Biochem Soc Trans* **34**(Pt 6):1103-1106.
- Patterson S.D. and Aebersold R. (1995). Mass spectrometric approaches for the identification of gel-separated proteins. *Electrophoresis* **16**(10):1791-1814.
- Pennington B.F., Moon J., Edgin J., Stedron J., and Nadel L. (2003). The neuropsychology of Down syndrome: evidence for hippocampal dysfunction. *Child Dev* **74**(1):75-93.
- Peretz H. and Elson D. (1976). Synthesis of a cleavable protein-crosslinking reagent for the investigation of ribosome structure. *Eur J Biochem* **63**(1):77-82.

- Peri S., Navarro J.D., Amanchy R., Kristiansen T.Z., Jonnalagadda C.K., Surendranath V., Niranjan V., Muthusamy B., Gandhi T.K., Gronborg M., *et al.* (2003). Development of human protein reference database as an initial platform for approaching systems biology in humans. *Genome Res* **13**(10):2363-2371.
- Persico M., Ceol A., Gavrilu C., Hoffmann R., Florio A., and Cesareni G. (2005). HomoMINT: an inferred human network based on orthology mapping of protein interactions discovered in model organisms. *BMC Bioinformatics* **6 Suppl 4**:S21.
- Peyrl A., Weitzdoerfer R., Gulesserian T., Fountoulakis M., and Lubec G. (2002). Aberrant expression of signaling-related proteins 14-3-3 gamma and RACK1 in fetal Down syndrome brain (trisomy 21). *Electrophoresis* **23**(1):152-157.
- Pheasant M. and Mattick J.S. (2007). Raising the estimate of functional human sequences. *Genome Res* **17**(9):1245-1253.
- Phizicky E.M. and Fields S. (1995). Protein-protein interactions: methods for detection and analysis. *Microbiol Rev* **59**(1):94-123.
- Pouli A.E., Kennedy H.J., Schofield J.G., and Rutter G.A. (1998). Insulin targeting to the regulated secretory pathway after fusion with green fluorescent protein and firefly luciferase. *Biochem J* **331 (Pt 2)**:669-675.
- Poustka A., Pohl T., Barlow D.P., Zehetner G., Craig A., Michiels F., Ehrich E., Frischauf A.M., and Lehrach H. (1986). Molecular approaches to mammalian genetics. *Cold Spring Harb Symp Quant Biol* **51 Pt 1**:131-139.
- Pozniak C.D., Barnabe-Heider F., Rymar V.V., Lee A.F., Sadikot A.F., and Miller F.D. (2002). p73 is required for survival and maintenance of CNS neurons. *J Neurosci* **22**(22):9800-9809.
- Prandini P., Deutsch S., Lyle R., Gagnebin M., Delucinge Vivier C., Delorenzi M., Gehrig C., Descombes P., Sherman S., Dagna Bricarelli F., *et al.* (2007). Natural gene-expression variation in Down syndrome modulates the outcome of gene-dosage imbalance. *Am J Hum Genet* **81**(2):252-263.
- Prenzel N., Fischer O.M., Streit S., Hart S., and Ullrich A. (2001). The epidermal growth factor receptor family as a central element for cellular signal transduction and diversification. *Endocr Relat Cancer* **8**(1):11-31.
- Pruitt K.D., Tatusova T., and Maglott D.R. (2007). NCBI reference sequences (RefSeq): a curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids Res* **35**(Database issue):D61-65.
- Pueschel S.M., Jackson I.M., Giesswein P., Dean M.K., and Pezzullo J.C. (1991a). Thyroid function in Down syndrome. *Res Dev Disabil* **12**(3):287-296.
- Pueschel S.M., Louis S., and McKnight P. (1991b). Seizure disorders in Down syndrome. *Arch Neurol* **48**(3):318-320.
- Pui C.H. (1995). Childhood leukemias. *N Engl J Med* **332**(24):1618-1630.
- Putkey J.A., Kleerekoper Q., Gaertner T.R., and Waxham M.N. (2003). A new role for IQ motif proteins in regulating calmodulin function. *J Biol Chem* **278**(50):49667-49670.
- Rabinowe S.L., Rubin I.L., George K.L., Adri M.N., and Eisenbarth G.S. (1989). Trisomy 21 (Down's syndrome): autoimmunity, aging and monoclonal antibody-defined T-cell abnormalities. *J Autoimmun* **2**(1):25-30.
- Rahmani Z., Blouin J.L., Creau-Goldberg N., Watkins P.C., Mattei J.F., Poissonnier M., Prieur M., Chettouh Z., Nicole A., Aurias A., *et al.* (1989). Critical role of the D21S55 region on chromosome 21 in the pathogenesis of Down syndrome. *Proc Natl Acad Sci U S A* **86**(15):5958-5962.
- Ralsler M., Nonhoff U., Albrecht M., Lengauer T., Wanker E.E., Lehrach H., and Krobitch S. (2005). Ataxin-2 and huntingtin interact with endophilin-A complexes to function in plastin-associated pathways. *Hum Mol Genet* **14**(19):2893-2909.
- Ramani A.K., Bunescu R.C., Mooney R.J., and Marcotte E.M. (2005). Consolidating the set of known human protein-protein interactions in preparation for large-scale mapping of the human interactome. *Genome Biol* **6**(5):R40.
- Rappsilber J., Ryder U., Lamond A.I., and Mann M. (2002). Large-scale proteomic analysis of the human spliceosome. *Genome Res* **12**(8):1231-1245.
- Rasmussen M.D., Deshpande M.S., Karypis G., Johnson J., Crow J.A., and Retzel E.F. (2003). wCLUTO: a Web-enabled clustering toolkit. *Plant Physiol* **133**(2):510-516.
- Rastan S. and Beeley L.J. (1997). Functional genomics: going forwards from the databases. *Curr Opin Genet Dev* **7**(6):777-783.
- Ratel D., Ravanat J.L., Charles M.P., Platet N., Breuillaud L., Lunardi J., Berger F., and Wion D. (2006). Undetectable levels of N6-methyl adenine in mouse DNA: Cloning and analysis of PRED28, a gene coding for a putative mammalian DNA adenine methyltransferase. *FEBS Lett* **580**(13):3179-3184.
- Raz N., Torres I.J., Briggs S.D., Spencer W.D., Thornton A.E., Loken W.J., Gunning F.M., McQuain J.D., Driesen N.R., and Acker J.D. (1995). Selective neuroanatomic abnormalities in Down's syndrome and their cognitive correlates: evidence from MRI morphometry. *Neurology* **45**(2):356-366.
- Reeves R.H. (2006). Down syndrome mouse models are looking up. *Trends Mol Med* **12**(6):237-240.
- Reeves R.H., Irving N.G., Moran T.H., Wohn A., Kitt C., Sisodia S.S., Schmidt C., Bronson R.T., and Davisson M.T. (1995). A mouse model for Down syndrome exhibits learning and behaviour deficits. *Nat Genet* **11**(2):177-184.
- Reimann F. and Ashcroft F.M. (1999). Inwardly rectifying potassium channels. *Curr Opin Cell Biol* **11**(4):503-508.

5. Bibliography

- Remm M., Storm C.E., and Sonnhammer E.L. (2001). Automatic clustering of orthologs and in-paralogs from pairwise species comparisons. *J Mol Biol* **314**(5):1041-1052.
- Rentero C. and Puigdomenech P. (2006). Specific use of start codons and cellular localization of splice variants of human phosphodiesterase 9A gene. *BMC Mol Biol* **7**:39.
- Rex A.P. and Preus M. (1982). A diagnostic index for Down syndrome. *J Pediatr* **100**(6):903-906.
- Reymond A., Marigo V., Yaylaoglu M.B., Leoni A., UCLA C., Scamuffa N., Caccioppoli C., Dermitzakis E.T., Lyle R., Banfi S., et al. (2002). Human chromosome 21 gene expression atlas in the mouse. *Nature* **420**(6915):582-586.
- Richtsmeier J.T., Baxter L.L., and Reeves R.H. (2000). Parallels of craniofacial maldevelopment in Down syndrome and Ts65Dn mice. *Dev Dyn* **217**(2):137-145.
- Richtsmeier J.T., Zumwalt A., Carlson E.J., Epstein C.J., and Reeves R.H. (2002). Craniofacial phenotypes in segmentally trisomic mouse models for Down syndrome. *Am J Med Genet* **107**(4):317-324.
- Rigaut G., Shevchenko A., Rutz B., Wilm M., Mann M., and Seraphin B. (1999). A generic protein purification method for protein complex characterization and proteome exploration. *Nat Biotechnol* **17**(10):1030-1032.
- Robinson S.W., Morris C.D., Goldmuntz E., Reller M.D., Jones M.A., Steiner R.D., and Maslen C.L. (2003). Missense mutations in CRELD1 are associated with cardiac atrioventricular septal defects. *Am J Hum Genet* **72**(4):1047-1052.
- Roper R.J., Baxter L.L., Saran N.G., Klinedinst D.K., Beachy P.A., and Reeves R.H. (2006). Defective cerebellar response to mitogenic Hedgehog signaling in Down [corrected] syndrome mice. *Proc Natl Acad Sci U S A* **103**(5):1452-1456.
- Rosen J., He M., Umbricht C., Alexander H.R., Dackiw A.P., Zeiger M.A., and Libutti S.K. (2005). A six-gene model for differentiating benign from malignant thyroid tumors on the basis of gene expression. *Surgery* **138**(6):1050-1056; discussion 1056-1057.
- Roth G.M., Sun B., Greensite F.S., Lott I.T., and Dietrich R.B. (1996). Premature aging in persons with Down syndrome: MR findings. *AJNR Am J Neuroradiol* **17**(7):1283-1289.
- Rothermel B., Vega R.B., Yang J., Wu H., Bassel-Duby R., and Williams R.S. (2000). A protein encoded within the Down syndrome critical region is enriched in striated muscles and inhibits calcineurin signaling. *J Biol Chem* **275**(12):8719-8725.
- Rual J.F., Hirozane-Kishikawa T., Hao T., Bertin N., Li S., Dricot A., Li N., Rosenberg J., Lamesch P., Vidalain P.O., et al. (2004). Human ORFeome version 1.1: a platform for reverse proteomics. *Genome Res* **14**(10B):2128-2135.
- Rual J.F., Venkatesan K., Hao T., Hirozane-Kishikawa T., Dricot A., Li N., Berriz G.F., Gibbons F.D., Dreze M., Ayivi-Guedehoussou N., et al. (2005). Towards a proteome-scale map of the human protein-protein interaction network. *Nature*.
- Ruiz-Leon Y. and Pascual A. (2001). Brain-derived neurotrophic factor stimulates beta-amyloid gene promoter activity by a Ras-dependent/AP-1-independent mechanism in SH-SY5Y neuroblastoma cells. *J Neurochem* **79**(2):278-285.
- Rumble B., Retallack R., Hilbich C., Simms G., Multhaup G., Martins R., Hockey A., Montgomery P., Beyreuther K., and Masters C.L. (1989). Amyloid A4 protein and its precursor in Down's syndrome and Alzheimer's disease. *N Engl J Med* **320**(22):1446-1452.
- Rzhetsky A., Iossifov I., Koike T., Krauthammer M., Kra P., Morris M., Yu H., Duboue P.A., Weng W., Wilbur W.J., et al. (2004). GeneWays: a system for extracting, analyzing, visualizing, and integrating molecular pathway data. *J Biomed Inform* **37**(1):43-53.
- Sago H., Carlson E.J., Smith D.J., Kilbridge J., Rubin E.M., Mobley W.C., Epstein C.J., and Huang T.T. (1998). Ts1Cje, a partial trisomy 16 mouse model for Down syndrome, exhibits learning and behavioral abnormalities. *Proc Natl Acad Sci U S A* **95**(11):6256-6261.
- Saini H.K., Griffiths-Jones S., and Enright A.J. (2007). Genomic analysis of human microRNA transcripts. *Proc Natl Acad Sci U S A* **104**(45):17719-17724.
- Sambrook J., Fritsch E.F., and Maniatis T. (1989). *Molecular Cloning. A Laboratory Manual*. Cold Spring Harbour, Cold Spring Harbor Laboratory Press.
- Sanger F., Air G.M., Barrell B.G., Brown N.L., Coulson A.R., Fiddes C.A., Hutchison C.A., Slocombe P.M., and Smith M. (1977). Nucleotide sequence of bacteriophage phi X174 DNA. *Nature* **265**(5596):687-695.
- Saran N.G., Pletcher M.T., Natale J.E., Cheng Y., and Reeves R.H. (2003). Global disruption of the cerebellar transcriptome in a Down syndrome mouse model. *Hum Mol Genet* **12**(16):2013-2019.
- Saunders M., Eldeen M.B., Valle L.D., Reiss K., Peruzzi F., Marnett G., Gelman B.B., Khalili K., Amini S., and Sawaya B.E. (2005). p73 modulates HIV-1 Tat transcriptional and apoptotic activities in human astrocytes. *Apoptosis* **10**(6):1419-1431.
- Savino T.M., Bastos R., Jansen E., and Hernandez-Verdun D. (1999). The nucleolar antigen Nop52, the human homologue of the yeast ribosomal RNA processing RRP1, is recruited at late stages of nucleologenesis. *J Cell Sci* **112** (Pt 12):1889-1900.
- Sawa A. (1999). Neuronal cell death in Down's syndrome. *J Neural Transm Suppl* **57**:87-97.
- Schena M., Shalon D., Davis R.W., and Brown P.O. (1995). Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science* **270**(5235):467-470.

- Schinzel A. (1976). Letter: Does full monosomy 21 exist? A comment to the paper: A male infant with monosomy 21 by Y. Kaneko, T. Ikeuchi, M. Sasaki, Y. Satake and S. Kuwajima *Humangenetik* 29, 1-7 (1975). *Hum Genet* **32**(1):105-107.
- Schwikowski B., Uetz P., and Fields S. (2000). A network of protein-protein interactions in yeast. *Nat Biotechnol* **18**(12):1257-1261.
- Seger R., Buchinger G., and Stroder J. (1977). On the influence of age on immunity in Down's syndrome. *Eur J Pediatr* **124**(2):77-87.
- Sethupathy P., Borel C., Gagnebin M., Grant G.R., Deutsch S., Elton T.S., Hatzigeorgiou A.G., and Antonarakis S.E. (2007). Human microRNA-155 on chromosome 21 differentially interacts with its polymorphic target in the AGTR1 3' untranslated region: a mechanism for functional single-nucleotide polymorphisms related to phenotypes. *Am J Hum Genet* **81**(2):405-413.
- Shetty J., Klotz K.L., Wolkowicz M.J., Flickinger C.J., and Herr J.C. (2007). Radial spoke protein 44 (human meichroacidin) is an axonemal alloantigen of sperm and cilia. *Gene* **396**(1):93-107.
- Shi Y.Y., Miller G.A., Qian H., and Bomsztyk K. (2006). Free-energy distribution of binary protein-protein binding suggests cross-species interactome differences. *Proc Natl Acad Sci U S A* **103**(31):11527-11532.
- Shibuya K., Kudoh J., Obayashi I., Shimizu A., Sasaki T., Minoshima S., and Shimizu N. (2004). Comparative genomics of the keratin-associated protein (KAP) gene clusters in human, chimpanzee, and baboon. *Mamm Genome* **15**(3):179-192.
- Simpson J.C., Wellenreuther R., Poustka A., Pepperkok R., and Wiemann S. (2000). Systematic subcellular localization of novel proteins identified by large-scale cDNA sequencing. *EMBO Rep* **1**(3):287-292.
- Simpson R.J. (2002). *Proteins and Proteomics: A Laboratory Manual*, Cold Spring Harbor Laboratory Press; Lab Manual edition.
- Sittler A., Walter S., Wedemeyer N., Hasenbank R., Scherzinger E., Eickhoff H., Bates G.P., Lehrach H., and Wanker E.E. (1998). SH3GL3 associates with the Huntingtin exon 1 protein and promotes the formation of polyglutamine-containing protein aggregates. *Mol Cell* **2**(4):427-436.
- Smith G.P. (1985). Filamentous fusion phage: novel expression vectors that display cloned antigens on the virion surface. *Science* **228**(4705):1315-1317.
- Soares L., Seroogy C., Skrenta H., Anandasabapathy N., Lovelace P., Chung C.D., Engleman E., and Fathman C.G. (2004). Two isoforms of otubain 1 regulate T cell anergy via GRAIL. *Nat Immunol* **5**(1):45-54.
- Sonnhammer E.L., von Heijne G., and Krogh A. (1998). A hidden Markov model for predicting transmembrane helices in protein sequences. *Proc Int Conf Intell Syst Mol Biol* **6**:175-182.
- Sprenger J., Fink J.L., and Teasdale R.D. (2006). Evaluation and comparison of mammalian subcellular localization prediction methods. *BMC Bioinformatics* **7 Suppl 5**:S3.
- Stanyon C.A., Liu G., Mangiola B.A., Patel N., Giot L., Kuang B., Zhang H., Zhong J., and Finley R.L., Jr. (2004). A Drosophila protein-interaction map centered on cell-cycle regulators. *Genome Biol* **5**(12):R96.
- Stelzl U. and Wanker E.E. (2006). The value of high quality protein-protein interaction networks for systems biology. *Curr Opin Chem Biol* **10**(6):551-558.
- Stelzl U., Worm U., Lalowski M., Haenig C., Brembeck F.H., Goehler H., Stroedicke M., Zenkner M., Schoenherr A., Koeppen S., et al. (2005). A human protein-protein interaction network: a resource for annotating the proteome. *Cell* **122**(6):957-968.
- Stertz S., Reichelt M., Krijnse-Locker J., Mackenzie J., Simpson J.C., Haller O., and Kochs G. (2006). Interferon-induced, antiviral human MxA protein localizes to a distinct subcompartment of the smooth endoplasmic reticulum. *J Interferon Cytokine Res* **26**(9):650-660.
- Stumpf M.P., Wiuf C., and May R.M. (2005). Subnets of scale-free networks are not scale-free: sampling properties of networks. *Proc Natl Acad Sci U S A* **102**(12):4221-4224.
- Sultan M., Piccini I., Balzereit D., Herwig R., Saran N.G., Lehrach H., Reeves R.H., and Yaspo M.L. (2007). Gene expression variation in Down's syndrome mice allows prioritization of candidate genes. *Genome Biol* **8**(5):R91.
- Suzuki H., Tashiro S., Hira S., Sun J., Yamazaki C., Zenke Y., Ikeda-Saito M., Yoshida M., and Igarashi K. (2004). Heme regulates gene expression by triggering Crm1-dependent nuclear export of Bach1. *Embo J* **23**(13):2544-2553.
- Swatton J.E., Sellers L.A., Faull R.L., Holland A., Iritani S., and Bahn S. (2004). Increased MAP kinase activity in Alzheimer's and Down syndrome but not in schizophrenia human brain. *Eur J Neurosci* **19**(10):2711-2719.
- Sweatt J.D. (2004). Mitogen-activated protein kinases in synaptic plasticity and memory. *Curr Opin Neurobiol* **14**(3):311-317.
- Takei Y., Assenberg M., Tsujimoto G., and Laskey R. (2002). The MCM3 acetylase MCM3AP inhibits initiation, but not elongation, of DNA replication via interaction with MCM3. *J Biol Chem* **277**(45):43121-43125.
- Takei Y. and Tsujimoto G. (1998). Identification of a novel MCM3-associated protein that facilitates MCM3 nuclear localization. *J Biol Chem* **273**(35):22177-22180.
- Tan Y.H., Schneider E.L., Tischfield J., Epstein C.J., and Ruddle F.H. (1974). Human chromosome 21 dosage: effect on the expression of the interferon induced antiviral state. *Science* **186**(4158):61-63.

5. Bibliography

- Tanaka T., Kurokawa M., Ueki K., Tanaka K., Imai Y., Mitani K., Okazaki K., Sagata N., Yazaki Y., Shibata Y., *et al.* (1996). The extracellular signal-regulated kinase pathway phosphorylates AML1, an acute myeloid leukemia gene product, and potentially regulates its transactivation ability. *Mol Cell Biol* **16**(7):3967-3979.
- Tessarollo L. (1998). Pleiotropic functions of neurotrophins in development. *Cytokine Growth Factor Rev* **9**(2):125-137.
- Teyssier C., Belguise K., Galtier F., Cavailles V., and Chalbos D. (2003). Receptor-interacting protein 140 binds c-Jun and inhibits estradiol-induced activator protein-1 activity by reversing glucocorticoid receptor-interacting protein 1 effect. *Mol Endocrinol* **17**(2):287-299.
- Thenot S., Charpin M., Bonnet S., and Cavailles V. (1999). Estrogen receptor cofactors expression in breast and endometrial human cancer cells. *Mol Cell Endocrinol* **156**(1-2):85-93.
- Tye B.K. (1999). MCM proteins in DNA replication. *Annu Rev Biochem* **68**:649-686.
- Uetz P. and Finley R.L., Jr. (2005). From protein networks to biological systems. *FEBS Lett* **579**(8):1821-1827.
- Uetz P., Giot L., Cagney G., Mansfield T.A., Judson R.S., Knight J.R., Lockshon D., Narayan V., Srinivasan M., Pochart P., *et al.* (2000). A comprehensive analysis of protein-protein interactions in *Saccharomyces cerevisiae*. *Nature* **403**(6770):623-627.
- Ule J., Jensen K.B., Ruggiu M., Mele A., Ule A., and Darnell R.B. (2003). CLIP identifies Nova-regulated RNA networks in the brain. *Science* **302**(5648):1212-1215.
- Van Lith J.M., Pratt J.J., Beekhuis J.R., and Mantingh A. (1992). Second-trimester maternal serum immunoreactive inhibin as a marker for fetal Down's syndrome. *Prenat Diagn* **12**(10):801-806.
- Van Valkenburgh H., Shern J.F., Sharer J.D., Zhu X., and Kahn R.A. (2001). ADP-ribosylation factors (ARFs) and ARF-like 1 (ARL1) have both specific and shared effectors: characterizing ARL1-binding proteins. *J Biol Chem* **276**(25):22826-22837.
- Vanhecke D. and Janitz M. (2004). High-throughput gene silencing using cell arrays. *Oncogene* **23**(51):8353-8358.
- Verner K. and Schatz G. (1988). Protein translocation across membranes. *Science* **241**(4871):1307-1313.
- von Mering C., Krause R., Snel B., Cornell M., Oliver S.G., Fields S., and Bork P. (2002). Comparative assessment of large-scale data sets of protein-protein interactions. *Nature* **417**(6887):399-403.
- Wald N.J., Kennard A., Hackshaw A., and McGuire A. (1997). Antenatal screening for Down's syndrome. *J Med Screen* **4**(4):181-246.
- Walhout A.J., Temple G.F., Brasch M.A., Hartley J.L., Lorson M.A., van den Heuvel S., and Vidal M. (2000). GATEWAY recombinational cloning: application to the cloning of large numbers of open reading frames or ORFeomes. *Methods Enzymol* **328**:575-592.
- Watanabe H., Fujiyama A., Hattori M., Taylor T.D., Toyoda A., Kuroki Y., Noguchi H., BenKahla A., Lehrach H., Sudbrak R., *et al.* (2004). DNA sequence and comparative analysis of chimpanzee chromosome 22. *Nature* **429**(6990):382-388.
- Watson J.D. (1990). The human genome project: past, present, and future. *Science* **248**(4951):44-49.
- Watson J.D. and Crick F.H.C. (1953). Molecular Structure of Nucleic Acids: A Structure for Deoxyribose Nucleic Acid. *Nature* **171**:737-738.
- Watts D.J. and Strogatz S.H. (1998). Collective dynamics of 'small-world' networks. *Nature* **393**(6684):440-442.
- Wei L.N., Hu X., Chandra D., Seto E., and Faroouqi M. (2000). Receptor-interacting protein 140 directly recruits histone deacetylases for gene silencing. *J Biol Chem* **275**(52):40782-40787.
- Weis S., Weber G., Neuhold A., and Rett A. (1991). Down syndrome: MR quantification of brain structures and comparison with normal control subjects. *AJNR Am J Neuroradiol* **12**(6):1207-1211.
- Wiemann S., Arlt D., Huber W., Wellenreuther R., Schlegler S., Mehrle A., Bechtel S., Sauer mann M., Korf U., Pepperkok R., *et al.* (2004). From ORFeome to biology: a functional genomics pipeline. *Genome Res* **14**(10B):2136-2144.
- Wilkins M.H.F., Stokes A.R., and Wilson H.R. (1953). Molecular Structure of Deoxyribose Nucleic Acids. *Nature* **171**:738-740.
- Wisniewski K.E. (1990). Down syndrome children often have brain with maturation delay, retardation of growth, and cortical dysgenesis. *Am J Med Genet Suppl* **7**:274-281.
- Wisniewski K.E., Laure-Kamionowska M., and Wisniewski H.M. (1984). Evidence of arrest of neurogenesis and synaptogenesis in brains of patients with Down's syndrome. *N Engl J Med* **311**(18):1187-1188.
- Wisniewski K.E., Wisniewski H.M., and Wen G.Y. (1985). Occurrence of neuropathological changes and dementia of Alzheimer's disease in Down's syndrome. *Ann Neurol* **17**(3):278-282.
- Wolvetang E.J., Wilson T.J., Sanij E., Busciglio J., Hatzistavrou T., Seth A., Hertzog P.J., and Kola I. (2003). ETS2 overexpression in transgenic models and in Down syndrome predisposes to apoptosis via the p53 pathway. *Hum Mol Genet* **12**(3):247-255.

- Wong C. and Naumovski L. (1997). Method to screen for relevant yeast two-hybrid-derived clones by coimmunoprecipitation and colocalization of epitope-tagged fragments--application to Bcl-xL. *Anal Biochem* **252**(1):33-39.
- Wong D., Bazopoulou D., Pujol N., Tavernarakis N., and Ewbank J.J. (2007). Genome-wide investigation reveals pathogen-specific and shared signatures in the response of *Caenorhabditis elegans* to infection. *Genome Biol* **8**(9):R194.
- Wu X., Zhu L., Guo J., Zhang D.Y., and Lin K. (2006). Prediction of yeast protein-protein interaction network: insights from the Gene Ontology and annotations. *Nucleic Acids Res* **34**(7):2137-2150.
- Xiao H., Chung J., Kao H.Y., and Yang Y.C. (2003). Tip60 is a co-repressor for STAT3. *J Biol Chem* **278**(13):11197-11204.
- Yang E.J., Ahn Y.S., and Chung K.C. (2001a). Protein kinase Dyrk1 activates cAMP response element-binding protein during neuronal differentiation in hippocampal progenitor cells. *J Biol Chem* **276**(43):39819-39824.
- Yang Q., Xie Y., Eriksson A.M., Nelson B.D., and DePierre J.W. (2001b). Further evidence for the involvement of inhibition of cell proliferation and development in thymic and splenic atrophy induced by the peroxisome proliferator perfluorooctanoic acid in mice. *Biochem Pharmacol* **62**(8):1133-1140.
- Yang W., Wang Q., Kaness S.J., Murray J.M., and Nishikura K. (2004). Altered RNA editing of serotonin 5-HT_{2C} receptor induced by interferon: implications for depression associated with cytokine therapy. *Brain Res Mol Brain Res* **124**(1):70-78.
- Yeger-Lotem E., Sattath S., Kashtan N., Itzkovitz S., Milo R., Pinter R.Y., Alon U., and Margalit H. (2004). Network motifs in integrated cellular networks of transcription-regulation and protein-protein interaction. *Proc Natl Acad Sci U S A* **101**(16):5934-5939.
- Yi Y.W., Kim D., Jung N., Hong S.S., Lee H.S., and Bae I. (2000). Gadd45 family proteins are coactivators of nuclear hormone receptors. *Biochem Biophys Res Commun* **272**(1):193-198.
- Yin L., Xiang Y., Zhu D.Y., Yan N., Huang R.H., Zhang Y., and Wang D.C. (2005). Crystal structure of human SH3BGR1 protein: the first structure of the human SH3BGR family representing a novel class of thioredoxin fold proteins. *Proteins* **61**(1):213-216.
- Yordy J.S. and Muise-Helmericks R.C. (2000). Signal transduction and the Ets family of transcription factors. *Oncogene* **19**(55):6503-6513.
- Yoshima T., Yura T., and Yanagi H. (1998). Novel testis-specific protein that interacts with heat shock factor 2. *Gene* **214**(1-2):139-146.
- Yuan Z.L., Guan Y.J., Chatterjee D., and Chin Y.E. (2005). Stat3 dimerization regulated by reversible acetylation of a single lysine residue. *Science* **307**(5707):269-273.
- Yuryev A., Patturajan M., Litingtung Y., Joshi R.V., Gentile C., Gebara M., and Corden J.L. (1996). The C-terminal domain of the largest subunit of RNA polymerase II interacts with a novel set of serine/arginine-rich proteins. *Proc Natl Acad Sci U S A* **93**(14):6975-6980.
- Zatyka M., Priestley M., Ladusans E.J., Fryer A.E., Mason J., Latif F., and Maher E.R. (2005). Analysis of CRELD1 as a candidate 3p25 atrioventricular septal defect locus (AVSD2). *Clin Genet* **67**(6):526-528.
- Zavzavadjian J.R., Couture S., Park W.S., Whalen J., Lyon S., Lee G., Fung E., Mi Q., Liu J., Wall E., *et al.* (2007). The alliance for cellular signaling plasmid collection: a flexible resource for protein localization studies and signaling pathway analysis. *Mol Cell Proteomics* **6**(3):413-424.
- Zhang L.V., King O.D., Wong S.L., Goldberg D.S., Tong A.H., Lesage G., Andrews B., Bussey H., Boone C., and Roth F.P. (2005). Motifs, themes and thematic maps of an integrated *Saccharomyces cerevisiae* interaction network. *J Biol* **4**(2):6.
- Zhou C., Cunningham L., Marcus A.I., Li Y., and Kahn R.A. (2006). Arl2 and Arl3 regulate different microtubule-dependent processes. *Mol Biol Cell* **17**(5):2476-2487.
- Zhou Z., Licklider L.J., Gygi S.P., and Reed R. (2002). Comprehensive proteomic analysis of the human spliceosome. *Nature* **419**(6903):182-185.
- Zhu C., Wang X., Deinum J., Huang Z., Gao J., Modjtahedi N., Neagu M.R., Nilsson M., Eriksson P.S., Hagberg H., *et al.* (2007). Cyclophilin A participates in the nuclear translocation of apoptosis-inducing factor in neurons after cerebral hypoxia-ischemia. *J Exp Med* **204**(8):1741-1748.
- Ziauddin J. and Sabatini D.M. (2001). Microarrays of cells expressing defined cDNAs. *Nature* **411**(6833):107-110.
- Zigman W., Schupf N., Haveman M., and Silverman W. (1997). The epidemiology of Alzheimer disease in intellectual disability: results and recommendations from an international conference. *J Intellect Disabil Res* **41** (Pt 1):76-80.
- Zolotukhin A.S., Tan W., Bear J., Smulevitch S., and Felber B.K. (2002). U2AF participates in the binding of TAP (NXF1) to mRNA. *J Biol Chem* **277**(6):3935-3942.
- Zou S., Meadows S., Sharp L., Jan L.Y., and Jan Y.N. (2000). Genome-wide study of aging and oxidative stress response in *Drosophila melanogaster*. *Proc Natl Acad Sci U S A* **97**(25):13726-13731.