

6. Literaturverzeichnis

Adams D.J., Barakeh J., Laskey R., Van Breemen C.. Ion channels and regulation of intracellular calcium in vascular endothelial cells. *FASEB J.* 3: 2389-400, 1989.

Bauersachs J., Popp R., Fleming I., Busse R.. Nitric oxide and endothelium-derived hyperpolarizing factor: formation and interactions. *Prostaglandins Leukot. Essent. Fatty Acids* 57: 439-446, 1997.

Bauersachs J., Popp R., Fleming I., Busse R.. Nitric oxide and endothelium-derived hyperpolarizing factor: formation and interactions. *Prostaglandins Leukot. Essent. Fatty Acids* 57: 439-446, 1997.

Ben-Ishay D., Yagil Y.. The sabra hypertension-prone and -resistant strains. *Experimental and Genetic Models of Hypertension*, edited by Ganten D., de Jong W., Amsterdam, Elsevier Science B.V.: 272, 1994.

Beny J.L.. Information networks in the arterial wall. *News Physiol. Sci.* 14: 68-73, 1999.

Bolz S.S., Fissthaler B., Pieperhoff S., de Wit C., Fleming I., Busse R., Pohl U.. Antisense oligonucleotides against cytochrome P450 2C8 attenuate EDHF-mediated Ca^{2+} changes and dilation in isolated resistance arteries. *FASEB J.* 14: 255-260, 2000.

Busse R., Fleming I., Hecker M.. Signal transduction in endothelium-dependent vasodilatation. *Eur. Heart J.* 14 (Suppl. I): 2-9, 1993.

Busse R., Fleming I.. Pulsatile stretch and shear stress: physical stimuli determining the production of endothelium-derived relaxing factors. *J. Vasc. Res.* 35: 73-84, 1998.

Busse R., Trogisch G., Bassenge E.. The role of endothelium in the control of vascular tone. *Basic Res. Cardiol.* 80: 475-90, 1985.

Cantiello, H.F., Stow, J.L., Prat., A.G., Ausiello, D.A.. Actin filaments regulate epithelial Na^+ channel activity. *Am. J. Physiol.* 261:C882-888, 1991.

Colden-Stanfield, M., Schilling, W.P., Possani, L.D., Kunze, D.. Bradykinin-induced potassium current in cultured bovine aortic endothelial cells. *J. Membr. Biol.* 116: 227-238, 1990.

Cooke, J.P., Rossitch, E., Andon, N.A., Loscalzo, J., Dzau, V.J.. Flow activates an endothelial potassium channel to release an endogenous nitrovasodilator. *J. Clin. Invest.* 88: 1663-1671, 1991.

Craelius W., Chen V., el-Sherif N.. Stretch activated ion channels in ventricular myocytes. *Biosci. Rep.* 8: 407-414, 1988.

Davies P.F., Barbee K.A., Volin M.V., Robotewskyj A., Chen J., Joseph L., Griem M.L., Wernick M.N., Jacobs E., Polacek D.C., de Paola N., Barakat A.I.. Spatial relationships in early signaling events of flow-mediated endothelial mechanotransduction. *Annu Rev. Physiol.* 59: 527-549, 1997.

Davies P.F.. Flow-mediated endothelial mechanotransduction. *Physiol. Rev.* 75: 519-560, 1995.

Davies, P.F., Tripathi, S.C.. Mechanical stress mechanisms and the cell. An endothelial paradigm. *Circ. Res.* 72: 239-245, 1993.

De Mendonca M., Garay R.P., Ben-Ishay D., Meyer P.. Abnormal erythrocyte cation transport in primary hypertension: Clinical and experimental studies. *Hypertension* 3: 179-183, 1981.

Dewey, C.F., Bussolari, S.R., Gimbrone, M.A., Davies, P.F.. The dynamic response of vascular endothelial cells to fluid shear stress. *J. Biomech. Eng.* 103: 177-185, 1981.

Diamond S.L., Sharefkin J.B., Dieffenbach C., Frasier-Scott K., McIntire L.V., Eskin S.G.. Tissue plasminogen activator messenger RNA levels increase in cultured human endothelial cells exposed to laminar shear stress. *J. Cell. Physiol.* 143: 364-371, 1990.

Dimmeler S., Fleming I., Fisslthaler B., Hermann C., Busse R., Zeiher A.M.. Activation of nitric oxide synthase in endothelial cells by Akt-dependent phosphorylation. *Nature* 399: 601-605, 1999.

Edwards G., Dora K.A., Gardener M.J., Garland C.J., Weston A.H.. K^+ is an endothelium-derived hyperpolarizing factor in rat arteries. *Nature* 396: 269-272, 1998.

Egashira K., Suzuki S., Hirooka Y., Kai H., Sugimachi M., Imaizumi T., Takeshita A.. Impaired endothelium-dependent vasodilation of large epicardial and resistance coronary arteries in patients with essential hypertension. Different responses to acetylcholine and substance P. *Hypertension* 25: 210-216, 1995.

Fasolato C., Nilius B.. Store depletion triggers the calcium release-activated calcium current (ICRAC) in macrovascular endothelial cells: a comparison with Jurkat and embryonic kidney cell lines. *Pflügers Arch.* 436: 69-74, 1998.

Fichtner, H., Fröbe, U., Busse, R., Kohlhardt, M.. Single non-selective cation channels and Ca^{2+} activated K^+ channels in aortic endothelial cells. *J. Membrane Biol.* 98: 125-133, 1987.

Fleming I., Bauersachs J., Schäfer A., Scholz D., Aldershvile J., Busse R.. Isometric contraction induces the Ca^{2+} -independent activation of the endothelial nitric oxide synthase. *Proc. Natl. Acad. Sci. USA* 96: 1123-1128, 1999.

Franco, A.; Lansman, J.B.. Calcium entry through stretch-inactivated ion channels in mdx myotubes. *Nature* 344: 670-673, 1990.

Frangos, J.A., McIntire, L.V., Eskin, S.G.. Shear stress induced stimulation of mammalian cell metabolism. *Biotech. Bioeng.* 32:1053-1060, 1988.

Gibbons G.H., Dzau VJ.. The emerging concept of vascular remodeling. *New Engl. J. Med.* 330: 1431-1438, 1994.

Griffith, T.M., Edwards, D.H., Newby, A.C., Lewis, M.J., Henderson, A.H.. Production of endothelium-derived relaxant factor is dependent on oxidative phosphorylation and extracellular calcium. *Cardiovasc. Res.* 20: 7-12, 1986.

Griffith, T.M., Edwards, D.H.. Myogenic autoregulation of flow may be inversely related to endothelium-derived relaxing-factor activity. *Am. J. Physiol.* 258: H1171-1180, 1990.

Guharay F., Sachs F.. Stretch-activated single ion channel currents in tissue-cultured embryonic chick skeletal muscle. *J. Physiol.* 352: 685-701, 1984.

Hamill, O.P., Marty, A., Neher, E., Sakmann, B., Sigworth, F.J.. Improved patch-clamp techniques with high-resolution current recordings from cells and cell-free membrane patches. *Pflügers Arch.* 391: 85-100, 1981.

Hanada T., Hashimoto M., Nosaka S., Sasaki T., Nakayama K., Masumura S., Yamauchi M., Tamura K.. Shear stress enhances prostacyclin release from endocardial endothelial cells. *Life Sci.* 66: 215-220, 2000.

Hille, B.: Ionic channels in excitable membranes. pp 1-19, Sinauer Ass. Inc. (editors), Sunderland, Massachusetts, 1984.

Hoyer J., Köhler R., Distler A.. Mechanosensitive Ca^{2+} oscillations and STOC activation in endothelial cells. *FASEB J.* 12: 359-366, 1998.

Hoyer J., Köhler R., Distler A.. Mechanosensitive cation channels in aortic endothelium of normotensive and hypertensive rats. *Hypertension* 30: 112-119, 1997.

Hoyer J., Köhler R., Haase W., Distler A.. Up-regulation of pressure-activated Ca^{2+} -permeable cation channel in intact vascular endothelium of hypertensive rats. *Proc. Natl. Acad. Sci. USA* 93: 11253-11258, 1996.

Hoyer, J., Distler, A., Haase, W., Gögelein, H.: Ca^{2+} -influx through stretch-activated cation channel activates maxi K^+ channels in porcine endocardial endothelium. *Proc. Natl. Acad. Sci. USA* 91: 2367-2371, 1994.

Hsieh, H.-J., Li, N.-Q., Frangos, J.A.. Shear stress induces endothelial platelet-derived growth factor mRNA levels. *Am. J. Physiol.* 260: H642-H646, 1991.

Huang, P.L., Huang, Z., Mashimo, H., Bloch, K.D., Moskowitz, M.A., Bevan, J.A., Fishman, M.C.. Hypertension in mice lacking the gene for endothelial nitric oxide synthase. *Nature* 377: 239-242, 1995.

Ikeuchi Y., Nishizaki T.. ATP activates the potassium channel and enhances cytosolic Ca^{2+} -release via a P2Y purinoceptor linked to pertussis toxin-insensitive G-protein in brain artery endothelial cells. *Biochem. Biophys. Res. Commun.* 215: 1022-1028, 1995.

Jacobs, E.R., Cheliakine, C., Gebremedhin, D., Birks, E.K., Davies, P.F., Harder, D.R.. Shear activated channels in cell-attached patches of cultured bovine aortic endothelial cells. *Pflügers Arch.* 431: 129-131, 1995.

Johns, A., Lategan, T.W., Lodge, N.J., Ryan, U.S., van Breemen, C., Adams, D.J.. Calcium entry through receptor-operated channels in bovine pulmonary artery endothelial cells. *Tissue Cell* 19: 733-745, 1987.

Köhler R., Distler A., Hoyer J.. Increased mechanosensitive currents in aortic endothelial cells from genetically hypertensive rats. *J. Hypertens.* 17: 365-371, 1999.

Köhler R., Distler A., Hoyer J.. Pressure-activated cation channel in intact rat endocardial endothelium. *Cardiovasc. Res.* 38: 433-440, 1998.

Koller, A., Huang, A.. Impaired flow-induced vasodilation in hypertension. *Circ. Res.* 74:416-421, 1994.

Kuo, L., Davis, M.J., Chilian, W.H.. Endothelium-dependent, flow-induced dilation of isolated coronary arterioles. *Am. J. Physiol.* 259: H1063-H1070, 1990.

Lamontagne D., Pohl U., Busse R.. Mechanical deformation of vessel wall and shear stress determine the basal release of endothelium-derived relaxing factor in the intact rabbit coronary vascular bed. *Circ. Res.* 70: 123-130, 1992.

Lansman, J.B., Hallam, T.J., Rink, T.J.. Single stretch-activated ion channels in vascular endothelial cells as mechanotransducers. *Nature* 325: 811-813, 1987.

Levesque M.J., Nerem RM.. The study of rheological effects on vascular endothelial cells in culture. *Biorheology* 26: 345-357, 1989.

Levesque, M.J., Nerem, R.M.. The elongation and orientation of cultured endothelial cells in response to shear stress. *J. Biomech. Eng.* 107: 341-347, 1985.

Lippoldt A., Gross V., Schneider K., Hansson A., Nadaud S., Schneider W., Bader M., Yagil C., Yagil Y., Luft F.. Nitric oxide synthase and renin-angiotensin system gene expression in salt-sensitive and salt-resistant sabra rats. *Hypertension* 30: 409-415, 1997.

Liu Y., Pleyte K.A., Knaus H.G., Rusch N.J.. Increased expression of Ca²⁺-sensitive K⁺ channels in aorta of hypertensive rats. *Hypertension* 30: 1403-1409, 1997.

Lückhoff A., Clapham D.E.. Inositol 1,3,4,5-tetrakisphosphate activates an endothelial Ca²⁺-permeable channel. *Nature* 355: 356-358, 1992.

Lückhoff, A., Busse, R.. Calcium influx into endothelial cells and formation of endothelium-derived relaxing factor is controlled by the membrane potential. *Pflügers Arch.* 416: 305-311, 1990.

Lückhoff, A., Pohl, U., Mülsch, A., Busse, R.. Differential role of extra- and intracellular calcium in the release of EDRF and prostacyclin from cultured endothelial cells. *Br. J. Pharmacol.* 95: 189-196, 1988.

Lüscher T.F., Aarhus L.L., Vanhoutte P.M.. Indomethacin improves the impaired endothelium-dependent relaxations in small mesenteric arteries of the spontaneously hypertensive rat. *Am. J. Hypertens.* 3: 55-58, 1990.

Lüscher T.F., Vanhoutte P.M., Raji L.. Antihypertensive treatment normalizes decreased endothelium-dependent relaxations in rats with salt-induced hypertension. *Hypertension* 9: 193-197, 1987.

Lüscher T.F.. Heterogeneity of endothelial dysfunction in hypertension. *Eur. Heart J.* 13 (Suppl. D): 50-55, 1992.

Lüscher T.F.. Platelet-vessel wall interaction: role of nitric oxide, prostaglandins and endothelins. *Baillieres Clin. Haematol.* 6: 609-627, 1993.

Lüscher, T.F., Vanhoutte, P.M.. Endothelium-dependent contractions to acetylcholine in the aorta of the spontaneous hypertensive rat. *Hypertension* 8: 344-348, 1986.

Lüscher, T.F., Vanhoutte, P.M.. Endothelium-dependent responses to platelets and serotonin in spontaneously hypertensive rats. *Hypertension* 8 (Suppl. II): II-55 - II-60, 1986.

Lüscher, T.F., Yang, Z., Diederich, D., Bühler, F.R.. Endothelium-derived vasoactive substances: potential role in hypertension, atherosclerosis, and vascular occlusion. *J. Cardiovasc. Pharmacol.* 14: S63-S69, 1989.

Lüscher, T.F.. Endotheliale Relaxation und Kontraktion. *Dtsch. Med. Wochenschr.* 116: 747-754, 1991.

Marchenko S.M., Sage S.O.. A novel mechanosensitive cationic channel from the endothelium of rat aorta. *J. Physiol.* 498: 419-425, 1997.

Mills J.W., Schwiebert E.M., Stanton B.A.. The cytoskeleton and membrane transport. *Curr. Opin. Nephrol. Hypertens.* 3: 529-534, 1994.

Moncada, S., Gryglewski, R., Bunting, S., Vane, J.R.. An enzyme isolated from arteries transforms prostaglandin endoperoxides to an unstable substance that inhibits platelet aggregation. *Nature* 263: 663-665, 1976.

Morris, C.E., Horn, R.. Technical comments. Single-channel mechanosensitive currents. *Science*, 253: 801-802, 1991.

Morris, C.E.. Mechanosensitive ion channels. *J. Membrane Biol.* 113: 93-107, 1990.

Nabel, E.G., Selwyn, A.P., Ganz, P.. Large coronary arteries in humans are responsive to changing blood flow: an endothelium-dependent mechanism that fails in patients with atherosclerosis. *J. Am. Coll. Cardiol.* 16: 349-356, 1990.

Nathan, C.. Nitric oxide as a secretory product of mammalian cells. *FASEB J.* 6: 3051-3064, 1993.

Neher E., Sakmann B.. Single-channel currents recorded from membrane of denervated frog muscle fibres. *Nature* 260: 799-802, 1976.

Ni Z., Oveisi F., Varizi N.D.. Nitric oxide synthase isotype expression in salt-sensitive and salt-resistant Dahl rats. *Hypertension* 34: 552-557, 1999.

Nilius B.. Signaltransduction in vascular endothelium: the role of intracellular calcium and ion channels. *Verh. K. Acad. Geneesk. Belg.* 60: 215-250, 1998.

Nilius, B.. Regulation of transmembrane calcium fluxes in endothelium. *NIPS* 6: 110-114, 1991.

Olesen, S.P., Clapham, D.E., Davies, P.F.. Hemodynamic shear stress activates a K^+ current in vascular endothelial cells. *Nature* 331: 168-170, 1988.

Panza, J.A., Quyyumi, A.A., Brush, J.E., Epstein, S.E.. Abnormal endothelium-dependent vascular relaxation in patients with essential hypertension. *New Engl. J. Med.* 323: 22-27, 1990.

Popp R., Fleming I., Busse R.. Pulsatile stretch in coronary arteries elicits release of endothelium-derived hyperpolarizing factor: a modulator of arterial compliance. *Circ. Res.* 82: 696-703, 1998.

Popp, R., Hoyer, J., Meyer, J., Galla, H.-J., Gögelein, H.. Stretch-activated channel in pig brain blood capillaries. *J. Physiol. (London)* 454, 435-449, 1992.

Prat, A.G., Bertorello, A.M., Ausiello, D.A., Cantiello, H.F.. Activation of epithelial Na^+ channels by protein kinase A requires actin filaments. *Am. J. Physiol.* 265: C224-C233, 1993.

Qiu H.Y., Henrion D., Benessiano J., Heymes C., Tournier B., Levy B.I.. Decreased flow-induced dilation and increased production of cGMP in spontaneously hypertensive rats. *Hypertension* 32: 1098-1103, 1998.

Rees D., Ben-Ishay D., Moncada S.. Nitric oxide and the regulation of blood pressure in the hypertension-prone and hypertension-resistant sabra rat. *Hypertension* 28: 367-371, 1996.

Resnik, N., Collins, T., Atkinson, W., Bonthron, D.T., Dewey, C.F., Gimbrone, M.A.. Platelet-derived growth factor B chain promoter contains a cis-acting fluid shear-stress-responsive element. *Proc. Natl. Acad. Sci. USA* 90: 4591-4595, 1993.

Rinaldi, G., Bohr.. Endothelium-mediated spontaneous response in aortic rings of deoxycorticosterone acetate-hypertensive rats. *Hypertension* 13: 256-261, 1989.

Rubanyi, G.M., Romero, J.C., Vanhoutte, P.M.. Flow-induced release of endothelium-derived relaxing factor. *Am. J. Physiol.* 219: H1145-H1149, 1986.

Rubanyi G.M., Freay A.D., Kauser K., Johns A., Harder DR.. Mechanoreception by the endothelium: mediators and mechanisms of pressure- and flow-induced vascular responses. *Blood Vessels* 27: 246-257, 1990.

Rusko J., Tanzi F., van Breemen C., Adams D.J.. Calcium-activated potassium channels in native endothelial cells from rabbit aorta: conductance, Ca^{2+} -sensitivity and block. *J. Physiol.* 455: 601-621, 1992.

Sackin, H.. Stretch-activated ion channels. *Kidney Int.* 48: 1134-1147, 1995.

Sato M, Levesque MJ, Nerem RM, Ohshima N.. Mechanical properties of cultured endothelial cells exposed to shear stress. *Front. Med. Biol. Eng.* 2: 171-175, 1990.

Sauvé, R., Chahine, M., Tremblay, J., Hamet, H.. Single-channel analysis of the electrical response of bovine aortic endothelial cells to bradykinin stimulation: contribution of a Ca^{2+} -dependent K^{+} -channel. *J. Hypertension* 8: S193-S201, 1990.

Sokabe M., Sachs F.. The structure and dynamics of patch-clamped membranes: a study using differential interference contrast light microscopy. *J. Cell. Biol.* 111: 599-606, 1991.

Van de Voorde, J., Vanheel, B., Leusen, I.. Endothelium-dependent relaxation and hyperpolarisation in aorta from control and renal hypertensive rats. *Circ. Res.* 70: 1-8, 1992.

Vanhoutte P.M.. The endothelium – modulator of vascular smooth-muscle tone. *New Engl. J. Med.* 319: 513-514, 1988.

Vaziri N.D., Oveisi F., Ni Z.. Upregulation of renal and vascular nitric oxide synthase in young spontaneously hypertensive rats. *Hypertension* 31: 1248-1254, 1999.

Yagil C., Katni G., Rubattu S., Stolpe C., Kreutz R., Lindpaintner K., Ganten D., Ben-Ishay D., Yagil Y.. Development, genotype and phenotype of a new colony of the sabra hypertension prone (SBH/y) and resistant (SBN/y) rat model of salt sensitivity and resistance. *J. Hypertens.* 14: 1175-1182, 1996.

Yagil Y., Ben-Ishay D., Wald H., Popovtzer M.M.. Water handling by the sabra hypertension prone (SBH) and resistant (SBN) rats. *Pflügers Arch.* 404: 61-66, 1985.

Yagil Y., Yagil C.. Does the sabra hypertension-prone rat represent a model of salt or mineralocorticoid sensitivity? *J. Hypertens.* 16: 1481-1484, 1998.

Yagil Y., Yagil C.. Genetic basis of salt-susceptibility in the sabra rat model of hypertension. *Kidney Int.* 53: 1493-1500, 1998.