

Literaturverzeichnis

- Abadir R und Hakami N. Ataxia telangiectasia with cancer. An indication for reduced radiotherapy and chemotherapy doses. *Br J Radiol* 1983, 56(665):343-5.
- Abbas AK und Lichtman AH. Basic Immunology - Functions and Disorders of the Immune System (2. Auflage 2004), Saunders, Philadelphia, USA.
- Alberts B, Bray D, Johnson A et al. Lehrbuch der Molekularen Zellbiologie (4. Auflage 2004), 6: Von der DNA zum Protein, S. 345-432, Wiley-VCH, Weinheim, New York, Chichester, Brisbane, Singapore, Toronto.
- Anderson DE, Trujillo KM, Sung P und Erickson HP. Structure of the Rad50 x Mre11 DNA repair complex from *Saccharomyces cerevisiae* by electron microscopy. *J Biol Chem* 2001, 276(40):37027-33.
- Antoccia A, Stumm M, Saar K, Ricordy R, Maraschio P und Tanzarella C. Impaired p53-mediated DNA damage response, cell-cycle disturbance and chromosome aberrations in Nijmegen breakage syndrome lymphoblastoid cell lines. *Int J Radiat Biol* 1999, 75(5):583-91.
- Athma P, Rappaport R, Swift M, Chase CL und Morrell D. Molecular genotyping shows that ataxia-telangiectasia heterozygotes are predisposed to breast cancer. *Cancer Genet Cytogenet* 1996, 92(2):130-4.
- Aurias A, Dutrillaux B, Buriot D und Lejeune L. High frequencies of inversions and translocations of chromosomes 7 and 14 in ataxia telangiectasia. *Mutat Res* 1980, 69(2):369-74.
- Avery OT, MacLeod CM und McCarty M. Studies on the chemical nature of the substance inducing transformation of pneumococcal types. *J Exp Med* 1944, 79:137-58.
- Bailey SM, Cornforth MN, Ullrich RL und Goodwin EH. Dysfunctional mammalian telomeres join with DNA double-strand breaks. *DNA Repair (Amst)* 2004, 3(4):349-57.
- Bailey SM, Meyne J, Chen DJ et al. DNA double-strand break repair proteins are required to cap the ends of mammalian chromosomes. *Proc Natl Acad Sci U S A* 1999, 96(26):14899-904.
- Bakhshi S, Cerosaletti KM, Concannon P et al. Medulloblastoma with adverse reaction to radiation therapy in nijmegen breakage syndrome. *J Pediatr Hematol Oncol* 2003, 25(3):248-51.
- Bakkenist CJ, Drissi R, Wu J, Kastan MB und Dome JS. Disappearance of the telomere dysfunction-induced stress response in fully senescent cells. *Cancer Res* 2004, 64(11):3748-52.
- Bakkenist CJ und Kastan MB. DNA damage activates ATM through intermolecular autophosphorylation and dimer dissociation. *Nature* 2003, 421(6922):499-506.
- Barbi G, Scheres JM, Schindler D et al. Chromosome instability and X-ray hypersensitivity in a

- microcephalic and growth-retarded child. Am J Med Genet 1991, 40(1):44-50.
- Barth E, Demori E, Pecile V, Zanazzo GA, Malorgio C und Tamaro P. Anthracyclines in Nijmegen breakage syndrome. Med Pediatr Oncol 2003, 40(2):122-4.
- Bekiesinska-Figatowska M, Chrzanowska KH, Sikorska J et al. Cranial MRI in the Nijmegen breakage syndrome. Neuroradiology 2000, 42(1):43-7.
- Böcker W, Denk H und Heitz U. Pathologie (3. Auflage 2004), Urban & Fischer, München, Jena.
- Boulton SJ und Jackson SP. Components of the Ku-dependent non-homologous end-joining pathway are involved in telomeric length maintenance and telomeric silencing. EMBO J 1998, 17(6):1819-28.
- Bressan DA, Baxter BK und Petrini JH. The Mre11-Rad50-Xrs2 protein complex facilitates homologous recombination-based double-strand break repair in *Saccharomyces cerevisiae*. Mol Cell Biol 1999, 19(11):7681-7.
- Burger S, Schindler D, Fehn M et al. Radiation-induced DNA damage and repair in peripheral blood mononuclear cells from Nijmegen breakage syndrome patients and carriers assessed by the Comet assay. Environ Mol Mutagen 2006, 47(4):260-70.
- Buscemi G, Savio C, Zannini L et al. Chk2 activation dependence on Nbs1 after DNA damage. Mol Cell Biol 2001, 21(15):5214-22.
- Buslov KG, Iyevleva AG, Chekmariova EV et al. NBS1 657del5 mutation may contribute only to a limited fraction of breast cancer cases in Russia. Int J Cancer 2005, 114(4):585-89.
- Carlomagno F, Chang-Claude J, Dunning AM und Ponder BA. Determination of the frequency of the common 657Del5 Nijmegen breakage syndrome mutation in the German population: no association with risk of breast cancer. Genes Chromosomes Cancer 1999, 25(4):393-5.
- Carney JP, Maser RS, Olivares H et al. The hMre11/hRad50 protein complex and Nijmegen breakage syndrome: linkage of double-strand break repair to the cellular DNA damage response. Cell 1998, 93(3):477-86.
- Carson CT, Schwartz RA, Stracker TH, Lilley CE, Lee DV und MD, W. The Mre11 complex is required for ATM activation and the G2/M checkpoint. EMBO J 2003, 22(24):6610-20.
- Cerosaletti K, Wright J und Concannon P. Active role for nibrin in the kinetics of atm activation. Mol Cell Biol 2005, 26(5):1691-9.
- Cerosaletti KM und Concannon P. Nibrin forkhead-associated domain and breast cancer C-terminal domain are both required for nuclear focus formation and phosphorylation. J Biol Chem 2003, 24(278):21944-51.
- Cerosaletti KM, Lange E, Stringham HM et al. Fine localization of the Nijmegen breakage syn-

- drome gene to 8q21: evidence for a common founder haplotype. Am J Hum Genet 1998, 63(1):125-34.
- Chen HT, Bhandoola A, Difilippantonio MJ et al. Response to RAG-mediated VDJ cleavage by NBS1 and gamma-H2AX. Sciece 2000, 290(5498):1962-5.
- Chien A, Edgar DB und Trela JM. Desoxyribonucleic acid polymerase from the extreme thermophile *Thermus aquaticus*. J Bact 1976, 127(3):1550-57.
- Chrzanowska K und Janninger CK. Nijmegen Breakage syndrome. eMedicine (aufgerufen am 05.07.2006 unter <http://www.emedicine.com/DERM/topic725.htm>).
- Chrzanowska KH, Kleijer WJ, Krajewska-Walasek M et al. Eleven Polish patients with microcephaly, immunodeficiency, and chromosomal instability: the Nijmegen breakage syndrome. Am J Med Genet 1995, 57(3):462-71.
- Chrzanowska KH, Piekutowska-Abramczuk D, Popowska E et al. Carrier frequency of mutation 657del5 in the NBS1 gene in a population of Polish pediatric patients with sporadic lymphoid malignancies. Int J Cancer 2006, 118(5):1269-74.
- Chrzanowska KH, Stumm M, Bekiesiska-Figatowska M et al. Atypical clinical picture of the Nijmegen breakage syndrome associated with developmental abnormalities of the brain. J Med Genet 2001, 38(1):E3.
- Clatworthy AE, Valencia-Burton MA, Haber JE und Oettinger MA. The MRE11-RAD50-XRS2 complex, in addition to other non-homologous end-joining factors, is required for V(D)J joining in yeast. J Biol Chem 2005, 280(21):20247-52.
- Cleary SP, Zhang W, Di Nicola N et al. Heterozygosity for the BLM(Ash) mutation and cancer risk. Cancer Res 2003, 63(8):1769-71.
- Concannon PJ und Gatti RA. Nijmegen Breakage Syndrom. geneReviews, (aufgerufen am 05.07.2006 unter www.geneclinics.org/profiles/nijmegen/details.html).
- Conley ME, Spinner NB, Emanuel BS, Nowell PC und Nichols WW. A chromosomal breakage syndrome with profound immunodeficiency. Blood 1986, 67(5):1251-56.
- Curry CJ, O'Lague P, Tsai J et al. ATfresno: a phenotype linking ataxia-telangiectasia with the Nijmegen breakage syndrome. Am J Hum Genet 1989, 45(2):270-5.
- Cybulski C, Gorski B, Debniak T et al. NBS1 is a prostate cancer susceptibility gene. Cancer Res 2004, 64(4):1215-9.
- D'Amours D und Jackson SP. The Mre11 complex: at the crossroads of dna repair and check-point signalling. Nat Rev Mol Cell Biol 2002, 3(5):317-27.
- De Jager M, van Noort J, van Gent DC, Dekker C, Kanaar R und Wyman C. Human Rad50/Mre11 is a flexible complex that can tether DNA ends. Mol Cell 2001,

- 8(5):1129-35.
- Debniak T, Gorski B, Cybulski C et al. Germline 657del5 mutation in the NBS1 gene in patients with malignant melanoma of the skin. *Melanoma Res* 2003, 13(4):365-70.
- Demuth I, Frappart PO, Hildebrand G et al. An inducible null mutant murine model of Nijmegen breakage syndrome proves the essential function of NBS1 in chromosomal stability and cell viability. *Hum Mol Gen* 2004, 13(20):2385-97.
- Der Kaloustian VM, Kleijer W, Booth A et al. Possible new variant of Nijmegen breakage syndrome. *Am J Med Genet* 1996, 65(1):21-6.
- Desai-Mehta A, Cerosaletti KM und Concannon P. Distinct functional domains of nibrin mediate Mre11 binding, focus formation, and nuclear localization. *Mol Cell Biol* 2001, 21(6):2184-91.
- Digweed M. Human genetic instability syndromes: single gene defects with increased risk of cancer. *Toxicol Lett* 1993, 67(1-3):259-81.
- Digweed M. Genomic Instability in Fanconi Anaemia and Nijmegen Breakage Syndrome. *Genome Dynamics & Stability*, Online-Dokument DOI 10.1007/7050_013 (2006), Springer-Verlag Berlin, Heidelberg
- Digweed M, Reis A und Sperling K. Nijmegen breakage syndrome: consequences of defective DNA double strand break repair. *Bioessays* 1999, 21(8):649-56.
- Digweed M und Sperling K. Nijmegen breakage syndrome: clinical manifestation of defective response to DNA double-strand breaks. *DNA Repair* 2004, 3(8-9):1207-17.
- Distel L, Neubauer S, Varon R, Holter W und Grabenbauer G. Fatal toxicity following radio- and chemotherapy of medulloblastoma in a child with unrecognized Nijmegen breakage syndrome. *Med Pediatr Oncol* 2003, 41(1):44-8.
- Dumon-Jones V, Frappart PO, Tong WM et al. Nbn heterozygosity renders mice susceptible to tumor formation and ionizing radiation-induced tumorigenesis. *Cancer Res* 2003, 63(21):7263-9.
- Elzouki AY, Harfi HA und Nazer H. *Textbook of Clinical Pediatrics* (1. Auflage 2001), 2: Cytogenetics, S. 20-32, Lippincott Williams & Wilkins, Philadelphia, Baltimore, New York, London.
- Erlich HA und Arnheim N. Genetic analysis with the polymerase chain reaction. *Ann Rev Genet* 1992, 26:479-506.
- Filipovich AH, Mathur A, Kamat D und Shapiro RS. Primary immunodeficiencies: genetic risk factors for lymphoma. *Cancer Res* 1994, 52(19 (Suppl)):5465s-5467s.
- Frappart PO, Tong WM, Demuth I et al. An essential function for NBS1 in the prevention of

- ataxia and cerebellar defects. *Nat Med* 2005, 11(5):538-44.
- Gassen HG, Sachse GE und Schulte A. PCR - Grundlagen und Anwendungen der Polymerase-Kettenreaktion. 1994, Gustav Fischer Verlag, Stuttgart, Jena, New York.
- Gatei M, Young D, Cerosaletti KM et al. ATM-dependent phosphorylation of nibrin in response to radiation exposure. *Nat Genet* 2000, 25(1):115-9.
- Goodarzi AA, Block WD und Lees-Miller SP. The role of ATM and ATR in DNA damage-induced cell cycle control. *Prog Cell Cycle Res* 2003, 5:393-411.
- Gorski B, Debniak T, Masojc B et al. Germline 657del5 mutation in the NBS1 gene in breast cancer patients. *Int J Cancer* 2003, 106(3):379-81.
- Green AJ, Yates JR, Taylor AM et al. Severe microcephaly with normal intellectual development: the Nijmegen breakage syndrome. *Arch Dis Child* 1995, 73(5):431-4.
- Gregorek H, Chrzanowska KH, Michalkiewicz J, Syczewska M und Madalinski K. Heterogeneity of humoral immune abnormalities in children with Nijmegen breakage syndrome: an 8-year follow-up study in a single centre. *Clin Exp Immunol* 2002, 130(2):319-24.
- Gruber SB, Ellis NA, Scott KK et al. BLM heterozygosity and the risk of colorectal cancer. *Science* 2002, 297(5589):2013.
- Haber JE. The many interfaces of Mre11. *Cell* 1998, 95(5):583-6.
- Hama S, Matsuura S, Tauchi H et al. Absence of mutations in the NBS1 gene in B-cell malignant lymphoma patients. *Anticancer Res* 2000, 20(3B):1897-900.
- Hart RM, Kimler BF, Evans RG und Park CH. Radiotherapeutic management of medulloblastoma in a pediatric patient with ataxia telangiectasia. *Int J Radiat Oncol Biol Phys* 1987, 13(8):1237-40.
- Heim RA, Lench NJ und Swift M. Heterozygous manifestations in four autosomal recessive human cancer-prone syndromes: ataxia telangiectasia, xeroderma pigmentosum, Fanconi anemia, and Bloom syndrome. *Mutat Res* 1992, 284(1):25-36.
- Hiel JA, Weemaes CM, van den Heuvel LP et al. Nijmegen breakage syndrome. *Arch Dis Child* 2000, 82:400-406.
- Hiel JA, Weemaes CM, van Engelen BG et al. Nijmegen breakage syndrome in a Dutch patient not resulting from a defect in NBS1. *J Med Genet* 2001, 38(6):E19.
- Hopfner KP, Karcher A, Craig L, Woo TT, Carney JP und Tainer JA. Structural biochemistry and interaction architecture of the DNA double-strand break repair Mre11 nuclease and Rad50-ATPase. *Cell* 2001, 105(4):473-85.
- Houldsworth J und Lavin MF. Effect of ionizing radiation on DNA synthesis in ataxia telangiectasia cells. *Nucl Acids Res* 1980, 8(16):3709-20.

- Hsu HL, Gilley D, Galande SA et al. Ku acts in a unique way at the mammalian telomere to prevent end joining. *Genes Dev* 2000, 14(22):2807-12.
- Huang SC, Koch CA, Vortmeyer AO et al. Duplication of the mutant RET allele in trisomy 10 or loss of the wild-type allele in multiple endocrine neoplasia type 2-associated pheochromocytomas. *Cancer Res* 2000, 60(22):6223-6.
- Innis MA, Myambo KB, Gelfand DH und Brow MA. DNA sequencing with *Thermus aquaticus* DNA polymerase and direct sequencing of polymerase chain reaction-amplified DNA. *Proc Natl Acad Sci USA* 1988, 85(24):9436-40.
- Inskip HM, Kinlen LJ, Taylor AM, Woods CG und Arlett CF. Risk of breast cancer and other cancers in heterozygotes for ataxia-telangiectasia. *Br J Cancer* 1999, 79(7-8):1304-7.
- Jahn W, Schumann J und Brockhoff E. Edel und Frei. Franken im Mittelalter. Katalog zur Landesausstellung. 2004, Augsburg.
- Janin N, Andrieu N, Ossian K et al. Breast cancer risk in ataxia telangiectasia (AT) heterozygotes: haplotype study in French AT families. *Br J Cancer* 1999, 80(7):1042-5.
- Jaspers NGJ und Bootsma D. Genetic heterogeneity in ataxia teleangiectasia studied by cell fusion. *Proc Natl Acad Sci USA* 1982a, 79(8):2641-4.
- Jaspers NGJ, De Wit JD und Bootsma D. The rate of DNA synthesis in ataxia teleangiectasia fibroblasts after exposure to DNA damaging agents. In: H.D. Bridges BA (Ed) Ataxia Teleangiectasia. Chichester, Wiley & Son, 1982b, S. 339-45.
- Jaspers NGJ, Gatti RA, Baan C, Linssen PCML und Bootsma D. Genetic complementation analysis of ataxia teleangiectasia and Nijmegen breakage syndrome: a survey of 50 patients. *Cytogenet Cell Genet* 1988b, 49(4):259-63.
- Jaspers NGJ, Painter RB, Paterson MC, Kidson C und Inoue T. Complementation analysis of ataxia teleangiectasia. *KROC Found Ser* 1985, 19:147-62.
- Jaspers NGJ, Taalman RDF und Baan C. Patients with an inherited syndrome characterized by immunodeficiency, microcephaly and chromosomal instability: genetic relationship to ataxia teleangiectasia. *Am J Hum Genet* 1988a, 42(1):66-73.
- Jongmans W, Vuillaume M, Chrzanowska K, Smeets D, Sperling K und Hall J. Nijmegen breakage syndrome cells fail to induce the p53-mediated DNA damage response following exposure to ionizing radiation. *Mol Cell Biol* 1997, 17(9):5016-22.
- Jorde LB, Carey JC, Banshad MJ und White RL. Human Genetics (3. Auflage 2003), 6: Clinical Cytogenetics: The Basis of Human Disease, S. 107-135; 14: Clinical Genetics and Genetic Counseling, S. 305-325, Mosby, St. Louis, Missouri, USA.
- Kang J, Ferguson D, Song H et al. Functional interaction of H2AX, NBS1, and p53 in ATM-

- dependent DNA damage responses and tumor suppression. *Mol Cell Biol* 2004, 25(2):661-70.
- Kleier S, Herrmann M, Wittwer B, Varon R, Reis A und Horst J. Clinical presentation and mutation identification in the NBS1 gene in a boy with Nijmegen breakage syndrome. *Clin Genet* 2000, 57(5):384-7.
- Knippers R. Molekulare Genetik (8. Auflage 2001), 2: DNA: Träger der genetischen Information, S. 9-36; 11: Exons und Introns. Struktur eukaryonter Gene Seiten 315-28; 17: Genomik: Forschungen über die Struktur und Expression komplexer Genome, Georg Thieme Verlag, Stuttgart, New York.
- Knudson AG. Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci USA* 1971, 68(4):820-3.
- Kobayashi J, Antoccia A, Tauchi H, Matsuura S und Komatsu K. NBS1 and its functional role in the DNA damage response. *DNA repair* 2004, 3(8-9):855-61.
- Kobayashi J, Tauchi H, Sakamoto S et al. NBS1 localizes the gamma-H2AX foci through interaction with the FHA/BRCT domain. *Curr Biol* 2002, 12(21):1846-51.
- Komatsu K, Matsuura S, Tauchi H et al. The gene for Nijmegen breakage syndrome (V2) is not located on chromosome 11. *Am J Hum Genet* 1996, 58(4):885-8.
- Kraakman-van der Zwet M, Overkamp WJ, Friedl AA et al. Immortalization and characterization of Nijmegen Breakage syndrome fibroblasts. *Mutat Res* 1999, 434(1):17-27.
- Kracker S, Bergmann Y, Demuth I et al. Nibrin functions in Ig class-switch recombination. *Proc Natl Acad Sci U S A* 2005, 102(5):1584-9.
- Lavin MF und Shiloh Y. The genetic defect in ataxia-teleangiectasia. *Annu Rev Immunol* 1997, 15:177-202.
- Le S, Moore JK, Haber JE und Greider CW. RAD50 and RAD51 define two pathways that collaborate to maintain telomeres in the absence of telomerase. *Genetics* 1999, 152(1):143-52.
- Lee JH, Ghirlando R, Bhaskara V, Hoffmeyer MR, Gu J und Paull TT. Regulation of Mre11/Rad50 by Nbs1: effects on nucleotide-dependent DNA binding and association with ataxia-telangiectasia-like disorder mutant complexes. *J Biol Chem* 2003, 278(14):45171-81.
- Lee JH und Paull TT. Direct activation of the ATM protein kinase by the Mre11/Rad50/Nbs1 complex. *Science* 2004, 304(5667):93-6.
- Lee JH und Paull TT. ATM activation by DNA double-strand breaks through the Mre11-Rad50-Nbs1 complex. *Science* 2005, 308(5721):551-4.
- Lentze MJ, Schaub J, Schulte FJ und Spranger J. Pädiatrie - Grundlagen und Praxis (2. Auflage

- 2003), 11: Immunologie, S. 583-629, Georg Thieme Verlag, Stuttgart, New York.
- Lim DS, Kim ST, Xu B et al. ATM phosphorylates p95/nbs1 in an S-phase checkpoint pathway. *Nature* 2000, 404(6778):613-7.
- Löffler G und Petrides PE. Biochemie & Pathobiochemie (7. Auflage 2003), 5: Nucleotide und Nucleinsäuren S. 141-168; 7: Replikation und Gentechnik S. 209-242, Springer Verlag, Berlin, Heidelberg, New York.
- Maraschio P, Danesino C, Antoccia A et al. A novel mutation and novel features in Nijmegen breakage syndrome. *J Med Genet* 2001, 38(2):113-7.
- Marcelain K, De La Torre C, Gonzalez P und Pincheira J. Roles of nibrin and AtM/ATR kinases on the G2 checkpoint under endogenous or radio-induced DNA damage. *Biol Res* 2005, 38(2-3):179-85.
- Maser RS, Zinkel R und Petrini JH. An alternative mode of translation permits production of a variant NBS1 protein from the common Nijmegen Breakage Syndrome allele. *Nat Genet* 2001, 27(4):417-22.
- Matsuura S, Kobayashi J, Tauchi H und Komatsu K. Nijmegen breakage syndrome and DNA double strand break repair by NBS1 complex. *Adv Biophys* 2004, 38:65-80.
- Matsuura S, Tauchi H, Nakamura A et al. Positional cloning of the gene for Nijmegen breakage syndrome. *Nat Genet* 1998, 19(2):171-81.
- Matsuura S, Weemaes C, Smeets D et al. Genetic mapping using microcell-mediated chromosome transfer suggests a locus for Nijmegen breakage syndrome at chromosome 8q21-24. *Am J Hum Genet* 1997, 60(6):1487-94.
- McConville CM, Byrd PJ, Ambrose HJ und Taylor AM. Genetic and physical mapping of the ataxia-telangiectasia locus on chromosome 11q22-q23. *Int J Radiat Biol* 1994, 66(6 Suppl):S45-56.
- Meyer S, Kingston H, Taylor AM et al. Rhabdomyosarcoma in Nijmegen breakage syndrome: strong association with perianal primary site. *Cancer Genet Cytogenet* 2004, 154(2):169-74.
- Meyn MS. Chromosome instability syndromes: lessons for carcinogenesis. *Curr Top Microbiol Immunol* 1997, 221:71-148.
- Michallet AS, Lesca G, Radford-Weiss I, Delarue R, Varet B und Buzyn A. T-cell prolymphocytic leukemia with autoimmune manifestations in Nijmegen breakage syndrome. *Ann Hemat* 2003, 82(8):515-517.
- Morrell D, Chase CL und Swift M. Cancers in 44 families with ataxia-telangiectasia. *Cancer Genet Cytogenet* 1990, 50(1):119-23.

- Morrell D, Cromartie E und Swift M. Mortality and cancer incidence in 263 patients with ataxiatelangiectasia. *J Natl Cancer Inst* 1986, 77(1):89-92.
- Neubauer S, Arutyunyan R, Stumm M et al. Radiosensitivity of ataxia telangiectasia and Nijmegen breakage syndrome homozygotes and heterozygotes as determined by three-color FISH chromosome painting. *Radiat Res* 2002, 157(3):312-21.
- New HV, Cale CM, Tischkowitz M et al. Nijmegen breakage syndrome diagnosed as Fanconi anaemia. *Pediatr Blood Cancer* 2005, 44(5):494-9.
- Nicholl DST. Gentechnische Methoden (2. Auflage 2002), 7: Die Polymerase-Kettenreaktion, S. 119-136, Spektrum Akademischer Verlag, Heidelberg, Berlin.
- Nove J, Little JB, Mayer PJ, Troilo P und Nichols WW. Hypersensitivity of cells from a new chromosomal-breakage syndrome to DNA-damaging agents. *Mutat Res* 1986, 163(3):255-62.
- Orita M, Suzuki Y, Sekiya T und Hayashi K. Rapid and sensitive detection of point mutations and DNA polymorphisms using the polymerase chain reaction. *Genomics* 1989, 5(4):874-9.
- Pan Q, Petit-Frere C, Lahdesmaki A, Gregorek H, Chrzanowska KH und Hammarstrom L. Alternative end joining during switch recombination in patients with ataxia-telangiectasia. *Euro J Immunol* 2002, 32(5):1300-8.
- Passarge E. Taschenatlas der Genetik (2. Auflage 2004), Georg Thieme Verlag, Stuttgart, New York.
- Paull TT und Gellert M. The 3' to 5' exonuclease activity of Mre11 facilitates repair of DNA double-strand breaks. *Mol Cell* 1998, 1(7):969-79.
- Paull TT und Gellert M. Nbs1 potentiates ATP-driven DNA unwinding and endonuclease cleavage by the Mre11/Rad50 complex. *Genes Dev* 1999, 13(10):1276-88.
- Paull TT und Lee JH. The Mre11/Rad50/Nbs1 complex and its role as a DNA double-strand break sensor for ATM. *Cell Cycle* 2005, 4(6):737-40.
- Paull TT, Rogakou EP, Yamazaki V, Kirchgessner CU, Gellert M und Bonner WM. A critical role for histone H2AX in recruitment of repair factors to nuclear foci after DNA damage. *Curr Biol* 2000, 10(15):886-95.
- Petersen S, Casellas R, Reina-San-Martin B et al. AID is required to initiate Nbs1/gamma-H2AX focus formation and mutations at sites of class switching. *Nature* 2001, 414(6864):660-5.
- Petrini JH und Stracker TH. The cellular response to DNA double-strand breaks: defining the sensors and mediators. *Trends Cell Biol* 2003, 13(9):458-62.
- Pippard EC, Hall AJ, Barker DJ und Bridges BA. Cancer in homozygotes and heterozygotes of

- ataxia-telangiectasia and xeroderma pigmentosum in Britain. *Cancer Res* 1988, 48(10):2929-32.
- Plattner H und Hentschel J. *Zellbiologie* (2. Auflage 2002), 22: Zellzyklus, Kernteilung und Zellteilung, S. 365-386, Georg Thieme Verlag, Stuttgart, New York.
- Plisiecka-Halasa J, Dansonka-Mieszkowska A, Rembiszewska A, Bidzinski M, Steffen J und Kupryjanczyk J. Nijmegen breakage syndrome gene (NBS1) alterations and its protein (nibrin) expression in human ovarian tumours. *Ann Hum Genet* 2002, 66(Pt 5-6):353-9.
- Ranganathan V, Heine WF, Ciccone DN et al. Rescue of a telomere length defect of Nijmegen breakage syndrome cells requires NBS and telomerase catalytic subunit. *Cell Biol* 2001, 11(12):962-6.
- Reis A. Genetics and B-cell leukaemia. *Lancet* 1999, 353(9146):3.
- Resnick IB, Kondratenko I, Togoev O et al. Nijmegen breakage syndrome: clinical characteristics and mutation analysis in eight unrelated Russian families. *J Pediatr* 2002, 140(3):355-61.
- Rischewski J, Bismarck P, Kabisch H, Janka-Schaub G, Obser T und Schneppenheim R. The common deletion 657del5 in the Nibrin gene is not a major risk factor for B or T cell non-Hodgkin lymphoma in a pediatric population. *Leukemia* 2000, 14(8):1528-9.
- Rogakou EP, Pilch DR, Orr AH, Ivanova VS und Bonner WM. DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *J Biol Chem* 1998, 273(10):5858-68.
- Saar K, Chrzanowska KH, Stumm M et al. The Gene for the Ataxia-Teleangiectasia Variant, Nijmegen Breakage Syndrome, Maps to a 1-cM Interval on Chromosome 8q21. *Am J Hum Genet* 1997, 60(3):605-610.
- Saiki RK, Gelfand DH, Stoffel S et al. Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 1988, 239:487-91.
- Saiki RK, Scharf S, Faloona F et al. Enzymatic amplification of beta-globulin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* 1985, 230(4732):1350-4.
- Sanger F und Coulson AR. A rapid method for determining sequences in DNA by primed synthesis with DNA polymerases. *J Mol Biol* 1975, 94:441-8.
- Savov A, Angelicheva D, Jordanova A, Eigel A und Kalaydjieva L. High percentage acrylamide gels improve resolution in SSCP analysis. *Nucleic Acids Res* 1992, 20(24):6741-2.
- Sedgwick RP und Boder E. Ataxia-teleangiectasia. In: Vinken PJ, Klawans HL, (Ed) *Handbook of Clinical Neurology*. Elsevier, Amsterdam, 1991, S. 347-423.
- Seeman P, Gebertová K, Paderová K, Sperling K und Seemanová E. Nijmegen Breakage Syn-

- drome in 13 % of age-matched Czech children with primary microcephaly. *Pediatr Neurol* 2004, 30(3):195-200.
- Seemanova E. An increased risk for malignant neoplasms in heterozygotes for a syndrome of microcephaly, normal intelligence, growth retardation, remarkable facies, immunodeficiency and chromosomal instability. *Mutat Res* 1990, 238(3):321-4.
- Seemanova E, Jarolim P, Seeman P, Varon R und Sperling K. Increased risk of malignancies in heterozygotes in families of patients with Nijmegen breakage syndrome. *Cas Lek Cesk* 2006, 145(2):138-43.
- Seemanova E, Passarge E, Beneskova D, Houstek J, Kasal P und Sevcikova M. Familial microcephaly with normal intelligence, immunodeficiency, and risk for lymphoreticular malignancies: a new autosomal recessive disorder. *Am J Med Genet* 1985, 20(4):639-48.
- Seemanova E, Pohanka V, Seeman P et al. Nijmegen breakage syndrome in Slovakia. *Cas Lek Cesk* 2004, 143(8):538-41.
- Seemanova E, Sperling K, Neitzel H et al. Nijmegen Breakage Syndrome (NBS) with neurological abnormalities and without chromosomal instability. *J Med Genet* 2005, 43(3):218-24.
- Seidemann K, Henze G, Beck JD et al. Non-Hodgkin's lymphoma in pediatric patients with chromosomal breakage syndromes (AT and NBS): experience from the BFM trials. *Ann Oncol* 2000, 11(Suppl 1):141-5.
- Shiloh Y. Ataxia-telangiectasia and the Nijmegen breakage syndrome: related disorders but genes apart. *Annu Rev Genet* 1997, 31:635-62.
- Sobel E, Lange E, Jaspers NG et al. Ataxia-telangiectasia: linkage evidence for genetic heterogeneity. *Am J Hum Genet* 1992, 50(6):1343-8.
- Soucek P, Gut I, Trneny M et al. Multiplex single-tube screening for mutations in the Nijmegen Breakage Syndrome (NBS1) gene in Hodgkin's and non-Hodgkin's lymphoma patients of Slavic origin. *Eur J Hum Genet* 2003, 11(5):416-9.
- Stankovic T, Weber P, Stewart G et al. Inactivation of ataxia telangiectasia mutated gene in B-cell chronic lymphocytic leukaemia. *Lancet* 1999, 353(9146):26-9.
- Stanulla M, Stumm M, Dieckvoss BO et al. No evidence for a major role of heterozygous deletion 657del5 within the NBS1 gene in the pathogenesis of non-Hodgkin's lymphoma of childhood and adolescence. *Br J Haematol* 2000, 109(1):117-20.
- Steffen J, Varon R, Mosor M et al. Increased cancer risk of heterozygotes with NBS1 germline mutations in Poland. *Int J Cancer* 2004, 111:67-71.
- Stiff T, Reis C, Alderton GK, Woodbine L, O'Driscoll M und Jeggo PA. Nbs1 is required for ATR-dependent phosphorylation events. *EMBO J* 2005, 24(1):199-208.

- Stockklausner C, Behnisch W, Mechtersheimer G, Moller P und Kulozik AE. Long-term remission of children with relapsed and secondary anaplastic large cell non-Hodgkin lymphoma (ALCL) following treatment with pulsed dexamethasone and low dose etoposide. *Pediatr Blood Cancer* 2006, 11(im Druck).
- Stoppa-Lyonnet D, Girault D, LeDeist F und Aurias A. Unusual T cell clones in a patient with Nijmegen breakage syndrome. *J Med Genet* 1992, 29(2):136-7.
- Stumm M, Gatti RA, Reis A et al. The ataxia-telangiectasia-variant genes 1 and 2 are distinct from the ataxia-telangiectasia gene on chromosome 11q23.1. *Am J Hum Genet* 1995, 57(4):960-2.
- Stumm M, Sperling K und Wegner RD. Noncomplementation of radiation-induced chromosome aberrations in ataxia-telangiectasia/ataxia-telangiectasia-variant heterodikaryons. *Am J Hum Genet* 1997, 60(5):1246-51.
- Stumm M, von Ruskowsky A, Siebert R et al. No evidence for deletions of the NBS1 gene in lymphomas. *Cancer Genet Cytogenet* 2001, 126(1):60-2.
- Sullivan KE, Veksler E, Lederman H und Lees-Miller SP. Cell cycle checkpoints and DNA repair in Nijmegen breakage syndrome. *Clin Immunol Immunopathol* 1997, 82(1):43-8.
- Swift M. Fanconi's anemia in the genetics of neoplasia. *Nature* 1971, 230:370-3.
- Swift M, Caldwell RJ und Chase C. Reassessment of cancer predisposition of Fanconi anemia heterozygotes. *J Natl Cancer Inst* 1980, 65(5):863-7.
- Swift M, Chase CL und Morrell D. Cancer predisposition of ataxia-telangiectasia heterozygotes. *Cancer Genet Cytogenet* 1990, 46(1):21-7.
- Swift M, Morrell D, Massey RB und Chase CK. Incidence of cancer in 161 families affected by ataxia teleangiectasia. *N Engl J Med* 1991, 325(26):1831-6.
- Swift M, Reitnauer PJ, Morrell D und Chase CL. Breast and other cancers in families with ataxia-telangiectasia. *N Engl J Med* 1987, 316(21):1289-94.
- Taalman RD, Jaspers NG, Scheres JM, de Wit J und Hustinx TW. Hypersensitivity to ionizing radiation, in vitro, in a new chromosomal breakage disorder, the Nijmegen Breakage Syndrome. *Mutat Res* 1983, 112(1):23-32.
- Taalman RDFM, Hustinx TWJ, Weemaes CMR et al. Further delination of the Nijmegen breakage syndrome. *Am J Med Genet* 1989, 32:425-31.
- Tanzanella C, Antoccia A, Spadoni E et al. Chromosome instability and nibrin protein variants in NBS heterozygotes. *Eur J Hum Genet* 2003, 11(4):297-303.
- Tariverdian G und Buselmaier W. Humangenetik (3. Auflage 2004), 3: Chromosomen des Menschen, S. 89-118; 9: Genetische Diagnostik und Beratung, S. 303-352, Springer-Verlag,

- Berlin, Heidelberg, New York.
- Tauchi H, Kobayashi J, Morishima K et al. The forkhead-associated domain of NBS1 is essential for nuclear foci formation after irradiation but not essential for hRAD50/hMRE11/NBS1 complex DNA repair activity. *J Biol Chem* 2001, 276(1):12-5.
- Tauchi H, Kobayashi J, Morishima K et al. Nbs1 is essential for DNA repair by homologous recombination in higher vertebrate cells. *Nature* 2002, 420(6911):93-98.
- Tauchi H, Matsuura S, Kobayashi J, Sakamoto S und Komatsu K. Nijmegen breakage syndrome gene, NBS1, and molecular links to factors for genome stability. *Oncogene* 2002, 21(58):8967-80.
- Taylor AMR, Groom A und Byrd PJ. Ataxia-teleangiectasia-like disorder (ATLD) - its clinical presentation and molecular basis. *DNA Repair* 2004, 3(8-9):1219-25.
- Taylor AMR, Metcalfe JA, Thick J und Mak YF. Leukemia and lymphoma in ataxia teleangiectasia. *Blood* 1996, 87(2):423-38.
- Taylor GM, O'Brien HP, Greaves MF, Ravetto PF und Eden OB. Correspondence re: R. Varon et al., Mutations in the Nijmegen breakage syndrome gene (NBS1) in childhood acute lymphoblastic leukemia. *Cancer Res.*, 61: 3570-3572, 2001. *Cancer Res* 2001, 63(19):6563-4.
- Teebi A, Al-Awadi SA und White AG. Autosomal recessive nonsyndromal microcephaly with normal intelligence. *Am J Med Genet* 1987, 26:355-59.
- Tekin M, Dogu F, Tacyildiz N et al. 657del5 mutation in the NBS1 gene is associated with Nijmegen breakage syndrome in a Turkish family. *Clin Genet* 2002, 62(1):84-8.
- The International Nijmegen Breakage Syndrome Study Group: Hiel JA, Weemaes CM, van den Heuvel LP et al. Nijmegen breakage syndrome. *Arch Dis Child* 2000, 82:400-406.
- Thompson D, Duedal S, Kirner J et al. Cancer risks and mortality in heterozygous ATM mutation carriers. *J Natl Cancer Inst* 2005, 97(11):813-22.
- Tindall KR und Kunkel TA. Fidelity of DNA synthesis by the *Thermus aquaticus* DNA polymerase. *Biochemistry* 1988, 27(16):6008-13.
- Tischkowitz M und Dokal I. Fanconi anemia and leukaemia - clinical and molecular aspects. *Br J Haematol* 2004, 126(2):176-91.
- Tupler R, Marseglia GL, Stefanini M et al. A variant of the Nijmegen breakage syndrome with unusual cytogenetic features and intermediate cellular radiosensitivity. *J Med Genet* 1997, 34(3):196-202.
- Ueno M, Nakazaki T, Akamatsu Y et al. Molecular characterization of the *Schizosaccharomyces pombe* nbs1+ gene involved in DNA repair and telomere maintenance. *Mol Cell Biol*

- 2003, 23(18):6553-63.
- Uhrhammer N, Bay JO, Gosse-Brun S et al. Allelic imbalance at NBS1 is frequent in both proximal and distal colorectal carcinoma. *Oncol Rep* 2000, 7(2):427-31.
- Uziel T, Lerenthal Y, Moyal L, Andegeko Y, Mittelman L und Shiloh Y. Requirement of the MRN complex for ATM activation by DNA damage. *EMBO J* 2003, 22(20):5612-21.
- Van de Kaa CA, Weemaes CM, Wesseling P, Schaafsma HE, Haraldsson A und De Weger RA. Postmortem findings in the Nijmegen breakage syndrome. *Pediatr Pathol* 1994, 14(5):787-96.
- Van der Burgt I, Chrzanowska KH, Smeets D und Weemaes C. Nijmegen breakage syndrome. *J Med Genet* 1996, 33(2):153-6.
- Van Engelen BG, Hiel JA, Gabreels FJ, van den Heuvel LP, van Gent DC und Weemaes CM. Decreased immunoglobulin class switching in Nijmegen Breakage syndrome due to the DNA repair defect. *Hum Immunol* 2001, 62(12):1324-7.
- Varon R, Gosse-Brun S, Bignon YJ, Sperling K und Uhrhammer N. Nijmegen breakage syndrome gene (NBS1) is not the tumor suppressor gene at 8q21.3 involved in colorectal carcinoma. *Oncol Rep* 2002, 9(4):709-11.
- Varon R, Reis A, Henze G, Graf von Einsiedel H, Sperling K und Seeger K. Mutations in the Nijmegen Breakage Syndrome Gene (NBS1) in Childhood Acute Lymphoblastic Leukemia (ALL). *Cancer Res* 2001, 61(9):3570-2.
- Varon R, Seemanova E, Chrzanowska K et al. Clinical ascertainment of Nijmegen breakage syndrome (NBS) and prevalence of the major mutation, 657del5, in three Slav populations. *Eur J Hum Genet* 2000, 8(11):900-2.
- Varon R, Vissinga C, Platzer M et al. Nibrin, a novel DNA double-strand break repair protein, is mutated in Nijmegen breakage syndrome. *Cell* 1998, 93(3):467-76.
- Volkenandt M, Löhr M und Dicker AP. Gen-Amplifikation durch Polymerase-Kettenreaktion. *Dtsch Med Wschr* 1990, 115(17):670-6.
- Vorechovsky I, Luo L, Dyer MJ et al. Clustering of missense mutations in the ataxiatelangiectasia gene in a sporadic T-cell leukaemia. *Nat Genet* 1997, 17(1):96-9.
- Watson JD und Crick FHC. Molecular structure of nucleic acid. *Nature* 1953, 171:737-738.
- Watson JD, Gilman M, Witkowski J und Zoller M. Rekombinate DNA (2. Auflage 1993), Spektrum Akademischer Verlag, Heidelberg, Berlin, Oxford.
- Weemaes CM, Smeets DF und van der Burgt CJ. Nijmegen Breakage syndrome: a progress report. *Int J Radiat Biol* 1994, 66(6 Suppl):S185-88.
- Weemaes CMR, Hustinx TWJ, Scheres JMJC, van Munster PJJ, Bakkeren JAJM und Taalman

- RDFM. A new chromosomal instability disorder: the Nijmegen Breakage Syndrome. *Acta Paediatr Scand* 1981, 70:557-64.
- Weemaes CMR, van de Kaa C, Wesseling P et al. Nijmegen Breakage Syndrome: clinical, immunological and pathological findings. In: L.R. Chapel HE, Webster ADB (Ed) *Progress in Immune Deficiency*. London Royal Society of Medicine Services, 1991, S. 191-93.
- Wegner RD, Chrzanowska KH, Sperling K und Stumm M. Ataxia-Teleangiectasia variants (Nijmegen Breakage Syndrome). In: S.C. Ochs HD, Puck JM (Ed) *Primary Immunodeficiency Diseases, a Molecular and Genetic Approach*. Oxford University Press, Oxford, UK, 1999, S. 324-34.
- Wegner RD, Metzger M, Hanefeld F et al. A new chromosomal instability disorder confirmed by complementation studies. *Clin Genet* 1988, 33(1):20-32.
- White TJ, Arnheim N und Erlich HA. The polymerase chain reaction. *Trends Genet* 1989, 5(6):185-9.
- Wilkins MFH, Stokes AR und Wilson HR. Molecular structure of DNA. *Nature* 1953, 171:738-40.
- Woods CG, Leversha M und Rogers JG. Severe intrauterine growth retardation with increased mitomycin C sensitivity: a further chromosome breakage syndrome. *J Med Genet* 1995, 32(4):301-5.
- Wu X, Ranganathan V, Weisman DS et al. ATM phosphorylation of Nijmegen breakage syndrome protein is required in a DNA damage response. *Nature* 2000, 405(6785):477-82.
- Xu B, Kim ST, Lim DS und Kastan MB. Two molecularly distinct G(2)/M checkpoints are induced by ionizing irradiation. *Mol Cell Biol* 2002, 22(4):1049-59.
- Yamazaki V, Wegner RD und Kirchgessner CU. Characterization of cell cycle checkpoint responses after ionizing radiation in Nijmegen breakage syndrome cells. *Cancer Res* 1998, 58(11):2316-22.
- Ying KL und Decoteau WE. Cytogenetic anomalies in a patient with ataxia, immune deficiency, and high alpha-fetoprotein in the absence of telangiectasia. *Cancer Genet Cytogenet* 1981, 4(4):311-7.
- You Z, Chahwan C, Bailis J, Hunter T und Russell P. ATM activation and its recruitment to damaged DNA require binding to the C terminus of Nbs1. *Mol Cell Biol* 2005, 25(13):5363-79.
- Zhang Y, Lim CU, Williams ES et al. NBS1 knockdown by small interfering RNA increases ionizing radiation mutagenesis and telomere association in human cells. *Cancer Res* 2005, 65(13):5544-53.

- Zhang Y, Zhou J und Lim CU. The role of NBS1 in DNA double strand break repair, telomere stability, and cell cycle checkpoint control. *Cell Res* 2006, 16(1):45-54.
- Zhao S, Weng YC, Yuan SS et al. Functional link between ataxia-telangiectasia and Nijmegen breakage syndrome gene products. *Nature* 2000, 405(6785):473-7.
- Zhou J, Lim CU, Li JJ, Cai L und Zhang Y. The role of NBS1 in the modulation of PIKK family proteins ATM and ATR in the cellular response to DNA damage. *Cancer Lett* 2006, (im Druck).
- Zhu J, Petersen S, Tessarollo L und Nussenzweig A. Targeted disruption of the Nijmegen breakage syndrome gene NBS1 leads to early embryonic lethality in mice. *Curr Biol* 2001, 11(2):105-9.