

Literatur

- ¹ Aasland, R., Abrams, C., Ampe, C., Ball, L.J., Bedford, M.T., Cesareni, G., et al. (2002). Normalization of nomenclature for peptide motifs as ligands of modular protein domains. *FEBS Lett* **513**: 141-4.
- ² Adams, M.E., Mueller, H.A. & Froehner, S.C. (2001). In vivo requirement of the alpha-syntrophin PDZ domain for the sarcolemmal localization of nNOS and aquaporin-4. *J Cell Biol* **155**: 113-22.
- ³ Adzhubei, A.A. & Sternberg, M.J. (1993). Left-handed polyproline II helices commonly occur in globular proteins. *J Mol Biol* **229**: 472-93.
- ⁴ Akamatsu, M. (2002). Current state and perspectives of 3D-QSAR. *Curr Top Med Chem* **2**: 1381-94.
- ⁵ Aloy, P., Bottcher, B., Ceulemans, H., Leutwein, C., Mellwig, C., Fischer, S., et al. (2004). Structure-based assembly of protein complexes in yeast. *Science* **303**: 2026-9.
- ⁶ Aloy, P. & Russell, R.B. (2002). Interrogating protein interaction networks through structural biology. *Proc Natl Acad Sci U S A* **99**: 5896-901.
- ⁷ Aloy, P. & Russell, R.B. (2002). The third dimension for protein interactions and complexes. *Trends Biochem Sci* **27**: 633-8.
- ⁸ Amber. (2000). Assisted model building with energy refinement. Version 7, Department of Pharmaceutical Chemistry, University of California, San Francisco.
- ⁹ Anfinsen, C.B. (1973). Principles that govern the folding of protein chains. *Science* **181**: 223-30.
- ¹⁰ Apweiler, R., Bairoch, A., Wu, C.H., Barker, W.C., Boeckmann, B., Ferro, S., et al. (2004). UniProt: the Universal Protein knowledgebase. *Nucleic Acids Res* **32**: D115-9.
- ¹¹ Aqeilan, R.I., Pekarsky, Y., Herrero, J.J., Palamarchuk, A., Letofsky, J., Druck, T., et al. (2004). Functional association between Wwox tumor suppressor protein and p73, a p53 homolog. *Proc Natl Acad Sci U S A* **101**: 4401-6.
- ¹² Bader, G.D. & Hogue, C.W. (2002). Analyzing yeast protein-protein interaction data obtained from different sources. *Nat Biotechnol* **20**: 991-7.
- ¹³ Ball, L.J., Jarchau, T., Oschkinat, H. & Walter, U. (2002). EVH1 domains: structure, function and interactions. *FEBS Lett* **513**: 45-52.
- ¹⁴ Ball, L.J., Kühne, R., Schneider-Mergener, J. & Oschkinat, H. (2004). Recognition of proline rich motifs (PRMs) by protein-protein interaction domains. *Angew. Chem. Int. Ed. Engl.* **44**: 2852-69.
- ¹⁵ Baroni, M., Costantino, G., Cruciani, G., Riganelli, D., Valigi, R. & Clementi, S. (1993). Generating optimal linear PLS estimations (GOLPE): an advanced chemometric tool for handling 3D-QSAR problems. *Quant Struct-Act Relat* **12**: 9-20.
- ¹⁶ Bateman, A., Birney, E., Durbin, R., Eddy, S.R., Howe, K.L. & Sonnhammer, E.L. (2000). The Pfam protein families database. *Nucleic Acids Res* **28**: 263-6.
- ¹⁷ Bates, D.M. & Pinheiro, J.C. (1998). *Computational methods for multilevel models*, (<http://franz.stat.wisc.edu/pub/NLME>).

- ¹⁸ Bedford, M.T., Reed, R. & Leder, P. (1998). WW domain-mediated interactions reveal a spliceosome-associated protein that binds a third class of proline-rich motif: the proline-glycine and methionine-rich motif. *Proc Natl Acad Sci U S A* **95**: 10602-7.
- ¹⁹ Bedford, M.T., Sarbassova, D., Xu, J., Leder, P. & Yaffe, M.B. (2000). A novel pro-Arg motif recognized by WW domains. *J Biol Chem* **275**: 10359-69.
- ²⁰ Berman, H.M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T.N., Weissig, H., et al. (2000). The Protein Data Bank. *Nucleic Acids Res* **28**: 235-42.
- ²¹ Bezprozvanny, I. & Maximov, A. (2001). Classification of PDZ domains. *FEBS Lett* **509**: 457-62.
- ²² Bezprozvanny, I. & Maximov, A. (2001). PDZ domains: More than just a glue. *Proc Natl Acad Sci U S A* **98**: 787-9.
- ²³ Birrane, G., Chung, J. & Ladis, J.A. (2003). Novel mode of ligand recognition by the Erbin PDZ domain. *J Biol Chem* **278**: 1399-402.
- ²⁴ Bitner-Glindzicz, M., Lindley, K.J., Rutland, P., Blaydon, D., Smith, V.V., Milla, P.J., et al. (2000). A recessive contiguous gene deletion causing infantile hyperinsulinism, enteropathy and deafness identifies the Usher type 1C gene. *Nat Genet* **26**: 56-60.
- ²⁵ Bock, J.R. & Gough, D.A. (2001). Predicting protein-protein interactions from primary structure. *Bioinformatics* **17**: 455-60.
- ²⁶ Boerkoel, C.F., Takashima, H., Stankiewicz, P., Garcia, C.A., Leber, S.M., Rhee-Morris, L., et al. (2001). Periaxin mutations cause recessive Dejerine-Sottas neuropathy. *Am J Hum Genet* **68**: 325-33.
- ²⁷ Boisguerin, P. (2004). Characterization of PDZ Domain/Ligand Specificity, Dissertation an der Freie Universität Berlin.
- ²⁸ Boisguerin, P., Leben, R., Ay, B., Radziwill, G., Moelling, K., Dong, L., et al. (2004). An improved method for the synthesis of cellulose membrane-bound peptides with free C termini is useful for PDZ domain binding studies. *Chem Biol* **11**: 449-59.
- ²⁹ Borg, J.P., Marchetto, S., Le Bivic, A., Ollendorff, V., Jaulin-Bastard, F., Saito, H., et al. (2000). ERBIN: a basolateral PDZ protein that interacts with the mammalian ERBB2/HER2 receptor. *Nat Cell Biol* **2**: 407-14.
- ³⁰ Bork, P. & Sudol, M. (1994). The WW domain: a signalling site in dystrophin? *Trends Biochem Sci* **19**: 531-3.
- ³¹ Brannetti, B., Via, A., Cestra, G., Cesareni, G. & Helmer-Citterich, M. (2000). SH3-SPOT: an algorithm to predict preferred ligands to different members of the SH3 gene family. *J Mol Biol* **298**: 313-28.
- ³² Brenman, J.E., Chao, D.S., Gee, S.H., McGee, A.W., Craven, S.E., Santillano, D.R., et al. (1996). Interaction of nitric oxide synthase with the postsynaptic density protein PSD-95 and alpha1-syntrophin mediated by PDZ domains. *Cell* **84**: 757-67.
- ³³ Bryant, P.J. & Huwe, A. (2000). LAP proteins: what's up with epithelia? *Nat Cell Biol* **2**: E141-3.
- ³⁴ Byrd, R.H., Lu, P., Nocedal, J. & Zhu, C. (1995). A limited memory algorithm for bound constrained optimization. *SIAM J Scientific Comp* **16**: 1190-208.
- ³⁵ Chang, A., Cheang, S., Espanol, X. & Sudol, M. (2000). Rsp5 WW domains interact directly with the carboxyl-terminal domain of RNA polymerase II. *J Biol Chem* **275**: 20562-71.
- ³⁶ Chen, H.I., Einbond, A., Kwak, S.J., Linn, H., Koepf, E., Peterson, S., et al. (1997). Characterization of the WW domain of human yes-associated protein and its polyproline-containing ligands. *J Biol Chem* **272**: 17070-7.
- ³⁷ Chen, H.I. & Sudol, M. (1995). The WW domain of Yes-associated protein binds a proline-rich ligand that differs from the consensus established for Src homology 3-binding modules. *Proc Natl Acad Sci U S A* **92**: 7819-23.
- ³⁸ Cho, K.O., Hunt, C.A. & Kennedy, M.B. (1992). The rat brain postsynaptic density fraction contains a homolog of the Drosophila discs-large tumor suppressor protein. *Neuron* **9**: 929-42.

- ³⁹ Connolly, M.L. (1983). Solvent-accessible surfaces of proteins and nucleic acids. *Science* **221**: 709-13.
- ⁴⁰ Cowan, P.M. & McGavin, S. (1955). Structure of Poly-L-Proline. *Nature* **176**: 501-3.
- ⁴¹ Cramer, P., Bushnell, D.A. & Kornberg, R.D. (2001). Structural basis of transcription: RNA polymerase II at 2.8 angstrom resolution. *Science* **292**: 1863-76.
- ⁴² Cramer, R.D., Patterson, D.E. & Bunce, J.D. (1988). Comparative Molecular-Field Analysis(CoMFA) .1. Effect of Shape on Binding of Steroids to Carrier Proteins. *J. Am. Chem. Soc.* **110**: 5959-67.
- ⁴³ Crooks, G.E., Hon, G., Chandonia, J.M. & Brenner, S.E. (2004). WebLogo: a sequence logo generator. *Genome Res* **14**: 1188-90.
- ⁴⁴ Dayhoff, M.O. & Orcutt, B.C. (1979). Methods for identifying proteins by using partial sequences. *Proc Natl Acad Sci U S A* **76**: 2170-4.
- ⁴⁵ Doyle, D.A., Lee, A., Lewis, J., Kim, E., Sheng, M. & MacKinnon, R. (1996). Crystal structures of a complexed and peptide-free membrane protein-binding domain: molecular basis of peptide recognition by PDZ. *Cell* **85**: 1067-76.
- ⁴⁶ Ebnet, K., Schulz, C.U., Meyer Zu Brickwedde, M.K., Pendl, G.G. & Vestweber, D. (2000). Junctional adhesion molecule interacts with the PDZ domain-containing proteins AF-6 and ZO-1. *J Biol Chem* **275**: 27979-88.
- ⁴⁷ Eddy, S.R. (1998). Profile hidden Markov models. *Bioinformatics* **14**: 755-63.
- ⁴⁸ Ermekova, K.S., Zambrano, N., Linn, H., Minopoli, G., Gertler, F., Russo, T., et al. (1997). The WW domain of neural protein FE65 interacts with proline-rich motifs in Mena, the mammalian homolog of Drosophila enabled. *J Biol Chem* **272**: 32869-77.
- ⁴⁹ Favre, B., Fontao, L., Koster, J., Shafaatian, R., Jaunin, F., Saurat, J.H., et al. (2001). The hemidesmosomal protein bullous pemphigoid antigen 1 and the integrin beta 4 subunit bind to ERBIN. Molecular cloning of multiple alternative splice variants of ERBIN and analysis of their tissue expression. *J Biol Chem* **276**: 32427-36.
- ⁵⁰ Felsenstein, J. (1989). PHYLIP - Phylogeny Inference Package (Version 3.2). *Cladistics* **5**: 164-6.
- ⁵¹ Feng, S., Chen, J.K., Yu, H., Simon, J.A. & Schreiber, S.L. (1994). Two binding orientations for peptides to the Src SH3 domain: development of a general model for SH3-ligand interactions. *Science* **266**: 1241-7.
- ⁵² Fields, S. & Song, O. (1989). A novel genetic system to detect protein-protein interactions. *Nature* **340**: 245-6.
- ⁵³ Fisher, R.A. (1935). *The design of experiments*, Oliver & Boyd, Edinburgh & London.
- ⁵⁴ Frank, R. (1992). Spot synthesis: an easy technique for the positionally addressable, parallel chemical synthesis on a membrane support. *Tetrahedron* **48**: 9217-32.
- ⁵⁵ Fuh, G., Pisabarro, M.T., Li, Y., Quan, C., Lasky, L.A. & Sidhu, S.S. (2000). Analysis of PDZ domain-ligand interactions using carboxyl-terminal phage display. *J Biol Chem* **275**: 21486-91.
- ⁵⁶ Gallivan, J.P. & Dougherty, D.A. (1999). Cation-pi interactions in structural biology. *Proc Natl Acad Sci U S A* **96**: 9459-64.
- ⁵⁷ Gavin, A.C., Bosche, M., Krause, R., Grandi, P., Marzioch, M., Bauer, A., et al. (2002). Functional organization of the yeast proteome by systematic analysis of protein complexes. *Nature* **415**: 141-7.
- ⁵⁸ Gavva, N.R., Gavva, R., Ermekova, K., Sudol, M. & Shen, C.J. (1997). Interaction of WW domains with hematopoietic transcription factor p45/NF-E2 and RNA polymerase II. *J Biol Chem* **272**: 24105-8.
- ⁵⁹ Gee, S.H., Madhavan, R., Levinson, S.R., Caldwell, J.H., Sealock, R. & Froehner, S.C. (1998). Interaction of muscle and brain sodium channels with multiple members of the syntrophin family of dystrophin-associated proteins. *J Neurosci* **18**: 128-37.

- ⁶⁰ Ghose, A.K. & Crippen, G.M. (1986). Atomic Physicochemical Parameters for Three-Dimensional Structure-Directed Quantitative Structure-Activity Relationships I. Partition Coefficients as a Measure of Hydrophobicity. *J Comp Chem* **7**: 565-77.
- ⁶¹ Gobel, U., Sander, C., Schneider, R. & Valencia, A. (1994). Correlated mutations and residue contacts in proteins. *Proteins* **18**: 309-17.
- ⁶² Grootjans, J.J., Reekmans, G., Ceulemans, H. & David, G. (2000). Syntenin-syndecan binding requires syndecan-synteny and the co-operation of both PDZ domains of syntenin. *J Biol Chem* **275**: 19933-41.
- ⁶³ Halperin, I., Ma, B., Wolfson, H. & Nussinov, R. (2002). Principles of docking: An overview of search algorithms and a guide to scoring functions. *Proteins* **47**: 409-43.
- ⁶⁴ Hannenhalli, S.S. & Russell, R.B. (2000). Analysis and prediction of functional sub-types from protein sequence alignments. *J Mol Biol* **303**: 61-76.
- ⁶⁵ Harris, B.Z. & Lim, W.A. (2001). Mechanism and role of PDZ domains in signaling complex assembly. *J Cell Sci* **114**: 3219-31.
- ⁶⁶ Hasegawa, M., Cuenda, A., Spillantini, M.G., Thomas, G.M., Buee-Scherrer, V., Cohen, P., et al. (1999). Stress-activated protein kinase-3 interacts with the PDZ domain of alpha1-syntrophin. A mechanism for specific substrate recognition. *J Biol Chem* **274**: 12626-31.
- ⁶⁷ Henikoff, S., Greene, E.A., Pietrokovski, S., Bork, P., Attwood, T.K. & Hood, L. (1997). Gene families: the taxonomy of protein paralogs and chimeras. *Science* **278**: 609-14.
- ⁶⁸ Higgs, H.N. & Pollard, T.D. (1999). Regulation of actin polymerization by Arp2/3 complex and WASp/Scar proteins. *J Biol Chem* **274**: 32531-4.
- ⁶⁹ Hillier, B.J., Christopherson, K.S., Prehoda, K.E., Bredt, D.S. & Lim, W.A. (1999). Unexpected modes of PDZ domain scaffolding revealed by structure of nNOS-syntrophin complex. *Science* **284**: 812-5.
- ⁷⁰ Ho, Y., Gruhler, A., Heilbut, A., Bader, G.D., Moore, L., Adams, S.L., et al. (2002). Systematic identification of protein complexes in *Saccharomyces cerevisiae* by mass spectrometry. *Nature* **415**: 180-3.
- ⁷¹ Hock, B., Bohme, B., Karn, T., Yamamoto, T., Kaibuchi, K., Holtrich, U., et al. (1998). PDZ-domain-mediated interaction of the Eph-related receptor tyrosine kinase EphB3 and the ras-binding protein AF6 depends on the kinase activity of the receptor. *Proc Natl Acad Sci U S A* **95**: 9779-84.
- ⁷² Hoffmann, R. & Valencia, A. (2003). Protein interaction: same network, different hubs. *Trends Genet* **19**: 681-3.
- ⁷³ Hoffmuller, U., Knaute, T., Hahn, M., Hohne, W., Schneider-Mergener, J. & Kramer, A. (2000). Evolutionary transition pathways for changing peptide ligand specificity and structure. *Embo J* **19**: 4866-74.
- ⁷⁴ Hoffmüller, U., Russwurm, M., Kleinjung, F., Ashurst, J., Oschkinat, H., Volkmer-Engert, R., et al. (1999). Interaktion einer PDZ-Proteindomäne mit einer synthetischen Bibliothek aller C-Termini humaner Proteine. *Angew Chemie* **11**: 2180-4.
- ⁷⁵ Hu, H., Columbus, J., Zhang, Y., Wu, D., Lian, L., Yang, S., et al. (2004). A map of WW domain family interactions. *Proteomics* **4**: 643-55.
- ⁷⁶ Huang, X., Poy, F., Zhang, R., Joachimiak, A., Sudol, M. & Eck, M.J. (2000). Structure of a WW domain containing fragment of dystrophin in complex with beta-dystroglycan. *Nat Struct Biol* **7**: 634-8.
- ⁷⁷ Huang, Y.Z., Zang, M., Xiong, W.C., Luo, Z. & Mei, L. (2003). Erbin suppresses the MAP kinase pathway. *J Biol Chem* **278**: 1108-14.
- ⁷⁸ Hubbard, T., Andrews, D., Caccamo, M., Cameron, G., Chen, Y., Clamp, M., et al. (2005). Ensembl 2005. *Nucleic Acids Res* **33 Database Issue**: D447-53.
- ⁷⁹ Hung, A.Y. & Sheng, M. (2002). PDZ domains: structural modules for protein complex assembly. *J Biol Chem* **277**: 5699-702.

- ⁸⁰ Ikeda, W., Nakanishi, H., Miyoshi, J., Mandai, K., Ishizaki, H., Tanaka, M., et al. (1999). Afadin: A key molecule essential for structural organization of cell-cell junctions of polarized epithelia during embryogenesis. *J Cell Biol* **146**: 1117-32.
- ⁸¹ Im, Y.J., Park, S.H., Rho, S.H., Lee, J.H., Kang, G.B., Sheng, M., et al. (2003). Crystal structure of GRIP1 PDZ6-peptide complex reveals the structural basis for class II PDZ target recognition and PDZ domain-mediated multimerization. *J Biol Chem* **278**: 8501-7.
- ⁸² Ito, T., Chiba, T., Ozawa, R., Yoshida, M., Hattori, M. & Sakaki, Y. (2001). A comprehensive two-hybrid analysis to explore the yeast protein interactome. *Proc Natl Acad Sci U S A* **98**: 4569-74.
- ⁸³ Ito, T., Tashiro, K., Muta, S., Ozawa, R., Chiba, T., Nishizawa, M., et al. (2000). Toward a protein-protein interaction map of the budding yeast: A comprehensive system to examine two-hybrid interactions in all possible combinations between the yeast proteins. *Proc Natl Acad Sci U S A* **97**: 1143-7.
- ⁸⁴ Jansen, R., Yu, H., Greenbaum, D., Kluger, Y., Krogan, N.J., Chung, S., et al. (2003). A Bayesian networks approach for predicting protein-protein interactions from genomic data. *Science* **302**: 449-53.
- ⁸⁵ Jaulin-Bastard, F., Arsanto, J.P., Le Bivic, A., Navarro, C., Vely, F., Saito, H., et al. (2002). Interaction between Erbin and a Catenin-related protein in epithelial cells. *J Biol Chem* **277**: 2869-75.
- ⁸⁶ Kabsch, W. & Sander, C. (1983). Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features. *Biopolymers* **22**: 2577-637.
- ⁸⁷ Kanelis, V., Rotin, D. & Forman-Kay, J.D. (2001). Solution structure of a Nedd4 WW domain-ENaC peptide complex. *Nat Struct Biol* **8**: 407-12.
- ⁸⁸ Kang, B.S., Cooper, D.R., Devedjiev, Y., Derewenda, U. & Derewenda, Z.S. (2003). Molecular roots of degenerate specificity in syntenin's PDZ2 domain: reassessment of the PDZ recognition paradigm. *Structure (Camb)* **11**: 845-53.
- ⁸⁹ Karthikeyan, S., Leung, T., Birrane, G., Webster, G. & Ladias, J.A. (2001). Crystal structure of the PDZ1 domain of human Na(+)/H(+) exchanger regulatory factor provides insights into the mechanism of carboxyl-terminal leucine recognition by class I PDZ domains. *J Mol Biol* **308**: 963-73.
- ⁹⁰ Karthikeyan, S., Leung, T. & Ladias, J.A. (2002). Structural determinants of the Na+/H+ exchanger regulatory factor interaction with the beta 2 adrenergic and platelet-derived growth factor receptors. *J Biol Chem* **277**: 18973-8.
- ⁹¹ Kato, Y., Nagata, K., Takahashi, M., Lian, L., Herrero, J.J., Sudol, M., et al. (2004). Common mechanism of ligand recognition by group II/III WW domains: redefining their functional classification. *J Biol Chem* **279**: 31833-41.
- ⁹² Kavsak, P., Rasmussen, R.K., Causing, C.G., Bonni, S., Zhu, H., Thomsen, G.H., et al. (2000). Smad7 binds to Smurf2 to form an E3 ubiquitin ligase that targets the TGF beta receptor for degradation. *Mol. Cell* **6**: 1365-75.
- ⁹³ Kay, B.K., Williamson, M.P. & Sudol, M. (2000). The importance of being proline: the interaction of proline-rich motifs in signaling proteins with their cognate domains. *FASEB J.* **14**: 231-41.
- ⁹⁴ Kim, E., Niethammer, M., Rothschild, A., Jan, Y.N. & Sheng, M. (1995). Clustering of Shaker-type K⁺ channels by interaction with a family of membrane-associated guanylate kinases. *Nature* **378**: 85-8.
- ⁹⁵ Kolch, W. (2003). Erbin: sorting out ErbB2 receptors or giving Ras a break? *Sci STKE* **2003**: pe37.
- ⁹⁶ Komuro, A., Saeki, M. & Kato, S. (1999). Npw38, a novel nuclear protein possessing a WW domain capable of activating basal transcription. *Nucleic Acids Res* **27**: 1957-65.
- ⁹⁷ Koradi, R., Billeter, M. & Wuthrich, K. (1996). MOLMOL: a program for display and analysis of macromolecular structures. *J. Mol. Graph.* **14**: 51-5, 29-32.

- ⁹⁸ Kornau, H.C., Schenker, L.T., Kennedy, M.B. & Seuberg, P.H. (1995). Domain interaction between NMDA receptor subunits and the postsynaptic density protein PSD-95. *Science* **269**: 1737-40.
- ⁹⁹ Kozlov, G., Banville, D., Gehring, K. & Ekiel, I. (2002). Solution structure of the PDZ2 domain from cytosolic human phosphatase hPTP1E complexed with a peptide reveals contribution of the beta2-beta3 loop to PDZ domain-ligand interactions. *J Mol Biol* **320**: 813-20.
- ¹⁰⁰ Kramer, A., Reineke, U., Dong, L., Hoffmann, B., Hoffmuller, U., Winkler, D., et al. (1999). Spot synthesis: observations and optimizations. *J. Pept. Res.* **54**: 319-27.
- ¹⁰¹ Kuntz, I.D. (1992). Structure-based strategies for drug design and discovery. *Science* **257**: 1078-82.
- ¹⁰² Laird, N.M. & Ware, J.H. (1982). Random-effects models for longitudinal data. *Biometrics* **38**: 963-74.
- ¹⁰³ Lander, E.S., Linton, L.M., Birren, B., Nusbaum, C., Zody, M.C., Baldwin, J., et al. (2001). Initial sequencing and analysis of the human genome. *Nature* **409**: 860-921.
- ¹⁰⁴ Landgraf, C., Panni, S., Montecchi-Palazzi, L., Castagnoli, L., Schneider-Mergener, J., Volkmer-Engert, R., et al. (2004). Protein interaction networks by proteome peptide scanning. *PLoS Biol* **2**: E14.
- ¹⁰⁵ Laskowski, R.A., Rullmann, J.A., MacArthur, M.W., Kaptein, R. & Thornton, J.M. (1996). AQUA and PROCHECK-NMR: programs for checking the quality of protein structures solved by NMR. *J. Biomol. NMR* **8**: 477-86.
- ¹⁰⁶ Lau, L.F., Mammen, A., Ehlers, M.D., Kindler, S., Chung, W.J., Garner, C.C., et al. (1996). Interaction of the N-methyl-D-aspartate receptor complex with a novel synapse-associated protein, SAP102. *J Biol Chem* **271**: 21622-8.
- ¹⁰⁷ Legrain, P., Wojcik, J. & Gauthier, J.M. (2001). Protein-protein interaction maps: a lead towards cellular functions. *Trends Genet* **17**: 346-52.
- ¹⁰⁸ Letunic, I., Copley, R.R., Schmidt, S., Ciccarelli, F.D., Doerks, T., Schultz, J., et al. (2004). SMART 4.0: towards genomic data integration. *Nucleic Acids Res* **32**: D142-4.
- ¹⁰⁹ Lim, W.A., Richards, F.M. & Fox, R.O. (1994). Structural determinants of peptide-binding orientation and of sequence specificity in SH3 domains. *Nature* **372**: 375-9.
- ¹¹⁰ Lu, P.J., Wulf, G., Zhou, X.Z., Davies, P. & Lu, K.P. (1999). The prolyl isomerase Pin1 restores the function of Alzheimer-associated phosphorylated tau protein. *Nature* **399**: 784-8.
- ¹¹¹ Ludes-Meyers, J.H., Kil, H., Bednarek, A.K., Drake, J., Bedford, M.T. & Aldaz, C.M. (2004). WWOX binds the specific proline-rich ligand PPXY: identification of candidate interacting proteins. *Oncogene* **23**: 5049-55.
- ¹¹² Macias, M.J., Gervais, V., Civera, C. & Oschkinat, H. (2000). Structural analysis of WW domains and design of a WW prototype. *Nat Struct Biol* **7**: 375-9.
- ¹¹³ Macias, M.J., Hyvonen, M., Baraldi, E., Schultz, J., Sudol, M., Saraste, M., et al. (1996). Structure of the WW domain of a kinase-associated protein complexed with a proline-rich peptide. *Nature* **382**: 646-9.
- ¹¹⁴ Madura, J.D., Briggs, J.M., Wade, R.C., Davis, M.E., Luty, B.A., Ilin, A., et al. (1995). Electrostatics and diffusion of molecules in solution: simulations with the university of Houston brownian dynamics program. *Comput Phys Commun* **91**: 57-95.
- ¹¹⁵ McGhee, J.D. & von Hippel, P.H. (1974). Theoretical aspects of DNA-protein interactions: co-operative and non-co-operative binding of large ligands to a one-dimensional homogeneous lattice. *J Mol Biol* **86**: 469-89.
- ¹¹⁶ Merrifield, B. (1986). Solid phase synthesis. *Science* **232**: 341-7.
- ¹¹⁷ Montell, C. (1998). TRP trapped in fly signaling web. *Curr. Opin. Neurobiol.* **8**: 389-97.
- ¹¹⁸ Morais Cabral, J.H., Petosa, C., Sutcliffe, M.J., Raza, S., Byron, O., Poy, F., et al. (1996). Crystal structure of a PDZ domain. *Nature* **382**: 649-52.

- ¹¹⁹ Morris, A.L., MacArthur, M.W., Hutchinson, E.G. & Thornton, J.M. (1992). Stereochemical quality of protein structure coordinates. *Proteins* **12**: 345-64.
- ¹²⁰ Murzin, A.G. (1998). How far divergent evolution goes in proteins. *Curr. Opin. Struct. Biol.* **8**: 380-7.
- ¹²¹ Nicholas, K.B. & Nicholas, H.B.J. (1997). *GeneDoc: a tool for editing and annotating multiple sequence alignments*, Distributed by the author.
- ¹²² Niethammer, M., Valschanoff, J.G., Kapoor, T.M., Allison, D.W., Weinberg, T.M., Craig, A.M., et al. (1998). CRIPT, a novel postsynaptic protein that binds to the third PDZ domain of PSD-95/SAP90. *Neuron* **20**: 693-707.
- ¹²³ Nishizawa, K., Freund, C., Li, J., Wagner, G. & Reinherz, E.L. (1998). Identification of a proline-binding motif regulating CD2-triggered T lymphocyte activation. *Proc Natl Acad Sci U S A* **95**: 14897-902.
- ¹²⁴ Noury, C., Grant, S.G. & Borg, J.P. (2003). PDZ domain proteins: plug and play! *Sci STKE* **2003**: RE7.
- ¹²⁵ Ohno, S. (2001). Intercellular junctions and cellular polarity: the PAR-aPKC complex, a conserved core cassette playing fundamental roles in cell polarity. *Curr. Opin. Cell Biol.* **13**: 641-8.
- ¹²⁶ Olalla, L., Aledo, J.C., Bannenberg, G. & Marquez, J. (2001). The C-terminus of human glutaminase L mediates association with PDZ domain-containing proteins. *FEBS Lett* **488**: 116-22.
- ¹²⁷ Ortiz, A.R., Pisabarro, M.T., Gago, F. & Wade, R.C. (1995). Prediction of drug binding affinities by comparative binding energy analysis. *J Med Chem* **38**: 2681-91.
- ¹²⁸ Oschkinat, H. (1999). A new type of PDZ domain recognition. *Nat Struct Biol* **6**: 408-10.
- ¹²⁹ Otte, L. (2004). Charakterisierung von WW-Domänen-Liganden-Wechselwirkungen mit Hilfe synthetischer Protein- und Peptidbibliotheken, Dissertation an der Humboldt-Universität Berlin.
- ¹³⁰ Otte, L., Wiedemann, U., Schlegel, B., Pires, J.R., Beyermann, M., Schmieder, P., et al. (2003). WW domain sequence activity relationships identified using ligand recognition propensities of 42 WW domains. *Protein Sci.* **12**: 491-500.
- ¹³¹ Page, R.D. (1996). TreeView: an application to display phylogenetic trees on personal computers. *Comput. Appl. Biosci.* **12**: 357-8.
- ¹³² Passani, L.A., Bedford, M.T., Faber, P.W., McGinnis, K.M., Sharp, A.H., Gusella, J.F., et al. (2000). Huntington's WW domain partners in Huntington's disease post-mortem brain fulfill genetic criteria for direct involvement in Huntington's disease pathogenesis. *Hum. Mol. Genet.* **9**: 2175-82.
- ¹³³ Patrie, K.M., Drescher, A.J., Welihinda, A., Mundel, P. & Margolis, B. (2002). Interaction of two actin-binding proteins, synaptopodin and alpha-actinin-4, with the tight junction protein MAGI-1. *J Biol Chem* **277**: 30183-90.
- ¹³⁴ Pawson, T. & Nash, P. (2003). Assembly of cell regulatory systems through protein interaction domains. *Science* **300**: 445-52.
- ¹³⁵ Pazos, F. & Valencia, A. (2002). In silico two-hybrid system for the selection of physically interacting protein pairs. *Proteins* **47**: 219-27.
- ¹³⁶ Penkert, R.R., DiVittorio, H.M. & Prehoda, K.E. (2004). Internal recognition through PDZ domain plasticity in the Par-6-Pals1 complex. *Nat Struct Mol Biol* **11**: 1122-7.
- ¹³⁷ Pires, J.R., Parthier, C., Aido-Machado, R.D., Wiedemann, U., Otte, L., Bohm, G., et al. (2005). Structural Basis for APPTPPPLPP Peptide Recognition by the FBP11WW1 Domain. *J Mol Biol* **348**: 399-408.
- ¹³⁸ Pires, J.R., Taha-Nejad, F., Toepert, F., Ast, T., Hoffmuller, U., Schneider-Mergener, J., et al. (2001). Solution structures of the YAP65 WW domain and the variant L30 K in complex with the peptides GTPPPPYTVG, N-(n-octyl)-GPPPY and PLPPY and the application of peptide libraries reveal a minimal binding epitope. *J Mol Biol* **314**: 1147-56.

- ¹³⁹ Pirozzi, G., McConnell, S.J., Uveges, A.J., Carter, J.M., Sparks, A.B., Kay, B.K., et al. (1997). Identification of novel human WW domain-containing proteins by cloning of ligand targets. *J Biol Chem* **272**: 14611-6.
- ¹⁴⁰ Pizzuti, A., Novelli, G., Mari, A., Ratti, A., Colosimo, A., Amati, F., et al. (1996). Human homologue sequences to the Drosophila dishevelled segment-polarity gene are deleted in the DiGeorge syndrome. *Am J Hum Genet* **58**: 722-9.
- ¹⁴¹ Ponting, C.P. (1997). Evidence for PDZ domains in bacteria, yeast, and plants. *Protein Sci.* **6**: 464-8.
- ¹⁴² Ponting, C.P., Schultz, J., Copley, R.R., Andrade, M.A. & Bork, P. (2000). Evolution of domain families. *Adv. Protein Chem.* **54**: 185-244.
- ¹⁴³ Pornillos, O., Alam, S.L., Rich, R.L., Myszka, D.G., Davis, D.R. & Sundquist, W.I. (2002). Structure and functional interactions of the Tsg101 UEV domain. *Embo J* **21**: 2397-406.
- ¹⁴⁴ Powell, M.J.D. (1977). Restart Procedures for the Conjugate Gradient Method. *Math Prog* **12**: 241-54.
- ¹⁴⁵ Press, W.H., Teukolsky, S.A., Vetterling, W.T. & Flannery, B.P. (1992). *Numerical recipies in C: the Art of Scientific Computing*. 2nd edit, Cambridge University Press, Cambridge & New York.
- ¹⁴⁶ Quilliam, L.A., Castro, A.F., Rogers-Graham, K.S., Martin, C.B., Der, C.J. & Bi, C. (1999). M-Ras/R-Ras3, a transforming ras protein regulated by Sos1, GRF1, and p120 Ras GTPase-activating protein, interacts with the putative Ras effector AF6. *J Biol Chem* **274**: 23850-7.
- ¹⁴⁷ Radziwill, G., Erdmann, R.A., Margelisch, U. & Moelling, K. (2003). The Bcr kinase downregulates Ras signaling by phosphorylating AF-6 and binding to its PDZ domain. *Mol. Cell. Biol.* **23**: 4663-72.
- ¹⁴⁸ R-Development-Core-Team. (2003). *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Wien.
- ¹⁴⁹ Reina, J., Lacroix, E., Hobson, S.D., Fernandez-Ballester, G., Rybin, V., Schwab, M.S., et al. (2002). Computer-aided design of a PDZ domain to recognize new target sequences. *Nat Struct Biol* **9**: 621-7.
- ¹⁵⁰ Rentschler, S., Linn, H., Deininger, K., Bedford, M.T., Espanol, X. & Sudol, M. (1999). The WW domain of dystrophin requires EF-hands region to interact with beta-dystroglycan. *Biol. Chem.* **380**: 431-42.
- ¹⁵¹ Richardson, J.S. (1981). The anatomy and taxonomy of protein structure. *Adv. Protein Chem.* **34**: 167-339.
- ¹⁵² Rognan, D., Lauemoller, S.L., Holm, A., Buus, S. & Tschinke, V. (1999). Predicting binding affinities of protein ligands from three-dimensional models: application to peptide binding to class I major histocompatibility proteins. *J Med Chem* **42**: 4650-8.
- ¹⁵³ Rubin, G.M., Yandell, M.D., Wortman, J.R., Gabor Miklos, G.L., Nelson, C.R., Hariharan, I.K., et al. (2000). Comparative genomics of the eukaryotes. *Science* **287**: 2204-15.
- ¹⁵⁴ Russo, T., Faraonio, R., Minopoli, G., De Candia, P., De Renzis, S. & Zambrano, N. (1998). Fe65 and the protein network centered around the cytosolic domain of the Alzheimer's beta-amyloid precursor protein. *FEBS Lett* **434**: 1-7.
- ¹⁵⁵ Saitou, N. & Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**: 406-25.
- ¹⁵⁶ Samanta, U. & Chakrabarti, P. (2001). Assessing the role of tryptophan residues in the binding site. *Protein Eng.* **14**: 7-15.
- ¹⁵⁷ Schleinkofer, K., Wiedemann, U., Otte, L., Wang, T., Krause, G., Oschkinat, H., et al. (2004). Comparative structural and energetic analysis of WW domain-peptide interactions. *J Mol Biol* **344**: 865-81.
- ¹⁵⁸ Schneider, T.D. & Stephens, R.M. (1990). Sequence logos: a new way to display consensus sequences. *Nucleic Acids Res* **18**: 6097-100.

- ¹⁵⁹ Schueler-Furman, O., Altuvia, Y., Sette, A. & Margalit, H. (2000). Structure-based prediction of binding peptides to MHC class I molecules: application to a broad range of MHC alleles. *Protein Sci.* **9**: 1838-46.
- ¹⁶⁰ Schultz, J., Hoffmuller, U., Krause, G., Ashurst, J., Macias, M.J., Schmieder, P., et al. (1998). Specific interactions between the syntrophin PDZ domain and voltage-gated sodium channels. *Nat Struct Biol* **5**: 19-24.
- ¹⁶¹ Schultz, J., Milpetz, F., Bork, P. & Ponting, C.P. (1998). SMART, a simple modular architecture research tool: identification of signaling domains. *Proc Natl Acad Sci U S A* **95**: 5857-64.
- ¹⁶² Schuster-Bockler, B., Schultz, J. & Rahmann, S. (2004). HMM Logos for visualization of protein families. *BMC Bioinformatics* **5**: 7.
- ¹⁶³ Setou, M., Nakagawa, T., Seog, D.H. & Hirokawa, N. (2000). Kinesin superfamily motor protein KIF17 and mLin-10 in NMDA receptor-containing vesicle transport. *Science* **288**: 1796-802.
- ¹⁶⁴ Shen, M., Stukenberg, P.T., Kirschner, M.W. & Lu, K.P. (1998). The essential mitotic peptidyl-prolyl isomerase Pin1 binds and regulates mitosis-specific phosphoproteins. *Genes Dev.* **12**: 706-20.
- ¹⁶⁵ Sherman, D.L., Fabrizi, C., Gillespie, C.S. & Brophy, P.J. (2001). Specific disruption of a schwann cell dystrophin-related protein complex in a demyelinating neuropathy. *Neuron* **30**: 677-87.
- ¹⁶⁶ Shi, Z., Olson, C.A., Rose, G.D., Baldwin, R.L. & Kallenbach, N.R. (2002). Polyproline II structure in a sequence of seven alanine residues. *Proc Natl Acad Sci U S A* **99**: 9190-5.
- ¹⁶⁷ Sidhu, S.S., Lowman, H.B., Cunningham, B.C. & Wells, J.A. (2000). Phage display for selection of novel binding peptides. *Methods Enzymol.* **328**: 333-63.
- ¹⁶⁸ Skelton, N.J., Koehler, M.F., Zobel, K., Wong, W.L., Yeh, S., Pisabarro, M.T., et al. (2003). Origins of PDZ domain ligand specificity. Structure determination and mutagenesis of the Erbin PDZ domain. *J Biol Chem* **278**: 7645-54.
- ¹⁶⁹ Smith, G.R. & Sternberg, M.J. (2002). Prediction of protein-protein interactions by docking methods. *Curr. Opin. Struct. Biol.* **12**: 28-35.
- ¹⁷⁰ Songyang, Z., Fanning, A.S., Fu, C., Xu, J., Marfatia, S.M., Chishti, A.H., et al. (1997). Recognition of unique carboxyl-terminal motifs by distinct PDZ domains. *Science* **275**: 73-7.
- ¹⁷¹ Sotgia, F., Lee, H., Bedford, M.T., Petrucci, T., Sudol, M. & Lisanti, M.P. (2001). Tyrosine phosphorylation of beta-dystroglycan at its WW domain binding motif, PPxY, recruits SH2 domain containing proteins. *Biochemistry (Mosc.)* **40**: 14585-92.
- ¹⁷² Staub, O., Dho, S., Henry, P., Correa, J., Ishikawa, T., McGlade, J., et al. (1996). WW domains of Nedd4 bind to the proline-rich PY motifs in the epithelial Na⁺ channel deleted in Liddle's syndrome. *Embo J* **15**: 2371-80.
- ¹⁷³ Staub, O., Gautschi, I., Ishikawa, T., Breitschopf, K., Ciechanover, A., Schild, L., et al. (1997). Regulation of stability and function of the epithelial Na⁺ channel (ENaC) by ubiquitination. *Embo J* **16**: 6325-36.
- ¹⁷⁴ Steffen, M., Petti, A., Aach, J., D'Haeseleer, P. & Church, G. (2002). Automated modelling of signal transduction networks. *BMC Bioinformatics* **3**: 34.
- ¹⁷⁵ Sudol, M. (1996). Structure and function of the WW domain. *Prog. Biophys. Mol. Biol.* **65**: 113-32.
- ¹⁷⁶ Sudol, M. (1998). From Src Homology domains to other signaling modules: proposal of the 'protein recognition code'. *Oncogene* **17**: 1469-74.
- ¹⁷⁷ Sudol, M. (2002). WW DOMAINS. In *Wiley Encyclopedia of Molecular Medicine*, Vol. **5**: 3405-8. John Wiley & Sons Inc., New York.
- ¹⁷⁸ Sudol, M. & Hunter, T. (2000). NeW wrinkles for an old domain. *Cell* **103**: 1001-4.

- ¹⁷⁹ Takahashi, K., Nakanishi, H., Miyahara, M., Mandai, K., Satoh, K., Satoh, A., et al. (1999). Nectin/PRR: an immunoglobulin-like cell adhesion molecule recruited to cadherin-based adherens junctions through interaction with Afadin, a PDZ domain-containing protein. *J Cell Biol* **145**: 539-49.
- ¹⁸⁰ Tassabehji, M., Metcalfe, K., Fergusson, W.D., Carette, M.J., Dore, J.K., Donnai, D., et al. (1996). LIM-kinase deleted in Williams syndrome. *Nat Genet* **13**: 272-3.
- ¹⁸¹ Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F. & Higgins, D.G. (1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **25**: 4876-82.
- ¹⁸² Toepert, F., Knaute, T., Guffler, S., Pires, J.R., Matzdorf, T., Oschkinat, H., et al. (2003). Combining SPOT synthesis and native peptide ligation to create large arrays of WW protein domains. *Angew. Chem. Int. Ed. Engl.* **42**: 1136-40.
- ¹⁸³ Toepert, F., Pires, J.R., Landgraf, C., Oschkinat, H. & Schneider-Mergener, J. (2001). Synthesis of an Array Comprising 837 Variants of the hYAP WW Protein Domain This work was supported by the DFG (INK 16/B1-1), by the Fonds der Chemischen Industrie, and by the Universitätsklinikum Charité Berlin. *Angew. Chem. Int. Ed. Engl.* **40**: 897-900.
- ¹⁸⁴ Tong, A.H., Drees, B., Nardelli, G., Bader, G.D., Brannetti, B., Castagnoli, L., et al. (2002). A combined experimental and computational strategy to define protein interaction networks for peptide recognition modules. *Science* **295**: 321-4.
- ¹⁸⁵ Traweger, A., Fang, D., Liu, Y.C., Stelzhammer, W., Krizbai, I.A., Fresser, F., et al. (2002). The tight junction-specific protein occludin is a functional target of the E3 ubiquitin-protein ligase itch. *J Biol Chem* **277**: 10201-8.
- ¹⁸⁶ Tsodikov, O.V., Holbrook, J.A., Shkel, I.A. & Record, M.T., Jr. (2001). Analytic binding isotherms describing competitive interactions of a protein ligand with specific and nonspecific sites on the same DNA oligomer. *Biophys. J.* **81**: 1960-9.
- ¹⁸⁷ Uetz, P., Giot, L., Cagney, G., Mansfield, T.A., Judson, R.S., Knight, J.R., et al. (2000). A comprehensive analysis of protein-protein interactions in *Saccharomyces cerevisiae*. *Nature* **403**: 623-7.
- ¹⁸⁸ Ullmer, C., Schmuck, K., Figge, A. & Lubbert, H. (1998). Cloning and characterization of MUPP1, a novel PDZ domain protein. *FEBS Lett* **424**: 63-8.
- ¹⁸⁹ Vaccaro, P. & Dente, L. (2002). PDZ domains: troubles in classification. *FEBS Lett* **512**: 345-9.
- ¹⁹⁰ Valencia, A. & Pazos, F. (2002). Computational methods for the prediction of protein interactions. *Curr. Opin. Struct. Biol.* **12**: 368-73.
- ¹⁹¹ Venables, W.N. & Ripley, B.D. (2001). *Modern Applied Statistics with S*, Springer, Berlin & New York.
- ¹⁹² Venter, J.C., Adams, M.D., Myers, E.W., Li, P.W., Mural, R.J., Sutton, G.G., et al. (2001). The sequence of the human genome. *Science* **291**: 1304-51.
- ¹⁹³ Verdecia, M.A., Bowman, M.E., Lu, K.P., Hunter, T. & Noel, J.P. (2000). Structural basis for phosphoserine-proline recognition by group IV WW domains. *Nat Struct Biol* **7**: 639-43.
- ¹⁹⁴ Verpy, E., Leibovici, M., Zwaenepoel, I., Liu, X.Z., Gal, A., Salem, N., et al. (2000). A defect in harmonin, a PDZ domain-containing protein expressed in the inner ear sensory hair cells, underlies Usher syndrome type 1C. *Nat Genet* **26**: 51-5.
- ¹⁹⁵ von Mering, C., Krause, R., Snel, B., Cornell, M., Oliver, S.G., Fields, S., et al. (2002). Comparative assessment of large-scale data sets of protein-protein interactions. *Nature* **417**: 399-403.
- ¹⁹⁶ Wallace, A.C., Laskowski, R.A. & Thornton, J.M. (1995). LIGPLOT: a program to generate schematic diagrams of protein-ligand interactions. *Protein Eng.* **8**: 127-34.
- ¹⁹⁷ Wang, G., Yang, J. & Huibregtse, J.M. (1999). Functional domains of the Rsp5 ubiquitin-protein ligase. *Mol. Cell. Biol.* **19**: 342-52.
- ¹⁹⁸ Wang, T. & Wade, R.C. (2001). Comparative binding energy (COMBINE) analysis of influenza neuraminidase-inhibitor complexes. *J Med Chem* **44**: 961-71.

- ¹⁹⁹ Wang, T. & Wade, R.C. (2002). Comparative binding energy (COMBINE) analysis of OppA-peptide complexes to relate structure to binding thermodynamics. *J Med Chem* **45**: 4828-37.
- ²⁰⁰ Wiedemann, U., Boisguerin, P., Leben, R., Leitner, D., Krause, G., Moelling, K., et al. (2004). Quantification of PDZ domain specificity, prediction of ligand affinity and rational design of super-binding peptides. *J Mol Biol* **343**: 703-18.
- ²⁰¹ Wiedemann, U., Labudde, D., Kleinau, G. & Krause, G. (2003). AnCoRe: Analysis of Complementary Residues. *Currents in Computational Molecular Biology (RECOMB)* **2003**: 105-6.
- ²⁰² Wiesner, S., Stier, G., Sattler, M. & Macias, M.J. (2002). Solution structure and ligand recognition of the WW domain pair of the yeast splicing factor Prp40. *J Mol Biol* **324**: 807-22.
- ²⁰³ Williamson, M.P. (1994). The structure and function of proline-rich regions in proteins. *Biochem. J.* **297** (Pt 2): 249-60.
- ²⁰⁴ Wood, J.D., Yuan, J., Margolis, R.L., Colomer, V., Duan, K., Kushi, J., et al. (1998). Atrophin-1, the DRPLA gene product, interacts with two families of WW domain-containing proteins. *Mol. Cell. Neurosci.* **11**: 149-60.
- ²⁰⁵ Woods, D.F. & Bryant, P.J. (1993). ZO-1, DlgA and PSD-95/SAP90: homologous proteins in tight, septate and synaptic cell junctions. *Mech. Dev.* **44**: 85-9.
- ²⁰⁶ Wulf, G., Ryo, A., Liou, Y.C. & Lu, K.P. (2003). The prolyl isomerase Pin1 in breast development and cancer. *Breast Cancer Res* **5**: 76-82.
- ²⁰⁷ Yaffe, M.B., Schutkowski, M., Shen, M., Zhou, X.Z., Stukenberg, P.T., Rahfeld, J.U., et al. (1997). Sequence-specific and phosphorylation-dependent proline isomerization: a potential mitotic regulatory mechanism. *Science* **278**: 1957-60.
- ²⁰⁸ Yagi, R., Chen, L.F., Shigesada, K., Murakami, Y. & Ito, Y. (1999). A WW domain-containing yes-associated protein (YAP) is a novel transcriptional co-activator. *Embo J* **18**: 2551-62.
- ²⁰⁹ Yamamoto, T., Harada, N., Kawano, Y., Taya, S. & Kaibuchi, K. (1999). In vivo interaction of AF-6 with activated Ras and ZO-1. *Biochem. Biophys. Res. Commun.* **259**: 103-7.
- ²¹⁰ Yu, J.W. & Lemmon, M.A. (2003). Genome-wide analysis of signaling domain function. *Curr. Opin. Chem. Biol.* **7**: 103-9.
- ²¹¹ Yusupov, M.M., Yusupova, G.Z., Baucom, A., Lieberman, K., Earnest, T.N., Cate, J.H., et al. (2001). Crystal structure of the ribosome at 5.5 Å resolution. *Science* **292**: 883-96.
- ²¹² Zarrinpar, A., Bhattacharyya, R.P. & Lim, W.A. (2003). The structure and function of proline recognition domains. *Sci STKE* **2003**: RE8.
- ²¹³ Zarrinpar, A. & Lim, W.A. (2000). Converging on proline: the mechanism of WW domain peptide recognition. *Nat Struct Biol* **7**: 611-3.
- ²¹⁴ Zhadanov, A.B., Provance, D.W., Jr., Speer, C.A., Coffin, J.D., Goss, D., Blixt, J.A., et al. (1999). Absence of the tight junctional protein AF-6 disrupts epithelial cell-cell junctions and cell polarity during mouse development. *Curr. Biol.* **9**: 880-8.
- ²¹⁵ Zhu, H., Bilgin, M., Bangham, R., Hall, D., Casamayor, A., Bertone, P., et al. (2001). Global analysis of protein activities using proteome chips. *Science* **293**: 2101-5.
- ²¹⁶ Zhu, H., Huang, S. & Dhar, P. (2004). The next step in systems biology: simulating the temporospatial dynamics of molecular network. *Bioessays* **26**: 68-72.
- ²¹⁷ Zimmermann, P., Meerschaert, K., Reekmans, G., Leenaerts, I., Small, J.V., Vandekerckhove, J., et al. (2002). PIP(2)-PDZ domain binding controls the association of syntenin with the plasma membrane. *Mol. Cell* **9**: 1215-25.