

# Literaturverzeichnis

- [1] B. Abdallah, A. Hassan, C. Benoist, D. Goula, J.P. Behr und B.A. Demeneix. A powerful nonviral vector for in vivo gene transfer into the adult mammalian brain: polyethylenimine. *Hum Gene Ther*, 7(16):1947–54, 1996.
- [2] T. Akao, T. Nakayama, K. Takeshia und A. Ito. Design of a new cationic amphiphile with efficient DNA-transfection ability. *Biochem Mol Biol Int*, 34(5):915–20, 1994.
- [3] W. Anderson. Human gene therapy. *Nature*, 392:25–30, 1998.
- [4] C.V. Arancibia-Carcamo, H.B. Oral, D.O. Haskard, D.F. Larkin und A.J.T. George. Lipoadenofection-mediated gene delivery to the corneal endothelium. *Transplant*, 65:62–67.1998.
- [5] S.M. Arcasoy, J.D. Latoche, M. Gondor, B.R. Pitt und J.M. Pilewski. Polycations increase the efficiency of adenovirus-mediated gene. *Gene Ther*, 4:32–38, 1997.
- [6] G. Ascadi, S.S. Jiao, A. Jani, D. Duke, P. Williams, W. Chon und J.A. Wolff. Direct gene transfer and expression into rat heart in vivo. *New Biol*, 3:71–81, 1991.
- [7] A. Baker, M. Saltik, H. Lehrmann, I. Killisch, V. Mautner, G. Lamm, G. Christofori und M. Cotten. Polyethylenimine (PEI) is a simple, inexpensive and effective reagent for condensing and linking plasmid-DNA to adenovirus for gene delivery. *Gene Ther*, 4:773–782.1997.
- [8] R.P. Balasubramaniam, M.J. Bennett, A.M. Aberle, J.G. Malone, M.H. Nantz und R.W. Malone. Structural and functional analysis of cationic transfection lipids: the hydrophobic domain. *Gene Ther*, 3:163–172, 1996.
- [9] A.D. Bangham und R.W. Horne. Negative staining of phospholipids and their structured modification by surface active agents as observed in the electron microscope. *J Mol Biol*, 8:660–668, 1964.
- [10] A.D. Bangham, M.M. Standish und J.C. Watkins. Diffusion of univalent ions across lamellae of swollen phospholipids. *J Mol Biol*, 13:238–252, 1965.
- [11] L.G. Barron, K.B. Meyer und F.C. Jr Szoka. Effects of complement depletion on the pharmacokinetics and gene delivery mediated by cationic lipid-DNA complexes. *Hum Gene Ther*, 9(3):315–23, 1998.

- [12] F. Barthel, J.S. Remy, J.P. Loeffler und J.P. Behr. Gene transfer optimization with lipospermine-coated DNA. *DNA Cell Biol*, 12(6):553–60, 1993.
- [13] M. Baru, J.H. Axelrod und I. Nur. Liposome-encapsulated DNA-mediated gene transfer and synthesis of human factor IX in mice. *Gene*, 161(2):143–50, 1995.
- [14] C. F. Bennett, M. Y. Chiang, H. Chan, J. E. Shoemaker und C. Mirabelli. Cationic lipids enhance cellular uptake and activity of phosphorothioate antisense oligonucleotides. *Mol Pharmacol*, 41(6):1023–33, 1992.
- [15] T. Blessing, J.S. Remy und J.P. Behr. Monomolecular collapse of plasmid DNA into stable virus-like particles. *Proc Natl Acad Sci USA*, 95(4):1427–31, 1998.
- [16] R.C. Boucher. *Harrisons Innere Medizin. (K.J.K. Schmailzl Hrsg.)*, Zystische Fibrose. S. 1404–07. Blackwell Wissenschafts Verlag, Berlin, Wien. 1995.
- [17] T. Boulikas. Status of gene therapy in 1997: molecular mechanism, disease targets, and clinical applications. *Gene Ther Mol Biol*, 1:1–172, 1998.
- [18] O. Boussif, F. Lezoualc'h, M.A. Zanta, M.D. Mergny, D. Scherman, B. Demeneix, und J.P. Behr. A versatile vector for gene and oligonucleotide transfer into cells in culture and in vivo: polyethylenimine. *Proc Natl Acad Sci USA*, 92(16):7297–301, 1995.
- [19] B. Buchberger, E. Fernholz, H. vd Eltz und M. Hinzpeter. DOSPER Liposomal Transfection Reagent: A Reagent with unique Transfection Properties. *Biochem Inf*, 98:27–29, 1996.
- [20] N.M. Calvin und P.C. Hanawalt. High-efficiency transformation of bacterial cells by electroporation. *J Bacteriol*, 170:2796–801, 1988.
- [21] MR Capecci. High efficiency transformation by direct microinjection of DNA into cultured mammalian cells. *Cell*, 22:479–488, 1980.
- [22] N.J. Caplen, E. Kinrade, F. Sorgi, X. Gao, D. Gruenert, D. Geddes, C. Coutelle, L. Huang, E.W. Alton und R. Williamson. In vitro liposome-mediated DNA transfection of epithelial cell lines using the cationic liposome DC-Chol/DOPE. *Gene Ther*, 2(9):603–13, 1995.
- [23] C. P. Chang, C. S. Lazar, B. J. Walsh, M. Komuro, J. F. Collawn, L. A. Kuhn, J. A. Tainer, I. S. Trowbridge, M. G. Farquhar, M. G. Rosenfeld und et Al. Ligand-induced internalization of the epidermal growth factor receptor is mediated by multiple endocytic codes analogous to the tyrosine motif found in constitutively internalized receptors. *J Biol Chem*, 268(26):19312–20, 1993.
- [24] J.Y. Cherng. *Pharmaceutical aspects of polymer-based gene delivery systems. Preparation, characterisation and stabilisation*. PhD thesis, Utrecht University, Faculty of Pharmacy, 1999.
- [25] J.Y. Cherng, P. Vandewetering, H. Talsma und D.J.A. Crommelin. Effect of size and serum proteins on transfection efficiency. *Pharm Res*, 13:1038–1042, 1996.

- [26] M. Y. Chiang, H. Chan, M. A. Zounes, S. M. Freier, W. F. Lima und C. Bennett. Antisense oligonucleotides inhibit intercellular adhesion molecule 1 expression by two distinct mechanisms. *J Biol Chem*, 266(27):18162–71, 1991.
- [27] M. Chillón, J.H. Lee, A. Fasbender und M.J. Welsh. Adenovirus complexed with polyethylene glycol and cationic lipid is shielded from neutralizing antibodies in vitro. *Gene Ther*, 5(7):995–1002, 1998.
- [28] C.T. Chung und R.H. Miller. Preparation and Storage of Competent Escherichia Coli Cells. *Meth Enzymol*, 218:621–7, 1993.
- [29] P.R. Cullis und B. de Kruijff. Lipid polymorphism and the functional roles of lipids in biological membranes. *Biochim Biophys Acta*, 559:399–420, 1979.
- [30] H.Z. Cummies und E.R. Pike. *Photon Correlations Spectroscopy and Light Beating Spectroscopy*. Plenum Press, New York. NATO Advanced Study Institutes. Series B, Vol 3.1974.
- [31] H.Z. Cummies und E.R. Pike. *Photon Correlations Spectroscopy and Velocimetry*. Plenum Press, New York. NATO Advanced Study Institutes. Series B, Vol 23.1977.
- [32] D.T. Curiel, S. Agarwal, E. Wagner und M. Cotten. Adenovirus enhancement of transferrin-polylysine-mediated gene delivery. *Proc Natl Acad Sci USA*, 88:8850–4, 1991.
- [33] A. Dathe. Circulardichroismus-Untersuchungen von Ncp7/DNA-Komplexen. Persönliche Mitteilung.
- [34] R. J. Debs, L. P. Freedman, S. Edmunds, K. L. Gaensler, N. Duzgunes und K. R. Yamamoto. Regulation of gene expression in vivo by liposome-mediated delivery of a purified transcription factor. *J Biol Chem*, 265(18):10189–92, 1990.
- [35] J.E. Diederichs, D. Groth und R. Reszka. Plasma protein adsorption patterns onto cationic liposomes and lipoplexes - influence of cationic lipid head group. *J Lipos Res*, 8:52–53, 1998.
- [36] S.J. Eastman, C. Siegel, J. Tousignant, A.E. Smith, S.H. Cheng und R.K. Scheule. Biophysical characterisation of cationic lipid:DNA complexes. *Biochim Biophys Acta*, 1325:41–62, 1997.
- [37] R.F. Epand, R. Kraayenhof, G.J. Sterk, Fong Sang H.W. Wong und R.M. Epand. Fluorescent probes of membrane surface properties. *Biochim Biophys Acta*, 1284(2):191–5, 1996.
- [38] V. Escriou, C. Ciolina, Leclerc A. Helbling, P. Wils und D. Scherman. Cationic lipid-mediated gene transfer: analysis of cellular uptake and nuclear import of plasmid DNA. *Cell Biol Toxicol*, 14(2):95–104, 1998.

- [39] V. Escriou, C. Ciolina, F. Lacroix, G. Byk, D. Scherman und P. Wils. Cationic lipid-mediated gene transfer: effect of serum on cellular uptake and intracellular fate of lipopolyamine/DNA complexes. *Biochim Biophys Acta.*, 1368(2):276–88, 1998.
- [40] H. Farhood, R. Bottega, R.M. Epand und L. Huang. Effect of cationic cholesterol derivatives on gene transfer and protein kinase C activity. *Biochim Biophys Acta*, 1111(2):239–46, 1992.
- [41] H. Farhood, N. Serbina und L. Huang. The role of dioleoyl phosphatidylethanolamine in cationic liposome mediated gene transfer. *Biochim Biophys Acta*, 1235(2):289–95, 1995.
- [42] A. Fasbender, J. Marshall, T.O. Moninger, T. Grunst, S. Cheng und M.J. Welsh. Effect of co-lipids in enhancing cationic lipid-mediated gene transfer in vitro and in vivo. *Gene Ther*, 4:716–25, 1997.
- [43] W.G. Feero, S. Li, J.D. Rosenblatt, N. Sirianni, J.E. Morgan, T.A. Partridge, L. Huang und E.P. Hoffman. Selection and use of ligands for receptor-mediated gene delivery to myogenic cells. *Gene Ther*, 4(7):664–74, 1997.
- [44] J.H. Felgner, R. Kumar, C.N. Sridhar, C.J. Wheeler, Y.J. Tsai, R. Border, P. Ramsey, M. Martin und P.L. Felgner. Enhanced gene delivery and mechanism studies with a novel series of cationic lipid formulations. *J Biol Chem*, 269(4):2550–61, 1994.
- [45] P.L. Felgner, Y. Barenholz, J.P. Behr, S.H. Cheng, P. Cullis, L. Huang, J.A. Jessee, L. Seymour, F. Szoka, A.R. Thierry, E. Wagner und G. Wu. Nomenclature for synthetic gene delivery systems [editorial]. *Hum Gene Ther*, 8(5):511–2, 1997.
- [46] P.L. Felgner, T.R. Gadek, M. Holm, R. Roman, H.W. Chan, M. Wenz, J.P. Northrop, G.M. Ringold und M. Danielsen. Lipofection: a highly efficient and lipid-mediated DNA-transfection procedure. *Proc Natl Acad Sci USA*, 84:7413–7, 1987.
- [47] T. Ferkol, J.C. Perales, F. Mularo und R.W. Hanson. Receptor-mediated gene transfer into macrophages. *Proc Natl Acad Sci USA*, 83:101–105, 1996.
- [48] S. Ferrari, E. Moro, A. Pettenazzo, J.P. Behr, F. Zacchello und M. Scarpa. ExGen 500 is an efficient vector for gene delivery to lung epithelial cells in vitro and in vivo. *Gene Ther*, 4(10):1100–6, 1997.
- [49] D. S. Friend, D. Papahadjopoulos und R. Debs. Endocytosis and intracellular processing accompanying transfection mediated by cationic liposomes. *Biochim Biophys Acta*, 1278(1):41–50, 1996.
- [50] J.D. Fritz, H. Herweijer, G. Zhang und J.A. Wolff. Gene transfer into mammalian cells using histone-condensed plasmid-DNA. *Hum Gene Ther*, 7:1395–1404, 1996.
- [51] X. Gao und L. Huang. A novel cationic liposome reagent for efficient transfection of mammalian cells. *Biochem Biophys Res Commun*, 179:280–5, 1991.

- [52] X. Gao und L. Huang. Cytoplasmic expression of a reporter gene by co-delivery of T7 RNA polymerase and T7 promoter sequence with cationic liposomes. *Nucleic Acids Res*, 21(12):2867–72, 1993.
- [53] X. Gao und L. Huang. Cationic liposome-mediated gene transfer. *Gene Ther*, 2:710–722, 1995.
- [54] X. Gao und L. Huang. Potentiation of cationic liposome-mediated gene delivery by polycations. *Biochemistry*, 35(3):1027–36, 1996.
- [55] H. Gershon, R. Ghirlando, S.B. Guttman und A. Minsky. Mode of formation of and structural features of DNA-cationic liposome complexes used for transfection. *Biochemistry*, 32:7143–7151, 1993.
- [56] S. Gottschalk, J.T. Sparrow, J. Hauer, M.P. Mims, F.E. Leland S.L. Woo und L.C. Smith. A novel DNA-peptide complex for efficient gene transfer and expression in mammalian cells. *Gene Ther*, 3:448–457, 1996.
- [57] F.L. Graham und A.J. van der Eb. A new technique for the assay of infectivity of human adenovirus 5 DNA. *Virology*, 52:456–467, 1973.
- [58] D. Groth, O. Keil, C. Lehmann, M. Schneider, M. Rudolph und R. Reszka. Preparation and characterisation of a new lipospermine for gene delivery into various cell lines. *Int J Pharm*, 162:143–157, 1998a.
- [59] D. Groth, O. Keil, M. Schneider und R. Reszka. Transfection assay for dual determination of toxicity and gene expression. *Anal Biochem*, 258:141–143, 1998.
- [60] D. Groth und R. Reszka. Verfahren zur dualen Bestimmung von Zellvitalität und Effizienz nach Gentransfer in eukaryontische Zellen. *Deut Pat Applik*, S. 197–50–790, 1997.
- [61] D. Groth, R. Reszka und J.A. Schenk. PEG mediated transformation of *E. coli* is increased by room temperature incubation,. *Anal Bioch*, 240:302–304, 1996.
- [62] J. Gustafsson, G. Arvidson, G. Karlsson und M. Almgren. Complexes between cationic liposomes and DNA visualized by cryo-TEM. *Biochim Biophys Acta*, 1235:305–12, 1995.
- [63] D. Hanahan. Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol*, 166:557–80, 1983.
- [64] R.P. Harbottle, R.G. Cooper, S.L. Hart, A. Ladhoff, T. McKay, A.M. Knight, E. Wagner, A.D. Miller und C. Coutelle. An RGD-oligolysine peptide: a prototype construct for integrin-mediated gene delivery [see comments]. *Hum Gene Ther*, 9(7):1037–47, 1998.
- [65] S.L. Hart, E. Mayall, M. Stern, F.M. Munkonge, A. Frost, L. Huang, M. Vasiliou, R. Williamson, E.W. Alton und C. Coutelle. The introduction of two silent mutations into a CFTR cDNA construct allows improved detection of exogenous mRNA in gene transfer experiments. *Hum Mol Gen*, 4(9):1597–602, 1995.

- [66] R. P. Haugland. *Handbook of Fluorescent Probes and Research Chemicals.*, Nucleic acid stains.S. 144–156. M.T.Z. Spence. Molecular Probes, Inc.1996.
- [67] P. Hawley-Nelson, V. Cicarrone, G. Gebeyehu und J. Jessee. LipofectAMINE reagent: a new, higher efficiency polycationic liposome transfection reagent. *Focus*, 15:73–79, 1993.
- [68] T. Heath. Covalent attachment of proteins to liposomes. *Meth Enzymol*, 149:111–9, 1987.
- [69] D. Hirsch-Lerner und Y. Barenholz. Probing DNA-cationic lipid interactions with the fluorophore trimethylammonium diphenyl-hexatriene (TMADPH). *Biochim Biophys Acta*, 1370(1):17–30, 1998.
- [70] R.M. Hoffmann, L.B. Margolis und L.D. Bergelson. Binding and Entrapment of high molecular weight DNA by lecithin liposomes. *FEBS Lett*, 93:229–232, 1978.
- [71] K. Hong, W. Zheng, A. Baker und D. Papahadjopoulos. Stabilization of cationic liposome-plasmid DNA complexes by polyamines and polyethyleneglycol-phospholipid conjugates for efficient in vivo gene delivery. *FEBS Lett*, 400(2):233–7, 1997.
- [72] L . Huang, A.L. Klibanov, A. Mori und X. Gao. Targeted delivery of drugs and DNA with liposomes. *J Lipos Res*, 3:505–515, 1993.
- [73] S.W. Hui, M. Langner, Y.L. Zhao, P. Ross, E. Hurley und K. Chan. The role of helper lipids in cationic liposome-mediated gene transfer. *Biophys J*, 71(2):590–9, 1996.
- [74] S.E. Jones, J.D. McHugh, C. Jomary, A. Shallal und M.J. Neal. Assessment of liposomal transfection of ocular tissues in vivo. *Gene Ther*, 1 Suppl 1:S61, 1994.
- [75] Y. Kaneda, K. Iwai und T. Uchida. Introduction and expression of the human insulin gene in adult rat liver. *J Biol Chem*, 264:12126–9, 1989.
- [76] Y. Kaneda, K. Kato, M. Nakanishi und T. Uchida. Introduction of plasmid DNA and nuclear protein into cells by using erythrocyte ghosts, liposomes and Sendai virus. *Meth Enzymol*, 221:317–27, 1993.
- [77] C. Kawaura, A. Noguchi, T. Furuno und M. Nakanashi. Atomic force microscopy for studying gene transfection mediated by cationic liposomes with a cationic cholesterol derivative. *FEBS Lett*, 421:69–72, 1998.
- [78] A. Kichler, W. Zauner, M. Ogris und E. Wagner. Influence of the DNA complexation medium on the transfection efficiency of lipospermine/DNA particles. *Gene Ther*, 5:855–860, 1998.
- [79] R. Kircheis, A. Kichler, G. Wallner, M. Kursa, M. Ogris, T. Felzmann, M. Buchberger und E. Wagner. Coupling of cell-binding ligands to polyethylenimine for targeted gene delivery. *Gene Ther*, 4:409–18, 1997.

- [80] L. Ko, A. Koestner und W. Wechsler. Characterisation of cell cycle and biological parameters of transplantation glioma cell clones. *Acta Neuropathol*, 51:107–111, 1980.
- [81] R. Kraayenhof, G.J. Sterk und H.W. Sang. Probing biomembrane interfacial potential and pH profiles with a new type of float-like fluorophores positioned at varying distance from the membrane surface. *Biochemistry*, 32(38):10057–66, 1993.
- [82] E. Krauss, D. Groth, S. Armeanu, R. Curth, J. Pelisek, O. Keil, B. Höfling, R. Reszka und S. Nikol. Optimization of non-viral gene transfer into vascular smooth muscle cells in vitro. *in preparation*, 1999.
- [83] J.F. Kukowska-Latallo, A.U. Bielinska, J. Johnson, R. Spindler, D.A. Tomalia, und J.R. Baker. Efficient transfer of genetic material into mammalian cells using starburst polyamidoamine dendrimers. *Proc Natl Acad Sci USA*, 93:4897–4902, 1996.
- [84] F. Labat-Moleur, A.M. Steffan, C. Brisson, H. Perron, O. Feugeas, P. Furstenberger, F. Oberling, E. Brambilla und J.P. Behr. An electron microscopy study into the mechanism of gene transfer with lipopolyamines. *Gene Ther*, 3(11):1010–7, 1996.
- [85] R.A. Lake und M.J. Owen. *Gene Transfer and Expression Protocols.*, Transfection of the Chloramphenicol-Acetyltransferase Gene into Eukaryotic cells using Diethyl-Aminoethyl (DEAE)-Dextran., Ed. Murray EJ. The Humana Press Inc. Clifton, NY, USA, 1991.
- [86] I. Lang, M. Scholz und R. Petrs. Molecular mobility and nucleocytoplasmic flux in hepatoma cells. *J Cell Biol*, 102:1183–1190, 1986.
- [87] D.D. Lasic. *Liposomes: from physics to applications*. Elsevier Science Publishers B.V. Amsterdam, 1993.
- [88] D.D. Lasic. *Liposomes: from physics to applications.*, Chemical stability of lipids: Oxidation.S. 35–37. Elsevier Science Publishers B.V. Amsterdam, 1993.
- [89] R.J. Lee und L. Huang. Folate-targeted, anionic liposome-entrapped polylysine-condensed DNA for tumor cell-specific gene transfer. *J Biol Chem*, 271(14):8481–7, 1996.
- [90] R.J. Lee und L. Huang. Lipidic vector systems for gene transfer. *Crit Rev Ther Drug Carr System*, 14(2):173–206, 1997.
- [91] C. Lehmann. Elektronenmikroskopische Untersuchungen von DOCSPER/DNA-Lipoplexen bei verschiedenen NaCl-Konzentrationen. Persönliche Mitteilung.
- [92] C. Lehmann. Untersuchungen zum Einfluß des Helferlipides auf die Liposomenformulierung mittels der Elektronenmikroskopie. Persönliche Mitteilung.

- [93] R. Leventis und J.R. Silvius. Interactions of mammalian cells with lipids dispersions containing novel metabolizable cationic amphiphiles. *Biochim Biophys Acta*, 1023:124–132, 1990.
- [94] S. Li und L. Huang. In vivo gene transfer via intravenous administration of cationic lipid-protamine-DNA (LPD) complexes. *Gene Ther*, 4(9):891–900, 1997.
- [95] S. Li, M.A. Rizzo, S. Bhattacharya und L. Huang. Characterization of cationic lipid-protamine-DNA (LPD) complexes for intravenous gene delivery. *Gene Ther*, 5(7):930–7, 1998.
- [96] W.W. Liang, X. Shi, D. Deshpande, C.J. Malanga und Y. Rojanasakul. Oligonucleotide targeting to alveolar macrophages by mannose receptor-mediated endocytosis. *Biochim Biophys Acta*, 1279:227–234, 1996.
- [97] J.P. Loeffler und J.P. Behr. Gene transfer into primary and established mammalian cell lines with lipopolyamine-coated DNA. *Meth Enzymol*, 217:599–618, 1993.
- [98] C.B. Lozzio und B.B. Lozzio. Cytotoxicity of a factor isolated from human spleen. *J Natl Cancer Inst*, 50:535–8, 1973.
- [99] R.H. Lyons, B.Q. Ferguson und M. Rosenberg. Pentapeptide nuclear localization signal in adenovirus E1a. *Mol Cell Biol*, 7:2451–2456, 1987.
- [100] R.W. Malone, P.L. Felgner und I. Verma. Cationic liposome-mediated RNA transfection. *Proc Natl Acad Sci USA*, 86:6077–6081, 1989.
- [101] R.J. Mannino und S. Gould-Fogerite. Liposome mediated gene transfer. *Bio-Techniques*, 6:682–690, 1988.
- [102] M. Marquet, N.A. Horn und J.A. Meek. Process development for the manufacture of plasmid DNA vectors for use in gene therapy. *Pharm*, 4:28–46, 1996.
- [103] F. Martin und T. Boulikas. The challenge of liposomes in gene therapy. *Gene Ther Mol Biol*, 1:173–214, 1998.
- [104] H. Matsui, L.G. Johnson, S.H. Randell und R.C. Boucher. Loss of binding and entry of liposome-DNA complexes decreases transfection efficiency in differentiated airway epithelial cells. *J Biol Chem*, 272(2):1117–26, 1997.
- [105] L.D. Mayer und L.C.L. Tai ... *Cancer Res*, 49:5922–5930, 1989.
- [106] R.A. McMasters, R.L. Sailors, K.E. Jones, M.E. Hendrix, M.P. Moyer und R.R. Drake. Lack of Bystander Killing in Herpes Simplex Virus Thymidine Kinase-Transduced Colon Cell Lines Due to Deficient Connexin43 Gap Junction Formation. *Hum Gene Ther*, 9:2253–61, 1998.
- [107] C. Meunier-Durmort, R. Picart, T. Ragot, M. Perricaudet, B. Hainque und C. Forest. Mechanism of adenovirus improvement of cationic liposome-mediated gene transfer. *Biochim Biophys Acta*, 1330(1):8–16, 1997.

- [108] O. Meyer, D. Kirpotin, K. Hong, B. Sternberg, J.W. Park, M.C. Woodle und D. Papahadjopoulos. Cationic liposomes coated with polyethylene glycol as carriers for oligonucleotides. *J Biol Chem*, 273(25):15621–7, 1998.
- [109] K.A. Mislick und J.D. Baldeschwieder. Evidence for a role of proteoglycans in cation-mediated gene transfer. *Proc Natl Acad Sci USA*, 93:12349–12354, 1996.
- [110] W.A. Mohler, C.A. Charlon und H.M. Blau. Spectrophotometric quantitation of tissue culture cell number in any medium. *BioTechniques*, 21:260–266, 1996.
- [111] R.H. Mueller. *Zetapotential und Partikelladung in der Laborpraxis. Rezepturen Für Pufferlösungen*. Wiss. Verlagsgesellschaft mbH Stuttgart.1996.
- [112] R.R.C. New. Liposomes a Practical Approach. Introduction. *Practical Approach Series*, S. 1–31, 1990.
- [113] C. Nicolau, A. Le-Pape, P. Soriano, F. Fargette und M. F. Juhel. In vivo expression of rat insulin after intravenous administration of the liposome-entrapped gene for rat insulin I. *Proc Natl Acad Sci USA*, 80(4):1068–1072, 1983.
- [114] B.S. Oler und E. Schenborn. Transfast Transfection Reagent Update. *Promega Notes*, 71:18–21, 1999.
- [115] A. Vaheri J.S. Pagano. *Virology*, 27:434–436, 1965.
- [116] R. Pal, W.A. Petri Jr, V. Ben-Yashar, R.R. Wagner und Y. Barenholz. Characterization of the fluorophore 4-heptadecyl-7-hydroxycoumarin: a probe for the head-group region of lipid bilayers and biological membranes. *Biochemistry*, 24:573–81, 1985.
- [117] B. Pitard, O. Aguerre, M. Airiau, A.M. Lachages, T. Boukhnikachvili, G. Byk, C. Dubertret, C. Herviou, D. Scherman, J.F. Mayaux und J. Crouzet. Virus-sized self-assembling lamellar complexes between plasmid DNA and cationic micelles promote gene transfer. *Proc Natl Acad Sci USA*, 94:14412–7, 1997.
- [118] H. Potter. Application of Electroporation in Recombinant DNA Technology. *Meth Enzymol*, 217:561–583, 1993.
- [119] C. Qiu, Young M.B. De, A. Finn und D.A. Dichek. Cationic liposomes enhance adenovirus entry via a pathway independent of the fiber receptor and alpha(v)-integrins. *Hum Gene Ther*, 9(4):507–20, 1998.
- [120] J.O. Raedler, I. Koltover, T. Salditt und Safinya. Structure of DNA-cationic liposome complexes: DNA intercalation in multilamellar membranes in distinct interhelical packing regimes. *Science*, 275:810–814, 1997.
- [121] J.S. Remy, A. Kichler, V. Mordvinov, F. Schuber und J.P. Behr. Targeted gene transfer into hepatoma cells with lipopolyamine-condensed DNA particles presenting galactose ligands: a stage toward artificial viruses. *Proc Natl Acad Sci USA* , 92(5):1744–8, 1995.

- [122] J.S. Remy, C. Sirlin, P. Vierling und J.P. Behr. Gene transfer with a series of lipophilic DNA-binding molecules. *Bioconj Chem*, 5(6):647–54, 1994.
- [123] R. Reszka, J. Zhu, F. Weber und W. Walther. Liposome mediated transfer of marker and cytokine genes into rat and human glioblastoma cells in vitro and in vivo. *J Liposom Res*, 5:149–167, 1995.
- [124] A.P. Rolland. From genes to gene medicines: Recent advances in nonviral gene delivery. *Crit Rev Ther Drug Carr System*, 15:143–198, 1998.
- [125] J.K. Rose, L. Buonocore und M.A. Whitt. A new cationic liposome reagent mediating nearly quantitative transfection of animal cells. *BioTechniques*, 4:520–525, 1991.
- [126] Y. Saeki, N. Matsumoto, Y. Nakano, M. Mori, K. Awai und Y. Kaneda. Development and characterization of cationic liposomes conjugated with HVJ (Sendai virus): reciprocal effect of cationic lipid for in vitro and in vivo gene transfer. *Hum Gene Ther*, 8(17):2133–41, 1997.
- [127] W.H. Gunzburg B. Salmons. *Gentransfer in Säugetierzellen*. Labor im Fokus. Spektrum Akademischer Verlag, Berlin, Heidelberg., 1997.
- [128] A. Schadendorf. Humane Melanomzelle Mewo. Persönliche Mitteilung.
- [129] M. Schaefer-Ridder, Y. Wang und P.H. Hofschneider. Liposomes as gene carriers: efficient transformation of mouse cells by thymidine kinase gene. *Science*, 215:166–168, 1982.
- [130] K.P. Schifferli und V. Ciccarone. Optimization of cationic lipid reagent-mediated transfection for suspension cell lines. *Focus*, 18:45–47, 1996.
- [131] M. Schneider, R. Reszka, D. Groth und O. Keil. Neuartige kationische Amphiphile für den liposomalen Gentransfer. *Deut Pat Applik 196.31.189.6.*, 1996.
- [132] D. Schutt, Schuhmann und R.H. Muller. Genauer durch Mehrwinkelanalyse. Teilchengrößenmessung im Submikronbereich. *Chemie-Technik*, S. 18–20, 1996.
- [133] B. Schwartz, C. Benoit, B. Abdallah, D. Sherman, J.P. Behr und B.A. Demeneix. Lipospermine-based gene transfer into the newborn mouse brain is optimized by a low lipospermine/DNA charge ratio. *Hum Gene Ther*, 6(12):1515–24, 1995.
- [134] M. Sioud, J.B. Natvig und O. Forre. Preformed ribozyme destroys tumour necrosis factor mRNA in human cells. *J Mol Biol*, 223:831–5, 1992.
- [135] S.J. Slater, M.B. Kelly, F.J. Taddeo, C. Ho, E. Rubin und C.D. Stubbs. The modulation of protein kinase C activity by membrane lipid bilayer structure. *J Biol Chem*, 269:4866–71, 1994.
- [136] J.G. Smith, R.L. Walzem und J.B. German. Liposomes as agents of DNA transfer. *Biochim Biophys Acta*, 1154:327–40, 1993.

- [137] J.G. Smith, T. Wedeking, J.H. Vernachio, H. Way und R.W. Niven. Characterization and in vivo testing of a heterogeneous cationic lipid-DNA formulation. *Pharm Res*, 15(9):1356–63, 1998.
- [138] F.L. Sorgi, S. Bhattacharya und L. Huang. Protamine sulfate enhances lipid-mediated gene transfer. *Gene Ther*, 4(9):961–8, 1997.
- [139] H.D. Soule, J. Vazquez, A. Long, S. Albert und M. Brennan. A human cell line from a pleural effusion derived from a breast carcinoma. *J Natl Cancer Inst.*, 51:1409–16, 1973.
- [140] J.C. Stavridis, G. Deliconstantinos, M.C. Psallidopoulos, N.A. Armenakas, D.J. Hadjiminas und J. Hadjiminas. Construction of transferrin-coated liposomes for in vivo transport of exogenous DNA to bone marrow erythroblasts in rabbits. *Exp Cell Res*, 164:568–72, 1986.
- [141] T. Stegmann und J.Y. Legendre. Gene transfer mediated by cationic lipids: lack of a correlation between lipid mixing and transfection. *Biochim Biophys Acta*, 1325(1):71–9, 1997.
- [142] B. Sternberg, F.L. Sorgi und L. Huang. New structures in complex formation between DNA and cationic liposomes visualized by freeze-fracture electron microscopy. *FEBS Lett*, 356:361–366, 1994.
- [143] W.P. Swaney, F.L. Sorgi, A.B. Bahnsen und J.A. Barranger. The effect of cationic liposome pretreatment and centrifugation on retrovirus-mediated gene transfer. *Gene Ther*, 4(12):1379–86, 1997.
- [144] K. Takeuchi, M. Ishihara, C. Kawaura, M. Noji, T. Furuno und M. Nakanishi. Effect of zeta potential of cationic liposomes containing cationic cholesterol derivatives on gene transfection. *FEBS Lett*, 397(2-3):207–9, 1996.
- [145] M.X. Tang, C.T. Redemann und F.C. Szoka FC Jr. In vitro gene delivery by degrades polyamidoamine dendrimers. *Bioconj Chem*, 7:703–714, 1996.
- [146] N.R. Taylor, B.E. Kaplan, P. Swiderski, H. Li und J. Rossi. Chimeric DNA-RNA hammerhead ribozymes have enhanced in vitro catalytic efficiency and increased stability in vivo. *Nucleic Acids Res*, 20:4559–65, 1992.
- [147] C.B. Thomas, R. Osieka und K.W. Kohn. DNA cross-linking by in vivo treatment with 1-(2-chloroethyl)-3-(4-methylcyclohexyl)-1-nitrosourea of sensitive and resistant human colon carcinoma xenograms in nude mice. *Cancer Res*, 38:2448–54, 1978.
- [148] J. Treat, A. Greenspan, D. Forst, J.A. Sanchez, V.J. Ferrans, L.A. Potkul, P.V. Woolley und A. Rahman. Antitumor activity of liposome encapsulated doxorubicin in advanced breast cancer: phase II study. *J Natl Cancer Inst*, 82:1706–1710, 1990.
- [149] A. Vainstain, A. Razin, A. Graessmann und A. Loyter. Fusogenic reconstituted Sendai virus envelopes as a vehicle for introducing DNA into viable mammalian cells. *Meth Enzymol*, 101:492–512, 1983.

- [150] I.M. Verma und N. Somia. Gene Therapy - Promises, Problems and Prospects. *Nature*, 389:239–42, 1997.
- [151] R.G. Vile, J.A. Nelson, S. Castleden, H. Chong und I.R. Hart. Systemic gene therapy of murine melanoma using tissue specific expression of the HSVtk gene involves an immune component. *Cancer Res*, 54(23):6228–34, 1994.
- [152] L. Vitiello, A. Chonn, J.D. Wasserman, C. Duff und R.G. Worton. Condensation of plasmid DNA with polylysine improves liposome-mediated gene transfer into established and primary muscle cells. *Gene Ther*, 3(5):396–404, 1996.
- [153] E. Wagner, C. Plank, K. Zatloukal, M. Cotten und M.L. Birnstiel. Influenza virus hemagglutinin HA-2 N-terminal fusogenic peptides augment gene transfer by transferrin-polylysine-DNA complexes: toward a synthetic virus-like gene-transfer vehicle. *Proc Natl Acad Sci USA*, 89(17):7934–8, 1992.
- [154] J. Wagner, H. Madry und R. Reszka. In Vivo Gene Transfer: Focus on the Kidney. *Nephrol Dial Transplant*, S. 1801–1807, 1995.
- [155] R. Wattiaux, M. Jadot, M.T. Warnier-Pirotte und S. Wattiaux de Coninck. Cationic lipids destabilize lysosomal membrane in vitro. *FEBS Lett*, 417:99–202, 1997.
- [156] M. Westphal, M. Haesel, R. Kunzmann und F. Hoelzel. Karyotype analysis of 20 human glioma cell lines. *Acta Neurochir*, 126:17–26, 1993.
- [157] C.J. Wheeler, L. Sukhu, G. Yang, Y. Tsai, C. Bustamente, P. Felgner, J. Norman, und M. Manthorpe. Converting an alcohol to an amine in a cationic lipid dramatically alters the co-lipid requirement, cellular transfection activity and the ultrastructure of DNA-cytosin complexes. *Biochim Biophys Acta*, 1280(1):1–11, 1996.
- [158] R. Widmaier, G.P. Wildner, G. Papsdorf und I. Graffi. A new infinite cell line, MaTu, of human mammary tumor-cells. *Arch Geschwulstforsch*, 44:1–10, 1974.
- [159] S. Wirsel. *Gentechnische Methoden*, Ausbeutebestimmung und analytische Überprüfung der DNA, S. 118–119. Gustav Fischer Verlag Stuttgart, S. Bertam und H. G. Gassen (Hrsg), 1991.
- [160] N. Wivel. Human gene transfer trials. *Adv Drug Del Rev*, 17:211–212, 1995.
- [161] J.A. Wolff, R.W. Malone, P. Williams, W. Chong, G. Ascadi, A. Jani und P.L. Felgner. Direct gene transfer into mouse muscle in vivo. *Science*, 247:1465–1468, 1990.
- [162] Y. Xu und F.C. Szoka Jr. Mechanism of DNA release from cationic liposome/DNA complexes used in cell transfection. *Biochemistry*, 35:5616–5623.1996.
- [163] J.P. Yang und L. Huang. Time-dependent maturation of cationic liposome-DNA complex for serum resistance. *Gene Ther*, 5(3):380–7, 1998.

- [164] K. Yang, G.L. Clifton und R.L. Hayes. Gene therapy for central nervous system injury: the use of cationic liposomes: an invited review. *J Neurotrauma*, 14(5):281–97, 1997.
- [165] N.S. Yang, J. Burkholder, B. Roberts, B. Martinell und D. McCabe. In vivo and in vitro gene transfer of mammalian somatic cells by particle bombardement. *Proc Natl Acad Sci USA*, 87:9568–9572, 1990.
- [166] Y. Yoshikawa, N. Emi, T. Kanbe, K. Yoshikawa und H. Saito. Folding and aggregation of DNA chains induced by complexation with lipospermine formation of a nucleosome-like structure and network assembly. *FEBS Lett*, 397:207–209.1996.
- [167] J. Zabner, A.J. Fasbender, T. Moninger, K. Poellinger und M.J. Welsh. Cellular and molecular barriers to gene transfer by a cationic lipid. *J Biol Chem*, 270:18997–19007, 1995.
- [168] O. Zelphati und F.C. Szoka Jr. Intracellular distribution and mechanism of delivery of oligonucleotides mediated by cationic lipids. *Pharm Res*, 13:367–1372, 1997.
- [169] O. Zelphati, C. Nguyen, M. Ferrari, J. Felgner, Y. Tsai und P.L. Felgner. Stable and monodisperse lipoplex formulations for gene delivery. *Gene Ther*, 5(9):1272–82, 1998.
- [170] X. Zhou und L. Huang. DNA transfection mediated by cationic liposomes containing lipopolylysine: characterization and mechanism of action. *Biochim Biophys Acta.*, 1189(2):195–203, 1994.
- [171] J. Zhu, L. Zhang und R. Reszka. Zinc finger motif nucleocapsid protein-facilitated gene transfer with lipoplexes and polyplexes. *submitted*, 1999.
- [172] N.J. Zuidam und Y. Barenholz. Electrostatic parameters of cationic liposomes commonly used for gene delivery as determined by 4-heptadecyl-7-hydroxycoumarin [published erratum appears in Biochem Biophys Acta 1998 Jun 24;1372(1):151]. *Biochim Biophys Acta*, 1329(2):211–22, 1997.