

7. Literaturverzeichnis

- AAEM:** American Association of Electrodiagnostic Medicine (AAEM) Somatosensory Evoked Potentials Subcommittee. Guidelines for somatosensory evoked potentials. Muscle Nerve 1999; 22: S123-S138
- AAN:** Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. Assessment: Neuropsychological testing of adults. Neurology 1996; 47: 592-599
- Abbruzzese** G, Dall'Agata D, Spadavecchia L, Abbruzzese M, Favale E. Parietal and prerolandic somatosensory evoked responses in carotid artery reversible ischemic attacks (RIA). Acta Neurol Scand 1987; 76: 480-485
- Abbruzzese** G, Reni L, Cocito L, Ratto S, Abbruzzese M, Favale E. Short-latency somatosensory evoked potentials in degenerative and vascular dementia. J Neurol Neurosurg Psychiatr 1984; 47: 1034-1037
- ACAS:** Toole JF, and the Executive Committee for the Asymptomatic Carotid Atherosclerosis Study. Endarterectomy for asymptomatic carotid artery stenosis. JAMA 1995; 273: 1421-1428
- ACST:** Halliday A, Mansfield A, Marro J, et al. Prevention of disabling and fatal strokes by successful carotid endarterectomy in patients without recent neurological symptoms: randomised controlled trial. Lancet 2004; 363: 1491-1502
- Allison** T, Mc Carthy G, Wood CC, Jones SJ. Potentials evoked in human and monkey cerebral cortex by stimulation of the median nerve. A review of scalp and intracranial recordings. Brain 1991; 114: 2465-2503
- Alvarez** JA, Emory E. Executive function and the frontal lobes: a meta-analytic review. Neuropsych Rev 2006; 16: 17-42
- Ancelin** M-L, De Roquetaeil G, Ledésert B, Bonnel F, Cheminal J-C, Ritchie K. Exposure to anaesthetic agents, cognitive functioning and depressive symptomatology in the elderly. Br J Psychiatry 2001; 178: 360-366
- Arning** C, Hammer E, Kortmann H et al. Quantifizierung von A. carotis interna-Stenosen: Welche Ultraschallkriterien sind geeignet? Ultraschall Med 2003; 24: 233-238
- Asken** MJ, Hobson RW. Intellectual change and carotid endarterectomy, subjective speculation or objective reality: a review. J Surg Res 1977; 23: 367-375
- Astarci** P, Guerit JM, Robert A, et al. Stump pressure and somatosensory evoked potentials for predicting the use of shunt during carotid surgery. Ann Vasc Surg 2007; 21: 312-317
- Astrup** J, Symon L, Branston NM, et al. Cortical evoked potential and extracellular K⁺ and H⁺ at critical levels of brain ischemia. Stroke 1977, 8: 51-57
- Auperin** A, Berr C, Bonithon-Kopp C, et al. Ultrasonographic assessment of carotid wall characteristics and cognitive functions in a community sample of 59 – 71-year-olds. Stroke 1996; 27: 1290-1295
- Bakker** FC, Klijn CJM, Jennekens-Schinkel A, et al. Cognitive disorders in patients with occlusive disease of the carotid artery: a systematic review of the literature. J Neurol 2000; 247: 669-676
- Bakker** FC, Klijn CJM, Jennekens-Schinkel A. Cognitive impairment in patients with carotid artery occlusion and ipsilateral transient ischemic attacks. J Neurol 2003; 250: 1340-1347
- Bakker** FC, Klijn CJM, van der Grond J, et al. Cognition and quality of life in patients with carotid artery occlusion. Neurology 2004; 62: 2230-2235
- Banoub** M, Tetzlaff JE, Schubert A. Pharmacologic and physiologic influences affecting sensory evoked potentials: implications for perioperative monitoring. Anesthesiology 2003; 99: 716-737
- Bardenheuer** HJ, Weigand MA. Nicht invasives und invasives Monitoring der zerebralen Perfusion. Anästhsiol Intensivmed Notfallmed Schmerzther 1997; 32(Suppl 2): 201-210
- Bender** R, Lange S, Ziegler A. Multiples Testen. Dtsch Med Wochenschr 2002; 127: T4-T7

- Benjamin** ME, Silva MB, Watt C, McCaffrey MT, Burford-Foggs A, Flinn WR. Awake patient monitoring to determine the need for shunting during carotid endarterectomy. *Surgery* 1993; 114: 673-681
- Benton** AL, Hamsher K, Sivan AB. Multilingual aphasia examination. 3rd ed., 1983, AJA Associates, Iowa.
- Berlit** P. Schlaganfall. Möglichkeiten der Primärprävention. *Nervenarzt* 2000; 71: 231-237
- Berman** L, Pietrzak RH, Mayes L. Neurocognitive changes after carotid revascularization: a review of the current literature. *J Psych Res* 2007; 63: 599-612
- Biller** J, Feinberg WM, Castaldo JE et al. Guidelines for carotid endarterectomy. A statement for healthcare professionals from a special writing group of the Stroke Council, American Heart Association. *Stroke* 1998; 29: 554-562
- Blume** WT, Ferguson GG, McNeill DK. Significance of EEG changes at carotid endarterectomy. *Stroke* 1986; 17: 891-897
- Bo** M, Massaia M, Speme S, et al. Risk of cognitive decline in older patients after carotid endarterectomy: an observational study. *J Am Geriatr Soc* 2006; 54: 932-936
- Boeke** S. The effect of carotid endarterectomy on mental functioning. *Clin Neurol Neurosurg* 1981; 83: 209-217
- Boisseau** N, Madany M, Staccini P, et al. Comparison of the effects of sevoflurane and propofol on cortical somatosensory evoked potentials. *Br J Anaesth* 2002; 88: 785-789
- Bond** R, Rerkasem K, Rothwell PM. Routine or selective carotid artery shunting for carotid endarterectomy (and different methods of monitoring in selective shunting). *Stroke* 2003a; 34: 824-825
- Bond** R, Rerkasem K, Rothwell PM. Systematic review of the risks of carotid endarterectomy in relation to the clinical indication for and timing of surgery. *Stroke* 2003b; 34: 2290-2303
- Borkowski** JG, Benton AL, Spreen O. Word fluency and brain damage. *Neuropsychologia* 1967; 5: 135-140
- Bornstein** RA, Benoit BG, Trites RL. Neuropsychological changes following carotid endarterectomy. *Can J Neurol Sci* 1981; 8: 127-132
- Borroni** B, Tiberio G, Bonardelli S, et al. Is mild vascular cognitive impairment reversible? Evidence from a study on the effect of carotid endarterectomy. *Neurol Res* 2004; 26: 594-597
- Bossema** ER, Brand AN, Geenen R, et al. Effect of carotid endarterectomy on patient evaluations of cognitive functioning and mental and physical health. *Ann Vasc Surg* 2005b; 19: 673-677
- Bossema** ER, Brand N, Moll FL et al. Cognitive functions in carotid artery disease before endarterectomy. *J Clin Exp Neuropsych* 2006; 28: 357-369
- Bossema** ER, Brand N, Moll FL, Ackerstaff RGA, van Doornen LJP. Does carotid endarterectomy improve cognitive functioning? *J Vasc Surg* 2005a; 41: 775-781
- Boysen** G, Engell HC, Henriksen H. The effect of induced hypertension on internal carotid artery pressure and regional cerebral blood flow during temporary carotid clamping for carotid endarterectomy. *Neurology* 1972; 22: 1133-1144
- Brand** N, Bossema ER, Ommen M, Moll FL, Ackerstaff RG. Left or right carotid endarterectomy in patients with atherosclerotic disease: ipsilateral effects on cognition? *Brain Cogn* 2004; 54: 117-123
- Branston** NM, Symon L, Crockard HA, et al. Relationship between the cortical evoked potential and local cortical blood flow following acute middle cerebral artery occlusion in the baboon. *Exp Neurol* 1974; 45: 195-208
- Brickenkamp** R. Test d2 : Aufmerksamkeits-Belastungs-Test. 1994, Hogrefe-Verlag für Psychologie, Göttingen.
- Brinkman** SD, Braun P, Ganji S, Morrell RM, Jacobs LA. Neuropsychological performance one week after carotid endarterectomy reflects intraoperative ischemia. *Stroke* 1984; 15: 497-503

- Bryson** GL, Wyand A. Evidence-based clinical update: General anesthesia and the risk of delirium and postoperative cognitive dysfunction. *Can J Anesth* 2006; 53: 669-677
- Canet** J, Raeder J, Rasmussen LS, et al. Cognitive dysfunction after minor surgery in the elderly. *Acta Anaesthesiol Scand* 2003; 47: 1204-1210
- Cao** P, De Rango P, Zannetti S. Eversion vs conventional carotid endarterectomy : a systematic review. *Eur J Vasc Endovasc Surg* 2002; 23: 195-201
- Carenini** L, Bottacchi E, Camerlingo M, et al. Considerations after intraoperative monitoring of somatosensory evoked potentials during carotid endarterectomy. *Ital J Neurol Sci* 1989; 10: 315-320
- Caza** N, Taha R, Qi Y, Blaise G. The effects of surgery and anesthesia on memory and cognition. *Progr Brain Res* 2008; 169: 409-422
- Cerhan** JR, Folsom AR, Mortimer JA, et al. Correlates of cognitive function in middle-aged adults. *Gerontology* 1998; 44: 95-105
- Chaturvedi** S, Bruno A, Feasby T, et al. Carotid endarterectomy – an evidence-based review. Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology* 2005; 65: 794-801
- Chaytor** N, Schmitter-Edgecombe M. The ecological validity of neuropsychological test: a review of the literature on everyday cognitive skills. *Neuropsych Rev* 2003; 13: 181-197
- Cho** H, Nemoto EM, Yonas H, et al. Cerebral monitoring by means of oxymetry and somatosensory evoked potentials during carotid endarterectomy. *J Neurosurg* 1998; 89 : 533-538
- Cinà** CS, Safar HA, Magissano R, Bailey R, Clase CM. Prevalence of significant carotid artery stenosis in patients with peripheral arterial occlusive disease. *J Vasc Surg* 2002; 36: 75-82
- Connolly** ES, Winfree CJ, Rampersad A, et al. Serum S100B levels are correlated with subclinical neurocognitive declines after carotid endarterectomy. *Neurosurgery* 2001; 49: 1076-1083
- Coulthard** E, Singh-Curry V, Husain M. Treatment of attention deficits in neurological disorders. *Curr Opin Neurol* 2006; 19: 613-618
- Crucu** G, Aminoff MJ, Curio G, et al. Recommendations for the clinical use of somatosensory-evoked potentials. *Clin Neurophysiol* 2008; 119: 1705-1719
- Cummings** JL. Anatomic and behavioural aspects of frontal-subcortical circuits. *Ann NY Acad Sci* 1995; 769: 1-13
- Cushman** L, Brinkman SD, Ganji S, Jacobs LA. Neuropsychological impairment after carotid endarterectomy correlates with intraoperative ischemia. *Cortex* 1984; 20: 403-412
- D'Addato** M, Pedrini L, Vitacchiano G. Intraoperative cerebral monitoring in carotid surgery. *Eur J Vasc Surg* 1993; 7 (Suppl A): 16-20
- Dahlöf** B, Devereux RB, Kjeldsen S, et al. Cardiovascular morbidity and mortality in the Losartan Intervention For Endpoint reduction in hypertension study (LIFE) : a randomised trial against atenolol. *Lancet* 2002; 359: 995-1003.
- de Haan** EH, Nys GM, van Zandvoort MJ. Cognitive function following stroke and vascular cognitive impairment. *Curr Opin Neurol* 2006; 19: 559-564
- De Rango** P, Caso V, Didier L, Paciaroni M, Lenti M, Cao P. The role of carotid artery stenting and carotid endarterectomy in cognitive performance. A systematic review. *Stroke* 2008; 39: 31116-3127
- De Vleeschauwer** P, Haupt WF, Heinrichs W, et al. Anwendung somatosensibel evoziert Potentiale in der Carotischirurgie. *Angio* 1985b; 7: 203-209
- De Vleeschauwer** P, Horsch S, Matamoros R. Monitoring of somatosensory evoked potentials in carotid surgery: results, usefulness and limitations of the method. *Ann Vasc Surg* 1988; 2: 63-68
- De Weerd** AW, Looijenga A, Veldhuizen RJ, Van Huffelen AC. Somatosensory evoked potentials in minor cerebral ischaemia: diagnostic significance and changes in serial records. *Electroencephalogr Clin Neurophysiol* 1985; 62: 45-55
- Debus** ES, Larena A, Wintzer C. Offene Therapie der Karotisstenose durch Endarterektomie. *Chirurg* 2004; 75: 658-666

- Delany** RC, Wallace JD, Egelko S. Transient cerebral ischemic attacks and neuropsychological deficit. *J Clin Exp Neuropsych* 1980; 2: 107-114
- DeLisa** J, Meckenzie K, Baran E. Manual of nerve conduction velocity and somatosensory evoked potentials. 2nd ed. New York, Raven 1987, 190-200
- Detsch** O, Kochs E. Perioperatives Neuromonitoring. *Anaesthetist* 1997; 46: 999-1014
- DGG: Deutsche Gesellschaft für Gefäßchirurgie** (Hrsg.). Leitlinie zu Stenosen der Arteria carotis. Gefässchirurgie 2003; 8: 25-28
- Di Legge** S, Di Piero V, Di Stani F, et al. Carotid endarterectomy and glial fibrillar S100b protein release. *Neurol Sci* 2003; 24: 351-356
- Dinkel** M, Kamp H-D, Schweiger H. Somatosensorisch evozierte Potentiale in der Karotischirurgie. *Anaesthetist* 1991; 40: 72-78
- Dinkel** M, Langer H, Schweiger H, Rügheimer E. Risikominimierung in der Gefäßchirurgie durch zentralnervöses Ischämemonitoring. *Anästh Intensivmed* 1994a; 35: 365-370
- Dinkel** M. In Rügheimer E, Dinkel M (Hrsg): Klinische Anästhesiologie und Intensivtherapie – Neuromonitoring in Anästhesie und Intensivmedizin, 1994b; Vol 46, 1. Auflage, Springer, Berlin, Heidelberg, New York
- Drummond** JC, Shapiro HM. Cerebral Physiology. In: Miller RD. Anesthesia. 4. Auflage. New York, Edinburgh, Melbourne: Churchill Livingstone, 1994: 689-729
- Duus** P. Blutversorgung des Gehirns. In: Duus P. Neurologisch-topische Diagnostik: Anatomie, Physiologie, Klinik. 5. Auflage. Stuttgart, New York: Thieme, 1990: 411-422
- Eckstein** HH, Allenberg JR. Einfluß evidenzbasierter Indikationen zur Karotis-TEA auf die Stadieneinteilung extrakranieller Karotisstenosen – Notwendigkeit einer aktualisierten Klassifikation. *Gefäßchirurgie* 2001; 6: 91 –97
- Eckstein** HH, Maeder N, Allenberg JR. Carotischirurgie als Apoplexprophylaxe. *Chirurg* 1999; 70: 353-363
- ECST** : The European Carotid Surgery Trialists' Collaborative Group. Randomised trial of endarterectomy for recently symptomatic carotid stenosis: final results of the MRC European Carotid Surgery Trial (ECST). *Lancet* 1998; 351: 1379-1387
- Ederle** J, Brown MM. The evidence for medicine versus surgery for carotid stenosis. *Eur J Radiol* 2006; 60: 3-7
- Engelhard** K, Werner C, Eberspacher E et al. Influence of propofol on neuronal damage and apoptotic factors after incomplete cerebral ischemia and reperfusion in rats: a long-term observation. *Anesthesiology* 2004; 101: 912-917
- Engelhard** K, Werner C, Möllenbergs O, Kochs E. Effects of remifentanil / propofol in comparison with isoflurane on dynamic cerebrovascular autoregulation in humans. *Acta Anaesthesiol Scand* 2001; 45: 971-976
- Engelhard** K, Werner C. Postoperatives kognitives Defizit. *Anaesthetist* 2005; 54: 588-594
- Epstein** CM, American Clinical Neurophysiology Society. Guidelines on short-latency somatosensory evoked potentials. *J Clin Neurophysiol* 2006; 23: 168-179
- Erasmi** H, Walter M, Löhr G. Läßt sich die Sicherheit des SEP-Monitorings während der Carotis-Chirurgie verbessern? *Langenbecks Arch Chir* 1997; Suppl II: 1293-1294
- Espaniola-Klein** C, Rupprecht HJ, Meyer J. Karotisstenose als Zusatzbefund bei koronarer Herzkrankheit. *Chirurg* 2004; 75: 667-671
- Fearn** SJ, Hutchinson S, Riding G, et al. Carotid endarterectomy improves cognitive function in patients with exhausted cerebrovascular reserve. *Eur J Endovasc Surg* 2003; 26: 529-536
- Ferguson** GG, Eliasziw M, Barr HW, et al. The North American Symptomatic Carotid Endarterectomy trial: surgical results in 1415 patients. *Stroke* 1999; 30: 1751-1758
- Fielmuth** S, Uhlig T. The role of somatosensory evoked potentials in detecting cerebral ischaemia during carotid endarterectomy. *Eur J Anaesthesiol* 2008; 25: 648-656

- Fisher RS, Raudzens P, Nunemacher M.** Efficacy of intraoperative neurophysiological monitoring. *J Clin Neurophysiol* 1995; 12: 97-109
- Florence G, Guérat JM, Gueguen B.** Electroencephalography (EEG) and somatosensory evoked potentials (SEP) to prevent cerebral ischaemia in the operating room. *Clin Neurophysiol* 2004; 34: 17-32
- Fredman B, Lahav M, Zohar E et al.** The effect of midazolam premedication on mental and psychomotor recovery in geriatric patients undergoing brief surgical procedures. *Anesth Analg* 1999; 89: 1161-1166
- Fukuda T, Maruno T, Ito H.** Neuropsychologic function in chronic carotid obstructive disease. *J Stroke Cerebrovasc Dis* 2001; 10: 49-54
- GALA: GALA Trial Collaborative Group.** General anaesthesia versus local anaesthesia for carotid surgery (GALA): a multicentre, randomised controlled trial. *Lancet* 2008; 372: 2132-2142
- Gallinat J, Möller H-J, Moser RL, Hegerl U.** Das postoperative Delir. Risikofaktoren, Prophylaxe und Therapie. *Anaesthesist* 1999; 48: 507-518
- Gao M-Y, Sillesen HH, Lorentzen JE, Schroeder TV.** Eversion carotid endarterectomy generates fewer microemboli than standard carotid endarterectomy. *Eur J Endovasc Surg* 2000; 20: 153-157
- Gentili F, Longheed WM, Ghate H, et al.** The role of intraoperative monitoring of sensory evoked potentials during cardiovascular surgery. In Suzuki J (ed.) *Advances in surgery for cerebral stroke*. 1997, Springer, Berlin, Heidelberg, New York
- Ghogawala Z, Westerveld M, Amin-Hanjani S.** Cognitive outcomes after carotid revascularization: the role of cerebral emboli and hypoperfusion. *Neurosurgery* 2008; 62: 385-395
- Gigli GL, Caramia M, Marciani MG, et al..** Monitoring of subcortical and cortical somatosensory evoked potentials during carotid endarterectomy: comparison with stump pressure levels. *Electroencephalogr Clin Neurophysiol* 1987; 68: 424-432
- Godet G, Reina M, Roux M et al.** Anaesthesia for carotid endarterectomy: comparison of hypnotic- and opioid-based technique. *Br J Anaesth* 2004; 92: 329-334
- Grubhofer G.** Systemic blood pressure and cerebral blood flow during carotid surgery. *Thorac Cardiovasc Surg* 1999; 47: 381-385
- Grundy BI.** Monitoring of somatosensory evoked potentials during neurosurgical operations: Methods and applications. *Neurosurgery* 1982; 11: 556-575
- Guérat JM, Fischer C, Facco E, et al.** Standards of clinical practice of EEG and EPs in comatose and other unresponsive states. The International Federation of Clinical Neurophysiology. *Electroencephalogr Clin Neurophysiol Suppl* 1999; 52: 117-131
- Guérat JM, Witdoeck C, de Tourtchaninoff M, et al.** Somatosensory evoked potentials in carotid endarterectomy. I. Relationships between qualitative SEP alterations and intraoperative events. *Electroencephalogr Clin Neurophysiol* 1997; 104: 459-469
- Hansen D.** Systemische und regionale Oxygenierung während tiefer kontrollierter Hypotension. 2000; Habilitationsschrift, Medizinische Fakultät der Freien Universität Berlin
- Hartmann A, Hupp T, Koch H-C, et al.** Prospective Study on the complication rate of carotid endarterectomy. *Cerebrovasc Dis* 1999; 9: 152-156
- Haupt WF, De Vleeschauwer P, Horsch S.** Veränderungen somatosensibel evozierter Potentiale während Karotis-Desobliteration. Diagnostische Bedeutung und mögliche Konsequenzen für die Therapie. *Z EEG-EMG* 1985; 16: 201-205
- Haupt WF, Erasmi-Körber H, Lanfermann H.** Intraoperative recording of parietal SEP can miss hemodynamic infarction during carotid endarterectomy: a case study. *Electroencephalogr Clin Neurophysiol* 1994; 92: 86-88
- Haupt WF, Horsch S.** Evoked potential monitoring in carotid surgery: a review of 994 cases. *Neurology* 1992; 42: 835-838
- Haynes CD, King GD, Dempsey RL.** Improvement of cognitive and personality changes after carotid endarterectomy. *Br J Surg* 1974; 61: 288-289

- Heckmann** JG, Erbguth FJ, Hilz MJ, Lang CJG, Neundörfer B. Die Hirndurchblutung aus klinischer Sicht. Historischer Überblick, Physiologie, Pathophysiologie, diagnostische und therapeutische Aspekte. *Med Klin* 2001; 96: 583-592
- Helmstaedter** C, Durwen HF. VLMT: Verbaler Lern- und Merkfähigkeitstest. Ein praktikables und differenziertes Instrumentarium zur Prüfung der verbalen Gedächtnisleistungen. *Schweiz Archiv für Neurologie und Psychiatrie* 1990; 141: 21-30
- Hemmingsen** R, Mejsholm B, Vorstrup S, et al. Carotid surgery, cognitive function, and cerebral blood flow in patients with transient ischemic attacks. *Ann Neurol* 1986; 20: 13-19
- Heyer** EJ, De La Paz R, Halazun HJ, et al.. Neuropsychological dysfunction in the absence of structural evidence for cerebral ischemia after uncomplicated carotid endarterectomy. *Neurosurgery* 2006; 58: 474-480
- Heyer** EJ, Sharma R, Rampersad A, et al. A controlled prospective study of neuropsychological dysfunction following carotid endarterectomy. *Arch Neurol* 2002; 59: 217-222
- Heyer** EJ, Sharma R, Winfree CJ, et al. Severe pain confounds neuropsychological test performance. *J Clin Neuropsychol* 2000; 22: 633-639
- Heyer** EJ, Wilson DA, Sahlein DH, et al. APOE-ε4 predisposes to cognitive dysfunction following uncomplicated carotid endarterectomy. *Neurology* 2005; 65: 1759-1763
- Hillen** T, Nieczaj R, Münzberg H, Schaub R, Borchelt M, Steinhagen-Thiessen E. Carotid atherosclerosis, vascular risk profile and mortality in a population-based sample of functionally healthy elderly subjects: the Berlin ageing study. *J Intern Med* 2000; 247: 679-688
- Hoffmann** A, Dinkel M, Schweiger H, Lang W. The influence of the state of the vertebral arteries on the peri- and postoperative risk in carotid surgery. *Eur J Vasc Endovasc Surg* 1998; 16: 329-333
- Horn** W. Leistungsprüfssystem LPS. 1983, Hogrefe Verlag, Göttingen
- Horsch** S, De Vleeschauwer P, Ktenidis K. Intraoperative assessment of cerebral ischemia during carotid surgery. *J Cardiovasc Surg* 1990; 31: 599-602
- Horsch** S, Ktenidis K. Intraoperative use of somatosensory evoked potentials for brain monitoring during carotid surgery. *Neurosurg Clin North Am* 1996; 7: 693-702
- Hossmann** KA. Viability thresholds and the penumbra of focal ischemia. *Ann Neurol* 1994; 36: 557-565
- Howell** SJ. Carotid endarterectomy. *Br J Anaesth* 2007; 99: 119-131
- Incalizi** RA, Gemma A, Landi F, et al. Neuropsychologic effects of carotid endarterectomy. *J Clin Exp Neuropsych* 1997; 19: 785-794
- Inzitari** D, Glazsiw M, Gates P, et al. The causes and risk of stroke in patients with asymptomatic internal-carotid-artery stenosis. North American Symptomatic Carotid Endarterectomy Trial Collaborators. *N Engl J Med* 2000; 342: 1693-1700
- Irvine** CD, Gardner FV, Davies AH, et al. Cognitive testing in patients undergoing carotid endarterectomy. *Eur J Vasc Endovasc Surg* 1998; 15: 195-204
- Jacobs** LA, Brinkman SD, Morrell RM, et al. Long-latency somatosensory evoked potentials during carotid endarterectomy. *The American Surgeon* 1983; 49: 338-344
- Jacobs** LA, Ganji S, Shirley JG, et al. Cognitive improvement after extracranial reconstruction for the low flow-endangered brain. *Surgery* 1983; 93: 683-687
- Jasper** HH. The ten-twenty electrode system of the International Federation. *Electroencephalogr Clin Neurophysiol* 1958; 10: 371-375
- Jellish** WS, Sheikh T, Baker WH, Louie EK, Slogoff S. Hemodynamic stability, myocardial ischemia, and perioperative outcome after carotid surgery with remifentanil / propofol or isoflurane / fentanyl anesthesia. *J Neurosurg Anaesthesiol* 2003; 15: 176-184
- Johnston** SC, O'Mara ES, Manolio TA, et al. Cognitive impairment and decline are associated with carotid artery disease in patients without clinically evident cerebrovascular disease. *Ann Intern Med* 2004; 140: 237-247

- Kearse** LA Jr., Brown EN, McPeck K. Somatosensory evoked potentials sensitivity relative to electroencephalography for cerebral ischemia during carotid endarterectomy. *Stroke* 1992; 23: 498-505
- Kelly** MP, Garron DC, Javid H. Carotid artery disease, carotid endarterectomy, and behavior. *Arch Neurol* 1980; 37: 743-748
- Keren** O, Ring H, Solzi P. Upper limb somatosensory evoked potentials as a predictor of rehabilitation progress in dominant hemisphere stroke patients. *Stroke* 1993; 24:1789-1793
- King** GD, Gideon DA, Haynes CD, et al. Intellectual and personality changes associated with carotid endarterectomy. *J Clin Psychol* 1977; 33: 215-220
- Kishikawa** K, Kamouchi M, Okada Y, Inoue T, Ibayashi S, Iida M. Effects of carotid endarterectomy on cerebral blood flow and neuropsychological test performance in patients with high-grade carotid stenosis. *J Neurol Sci* 2003; 213: 19-24
- Klijn** CJM, Kappelle LJ, Tulleken CAF, et al. Symptomatic carotid artery occlusion: a reappraisal of hemodynamic factors. *Stroke* 1997; 28: 2084-2093
- Kodl** CT, Seaquist ER. Cognitive dysfunction and diabetes mellitus. *Endocrine Reviews* 2008; 29: 494-511
- Kostopanagiotou** G, Markantonis SL, Polydoron M, Pandazi A, Kottis G. Recovery and cognitive function after fentanyl or remifentanyl administration for carotid endarterectomy. *J Clin Anesth* 2005; 17: 16-20
- Krul** JMJ, van Gijn J, Ackerstaff RGA, Eikelboom BC, Theodorides T, Vermeulen FEE. Site and pathogenesis of infarcts associated with carotid endarterectomy. *Stroke* 1989; 20: 324-328
- Kubale** R, Arning C: Stellenwert der Doppler-sonographischen Verfahren zur Diagnose der Karotisstenosen. *Radiologe* 2004; 44: 946-959
- Kuroda** S, Houkin K, Abe H, Hoshi Y, Tamura M. Near-infrared monitoring of cerebral oxygenation state during carotid endarterectomy. *Surg Neurol* 1996; 45: 450-458
- Lal** BK. Cognitive function after carotid artery revascularization. *Vasc Endovasc Surg* 2007; 41: 5-13
- Lam** AM, Manninen PH, Ferguson GG, et al. Monitoring electrophysiologic function during carotid endarterectomy: a comparison of somatosensory evoked potentials and conventional electroencephalogram. *Anesthesiology* 1991; 75: 15-21
- Larsen** B, Seitz A, Larsen R. Recovery of cognitive function after remifentanil-propofol anesthesia: a comparison with desflurane and sevoflurane anesthesia. *Anesth Analg* 2000; 90: 168-174
- Lazar** RM, Marshall RS, Pile-Spellmann J, et al. Continuous time estimation as a behavioural index of human cerebral ischemia during temporary occlusion of the internal carotid artery. *J Neurol Neurosurg Psychiatry* 1996; 60: 559-563
- Lee** EK, Seyal M. Generators of short latency human somatosensory-evoked potentials recorded over the spine and scalp. *J Clin Neurophysiol* 1998; 15: 227-234
- Levin** HS. A guide to clinical neuropsychological testing. *Arch Neurol* 1994; 51: 854-859
- Levine** B, Milbert WB, Stuss D. The Boston Examination of Executive Functioning (BEEF). Unpublished data, 1992. In: Albert ML, Knofel JE. Clinical neurology of ageing. 1997. Oxford University Press, Oxford, New York.
- Lewis** M, Maruff P, Silbert B. Statistical and conceptual issues in defining postoperative cognitive dysfunction. *Neuroscience and Behavioral Reviews* 2004; 28: 433-440
- Liebeskind** DS. Collateral circulation. *Stroke* 2003; 34: 2279-2284
- Lind** C, Wimmer A, Magometschnigg H, et al. Einflüsse der Karotisendarterektomie auf verschiedene Hirnleistungsparameter. Eine neuropsychologische Längsschnittstudie. *Langenbecks Arch Chir* 1993; 378: 345-352
- Linstedt** U, Maier C, Petry A. Intraoperative monitoring with somatosensory evoked potentials in carotid artery surgery – less reliable in patients with preoperative neurologic deficiency? *Acta Anaesthesiol Scand* 1998; 42: 13-16

- Liu** EHC, Wong HK, Chia CP, Lim HJ, Chen ZY, Lee TL. Effects of isoflurane and propofol on cortical somatosensory evoked potentials during comparable depth of anaesthesia guided by bispectral index. *Br J Anaesth* 2005; 94: 193-197
- López** J. The use of evoked potentials in intraoperative neurophysiologic monitoring. *Phys Med Rehabil Clin N Am* 2004; 15: 63-84
- Lunn** S, Crawley F, Harrison MJG, et al. Impact of carotid endarterectomy upon cognitive functioning. *Cerebrovasc Dis* 1999; 9: 74-81
- Mandell** AM, Knofel JE, Albert ML. Mental status examination in the elderly. In: Albert ML, Knofel JE. *Clinical neurology of ageing*. 1997. Oxford University Press, Oxford, New York.
- Manninen** P, Srjeant R, Joshi M. Posterior tibial nerve and median nerve somatosensory evoked potential monitoring during carotid endarterectomy. *Can J Anesth* 2004; 51: 937-941
- Manninen** PH, Tan TK, Sarjeant RM. Somatosensory evoked potential monitoring during carotid endarterectomy in patients with a stroke. *Anesth Analg* 2001; 93: 39-44
- Marrocco-Trischitta** MM, Tiezzi A, Svampa MG et al. Perioperative stress response to carotid endarterectomy: the impact of anaesthetic modality. *J Vasc Surg* 2004; 39: 1295-1304
- Marshall** KE, Vaitkeviciute I. Carotid endarterectomy, carotid artery shunting and outcome: an historical perspective. *Curr Opin Anaesthesiol* 2004; 17: 183-187
- Marshall** RS, Lazar RM, Mohr JP, et al. Higher cerebral function and hemispheric blood flow during awake carotid artery balloon test occlusions. *J Neurol Neurosurg Psychiatry* 1999; 66: 734-738
- Marshall** RS, Lazar RM, Pile-Spellman J, et al. Recovery of brain function during induced cerebral hypoperfusion. *Brain* 2001; 124: 1208-1217
- Marshall** RS. The functional relevance of cerebral hemodynamics: why blood flow matters to the injured and recovering brain. *Curr Opin Neurol* 2004; 17: 705-709
- Mathiesen** EB, Waterloo K, Joakimsen O, et al. Reduced neuropsychological test performance in asymptomatic carotid stenosis. *Neurology* 2004; 62: 695-701
- McCarthy** RJ, Nasr MK, Mc Ateer MK. Physiological advantages of cerebral blood flow during carotid endarterectomy under local anaesthesia: a randomised clinical trial. *Eur J Vasc Endovasc Surg* 2002; 24: 215-221
- McCleary** AJ, Maritati G, Gough MJ. Carotid endarterectomy: local or general anaesthesia? *Eur J Vasc Endovasc Surg* 2001; 22: 1-12
- McGirt** MJ, Perler BA, Brooke SB et al. 3-Hydroxy-3-methylglutaryl coenzyme A reductase inhibitors reduce the risk of perioperative stroke and mortality after carotid endarterectomy. *J Vasc Surg* 2005; 42 : 829-836
- Mehta** M, Rahmani O, Dietzek AM, et al. Eversion technique increases the risk for post-carotid endarterectomy hypertension. *J Vasc Surg* 2001; 34: 839-845
- Meyer** JS, Rauch G, Rauch RA, et al. Risk factors for cerebral hypoperfusion, mild cognitive impairment, and dementia. *Neurobiol Ageing* 2000; 21: 161-169
- Mocco** J, Wilson DA, Ducruet AF, et al. Elevations in preoperative monocyte predispose to acute neurocognitive decline after carotid endarterectomy for asymptomatic carotid artery stenosis. *Stroke* 2006b; 37: 240-242
- Mocco** J, Wilson DA, Komotar RJ, et al. Predictors of neurocognitive decline after carotid endarterectomy. *Neurosurgery* 2006a; 58: 844-850
- Momjian-Mayor** I, Baron J-C. The pathophysiology of watershed infarction in internal carotid artery disease. Review of cerebral perfusion studies. *Stroke* 2005; 36: 567-577
- Monk** TG, Weldon BC, Garvan CW, et al. Predictors of cognitive dysfunction after major noncardiac surgery. *Anesthesiology* 2008; 108: 18-30
- Morawetz** RB, Zeiger EH, McDowell HA, et al. Correlation of cerebral blood flow and EEG during carotid occlusion for endarterectomy (without shunt) and neurologic outcome. *Surgery* 1984; 96: 184-189

- Moritz** S, Kasprzak P, Arlt M, Taeger K, Metz C. Accuracy of cerebral monitoring in detecting cerebral ischemia during carotid endarterectomy: a comparison of transcranial Doppler sonography, near-infrared spectroscopy, stump pressure, and somatosensory evoked potentials. *Anesthesiology* 2007; 107: 563-569
- Mosso** M, Baumgartner RW. Karotisstenose: Epidemiologie und Symptomatologie. *Schweiz Med Wochenschr* 2000; 130: 1226-1230
- Müller** SV, Harth S, Hildebrandt H, Münte TF. Evidence based treatment of executive dysfunction. *Fortschr Neurol Psychiatr* 2006; 74: 10-18
- Murkin** JM, Newman SP, Stump DA, Blumenthal JA. Statement of consensus on assessment of neurobehavioral outcomes after cardiac surgery. *Ann Thorac Surg* 1995; 59: 1289-1295
- Mussak** T, Biberthaler P, Geisenberger T, et al. Assessment of early brain damage in carotid endarterectomy: evaluation of S-100B serum levels and somatosensory evoked potentials in a pilot study. *World J Surg* 2002; 26: 1251-1255
- NASCET**: Barnett HJ, Taylor DW, Eliasziw M, et al. Benefit of carotid endarterectomy in patients with symptomatic moderate or severe stenosis. North American Symptomatic Carotid Endarterectomy Trial Collaborators (NASCET). *N Engl J Med* 1998; 339: 1415-1425
- NASCET**: North American Symptomatic Carotid Endarterectomy Trial Collaborators. Beneficial effect of carotid endarterectomy in symptomatic patients with high-grade carotid stenosis. *N Engl J Med* 1991; 325: 445 – 453
- Nederkoorn** PJ, Mali WP, Eikelboom BC, et al. Preoperative diagnosis of cerebral artery stenosis. Accuracy of noninvasive testing. *Stroke* 2002; 33: 2003-2008
- Newman** MF, Grocott HP, Mathew JP, et al. Report of the substudy assessing the impact of neurocognitive function on quality of life 5 years after cardiac surgery. *Stroke* 2001; 32: 2874-2881
- Newman** S, Stygall J, Hirani S, Shaefi S, Maze M. Postoperative cognitive dysfunction after noncardiac surgery. *Anesthesiology* 2007; 106: 572-590
- Nuwer** MR, Daube J, Fischer C, et al. Neuromonitoring during surgery. Report of an IFCN committee. *Electroencephalogr Clin Neurophysiol* 1993; 87: 263-276
- Nys** GMS, van Zandvoort MJE, de Kort PLM, Jansen BPW, de Haan EHF, Kappelle LJ. Cognitive disorders in acute stroke: prevalence and clinical determinants. *Cerebrovasc Dis* 2007; 23: 408-416
- Ogasawara** K, Inoue T, Kobayashi M, et al. Cognitive impairment associated with intraoperative and postoperative hypoperfusion without neurologic deficits in a patient undergoing carotid endarterectomy. *Surg Neurol* 2006; 65: 577-581
- Oswald** WD, Fleischmann UM. Nürnberg-Alters-Inventar NAI. 1986. Psychologisches Institut II, Universität Erlangen-Nürnberg, Nürnberg.
- Oswald** WD. Das Nürnberg-Alters-Inventar NAI als psychometrische Methode der klinischen Pharmakologie. In: Platt D. Funktionsstörungen des Gehirns im Alter. 1981. Schattauer Verlag, Stuttgart, New York, S. 129-137
- Paciaroni** M, Eliasziw M, Kappelle LJ, et al. Medical complications associated with carotid endarterectomy. *Stroke* 1999; 30: 1759-1763
- Parker** JC, Granberg BW, Nichols WK, et al. Mental status outcomes following carotid endarterectomy: a six-month analysis. *J Clin Neuropsych* 1983; 5: 345-353
- Parker** JC, Nichols WK, Smarr KL, et al. Neuropsychological parameters of carotid endarterectomy: a two-year prospective analysis. *J Consult Clin Psychology* 1986; 54: 676-681
- Parton** A, Coulthard E, Husain M. Neuropharmacological modulation of cognitive deficits after brain damage. *Curr Opin Neurol* 2005; 18: 675-680
- Pearson** S, Maddern G, Fitridge R. Cognitive performance in patients after carotid endarterectomy. *J Vasc Surg* 2003; 38: 1248-1252
- Pedrini** L, Tarantini S, Cirelli MR, et al. Intraoperative assessment of cerebral ischaemia during carotid surgery. *Int Angiol* 1998; 17: 10-14

- Phillips** NA, Mate-Kole CC. Cognitive deficits in peripheral vascular disease. A comparison of mild stroke patients and normal control subjects. *Stroke* 1997; 28: 777-784
- Powers** WJ. Cerebral hemodynamics in ischemic cerebrovascular disease. *Ann Neurol* 1991; 29: 231-240
- Press** MJ, Chassin MR, Wang J, Tuhrim S, Halm EA. Predicting medical and surgical complications of carotid endarterectomy. *Arch Intern Med* 2006; 166: 914-920
- Prior** PF. EEG monitoring and evoked potentials in brain ischemia. *Br J Anaesth* 1985; 57: 63-81
- Prokop** A, Meyer GP, Walter M, et al. Validity of SEP monitoring in carotid surgery. *J Cardiovasc Surg* 1996; 37: 337-342
- Pugh** KG, Kiely DK, Milberg WP, Lipsitz LA. Selective impairment of frontal-executive function in african americans with cardiovascular risk factors. *J Am Geriatr Soc* 2003; 51: 1439-1444
- Pulli** R, Dorigo W, Barbanti E, et al. Carotid endarterectomy with contralateral carotid artery occlusion: is this a higher risk subgroup? *Eur J Endovasc Surg* 2002; 24: 63-68
- Rankin** KP, Kochamba GS, Boone KB, et al. Presurgical cognitive deficits in patients receiving coronary artery bypass graft surgery. *J Int Neuropsychol Soc* 2003; 9: 913 – 924
- Rao** R, Jackson S, Howard R. Neuropsychological impairment in stroke, carotid stenosis and peripheral vascular disease. *Stroke* 1999; 30: 2167-2173
- Rao** R. The role of carotid stenosis in vascular cognitive impairment. *Eur Neurol* 2001; 46: 63-69
- Rao** R. The role of carotid stenosis in vascular cognitive impairment. *J Neurol Sci* 2002; 203/204: 103-107
- Rasmussen** LS, Siersma VD, the ISPOCD group. Postoperative cognitive dysfunction: true detoriation versus random variation. *Acta Anaesthesiol Scand* 2004; 48: 1137-1143
- Rasmussen** LS, Steentoft A, Rasmussen H, et al. Benzodiazepines and postoperative cognitive dysfunction in the elderly. *Br J Anaesth* 1999; 83: 585-589
- Rasmussen** LS, Larsen K, Houx P, et al. The assessment of postoperative cognitive function. *Acta Anaesthesiol Scand* 2001; 45: 275-289
- Reischies** FM, Neu P. Comorbidity of mild cognitive disorder and depression: a neuropsychological analysis. *Eur Arch Psychiatry Clin Neurosci* 2000; 250: 186-193
- Reischies** FM. Neuropsychologisches Defizit-Screening. *Nervenarzt* 1987; 58: 219-226
- Reisecker** F, Witzmann A, Deisenhammer E. Somatosensory evoked potentials (SSEPs) in various groups of cerebro-vascular ischaemic disease. *Electroencephalogr Clin Neurophysiol* 1986; 65: 260-268
- Rey** A. L'examen clinique en psychologie. 1964, Presses Universitaires de France, Paris.
- Riles** TS, Imparato AM, Jacobowitz GR, et al. The cause of perioperative stroke after carotid endarterectomy. *J Vasc Surg* 1994; 19: 206-216
- Rockwood** K. Vascular cognitive impairment and vascular dementia. *J Neurol Sci* 2002; 203/204: 23-27
- Rohan** D, Buggy DJ, Crowley S, et al. Increased incidence of postoperative cognitive dysfunction 24hr after minor surgery in the elderly. *Can J Anesth* 2005; 52: 137-142
- Román** GC. Brain hypoperfusion: a critical factor in vascular dementia. *Neurol Res* 2004; 26: 454-458
- Ropper** AH. Evoked potentials in cerebral ischemia. *Stroke* 1986; 17: 3-5
- Rössel** T, Litz RJ, Heller AR, Koch T. Anästhesie zur Karotischirurgie. Gibt es einen Goldstandard? *Anaesthesia* 2008; 57: 115-130
- Rothwell** PM, Eliasziw M, Gutnikov SA, et al. for the Carotid endarterectomy Trialists Collaboration. Analysis of pooled data from the randomised controlled trials of endarterectomy for symptomatic carotid stenosis. *Lancet* 2003; 361: 107-116

- Rothwell PM**, Eliasziw M, Gutnikov SA, Fox AJ, Warlow CP, Barnett HJM. Endarterectomy for symptomatic carotid stenosis in relation to clinical subgroups and timing of surgery. *Lancet* 2004; 363: 915-924
- Rothwell PM**, Gutnikov SA, Myberg M, et al. A pooled analysis of individual patient data from trials of endarterectomy for symptomatic carotid stenosis. *Stroke* 2001; 32: 328
- Rothwell PM**, Pendlebury ST, Wardlaw J, Warlow CP. Critical appraisal of the design and reporting of studies of imaging and measurement of carotid stenosis. *Stroke* 2000; 31: 1444-1450
- Rundshagen I**. Postoperative kognitive Dysfunktion. *Anästh Intensivmed* 2008; 49: 566-580
- Sachdev PS**, Brodaty H, Valenzuela MJ, et al. The neuropsychological profile of vascular cognitive impairment in stroke and TIA patients. *Neurology* 2004; 62: 912-919
- Sachdev PS**, Brodaty H, Valenzuela MJ, Lorentz LM, Koschera A. Progression of cognitive impairment in stroke patients. *Neurology* 2004; 63: 1618-1623
- Samra SK**, Dy EA, Welch KB, Lovely LK, Graziano GP. Remifentanil- and fentanyl-based anesthesia for intraoperative monitoring of somatosensory evoked potentials. *Anesth Analg* 2001; 92: 1510-1515
- Saxton J**, Ratcliff G, Newman A, et al. Cognitive test performance and presence of subclinical cardiovascular disease in the Cardiovascular Health Study. *Neuroepidemiology* 2000; 19: 312-319
- Sbarigia E**, schioppa A, Misuraca M, et al. Somatosensory evoked potentials versus locoregional anaesthesia in the monitoring of cerebral function during carotid artery surgery: preliminary results of a prospective study. *Eur J Vasc Endovasc Surg* 2001; 21: 413-416
- Schneemilch CE**, Ludwig S, Ulrich A, Halloul Z, Hachenberg T. Somatosensory evoked potentials and biochemical markers of neuronal deficits in patients undergoing carotid endarterectomy under regional anesthesia. *Zentralbl Chir* 2007; 132: 176-182
- Schwarz G**, Pfurtscheller G, Tritthart H, Litscher G, List WF. Begleitphänomene beim EP-Monitoring auf der Intensivstation. *Intensivbehandlung* 1989; 14: 28-32
- Schweiger H**, Dinkel M, Lang W. Risk reduction in carotid surgery by monitoring somatosensory evoked potentials. *Vasa* 1992; 35 (Suppl): 21
- Schweiger H**, Kamp HD, Dinkel M. Somatosensory-evoked potentials during carotid artery surgery: experience in 400 operations. *Surgery* 1991; 109: 602-609
- Sciarroni L**, Gremigni P, Pedrini L. Psychological impact of carotid endarterectomy: a review of the studies. *Monaldi Arch Chest Dis* 2007; 68: 170-177
- Seidenberg M**, Parker JC, Nichols WK, Davenport J, Hewett JE. Carotid stenosis and arteriosclerotic heart disease: interactive effects on cognitive status. *Int J Clin Neuropsych* 1985; 7: 45-48
- Shah DM**, Leather RP, Darling RC, Chang BB, Paty PSK, Kreienberg PB. Technique of eversion carotid endarterectomy and contemporary results. *Perspect Vasc Surg Endovasc Ther* 1998; 9: 49-62
- Sheehan MK**, Baker WH, Littooy FN, et al. Timing of postcarotid complications: a guide to safe discharge planning. *J Vasc Surg* 2001; 34: 13-16
- Sillesen H**. The natural history of patients with carotid stenosis. *Pathophysiol Haemost Thromb* 2002; 32: 378-380
- Silverstein JH**, Steinmetz J, Reichenberg A, et al. Postoperative cognitive dysfunction in patients with preoperative cognitive impairment. *Anesthesiology* 2007b; 106: 431-435
- Silverstein JH**, Timberger M, Reich DL, Uysal S. Central nervous system dysfunction after noncardiac surgery and anesthesia in the elderly. *Anesthesiology* 2007a; 106: 622-628
- Sloan TB**. Anesthetic effects on electrophysiologic recordings. *J Clin Neurophysiol* 1998; 15: 217-226
- Sloan TB**. Evoked potential monitoring. In: Vender JS (Hrsg) *International Anaesthesiology Clinics*. Boston, MA. Little, Brown and Co 1996; 109-136
- Soinne L**, Helenius J, Tatlisumak T, et al. Cerebral hemodynamics in asymptomatic and symptomatic patients with high-grade carotid stenosis undergoing carotid endarterectomy. *Stroke* 2003; 34: 1655-1661

- Stöhr** M. Somatosensorisch evozierte Potentiale (SEP). In: Maurer K, Lowitzsch K, Stöhr M. Evozierte Potentiale: AEP-VEP-SEP. Stuttgart: Ferdinand Encke Verlag, 1988: 130-178
- Stroop** J. Studies of interference in serial verbal reactions. *J Exper Psych* 1935; 18: 643-662
- Stump** DA. Selection and clinical significance of neuropsychologic tests. *Ann Thorac Surg* 1995; 59: 1340-1344
- Sturm** W, Willmes K, Horn W. Leistungsprüfsystem für 50 – 90jährige (LPS 50+). 1993, Hogrefe Verlag, Göttingen
- Taniguchi** M, Nadstawek J, Pechstein U, Schremm J. Total intravenous anesthesia for improvement of intraoperative monitoring of somatosensory evoked potentials during aneurysm surgery. *Neurosurgery* 1992; 31: 891-897
- Taylor** DW, Barnett HJ, Haynes RB. Low-dose and high-dose acetylsalicylic acid for patients undergoing carotid endarterectomy: a randomized controlled trial. ASA and Carotid Endarterectomy (ACE) Trial Collaborators. *Lancet* 1999; 353: 2179-2184
- Telman** G, Kouperberg E, Nitecki S, et al. Cerebral hemodynamics in symptomatic and asymptomatic patients with severe unilateral carotid stenosis before and after carotid endarterectomy. *Eur J Vasc Endovasc Surg* 2006 ; 32 : 375-378
- Thiel** A, Ritzka M. Zerebrale Überwachungsmaßnahmen in der Karotischirurgie. Ergebnisse einer Umfrage in der Bundesrepublik Deutschland. *Anästhesiol Intensivmed Notfallmed Schmerzther* 2001; 36: 693-697
- Tolekis** JR. Intraoperative monitoring using somatosensory evoked potentials: A position statement by the American Society of Neurophysiological Monitoring. *J Clin Monit Comput* 2005; 19: 241-258
- Trivedi** RA, Green HAL, U-King-Im J et al. Cerebral haemodynamic disturbances in patients with moderate carotid artery stenosis. *Eur J Vasc Endovasc Surg* 2005; 29: 52-57
- Tsiptsios** I, Fountoulakis KN, Sitzoglou K, et al. Clinical and neuroimaging correlates of abnormal short-latency somatosensory evoked potentials in elderly vascular dementia patients: a psychophysiological exploratory study. *Ann Gen Hosp Psychiatry* 2003; 2: 8
- Tzvetanov** P, Rousseff RT, Atanassova P. Prognostic value of median and tibial somatosensory evoked potentials in acute stroke. *Neurosci Lett* 2005; 380: 99-104
- Tzvetanov** P, Rousseff RT. Predictive value of median-SSEP in early phase of stroke: a comparison in supratentorial infarction and hemorrhage. *Clin Neurol Neurosurg* 2005; 107: 475-481
- Uno** M, Suzne A, Nishi K, et al. Hemodynamic cerebral ischemia during carotid endarterectomy evaluated by intraoperative monitoring and post-operative diffusion-weighted imaging. *Neurol Res* 2007; 29: 70-77
- Van den Burg** W, Saan RJ, van Zomeren AH, et al. Carotid endarterectomy: does it improve cognitive or motor functioning? *Psychological Medicine* 1985; 15: 341-346
- van Mook** WNKA, Rennenberg RJMW, Schurink GW, et al. Cerebral hyperperfusion syndrome. *Lancet Neurol* 2005; 4: 877-888
- Vanninen** E, Vanninen R, Aikia M, et al. Frequency of carotid endarterectomy-related subclinical cerebral complications. *Cerebrovasc Dis* 1996; 6: 272-280
- VA-Study:** Mayberg MR, Wilson SE, Yatsu F, et al. Carotid endarterectomy and prevention of cerebral ischemia in symptomatic carotid stenosis. Veterans Affairs Cooperative Studies Program 309 Trialist Group. *JAMA* 1991; 266: 3289-3294
- Verborgh** C. Anaesthesia in patients with dementia. *Curr Opin Anaesthetol* 2004; 17: 277-283
- Wiedemayer** H, Fauser B, Sandalciogh IE, Schäfer H, Stolke D. The impact of neurophysiological intraoperative monitoring on surgical decisions: a critical analysis of 423 cases. *J Neurosurg* 2002; 96: 255-262
- Wilson** SE, Bennion RS, Owens L. Effect of carotid endarterectomy on postoperative psychological test performance. *J Cardiovasc Surg* 1983; 24: 428

Witdoeckt C, Chariani S. Somatosensory evoked potentials in carotid surgery. II. Comparison between qualitative and quantitative scoring systems. *Electroenceph Clin Neurophysiol* 1997; 104: 328-332

Wöber C, Zeitlhofer J, Asenbaum S, et al. Monitoring of median nerve somatosensory evoked potentials in carotid surgery. *J Clin Neurophysiol* 1998; 15: 429-438

Yamada T, Yeh M, Kimura J. Fundamental principles of somatosensory evoked potentials. *Phys Med Rehabil Clin N Am* 2004; 15: 19-42

Zinn S, Bosworth HB, Hoenig HM, Swartzwelder HS. Executive function in acute stroke. *Arch Phys Med Rehabil* 2007; 88: 173-180