

7. LITERATURVERZEICHNIS

- Abdulaev, N.G. and K.D. Ridge. 1998. Light-induced exposure of the cytoplasmic end of transmembrane helix seven in rhodopsin. *Proc Natl Acad Sci U S A.* 95:12854-9.
- Ala, Y., D. Morin, B. Mouillac, N. Sabatier, R. Vargas, N. Cotte, M. Dechaux, C. Antignac, M.F. Arthus, M. Lonergan, M.S. Turner, M.N. Balestre, G. Alonso, M. Hibert, C. Barberis, G.N. Hendy, D.G. Bichet and S. Jard. 1998. Functional studies of twelve mutant V2 vasopressin receptors related to nephrogenic diabetes insipidus: molecular basis of a mild clinical phenotype. *J Am Soc Nephrol.* 9:1861-72.
- Andersen-Beckh, B., M. Dehe, R. Schülein, B. Wiesner, C. Rutz, U. Liebenhoff, W. Rosenthal and A. Oksche. 1999. Polarized expression of the vasopressin V2 receptor in Madin-Darby canine kidney cells. *Kidney Int.* 56:517-27.
- Anukanth, A. and H.G. Khorana. 1994. Structure and function in rhodopsin. Requirements of a specific structure for the intradiscal domain. *J Biol Chem.* 269:19738-44.
- Appleyard, S.M., J. Celver, V. Pineda, A. Kovoov, G.A. Wayman and C. Chavkin. 1999. Agonist-dependent desensitization of the kappa opioid receptor by G protein receptor kinase and beta-arrestin. *J Biol Chem.* 274:23802-7.
- Beau, I., M.T. Groyer-Picard, A. Le Bivic, B. Vannier, H. Loosfelt, E. Milgrom and M. Misrahi. 1998. The basolateral localization signal of the follicle-stimulating hormone receptor. *J Biol Chem.* 273:18610-6.
- Beau, I., M. Misrahi, B. Gross, B. Vannier, H. Loosfelt, M.T. Hai, C. Pichon and E. Milgrom. 1997. Basolateral localization and transcytosis of gonadotropin and thyrotropin receptors expressed in Madin-Darby canine kidney cells. *J Biol Chem.* 272:5241-8.
- Bermak, J.C., M. Li, C. Bullock, P. Weingarten and Q.Y. Zhou. 2002. Interaction of gamma-COP with a transport motif in the D1 receptor C-terminus. *Eur J Cell Biol.* 81:77-85.
- Bermak, J.C., M. Li, C. Bullock and Q.Y. Zhou. 2001. Regulation of transport of the dopamine D1 receptor by a new membrane-associated ER protein. *Nat Cell Biol.* 3:492-8.

- Bichet, D.G., M. Birnbaumer, M. Lonergan, M.F. Arthus, W. Rosenthal, P. Goodyer, H. Nivet, S. Benoit, P. Giampietro, S. Simonetti and et al. 1994. Nature and recurrence of AVPR2 mutations in X-linked nephrogenic diabetes insipidus. *Am J Hum Genet.* 55:278-86.
- Bichet, D.G., A. Oksche and W. Rosenthal. 1997. Congenital nephrogenic diabetes insipidus. *J Am Soc Nephrol.* 8:1951-8.
- Bichet, D.G., M. Turner and D. Morin. 1998. Vasopressin receptor mutations causing nephrogenic diabetes insipidus. *Proc Assoc Am Physicians.* 110:387-94.
- Bockaert, J. and J.P. Pin. 1999. Molecular tinkering of G protein-coupled receptors: an evolutionary success. *Embo J.* 18:1723-9.
- Bowen-Pidgeon, D., G. Innamorati, H.M. Sadeghi and M. Birnbaumer. 2001. Arrestin effects on internalization of vasopressin receptors. *Mol Pharmacol.* 59:1395-401.
- Chapple, J.P. and M.E. Cheetham. 2003. The chaperone environment at the cytoplasmic face of the endoplasmic reticulum can modulate rhodopsin processing and inclusion formation. *J Biol Chem.*
- Chazenbalk, G.D., Y. Nagayama, D. Russo, H.L. Wadsworth and B. Rapoport. 1990. Functional analysis of the cytoplasmic domains of the human thyrotropin receptor by site-directed mutagenesis. *J Biol Chem.* 265:20970-5.
- Chuang, J.Z. and C.H. Sung. 1998. The cytoplasmic tail of rhodopsin acts as a novel apical sorting signal in polarized MDCK cells. *J Cell Biol.* 142:1245-56.
- Cyr, C.R., B. Rudy and R.M. Kris. 1993. Prolonged desensitization of the human endothelin A receptor in *Xenopus* oocytes. Comparative studies with the human neurokinin A receptor. *J Biol Chem.* 268:26071-4.
- Deretic, D., S. Schmerl, P.A. Hargrave, A. Arendt and J.H. McDowell. 1998. Regulation of sorting and post-Golgi trafficking of rhodopsin by its C-terminal sequence QVS(A)PA. *Proc Natl Acad Sci U S A.* 95:10620-5.
- Ellgaard, L. and A. Helenius. 2003. Quality control in the endoplasmic reticulum. *Nat Rev Mol Cell Biol.* 4:181-91.

- Ellis, R.J. 1990. The molecular chaperone concept. *Semin Cell Biol.* 1:1-9.
- Fiedler, K., M. Veit, M.A. Starnes and J.E. Rothman. 1996. Bimodal interaction of coatamer with the p24 family of putative cargo receptors. *Science.* 273:1396-9.
- Franzoni, L., G. Nicastro, T.A. Pertinhez, M. Tato, C.R. Nakaie, A.C. Paiva, S. Schreier and A. Spisni. 1997. Structure of the C-terminal fragment 300-320 of the rat angiotensin II AT1A receptor and its relevance with respect to G-protein coupling. *J Biol Chem.* 272:9734-41.
- Fritze, O., S. Filipek, V. Kuksa, K. Palczewski, K.P. Hofmann and O.P. Ernst. 2003. Role of the conserved NPxxY(x)5,6F motif in the rhodopsin ground state and during activation. *Proc Natl Acad Sci U S A.* 100:2290-5.
- Fukushima, Y., T. Asano, K. Takata, M. Funaki, T. Ogihara, M. Anai, K. Tsukuda, T. Saitoh, H. Katagiri, M. Aihara, N. Matsuhashi, Y. Oka, Y. Yazaki and K. Sugano. 1997. Role of the C terminus in histamine H2 receptor signaling, desensitization, and agonist-induced internalization. *J Biol Chem.* 272:19464-70.
- Gabilondo, A.M., J. Hegler, C. Krasel, V. Boivin-Jahns, L. Hein and M.J. Lohse. 1997. A dileucine motif in the C terminus of the beta2-adrenergic receptor is involved in receptor internalization. *Proc Natl Acad Sci U S A.* 94:12285-90.
- Gether, U. 2000. Uncovering molecular mechanisms involved in activation of G protein-coupled receptors. *Endocr Rev.* 21:90-113.
- Gilbert, A., M. Jadot, E. Leontieva, S. Wattiaux-De Coninck and R. Wattiaux. 1998. Delta F508 CFTR localizes in the endoplasmic reticulum-Golgi intermediate compartment in cystic fibrosis cells. *Exp Cell Res.* 242:144-52.
- Gut, A., F. Kappeler, N. Hyka, M.S. Balda, H.P. Hauri and K. Matter. 1998. Carbohydrate-mediated Golgi to cell surface transport and apical targeting of membrane proteins. *Embo J.* 17:1919-29.
- Hauri, H., C. Appenzeller, F. Kuhn and O. Nufer. 2000a. Lectins and traffic in the secretory pathway. *FEBS Lett.* 476:32-7.
- Hauri, H.P., F. Kappeler, H. Andersson and C. Appenzeller. 2000b. ERGIC-53 and traffic in the secretory pathway. *J Cell Sci.* 113 (Pt 4):587-96.

- Helenius, A. and M. Aebi. 2001. Intracellular functions of N-linked glycans. *Science*. 291:2364-9.
- Hermosilla, R. and R. Schülein. 2001. Sorting functions of the individual cytoplasmic domains of the G protein-coupled vasopressin V(2) receptor in Madin Darby canine kidney epithelial cells. *Mol Pharmacol*. 60:1031-9.
- Heyman, J.A. and S. Subramaniam. 1997. Expression, stability and membrane integration of truncation mutants of bovine rhodopsin. *Proc. Natl. Acad. Sci. USA*. 94:4966-4971.
- Horstmeyer, A., H. Cramer, T. Sauer, W. Muller-Esterl and C. Schroeder. 1996. Palmitoylation of endothelin receptor A. Differential modulation of signal transduction activity by post-translational modification. *J Biol Chem*. 271:20811-9.
- Huet, G., V. Gouyer, D. Delacour, C. Richet, J.P. Zanetta, P. Delannoy and P. Degand. 2003. Involvement of glycosylation in the intracellular trafficking of glycoproteins in polarized epithelial cells. *Biochimie*. 85:323-30.
- Hunziker, W. and C. Fumey. 1994. A di-leucine motif mediates endocytosis and basolateral sorting of macrophage IgG Fc receptors in MDCK cells. *Embo J*. 13:2963-7.
- Innamorati, G., C. Le Gouill, M. Balamotis and M. Birnbaumer. 2001. The long and the short cycle. Alternative intracellular routes for trafficking of G-protein-coupled receptors. *J Biol Chem*. 276:13096-103.
- Innamorati, G., H. Sadeghi and M. Birnbaumer. 1996. A fully active nonglycosylated V2 vasopressin receptor. *Mol Pharmacol*. 50:467-73.
- Innamorati, G., H. Sadeghi and M. Birnbaumer. 1999. Phosphorylation and recycling kinetics of G protein-coupled receptors. *J Recept Signal Transduct Res*. 19:315-26.
- Innamorati, G., H. Sadeghi, A.N. Eberle and M. Birnbaumer. 1997. Phosphorylation of the V2 vasopressin receptor. *J Biol Chem*. 272:2486-92.
- Innamorati, G., H.M. Sadeghi, N.T. Tran and M. Birnbaumer. 1998. A serine cluster prevents recycling of the V2 vasopressin receptor. *Proc Natl Acad Sci U S A*. 95:2222-6.

- Jung, H., R. Windhaber, D. Palm and K.D. Schnackerz. 1996. Conformation of a beta-adrenoceptor-derived signal transducing peptide as inferred by circular dichroism and ¹H NMR spectroscopy. *Biochemistry*. 35:6399-405.
- Kappeler, F., D.R. Klopfenstein, M. Foguet, J.P. Paccaud and H.P. Hauri. 1997. The recycling of ERGIC-53 in the early secretory pathway. ERGIC-53 carries a cytosolic endoplasmic reticulum-exit determinant interacting with COPII. *J Biol Chem*. 272:31801-8.
- Karpa, K.D., M.S. Lidow, M.T. Pickering, R. Levenson and C. Bergson. 1999. N-linked glycosylation is required for plasma membrane localization of D5, but not D1, dopamine receptors in transfected mammalian cells. *Mol Pharmacol*. 56:1071-8.
- Kaushal, S., K.D. Ridge and H.G. Khorana. 1994. Structure and function in rhodopsin: the role of asparagine-linked glycosylation. *Proc Natl Acad Sci U S A*. 91:4024-8.
- Kim, G.H., C.A. Ecelbarger, C. Mitchell, R.K. Packer, J.B. Wade and M.A. Knepper. 1999. Vasopressin increases Na-K-2Cl cotransporter expression in thick ascending limb of Henle's loop. *Am J Physiol*. 276:F96-F103.
- Kishore, B.K., B. Mandon, N.B. Oza, S.R. DiGiovanni, R.A. Coleman, N.L. Ostrowski, J.B. Wade and M.A. Knepper. 1996. Rat renal arcade segment expresses vasopressin-regulated water channel and vasopressin V2 receptor. *J Clin Invest*. 97:2763-71.
- Klussmann, E., K. Maric and W. Rosenthal. 2000. The mechanisms of aquaporin control in the renal collecting duct. *Rev Physiol Biochem Pharmacol*. 141:33-95.
- Krause, G., R. Hermosilla, A. Oksche, C. Rutz, W. Rosenthal and R. Schülein. 2000. Molecular and conformational features of a transport-relevant domain in the C-terminal tail of the vasopressin V(2) receptor. *Mol Pharmacol*. 57:232-42.
- Krishna, A.G., S.T. Menon, T.J. Terry and T.P. Sakmar. 2002. Evidence that helix 8 of rhodopsin acts as a membrane-dependent conformational switch. *Biochemistry*. 41:8298-309.
- Laemmli, U.K. 1970. Cleavage of structural proteins during assembly of the head of bacteriophage T4. *Nature*. 227: 680-685.

- Lattion, A.L., D. Diviani and S. Cotecchia. 1994. Truncation of the receptor carboxyl terminus impairs agonist-dependent phosphorylation and desensitization of the alpha 1B-adrenergic receptor. *J Biol Chem.* 269:22887-93.
- Letourneur, F., E.C. Gaynor, S. Hennecke, C. Demolliere, R. Duden, S.D. Emr, H. Riezman and P. Cosson. 1994. Coatamer is essential for retrieval of dilysine-tagged proteins to the endoplasmic reticulum. *Cell.* 79:1199-207.
- Liu, J., T. Schöneberg, M. van Rhee and J. Wess. 1995. Mutational analysis of the relative orientation of transmembrane helices I and VII in G-protein-coupled Receptors. *J Biol Chem.* 270:19532-9.
- Liu, X., D. Davis and D.L. Segaloff. 1993. Disruption of potential sites for N-linked glycosylation does not impair hormone binding to the lutropin/choriogonadotropin receptor if Asn-173 is left intact. *J Biol Chem.* 268:1513-6.
- Luttrell, L.M. and R.J. Lefkowitz. 2002. The role of beta-arrestins in the termination and transduction of G-protein-coupled receptor signals. *J Cell Sci.* 115:455-65.
- Ma, D., N. Zerangue, Y.F. Lin, A. Collins, M. Yu, Y.N. Jan and L.Y. Jan. 2001. Role of ER export signals in controlling surface potassium channel numbers. *Science.* 291:316-9.
- Marin, E.P., A.G. Krishna, T.A. Zvyaga, J. Isele, F. Siebert and T.P. Sakmar. 2000. The amino terminus of the fourth cytoplasmic loop of rhodopsin modulates rhodopsin-transducin interaction. *J Biol Chem.* 275:1930-6.
- Martinez-Ménarguez, J.A., H.J. Geuze, J.W. Slot and J. Klumperman. 1999. Vesicular tubular clusters between the ER and Golgi mediate concentration of soluble secretory proteins by exclusion from COPI-coated vesicles. *Cell.* 98:81-90.
- Matter, K., W. Hunziker and I. Mellman. 1992. Basolateral sorting of LDL receptor in MDCK cells: the cytoplasmic domain contains two tyrosine-dependent targeting determinants. *Cell.* 71:741-53.
- McBee, J.K., K. Palczewski, W. Baehr and D.R. Pepperberg. 2001. Confronting complexity: the interlink of phototransduction and retinoid metabolism in the vertebrate retina. *Prog Retin Eye Res.* 20:469-529.

- McPherson, J.D., M. Marra, L. Hillier, R.H. Waterston, A. Chinwalla, J. Wallis, M. Sekhon, K. Wylie, E.R. Mardis, R.K. Wilson, R. Fulton, T.A. Kucaba, C. Wagner-McPherson, W.B. Barbazuk, S.G. Gregory, S.J. Humphray, L. French, R.S. Evans, G. Bethel, A. Whittaker, J.L. Holden, O.T. McCann, A. Dunham, C. Soderlund, C.E. Scott, D.R. Bentley, G. Schuler, H.C. Chen, W. Jang, E.D. Green, J.R. Idol, V.V. Maduro, K.T. Montgomery, E. Lee, A. Miller, S. Emerling, Kucherlapati, R. Gibbs, S. Scherer, J.H. Gorrell, E. Sodergren, K. Clerc-Blankenburg, P. Tabor, S. Naylor, D. Garcia, P.J. de Jong, J.J. Catanese, N. Nowak, K. Osoegawa, S. Qin, L. Rowen, A. Madan, M. Dors, L. Hood, B. Trask, C. Friedman, H. Massa, V.G. Cheung, I.R. Kirsch, T. Reid, R. Yonescu, J. Weissenbach, T. Bruls, R. Heilig, E. Branscomb, A. Olsen, N. Doggett, J.F. Cheng, T. Hawkins, R.M. Myers, J. Shang, L. Ramirez, J. Schmutz, O. Velasquez, K. Dixon, N.E. Stone, D.R. Cox, D. Haussler, W.J. Kent, T. Furey, S. Rogic, S. Kennedy, S. Jones, A. Rosenthal, G. Wen, M. Schilhabel, G. Gloeckner, G. Nyakatura, R. Siebert, B. Schlegelberger, J. Korenberg, X.N. Chen, A. Fujiyama, M. Hattori, A. Toyoda, T. Yada, H.S. Park, Y. Sakaki, N. Shimizu, S. Asakawa, et al. 2001. A physical map of the human genome. *Nature*. 409:934-41.
- Meacham, G.C., Z. Lu, S. King, E. Sorscher, A. Tousson and D.M. Cyr. 1999. The Hdj-2/Hsc70 chaperone pair facilitates early steps in CFTR biogenesis. *Embo J*. 18:1492-505.
- Mizobe, T., M. Maze, V. Lam, S. Suryanarayana and B.K. Kobilka. 1996. Arrangement of transmembrane domains in adrenergic receptors. Similarity to bacteriorhodopsin. *J Biol Chem*. 271:2387-9.
- Molinari, M. and A. Helenius. 2000. Chaperone selection during glycoprotein translocation into the endoplasmic reticulum. *Science*. 288:331-333.
- Morello, J.P., A. Salahpour, A. Laperriere, V. Bernier, M.F. Arthus, M. Lonergan, U.E. Petaja-Repo, S. Angers, D. Morin, D.G. Bichet and M. Bouvier. 2000. Pharmacological chaperones rescue cell-surface expression and function of misfolded V2 vasopressin receptor mutants. *J. Clin. Invest*. 105:887-895.
- Morello, J.P., A. Salahpour, U.E. Petaja-Repo, A. Laperriere, M. Lonergan, M.F. Arthus, I.R. Nabi, D.G. Bichet and M. Bouvier. 2001. Association of calnexin with wild type and mutant AVPR2 that causes nephrogenic diabetes insipidus. *Biochemistry*. 40:6766-75.

- Morin, D., Y. Ala, N. Sabatier, N. Cotte, G. Hendy, R. Vargas, M. Dechaux, C. Antignac, M. Hibert, D. Bichet and C. Barberis. 1998. Functional study of two V2 vasopressin mutant receptors related to NDI. P322S and P322H. *Adv Exp Med Biol.* 449:391-3.
- Munshi, U.M., H. Peegel and K.M. Menon. 2001. Palmitoylation of the luteinizing hormone/human chorionic gonadotropin receptor regulates receptor interaction with the arrestin-mediated internalization pathway. *Eur J Biochem.* 268:1631-9.
- Nakagawa, M., T. Miyamoto, R. Kusakabe, S. Takasaki, T. Takao, Y. Shichida and M. Tsuda. 2001. O-Glycosylation of G-protein-coupled receptor, octopus rhodopsin. Direct analysis by FAB mass spectrometry. *FEBS Lett.* 496:19-24.
- Nakamura, K. and M. Ascoli. 1999. A dileucine-based motif in the C-terminal tail of the lutropin/choriogonadotropin receptor inhibits endocytosis of the agonist-receptor complex. *Mol Pharmacol.* 56:728-36.
- Nguyen, T.T. and S.G. Amara. 1996. N-linked oligosaccharides are required for cell surface expression of the norepinephrine transporter but do not influence substrate or inhibitor recognition. *J Neurochem.* 67:645-55.
- Nonoguchi, H., A. Owada, N. Kobayashi, M. Takayama, Y. Terada, J. Koike, K. Ujiie, F. Marumo, T. Sakai and K. Tomita. 1995. Immunohistochemical localization of V2 vasopressin receptor along the nephron and functional role of luminal V2 receptor in terminal inner medullary collecting ducts. *J Clin Invest.* 96:1768-78.
- Nufer, O., S. Guldbrandsen, M. Degen, F. Kappeler, J.P. Paccaud, K. Tani and H.P. Hauri. 2002. Role of cytoplasmic C-terminal amino acids of membrane proteins in ER export. *J Cell Sci.* 115:619-28.
- Oakley, R.H., S.A. Laporte, J.A. Holt, M.G. Caron and L.S. Barak. 2000. Differential affinities of visual arrestin, beta arrestin1, and beta arrestin2 for G protein-coupled receptors delineate two major classes of receptors. *J Biol Chem.* 275:17201-10.
- O'Dowd, B.F., M. Hnatowich, M.G. Caron, R.J. Lefkowitz and M. Bouvier. 1989. Palmitoylation of the human beta 2-adrenergic receptor. Mutation of Cys341 in the carboxyl tail leads to an uncoupled nonpalmitoylated form of the receptor. *J Biol Chem.* 264:7564-9.

- O'Hara, P.J., P.O. Sheppard, H. Thogersen, D. Venezia, B.A. Haldeman, V. McGrane, K.M. Houamed, C. Thomsen, T.L. Gilbert and E.R. Mulvihill. 1993. The ligand-binding domain in metabotropic glutamate receptors is related to bacterial periplasmic binding proteins. *Neuron*. 11:41-52.
- Oksche, A., G. Boese, A. Horstmeyer, J. Furkert, M. Beyermann, M. Bienert and W. Rosenthal. 2000. Late endosomal/lysosomal targeting and lack of recycling of the ligand-occupied endothelin B receptor. *Mol Pharmacol*. 57:1104-13.
- Oksche, A., M. Dehe, R. Schülein, B. Wiesner and W. Rosenthal. 1998. Folding and cell surface expression of the vasopressin V2 receptor: requirement of the intracellular C-terminus. *FEBS Lett*. 424:57-62.
- Oksche, A. and W. Rosenthal. 1998. The molecular basis of nephrogenic diabetes insipidus. *J Mol Med*. 76:326-37.
- Oksche, A., R. Schülein, C. Rutz, U. Liebenhoff, J. Dickson, H. Muller, M. Birnbaumer and W. Rosenthal. 1996. Vasopressin V2 receptor mutants that cause X-linked nephrogenic diabetes insipidus: analysis of expression, processing, and function. *Mol Pharmacol*. 50:820-8.
- Orci, L., M. Stames, M. Ravazzola, M. Amherdt, A. Perrelet, T.H. Sollner and J.E. Rothman. 1997. Bidirectional transport by distinct populations of COPI-coated vesicles. *Cell*. 90:335-49.
- Otte, S and C. Barlowe. 2003. The Erv41p-Erv46p complex: multiple export signals are required in trans for COPII-dependent transport from the ER. *EMBO J*. 21:6095-104.
- Palczewski, K., T. Kumasaka, T. Hori, C.A. Behnke, H. Motoshima, B.A. Fox, I. Le Trong, D.C. Teller, T. Okada, R.E. Stenkamp, M. Yamamoto and M. Miyano. 2000. Crystal structure of rhodopsin: A G protein-coupled receptor. *Science*. 289:739-45.
- Parodi, A.J. 2000. Protein glycosylation and its role in protein folding. *Annu Rev Biochem*. 69:69-93.
- Peleg, G., P. Ghanouni, B.K. Kobilka and R.N. Zare. 2001. Single-molecule spectroscopy of the beta(2) adrenergic receptor: observation of conformational substates in a membrane protein. *Proc Natl Acad Sci U S A*. 98:8469-74.

- Petäjä-Repo, U.E., M. Hogue, A. Laperriere, S. Bhalla, P. Walker and M. Bouvier. 2001. Newly synthesized human delta opioid receptors retained in the endoplasmic reticulum are retrotranslocated to the cytosol, deglycosylated, ubiquitinated, and degraded by the proteasome. *J Biol Chem.* 276:4416-23.
- Petäjä-Repo, U.E., M. Hogue, A. Laperriere, P. Walker and M. Bouvier. 2000. Export from the endoplasmic reticulum represents the limiting step in the maturation and cell surface expression of the human delta opioid receptor. *J Biol Chem.* 275:13727-36.
- Pfeiffer, R., J. Kirsch and F. Fahrenholz. 1998. Agonist and antagonist-dependent internalization of the human vasopressin V2 receptor. *Exp Cell Res.* 244:327-39.
- Prioleau, C., I. Visiers, B.J. Ebersole, H. Weinstein and S.C. Sealfon. 2002. Conserved helix 7 tyrosine acts as a multistate conformational switch in the 5HT2C receptor. Identification of a novel "locked-on" phenotype and double revertant mutations. *J Biol Chem.* 277:36577-84.
- Qanbar, R. and M. Bouvier. 2003. Role of palmitoylation/depalmitoylation reactions in G-protein-coupled receptor function. *Pharmacol Ther.* 97:1-33.
- Rands, E., M.R. Candelore, A.H. Cheung, W.S. Hill, C.D. Strader and R.A. Dixon. 1990. Mutational analysis of beta-adrenergic receptor glycosylation. *J Biol Chem.* 265:10759-64.
- Risler, J.L., M.O. Delorme, H. Delacroix and A. Henaut. 1988. Amino acid substitutions in structurally related proteins. A pattern recognition approach. Determination of a new and efficient scoring matrix. *J Mol Biol.* 204:1019-29.
- Rodriguez, M.C., Y.B. Xie, H. Wang, K. Collison and D.L. Segaloff. 1992. Effects of truncations of the cytoplasmic tail of the luteinizing hormone/chorionic gonadotropin receptor on receptor-mediated hormone internalization. *Mol Endocrinol.* 6:327-36.
- Rodriguez-Boulan, E. and C. Zurzolo. 1993. Polarity signals in epithelial cells. *J Cell Sci Suppl.* 17:9-12.
- Rohl, C.A., W. Fiori and R.L. Baldwin. 1999. Alanine is helix-stabilizing in both template-nucleated and standard peptide helices. *Proc Natl Acad Sci U S A.* 96:3682-7.

- Roth, A., H.J. Kreienkamp, W. Meyerhof and D. Richter. 1997. Phosphorylation of four amino acid residues in the carboxyl terminus of the rat somatostatin receptor subtype 3 is crucial for its desensitization and internalization. *J Biol Chem.* 272:23769-74.
- Rothman, J.E. 1994. Mechanisms of intracellular protein transport. *Nature.* 372:55-63.
- Rozell, T.G., D.P. Davis, Y. Chai and D.L. Segaloff. 1998. Association of gonadotropin receptor precursors with the protein folding chaperone calnexin. *Endocrinology.* 139:1588-93.
- Rudiger, S., A. Buchberger and B. Bukau. 1997. Interaction of Hsp70 chaperones with substrates. *Nat Struct Biol.* 4:342-9.
- Sadeghi, H., and M. Birnbaumer. 1999. O-Glycosylation of the V2 vasopressin receptor. *Glycobiology.* 9:731-7.
- Sadeghi, H.M., G. Innamorati and M. Birnbaumer. 1997. Maturation of receptor proteins in eukaryotic expression systems. *J Recept Signal Transduct Res.* 17:433-45.
- Sadeghi, H.M., G. Innamorati and M. Birnbaumer. 1997. An X-linked NDI mutation reveals a requirement for cell surface V2R expression. *Mol Endocrinol.* 11:706-13.
- Sadeghi, H.M., G. Innamorati, M. Dagarag and M. Birnbaumer. 1997. Palmitoylation of the V2 vasopressin receptor. *Mol Pharmacol.* 52:21-9.
- Sakmar, T.P., S.T. Menon, E.P. Marin and E.S. Awad. 2002. Rhodopsin: insights from recent structural studies. *Annu Rev Biophys Biomol Struct.* 31:443-84.
- Saliba, R.S., P.M. Munro, P.J. Luthert and M.E. Cheetham. 2002. The cellular fate of mutant rhodopsin: quality control, degradation and aggresome formation. *J Cell Sci.* 115:2907-18.
- Sasakawa, N., M. Sharif and M.R. Hanley. 1994. Attenuation of agonist-induced desensitization of the rat substance P receptor by progressive truncation of the C-terminus. *FEBS Lett.* 347:181-4.
- Saunders, C., J.R. Keefer, C.A. Bonner and L.E. Limbird. 1998. Targeting of G protein-coupled receptors to the basolateral surface of polarized renal epithelial cells involves multiple, non-contiguous structural signals. *J Biol Chem.* 273:24196-206.

- Saunders, C., J.R. Keefer, A.P. Kennedy, J.N. Wells and L.E. Limbird. 1996. Receptors coupled to pertussis toxin-sensitive G-proteins traffic to opposite surfaces in Madin-Darby canine kidney cells. A1 adenosine receptors achieve apical and alpha 2A adrenergic receptors achieve basolateral localization. *J Biol Chem.* 271:995-1002.
- Scheer, A., F. Fanelli, T. Costa, P.G. De Benedetti and S. Cotecchia. 1996. Constitutively active mutants of the alpha 1B-adrenergic receptor: role of highly conserved polar amino acids in receptor activation. *EMBO J.* 15:3566-3578.
- Scheiffele, P. and J. Fullekrug. 2000. Glycosylation and protein transport. *Essays Biochem.* 36:27-35.
- Schneider, H., J.H. Feyen and K. Seuwen. 1994. A C-terminally truncated human parathyroid hormone receptor is functional and activates multiple G proteins. *FEBS Lett.* 351:281-5.
- Schubert, U., L.C. Anton, J. Gibbs, C.C. Norbury, J.W. Yewdell and J.R. Bennink. 2000. Rapid degradation of a large fraction of newly synthesized proteins by proteasomes. *Nature.* 404:770-4.
- Schülein, R., R. Hermosilla, A. Oksche, M. Dehe, B. Wiesner, G. Krause and W. Rosenthal. 1998. A dileucine sequence and an upstream glutamate residue in the intracellular carboxyl terminus of the vasopressin V2 receptor are essential for cell surface transport in COS.M6 cells. *Mol Pharmacol.* 54:525-35.
- Schülein, R., U. Liebenhoff, H. Müller, M. Birnbaumer and W. Rosenthal. 1996. Properties of the human arginine vasopressin V2 receptor after site-directed mutagenesis of its putative palmitoylation site. *Biochem J.* 313 (Pt 2):611-6.
- Schülein, R., D. Lorenz, A. Oksche, B. Wiesner, R. Hermosilla, J. Ebert and W. Rosenthal. 1998. Polarized cell surface expression of the green fluorescent protein-tagged vasopressin V2 receptor in Madin Darby canine kidney cells. *FEBS Lett.* 441:170-6.
- Schülein, R., C. Rutz and W. Rosenthal. 1996. Membrane targeting and determination of transmembrane topology of the human vasopressin V2 receptor. *J Biol Chem.* 271:28844-52.

- Schülein, R., K. Zühlke, A. Oksche, R. Hermosilla, J. Furkert and W. Rosenthal. 2000. The role of conserved extracellular cysteine residues in vasopressin V2 receptor function and properties of two naturally occurring mutant receptors with additional extracellular cysteine residues. *FEBS Lett.* 466:101-6.
- Sevier, C.S., O.A. Weisz, M. Davies and C.E. Machamer. 2000. Efficient export of the vesicular stomatitis virus G protein from the endoplasmic reticulum requires a signal in the cytoplasmic tail that includes both tyrosine-based and di-acidic motifs. *Mol Biol Cell.* 11:13-22
- Sheikh, H. and C.M. Isacke. 1996. A di-hydrophobic Leu-Val motif regulates the basolateral localization of CD44 in polarized Madin-Darby canine kidney epithelial cells. *J Biol Chem.* 271:12185-90.
- Spiess, M. 1995. Head or tails-what determines the orientation of proteins in the membrane. *FEBS Lett.* 369:76-79.
- Strickland, E., B.H. Qu, L. Millen and P.J. Thomas. 1997. The molecular chaperone Hsc70 assists the in vitro folding of the N-terminal nucleotide-binding domain of the cystic fibrosis transmembrane conductance regulator. *J Biol Chem.* 272:25421-4.
- Thomas, W.G., K.M. Baker, T.J. Motel and T.J. Thekkumkara. 1995. Angiotensin II receptor endocytosis involves two distinct regions of the cytoplasmic tail. A role for residues on the hydrophobic face of a putative amphipathic helix. *J Biol Chem.* 270:22153-9.
- Tohgo, A., E.W. Choy, D. Gesty-Palmer, K.L. Pierce, S. Laporte, R.H. Oakley, M.G. Caron, R.J. Lefkowitz and L.M. Luttrell. 2003. The stability of the G protein-coupled receptor-beta-arrestin interaction determines the mechanism and functional consequence of ERK activation. *J Biol Chem.* 278:6258-67.
- Ulrich, C.D., 2nd, M. Holtmann and L.J. Miller. 1998. Secretin and vasoactive intestinal peptide receptors: members of a unique family of G protein-coupled receptors. *Gastroenterology.* 114:382-97.
- Unger, V.M., P.A. Hargrave, J.M. Baldwin and G.F. Schertler. 1997. Arrangement of rhodopsin transmembrane alpha-helices. *Nature.* 389:203-6.

- Unson, C.G., A.M. Cypess, H.N. Kim, P.K. Goldsmith, C.J. Carruthers, R.B. Merrifield and T.P. Sakmar. 1995. Characterization of deletion and truncation mutants of the rat glucagon receptor. Seven transmembrane segments are necessary for receptor transport to the plasma membrane and glucagon binding. *J Biol Chem.* 270:27720-27.
- VanLeuwen, D., M.E. Steffey, C. Donahue, G. Ho and R.G. MacKenzie. 2003. Cell surface expression of the Melanocortin-4 Receptor is dependent on a C-terminal Di-isoleucine sequence at codons 316/317. *J Biol Chem.* 278:15935-15940.
- Wang, J., L. Wang, J. Zheng, J.L. Anderson and M.L. Toews. 2000. Identification of distinct carboxyl-terminal domains mediating internalization and down-regulation of the hamster alpha(1B)- adrenergic receptor. *Mol Pharmacol.* 57:687-94.
- Warne, T., J. Chirnside and G.F. Schertler. 2003. Expression and purification of truncated, non-glycosylated turkey beta-adrenergic receptors for crystallization. *Biochim Biophys Acta.* 1610:133-40.
- Weiss, E.R., S. Osawa, W. Shi and C.D. Dickerson. 1994. Effects of carboxyl-terminal truncation on the stability and G protein-coupling activity of bovine rhodopsin. *Biochemistry.* 33:7587-93.
- Wieland, F. and C. Harter. 1999. Mechanisms of vesicle formation: insights from the COP system. *Curr Opin Cell Biol.* 11:440-6.
- Wieland, F.T., M.L. Gleason, T.A. Serafini and J.E. Rothman. 1987. The rate of bulk flow .from the endoplasmic reticulum to the cell surface. *Cell.* 50:289-300.
- Willbanks, A.M., S.A. LaPorte, L.M. Bohn, L.S. Barak and M.G. Caron. 2002. Apparent loss-of-function mutant GPCRs revealed as constitutively desensitized receptors. *Biochemistry.* 40:11981-89.
- Wormald, M.R. and R.A. Dwek. 1999. Glycoproteins: glycan presentation and protein-fold stability. *Structure Fold Des.* 7:155-60
- Wozniak, M. and L.E. Limbird. 1996. The three alpha 2-adrenergic receptor subtypes achieve basolateral localization in Madin-Darby canine kidney II cells via different targeting mechanisms. *J Biol Chem.* 271:5017-24.

- Wright, P.E., H.J. Dyson and R.A. Lerner. 1988. Conformation of peptide fragments of proteins in aqueous solution: implications for initiation of protein folding. *Biochemistry*. 27:7167-75.
- Yamamoto, K., R. Fujii, Y. Toyofuku, T. Saito, H. Koseki, V.W. Hsu and T. Aoe. 2001. The KDEL receptor mediates a retrieval mechanism that contributes to quality control at the endoplasmic reticulum. *Embo J*. 20:3082-91.
- Yeagle, P.L., J.L. Alderfer and A.D. Albert. 1997. Three-dimensional structure of the cytoplasmic face of the G protein receptor rhodopsin. *Biochemistry*. 36:9649-54.
- Yeagle, P.L., C. Danis, G. Choi, J.L. Alderfer and A.D. Albert. 2000. Three dimensional structure of the seventh transmembrane helical domain of the G-protein receptor, rhodopsin. *Mol Vis*. 6:125-31.