## **Table of contents**

1 Dire	ctory	1
2 Intro	oduction	2
3 Mate	erials and methods	3
3.1 Det	termination of soil parameters	3
3.1.1	Water content	3
3.1.2	Maximum water holding capacity (MWC)	3
3.1.3	Soil acidity	3
3.1.4	Content of organic matter and organic carbon	4
3.1.5	Grain size	4
3.2 Tes	st organisms	4
3.3 Tes	sts	5
3.3.1	Soil tests	5
3.3.2	Non-soil tests	8
3.3.3	Statistics	9
4 Pure	substances	11
4.1 Ab	stract	11
4.2 The	eoretical background	11
4.2.1	TNT	12
4.2.2	Hexyl	14
4.2.3	Hexogen (RDX)	15
4.2.4	Octogen (HMX)	16
4.2.5	Triaminotoluene (TAT)	17
4.3 Ma	terials and methods	18
4.3.1	Contaminants	18
4.3.2	Tests	19
4.4 Res	sults	20
4.4.1	TNT	20

4.4.2	Hexyl	27
4.4.3	Hexogen (RDX)	36
4.4.4	Octogen (HMX)	38
4.4.5	Triaminotoluene (TAT)	39
4.4.6	Summary of Results	42
4.5 Dis	cussion	43
4.5.1	TNT	43
4.5.2	Hexyl	48
4.5.3	Hexogen (RDX) and Octogen (HMX)	50
4.5.4	TAT	51
4.5.5	Comparison of the toxicity to different organisms	52
4.5.6	Comparison of tests	55
5 Toxi	city of TNT in different reference soil materials	<b>57</b>
5.1 Abs	stract	57
5.2 The	eoretical background	57
5.2.1	Organic matter	57
5.2.2	Clay	58
5.2.3	Sorption to organic matter and clay	58
5.2.4	Sorption of TNT	58
5.3 Ma	terials and methods	59
5.3.1	Reference soil materials	59
5.3.2	Treatment of soil materials and tests	61
5.3.3	Validity of tests and treatment of results	62
5.4 Res	sults	62
5.4.1	Comparison of the uncontaminated reference soil materials	62
5.4.2	Lufa 2.3	63
5.4.3	Lufa 3	64
5.4.4	Lufa 4	64
5.4.5	Potting soil	65
5.4.6	Comparison of the toxicity of TNT in different reference soil materials	65

5.5	Dis	cussion	66
5	.5.1	Comparison of uncontaminated reference soil materials	66
5.	.5.2	Comparison of the toxicity of TNT in the reference soil materials	68
6	Age	ing of TNT-contaminated soil materials	71
6.1	Ab	stract	71
6.2	The	eoretical background	71
6	.2.1	Sorption	72
6	.2.2	Microbial degradation of TNT	72
6.3	Ma	terials and methods	74
6	.3.1	Ageing at different temperatures	74
6.4	Res	sults	75
6	.4.1	Ageing at different temperatures	75
6	.4.2	More detailed assessment of ageing at 20°C	82
6	.4.3	Summary of results	85
6.5	Dis	cussion	86
6	.5.1	Ageing at different temperatures	86
6	.5.2	More detailed assessment of ageing at 20°C	90
7	Con	taminated soil materials	92
7.1	Ab	stract	92
7.2		eoretical background	
	.2.1	"Werk Tanne" Clausthal-Zellerfeld	
	.2.2	Elsnig near Torgau	
	.2.3	Hambühren	
	.2.4	Stadtallendorf	
7.3	Ma	terials and methods	95
	.3.1	Properties of the soil materials	
	.3.2	Analysis of contaminated soil samples	
	.3.3	Evaluation of the toxicity of heavy metals in the contaminated soil materials	
7.4	Res	sults	
	.4.1	"Werk Tanne" Clausthal-Zellerfeld	

7.4.2	Elsnig near Torgau	100
7.4.3	Hambühren	103
7.4.4	Stadtallendorf	103
7.4.5	Toxicity of TNT-contaminated soil materials	104
7.4.6	Summary of results	104
7.5 Dis	scussion	105
7.5.1	Toxicity of the soil materials	106
7.5.2	Comparison of the toxicity of contaminated soil materials	107
7.5.3	Comparison with literature	108
8 Ren	nediation	113
8.1 Ab	stract	113
8.2 Th	eoretical background	113
8.2.1	Physico-chemical remediation	114
8.2.2	Biological methods	115
8.3 Ma	iterials and methods	119
8.3.1	"Werk Tanne" at Clausthal-Zellerfeld: CTNT2, CTNT3, CTNT4	119
8.3.2	Elsnig near Torgau: ETNT	121
8.3.3	Hambühren: LTNT	122
8.4 Re	sults	123
8.4.1	"Werk Tanne" at Clausthal-Zellerfeld	123
8.4.2	Elsnig near Torgau	126
8.4.3	Hambühren	127
8.4.4	Summary of results	129
8.5 Dis	scussion	130
8.5.1	Evaluation of the remediation processes	130
8.5.2	Comparison with literature	135
9 Sun	nmary	138
10 Futu	ure research	140
11 Refu	erences	141

Appe	ndix	i
A D	ata sheets of the reference soil materials	i
A.1	Lufa 2.2	i
A.2	Lufa 2.3	iii
A.3	Lufa 3	v
A.4	Lufa 4	vi
В А	nalysis of aged soil materials	vii
B.1	Soil samples	vii
B.2	Reports	vii
СС	haracteristics of the different soil materials	xi
D O	riginal Data	xii
D.1	Betanal	ii
D.2	Metals in the contaminated and remediated soil materials	xiv
E P	ersonal data	xxiii
E.1	Curriculum Vitae	xxiii
E.2	List of publications	xxiv
E.3	Acknowledgement	xxv
E.4	Erklärung	xxvii

## List of figures

Fig. 4.2-1:	Structural formula of TNT	. 12
Fig. 4.2-2:	Structural formula of Hexyl	. 14
Fig. 4.2-3:	Structural formula of Hexogen	. 15
Fig. 4.2-4:	Structural formula of Octogen	. 16
Fig. 4.2-5:	Structural formula of TAT	. 17
Fig. 4.4-1:	Concentration-effect relationship for TNT in standard soil material Lufa 2.2	. 20
Fig. 4.4-2:	Surviving <i>E. crypticus</i> of a mortality test with various concentrations of TNT-contaminated soil materials.	. 20
Fig. 4.4-3:	Cocoon rates for <i>E. crypticus</i> transferred from different concentrations of TNT-contaminated soil materials onto agar-agar	. 22
Fig 4.4-4::	Hatching rates for the cocoons laid 1-3 days after the transfer of the adult E crypticus from TNT-contaminated soil materials	. 23
Fig. 4.4-5:	Concentration-effect relationship for TNT in the contact test with <i>F. candida</i> and in the water test with <i>E. crypticus</i>	. 23
Fig. 4.4-6:	Distribution of <i>E. crypticus</i> in a separated test set of a choice test with TNT	. 26
Fig. 4.4-7:	Distribution of <i>E. crypticus</i> in a separated test set of a choice test with TNT after 7 days of ageing	. 27
Fig. 4.4-8:	Concentration-effect relationship with Hexyl for <i>F. candida</i> and <i>E. crypticus</i> . The data for mortality and reproduction are given in relation to the values in the standard soil material Lufa 2.2	. 28
Fig. 4.4-9:	Reproduction rates of <i>F. candida</i> transferred from a mortality test in various Hexyl-contaminated soil materials into uncontaminated soil material	. 29
Fig. 4.4-10:	Photos of surviving <i>E. crypticus</i> from a mortality test in various concentrations of Hexyl-contaminated soil materials	. 30
Fig. 4.4-11:	Cocoons laid by worms out of the control and cocoons laid by worms out of Hexyl-contaminated soil material immediately after the transfer	. 30
Fig. 4.4-12:	Cocoon rates for <i>E. crypticus</i> transferred from different concentrations of Hexyl-contaminated soil materials onto agar-agar	. 31
Fig. 4.4-13:	Hatching rates of cocoons laid 1-3 days after the transfer of adult <i>E. crypticus</i> from various Hexyl-contaminated soil materials onto agar-agar	. 32
Fig. 4.4-14:	Hatching rate of cocoons laid 4-6 days after the transfer of adult <i>E. crypticus</i> from various Hexyl-contaminated soil materials onto agar-agar	. 33

Fig. 4.4-15:	Hatching rates of cocoons laid 7-9 days after the transfer of adult <i>E. crypticus</i>
	from various Hexyl-contaminated soil materials onto agar-agar
Fig. 4.4-16:	Distribution of <i>F. candida</i> in a separated test set of a choice test with Hexyl 35
Fig. 4.4-17:	Distribution of <i>E. crypticus</i> in a separated test set of a choice test with Hexyl 36
Fig. 4.4-18:	Concentration-effect relationship for TAT with F. candida and E. crypticus 40
Fig. 4.4-19:	Concentration-effect relationship for TAT in the water tests with F. candida
	and E. crypticus40
Fig. 4.4-20:	Distribution of <i>E. crypticus</i> in a separated test set of a choice test with TAT42
Fig. 5.3-1:	Distribution of grain size in the Lufa soil materials61
Fig. 5.3-2:	Content of organic carbon in the reference soil materials61
Fig. 5.4-1:	Mortality and reproduction rate in different reference soil materials for <i>F. candida</i> and <i>E. crypticus</i> .
Fig. 5.4-2:	Concentration-effect relationship for TNT in the reference soil material Lufa 2.3 with <i>F. candida</i> and <i>E. crypticus</i>
Fig. 5.4-3:	Concentration-effect relationship for TNT in the reference soil material Lufa 3 with <i>F. candida</i> and <i>E. crypticus</i>
Fig. 5.4-4:	Concentration-effect relationship for TNT in the reference soil material Lufa 4 with <i>F. candida</i> and <i>E. crypticus</i>
Fig. 5.4-5:	Concentration-effect relationship for TNT in the potting soil with <i>F. candida</i> and <i>E. crypticus</i>
Fig. 5.4-6:	LC50(7d)- and EC50(28d)-values in the different reference soil materials with  F. candida and E. crypticus
Fig. 6.2-1:	Possible intermediates and condensation products of TNT originating from gratuitous reduction
Fig. 6.4-1:	Reproduction rates for <i>F. candida</i> in soil materials aged at different temperatures for the two lowest tested concentrations
Fig. 6.4-2:	LC50(7d)-values for <i>F. candida</i> in TNT-contaminated soil materials aged at 4°C and 20°C in comparison to unaged soil material indicated by the line 80
Fig. 6.4-3:	Reproduction rates for <i>F. candida</i> in TNT-contaminated soil materials aged at different temperatures for the two lowest tested concentrations
Fig. 6.4-4:	Mortality and reproduction rates for <i>F. candida</i> in TNT-contaminated soil materials aged at 20°C
Fig. 6.4-5:	LC50(7d)- and EC50(28d)-values for <i>F. candida</i> in TNT-contaminated soil materials aged at 20°C85

Fig. 7.4-1:	Concentration-effect relationship with CTNT1a in dilution with the standard soil material Lufa 2.2 for <i>F. candida</i> and <i>E. crypticus</i>	98
Fig. 7.4-2:	Concentration-effect relationship with CTNT2a in dilution with the standard soil material Lufa 2.2 for <i>F. candida</i> and <i>E. crypticus</i>	99
Fig. 7.4-3:	Concentration-effect relationship with CTNT4a in dilution with the standard soil material Lufa 2.2 for <i>F. candida</i> and <i>E. crypticus</i>	.100
Fig. 7.4-4:	Concentration-effect relationship with ETNTa in dilution with the standard soil material Lufa 2.2 for <i>F. candida</i> and <i>E. crypticus</i>	.102
Fig. 7.4-5:	Concentration-effect relationship with LTNT1a for F. candida and E. crypticus	103
Fig. 7.4-6:	LC50(7d)- and EC50(28d)-values for the TNT-contaminated soil materials in dilution with Lufa 2.2 for <i>F. candida</i> and <i>E. crypticus</i> .	.104
Fig. 8.2-1:	Flow chart of a soil washing process by the AB Umwelttechnik GmbH	115
Fig. 8.2-2:	Microbial reduction of TNT	117
Fig. 8.4-1:	Mortality and reproduction rates of <i>F. candida</i> and <i>E. crypticus</i> in the contaminated soil material CTNT2a and the remediated soil material CTNT2aVA from the ammunition plant "Werk Tanne" at Clausthal-Zellerfeld (Lower Saxony, Germany)	.124
Fig. 8.4-2:	Mortality and reproduction rates of <i>F. candida</i> and <i>E. crypticus</i> in the contaminated soil material CTNT2a and the remediated soil material CTNT3b from the ammunition plant "Werk Tanne" at Clausthal-Zellerfeld (Lower Saxony, Germany)	.125
Fig. 8.4-3:	Mortality and reproduction rates of <i>F. candida</i> and <i>E. crypticus</i> in the contaminated soil material CTNT4a and the remediated soil materials CTNT4b, CTNT7b as well as in the uncontaminated soil material CTNT04a before and after the remediation (CTNT04b) from the munition plant "Werk Tanne" (Lower Saxony, Germany)	126
Fig. 8.4-4:	Mortality and reproduction rates of the original soil material LTNTa from Hambühren (Lower Saxony, Germany) and the two soil fractions LTNT1b and LTNT1c after the washing process in the stationary soil washing plant of the	120
	ABU at Lägersdorf	.128
Fig. 8.4-5:	Concentration-effect relationship with LTNT1b for F. candida and E. crypticus	129

## List of tables

Table 3.3-1: Overview of the used statistical tests	9
Table 3.3-2: Overview of the biotest-systems with <i>E. crypticus</i> and <i>F. candida</i>	10
Table 4.3-1: Properties of explosives	18
Table 4.4-1: Cocoon rates for E. crypticus transferred from different concentrations of	
TNT-contaminated soil materials onto agar-agar	21
Table 4.4-2: Hatching rates for the cocoons laid 1-3 days after the transfer of adult	
E. crypticus from different concentrations of TNT-contaminated soil materials	22
Table 4.4-3: LC 50 values of TNT for <i>F. candida</i> in the contact test and <i>E. crypticus</i> in the	
water test after 24, 48 and 72 hours	24
Table 4.4-4: Distribution of <i>F. candida</i> in a separated test set with TNT	25
Table 4.4-5: Distribution of <i>E. crypticus</i> in a separated test set with TNT	25
Table 4.4-6: Distribution of <i>E. crypticus</i> in a separated test set with TNT after an ageing	
period of 7 days	26
Table 4.4-7: Mortality for <i>F. candida</i> with Hexyl	28
Table 4.4-8: Number of adults transferred from the Hexyl-contaminated soil materials onto	
the uncontaminated soil material, number of adults found after a test period	
of three weeks and the reproduction rate after 3 weeks	29
Table 4.4-9: Cocoon rate for <i>E. crypticus</i> transferred from different concentrations of	
Hexyl-contaminated soil materials onto agar-agar	31
Table 4.4-10: Hatching rates for the cocoons laid 1-3 days after the transfer of adult	
E. crypticus from different concentrations of Hexyl-contaminated soil	
materials	32
Table 4.4-11: Hatching rates for the cocoons laid 4-6 days after the transfer of adult	
E crypticus from different concentrations of Hexyl-contaminated soil	00
materials	33
Table 4.4-12: Hatching rates for the cocoons laid 7-9 days after the transfer of adult	
E. crypticus from different concentrations of Hexyl-contaminated soil materials	33
Table 4.4-13: Distribution of <i>F. candida</i> in a separated test set with Hexyl	
Table 4.4-14: Distribution of <i>E. crypticus</i> in a separated test set with Hexyl	
Table 4.4-15: Mortalitly and reproduction rates for <i>F. candida</i> and <i>E. crypticus</i> with Hexogen	
Table T.T- 10. Mortality and reproduction rates for 1. Candida and L. Crypticus With Nexogen	01

Table 4.4-16:	Distribution of <i>F. candida</i> and <i>E. crypticus</i> in a combined test set with	
	Hexogen	37
Table 4.4-17:N	Mortality and reproduction rates for F. candida and E. crypticus with Octogen	38
Table 4.4-18:[	Distribution of F. candida and E. crypticus in a combined test set with	
	Octogen	38
Table 4.4-19:N	Mortality and reproduction rates for <i>F. candida</i> and <i>E. crypticus</i> with TAT	39
Table 4.4-20:L	LC 50 values of TAT for <i>F. candida</i> and <i>E. crypticus</i> in the water test after 24,	
	48 and 72 hours	
Table 4.4-21:[	Distribution of <i>F. candida</i> in a separated test set with TAT	41
Table 4.4-22:[	Distribution of <i>E. crypticus</i> in a separated test set with TAT	41
Table 4.4-23:5	Summary of the results with the soil and non-soil tests	43
Table 4.5-1: (	Overview of the toxicities of TNT, Hexyl, Hexogen, Octogen and TAT to different test organisms in respect to mortality and reproduction	54
Table 4.5-2: (	Comparison of the sensitivity of the tests for the two test organisms for the tested explosives and TAT	56
Table 5.3-1: (	Comparison of chemical and physical properties of the different reference soil materials with the usual standard soil material Lufa 2.2 indicated by shading	60
Table 5.4-1:(	Comparison of the mortality and the reproduction rate in uncontaminated reference soil materials with the usual standard soil material Lufa 2.2 indicated by shading	62
Table 5.4-2: L	LC50(7d)- and EC50(28d)-values for TNT in different reference soil materials for <i>F. candida</i> and <i>E. crypticus</i> in comparison to the usually used standard soil material Lufa 2.2 indicated by shading	65
Table 5.5-1: §	Soil properties of different uncontaminated soil materials and mean of the reproduction rate in percent in relation to the usual standard soil material Lufa 2.2 for <i>F. candida</i> and <i>E. crypticus</i>	68
Table 6.4-1:(	Concentrations of TNT and the ADNTs detected in the spiked soil samples after different ageing periods	75
Table 6.4-2: (	Concentrations of TNT and the ADNTs detected in the spiked soil samples after different ageing periods at the beginning and the end of the reproduction test	76
Table 6.4-3: N	Mortality for <i>F. candida</i> with TNT-contaminated soil material stored at 4°C after different periods of ageing	

Table 6.4-4:	Mortality for <i>F. candida</i> with TNT-contaminated soil materials stored at 20°C	70
	after different periods of ageing	78
Table 6.4-5:	LC50(7d)-values for <i>F. candida</i> in TNT-contaminated soil materials aged at 4°C and 20°C in comparison to the one without ageing indicated by shading7	79
Table 6 4 6:		•
1 able 6.4-6:	Reproduction rates for <i>F. candida</i> with TNT-contaminated soil materials stored at 4°C after different periods of ageing	30
Table 6.4-7:	Reproduction rates for <i>F. candida</i> with TNT-contaminated soil materials	
	stored at 20°C after different periods of ageing	31
Table 6.4-8:	Mortality for <i>F. candida</i> with TNT-contaminated soil materials stored at 20°C	
	after different periods of ageing	32
Table 6.4-9:	Reproduction rates for <i>F. candida</i> with TNT-contaminated soil materials stored at 20°C after different periods of ageing	33
Table 6 4-10	:LC50(7d)-and EC50(28d)-values for <i>F. candida</i> in TNT-contaminated soil	
14516 6.1 16	materials aged at 20°C in comparison to the one without ageing indicated by	
	shading	35
Table 6.4-11	:LC50(7d)- and EC50(28d)-values for <i>F. candida</i> determined in both test sets	35
Table 7.3-1:	Soil Parameters and contaminations above the German intervention values	
	for the contaminated soils materials from the four sites	96
Table 7.3-2:	Reference values and the soil reference value formulas for some metals	97
Table 7.3-3:	Calculated EC50-values for some metals in the tested soil materials	97
Table 7.4-1:	Mortality and reproduction rate with CTNT1 for F. candida and E. crypticus in	
	relation to control with Lufa 2.2	98
Table 7.4-2:	Mortality and reproduction rate with CTNT2a for <i>F. candida</i> and <i>E. crypticus</i> in relation to control with Lufa 2.2	20
		19
Table 7.4-3:	Mortality and reproduction rate with CTNT4a for <i>F. candida</i> and <i>E. crypticus</i> in relation to control with Lufa 2.2	00
Table 7.4-4:	Mortality and reproduction rate of TNT in combination with other explosives	
	for F. candida and E. crypticus in relation to control with Lufa 2.2	)1
Table 7.4-5:	Mortality of the LC10(7d) of TNT in combination with different Hexyl-	
	concentrations for <i>F. candida</i> and <i>E. crypticus</i> in relation to control with Lufa	
	2.2	J1
Table 7.4-6:	Mortality and reproduction rate with ETNTa for <i>F. candida</i> and <i>E. crypticus</i> in relation to control with Lufa 2.2	าว
	101011011 tO 00111101 WILLI EUIG E.E	

Table 7.4-7:	Mortality and reproduction rate with LTNT1a for <i>F. candida</i> and <i>E. crypticus</i> in relation to control with Lufa 2.2	.103
Table 7.4-8:	Mortality and reproduction rate with STNTa for <i>F. candida</i> and <i>E. crypticus</i> in relation to control with Lufa 2.2	.104
Table 7.4-9:	Toxicity of the TNT-contaminated soil materials in reference to their LC50(7d)- and EC50(28d)-values in dilution with Lufa 2.2	.104
Table 7.4-10	:Toxicity criteria for the evaluation of soils in relation to a control with the reference soil material Lufa 2.2	.105
Table 7.4-11	:Classification of the contaminated soil materials according to the guidelines established for contaminated soils	.105
Table 7.5-1:	Toxicity of TNT-contaminated soil materials form various sites at JAAP	.110
Table 7.5-2:	Evaluation of the tested soil materials with other biotests; the results with the collembola and enchytraeid-biotest are given as a comparison and indicated by shading	112
Table 8.3-1:	Soil properties and contaminations above the German intervention values for the contaminated and the remediated soil materials from the ammuntion plant Werk Tanne at Clausthal-Zellerfeld (Lower Saxony, Germany)	
Table 8.3-2:	Soil properties and contaminations above the German intervention values for the contaminated and remediated soil materials from the open-burning site in Elsnig near Torgau (Saxony, Germany)	.122
Table 8.3-3:	Soil properties and contaminations above the German intervention values for the contaminated and remediated soil materials from the washing process at Lägersdorf (Schleswig-Holstein, Germany)	.123
Table 8.4-1:	Mortality and reproduction rate for <i>F. candida</i> and <i>E. crypticus</i> in CTNT2a and the remediated soil material CTNT2aVA in relation to control with Lufa 2.2	.124
Table 8.4-2:	Mortality and reproduction rate for <i>F. candida</i> and <i>E. crypticus</i> in CTNT2a and the remediated soil material CTNT 3b in relation to control with Lufa 2.2	.124
Table 8.4-3:	Mortality and reproduction rate for <i>F. candida</i> and <i>E. crypticus</i> in CTNT4a, the remediated soil materials CTNT4b, CTNT7b and the uncontaminated soil materials CTNT04a and CTNT04b before and after the remediation in relation to control with Lufa 2.2	.126
Table 8.4-4:	Reproduction rates for <i>F. candida</i> and <i>E. crypticus</i> in the ETNTa and the remediated soil materials ETNTb1% and ETNTb2% in relation to control with Lufa 2.2	127

Table 8.4-5:	Mortality and reproduction rate for <i>F. candida</i> and <i>E. crypticus</i> with the additives of the remediation process for ETNTa	127
Table 8.4-6:	Mortality and reproduction rate for <i>F. candida</i> and <i>E. crypticus</i> in LTNT1a and in the fractions of the washing process LTNT1b and LTNT1c in relation to control with Lufa 2.2	128
Table 8.4-7:	Toxicity of the contaminated and remediated soil materials according to the guideline established for contaminated soils	129
Table 8.5-1:	Evaluation of the tested contaminated and remediated soil materials with other biotests: results with the collembola- and enchytraeid-biotest are given as a comparison and indicated by shading	137
Table-C-1:	Characteristics of the different contaminated and remediated soil materials	xi
Table D-1:	Original Data of the contact test with <i>F. candida</i> for Betanal	xii
Table D-2:	Original Data of the water test with <i>E.crypticus</i> for Betanal	. xiii
Table D-3:	Calculation of standardised EC50 for <i>F. candida</i> in Cd-contaminated soil materials	.xiv
Table-D-4:	Calculation of EC50-values of Cd in different contaminated and remediated soil materials for <i>F. candida</i>	.xiv
Table D-5:	Calculation of standardised EC50 for <i>E. crypticus</i> in Cd-contaminated soil materials	xv
Table D-6:	Calculation of EC50-values of Cd in different contaminated and remediated soil materials for <i>E. crypticus</i>	xv
Table D-7:	Calculation of standardised EC50 for <i>F. candida</i> in Cu-contaminated soil materials	.xvi
Table D-8:	Calculation of EC50-values of Cu in different contaminated and remediated soil materials for <i>F. candida</i>	.xvi
Table D-9:	Calculation of standardised EC50 for <i>E. crypticus</i> in Cu-contaminated soil materials	xvii
Table D-10:	Calculation of EC50-values of Cu in different contaminated and remediated soil materials for <i>E. crypticus</i>	xvii
Table D-11:	Calculation of standardised EC50 for <i>F. candida</i> in Pb-contaminated soil materials	xviii
Table D-12:	Calculation of EC50-values of Pb in different contaminated and remediated soil materials for <i>F. candida</i>	xviii

Table D-13:	Calculation of standardised EC50 for <i>E. crypticus</i> in Pb-contaminated soil	
	materials	. xix
Table D-14:	Calculation of EC50-values of Pb in different contaminated and remediated soil materials for <i>E. crypticus</i>	. xix
Table D-15:	Calculation of standardised EC50 for <i>F. candida</i> in Zn-contaminated soil materials	xx
Table D-16:	Calculation of EC50-values of Zn in different contaminated and remediated soil materials for <i>F. candida</i>	xx
Table D-17:	Calculation of standardised EC50 for <i>E. crypticus</i> in Zn-contaminated soil materials	. xxi
Table D-18:	Calculation of EC50-values of Zn in different contaminated and remediated soil materials for <i>E. crypticus</i>	. xxi