

## **6 LITERATURVERZEICHNIS**

---

## 6 LITERATURVERZEICHNIS

- Ademola, J.I., Wester, R.C. and Maibach, H.I. (1992) Cutaneous metabolism of theophylline by the human skin. *J Invest Dermatol*, **98**, 310-314.
- Aeby, P., Wyss, C., Beck, H., Griem, P., Scheffler, H. and Goebel, C. (2004) Characterization of the sensitizing potential of chemicals by in vitro analysis of dendritic cell activation and skin penetration. *J Invest Dermatol*, **122**, 1154-1164.
- Ahmed, S., Imai, T. and Otagiri, M. (1996) Evaluation of stereoselective transdermal transport and concurrent cutaneous hydrolysis of several ester prodrugs of propranolol: mechanism of stereoselective permeation. *Pharm Res*, **13**, 1524-1529.
- Al Masaoudi, T., Sieben, S. and Callahan, C.P. (2001) Eugenol but not isoeugenol induces CYP1A1 mRNA in human keratinocytes. *International archives of allergy and immunology*, **124**, 309-311.
- Allombert-Blaise, C., Tamiji, S., Mortier, L., Fauvel, H., Tual, M., Delaporte, E., Piette, F., DeLassale, E.M., Formstecher, P., Marchetti, P. and Polakowska, R. (2003) Terminal differentiation of human epidermal keratinocytes involves mitochondria- and caspase-dependent cell death pathway. *Cell Death Differ*, **10**, 850-852.
- Amato, M., Isenschmid, M. and Huppi, P. (1991) Percutaneous caffeine application in the treatment of neonatal apnoea. *Eur J Pediatr*, **150**, 592-594.
- An, N.M., Kim, D.D., Shin, Y.H. and Lee, C.H. (2003) Development of a novel soft hydrogel for the transdermal delivery of testosterone. *Drug Dev Ind Pharm*, **29**, 99-105.
- Anderson, B.J., Gunn, T.R., Holford, N.H. and Johnson, R. (1999) Caffeine overdose in a premature infant: clinical course and pharmacokinetics. *Anaesth Intensive Care*, **27**, 307-311.
- Arnaud, M.J. (1985) Comparative metabolic disposition of [1-Me<sup>14</sup>C]caffeine in rats, mice, and Chinese hamsters. *Drug Metab Dispos*, **13**, 471-478.
- Asselineau, D., Bernhard, B., Bailly, C. and Darmon, M. (1985) Epidermal morphogenesis and induction of the 67 kD keratin polypeptide by culture of human keratinocytes at the liquid-air interface. *Exp Cell Res*, **159**, 536-539.
- Bando, H., Mohri, S., Yamashita, F., Takakura, Y. and Hashida, M. (1997) Effects of skin metabolism on percutaneous penetration of lipophilic drugs. *J Pharm Sci*, **86**, 759-761.
- Baron, J.M., Goh, L.B., Yao, D., Wolf, C.R. and Friedberg, T. (2001a) Modulation of P450 CYP3A4-dependent metabolism by P-glycoprotein: implications for P450 phenotyping. *J Pharmacol Exp Ther*, **296**, 351-358.
- Baron, J.M., Holler, D., Schiffer, R., Frankenberg, S., Neis, M., Merk, H.F. and Jugert, F.K. (2001b) Expression of multiple cytochrome p450 enzymes and multidrug resistance-associated transport proteins in human skin keratinocytes. *J Invest Dermatol*, **116**, 541-548.
- Barrett, D.A. and Rutter, N. (1994) Transdermal delivery and the premature neonate. *Crit Rev Ther Drug Carrier Syst*, **11**, 1-30.
- Bartek, M.J., LaBudde, J.A. and Maibach, H.I. (1972) Skin permeability in vivo: comparison in rat, rabbit, pig and man. *J Invest Dermatol*, **58**, 114-123.

- Barth, J., Lehr, K.H., Derendorf, H., Mollmann, H.W., Hohler, T. and Hochhaus, G. (1993) Studies on the pharmacokinetics and metabolism of prednicarbate after cutaneous and oral administration. *Skin Pharmacol*, **6**, 179-186.
- Baskettter, D.A., Pendlington, R.U. and Sarll, A. (1996) The role of P450 1A in the activation of prohaptens in skin sensitization. *Journal of Investigative Dermatology*, **106**, 915.
- Bauerova, K., Matusova, D. and Kassai, Z. (2001) Chemical enhancers for transdermal drug transport. *Eur J Drug Metab Pharmacokinet*, **26**, 85-94.
- Baynes, R.E., Halling, K.B. and Riviere, J.E. (1997) The influence of diethyl-m-toluamide (DEET) on the percutaneous absorption of permethrin and carbaryl. *Toxicol Appl Pharmacol*, **144**, 332-339.
- Bell, E., Ivarsson, B. and Merrill, C. (1979) Production of a tissue-like structure by contraction of collagen lattices by human fibroblasts of different proliferative potential in vitro. *Proc Natl Acad Sci U S A*, **76**, 1274-1278.
- Benfeldt, E. (1999) In vivo microdialysis for the investigation of drug levels in the dermis and the effect of barrier perturbation on cutaneous drug penetration. Studies in hairless rats and human subjects. *Acta Derm Venereol Suppl (Stockh)*, **206**, 1-59.
- Bertaux, B., Morliere, P., Moreno, G., Courtalon, A., Masse, J.M. and Dubertret, L. (1988) Growth of melanocytes in a skin equivalent model in vitro. *Br J Dermatol*, **119**, 503-512.
- Bertin, C., Zunino, H., Pittet, J.C., Beau, P., Pineau, P., Massonneau, M., Robert, C. and Hopkins, J. (2001) A double-blind evaluation of the activity of an anti-cellulite product containing retinol, caffeine, and ruscogenine by a combination of several non-invasive methods. *J Cosmet Sci*, **52**, 199-210.
- Bodde, H.E., Holman, B., Spies, F., Weerheim, A., Kempenaar, J., Mommaas, M. and Ponec, M. (1990) Freeze-fracture electron microscopy of in vitro reconstructed human epidermis. *J Invest Dermatol*, **95**, 108-116.
- Boderke, P., Schittkowski, K., Wolf, M. and Merkle, H.P. (2000) Modeling of diffusion and concurrent metabolism in cutaneous tissue. *J Theor Biol*, **204**, 393-407.
- Boehnlein, J., Sakr, A., Lichtin, J.L. and Bronaugh, R.L. (1994) Characterization of esterase and alcohol dehydrogenase activity in skin. Metabolism of retinyl palmitate to retinol (vitamin A) during percutaneous absorption. *Pharm Res*, **11**, 1155-1159.
- Boelsma, E., Gibbs, S., Faller, C. and Ponec, M. (2000) Characterization and comparison of reconstructed skin models: morphological and immunohistochemical evaluation. *Acta Derm Venereol*, **80**, 82-88.
- Boogaard, P.J., van Elburg, P.A., de Kloe, K.P., Watson, W.P. and van Sittert, N.J. (1999) Metabolic inactivation of 2-oxiranylmethyl 2-ethyl-2,5-dimethylhexanoate (C10GE) in skin, lung and liver of human, rat and mouse. *Xenobiotica*, **29**, 987-1006.
- Bosman, I.J., Avegaart, S.R., Lawant, A.L., Ensing, K. and de Zeeuw, R.A. (1998) Evaluation of a novel diffusion cell for in vitro transdermal permeation: effects of injection height, volume and temperature. *J Pharm Biomed Anal*, **17**, 493-499.
- Bowen, A.R., Hanks, A.N., Allen, S.M., Alexander, A., Diedrich, M.J. and Grossman, D. (2003) Apoptosis regulators and responses in human melanocytic and keratinocytic cells. *J Invest Dermatol*, **120**, 48-55.
- Boyce, S., Michel, S., Reichert, U., Shroot, B. and Schmidt, R. (1990) Reconstructed skin from cultured human keratinocytes and fibroblasts on a collagen-glycosaminoglycan biopolymer substrate. *Skin Pharmacol*, **3**, 136-143.

- Bronaugh, R.L., Stewart, R.F. and Simon, M. (1986) Methods for in vitro percutaneous absorption studies. VII: Use of excised human skin. *J Pharm Sci*, **75**, 1094-1097.
- Bronaugh, R.L., Stewart, R.F. and Storm, J.E. (1989) Extent of cutaneous metabolism during percutaneous absorption of xenobiotics. *Toxicol Appl Pharmacol*, **99**, 534-543.
- Brunskill, N.J. (1998) Molecular interactions between albumin and proximal tubular cells. *Exp Nephrol*, **6**, 491-495.
- Casciola-Rosen, L.A., Anhalt, G. and Rosen, A. (1994) Autoantigens targeted in systemic lupus erythematosus are clustered in two populations of surface structures on apoptotic keratinocytes. *J Exp Med*, **179**, 1317-1330.
- Castro-Munozledo, F., Hernandez-Quintero, M., Marsch-Moreno, M. and Kuri-Harcuch, W. (1997) Cultivation, serial transfer, and differentiation of epidermal keratinocytes in serum-free medium. *Biochem Biophys Res Commun*, **236**, 167-172.
- Caubet, M.S., Elbast, W., Dubuc, M.C. and Brazier, J.L. (2002) Analysis of urinary caffeine metabolites by HPLC-DAD: the use of metabolic ratios to assess CYP1A2 enzyme activity. *J Pharm Biomed Anal*, **27**, 261-270.
- Christensen, E.I. and Birn, H. (2001) Megalin and cubilin: synergistic endocytic receptors in renal proximal tubule. *Am J Physiol Renal Physiol*, **280**, F562-573.
- COLIPA. (1995) Cosmetic Ingredients: Guidelines for Percutaneous Absorption / Penetration. COLIPA.
- Collier, S.W., Sale, M.E., Gastonguay, M.R., Bronaugh, R.L. and Schwartz, S.L. (1993) Role of stratum corneum, viable epidermis and saturable metabolism on percutaneous penetration of phenylazophenol. *Pharmaceutical Research*, **10 (Suppl.)**, S412.
- Collier, S.W., Sheikh, N.M., Sakr, A., Lichtin, J.L., Stewart, R.F. and Bronaugh, R.L. (1989) Maintenance of skin viability during in vitro percutaneous absorption/metabolism studies. *Toxicol Appl Pharmacol*, **99**, 522-533.
- Cooper, E.R. and Berner, B. (1985) *Methods In Skin Research*. John Wiley and Sons.
- Coulomb, B., Lebreton, C. and Dubertret, L. (1989) Influence of human dermal fibroblasts on epidermalization. *J Invest Dermatol*, **92**, 122-125.
- Cozzani, E., Cacciapuoti, M., Parodi, A., Ghoshestani, R. and Rebora, A. (2000) Desmosomes and their autoimmune pathologies. *Eur J Dermatol*, **10**, 255-261.
- Cross, S.E., Magnusson, B.M., Winckle, G., Anissimov, Y. and Roberts, M.S. (2003) Determination of the effect of lipophilicity on the in vitro permeability and tissue reservoir characteristics of topically applied solutes in human skin layers. *J Invest Dermatol*, **120**, 759-764.
- de Heer, C., Bosman-Hoefakker, S. and Hakkert, B.C. (1999) Principles for study protocols addressing the dermal absorption of pesticides. *TNO report, V98.356*.
- Dias, M., Farinha, A., Faustino, E., Hadgraft, J., Pais, J. and Toscano, C. (1999) Topical delivery of caffeine from some commercial formulations. *Int J Pharm*, **182**, 41-47.
- Diembeck, W., Beck, H., Benech-Kieffer, F., Courtellemont, P., Dupuis, J., Lovell, W., Paye, M., Spengler, J. and Steiling, W. (1999) Test guidelines for in vitro assessment of dermal absorption and percutaneous penetration of cosmetic

- ingredients. European Cosmetic, Toiletry and Perfumery Association. *Food Chem Toxicol*, **37**, 191-205.
- Doucet, O., Garcia, N., Rosdy, M., Fartasch, M. and Zastrow, L. (1997) Critical events in the barrier development of reconstituted epidermis. *Perspectives in Percutaneous Penetration*, **5B**, 141-144.
- Dreher, F., Fouchard, F., Patouillet, C., Andrian, M., Simonnet, J.T. and Benech-Kieffer, F. (2002a) Comparison of cutaneous bioavailability of cosmetic preparations containing caffeine or alpha-tocopherol applied on human skin models or human skin ex vivo at finite doses. *Skin Pharmacol Appl Skin Physiol*, **15 Suppl 1**, 40-58.
- Dreher, F., Patouillet, C., Fouchard, F., Zanini, M., Messager, A., Roguet, R., Cottin, M., Leclaire, J. and Benech-Kieffer, F. (2002b) Improvement of the experimental setup to assess cutaneous bioavailability on human skin models: dynamic protocol. *Skin Pharmacol Appl Skin Physiol*, **15 Suppl 1**, 31-39.
- Drexler, H.G., Quentmeier, H., Dirks, W.G. and MacLeod, R.A. (2002) Bladder carcinoma cell line ECV304 is not a model system for endothelial cells. *In Vitro Cell Dev Biol Anim*, **38**, 185-186; author reply 187.
- Du, L., Neis, M.M., Ladd, P.A., Lanza, D.L., Yost, G.S. and Keeney, D.S. (2005) Effects of the differentiated keratinocyte phenotype on expression levels of CYP1-4 family genes in human skin cells. *Toxicol Appl Pharmacol*.
- Duvall, E., Wyllie, A.H. and Morris, R.G. (1985) Macrophage recognition of cells undergoing programmed cell death (apoptosis). *Immunology*, **56**, 351-358.
- EC. (1998) EC Directive concerning the placing of biocidal products on the market (L123/1-63).
- EC. (2003a) 7th Amendment to the EC Cosmetic Directive (L66/26-35).
- EC. (2003b) Amendment to the EC Directive relating to medicinal products for human use (L159/46-94).
- Eckert, R.L., Yaffe, M.B., Crish, J.F., Murthy, S., Rorke, E.A. and Welter, J.F. (1993) Involucrin--structure and role in envelope assembly. *J Invest Dermatol*, **100**, 613-617.
- Eckhart, L., Declercq, W., Ban, J., Rendl, M., Lengauer, B., Mayer, C., Lippens, S., Vandenabeele, P. and Tschachler, E. (2000) Terminal differentiation of human keratinocytes and stratum corneum formation is associated with caspase-14 activation. *J Invest Dermatol*, **115**, 1148-1151.
- EHSCLC. (2002) Guidelines for analysis of samples from pharmacokinetic studies - Method Validation. *Guidelines of European Horserace Scientific Liaison Committee*.
- Elias, P.M. and Menon, G.K. (1991) Structural and lipid biochemical correlates of the epidermal permeability barrier. *Adv Lipid Res*, **24**, 1-26.
- Erkan, E., De Leon, M. and Devarajan, P. (2001) Albumin overload induces apoptosis in LLC-PK(1) cells. *Am J Physiol Renal Physiol*, **280**, F1107-1114.
- Fartasch, M., Bassukas, I.D. and Diepgen, T.L. (1993) Structural relationship between epidermal lipid lamellae, lamellar bodies and desmosomes in human epidermis: an ultrastructural study. *Br J Dermatol*, **128**, 1-9.
- Fasano, W.J., Manning, L.A. and Green, J.W. (2002) Rapid integrity assessment of rat and human epidermal membranes for in vitro dermal regulatory testing: correlation of electrical resistance with tritiated water permeability. *Toxicol In Vitro*, **16**, 731-740.
- Feldmann, R.J. and Maibach, H.I. (1969) Percutaneous penetration of steroids in man. *J Invest Dermatol*, **52**, 89-94.

- Fentem, J.H. and Botham, P.A. (2002) ECVAM's activities in validating alternative tests for skin corrosion and irritation. *Altern Lab Anim*, **30 Suppl 2**, 61-67.
- Finnen, M.J., Herdman, M.L. and Shuster, S. (1985) Distribution and sub-cellular localization of drug metabolizing enzymes in the skin. *Br J Dermatol*, **113**, 713-721.
- Forestier, J.P. (1992) Les enzymes de l'espace extra cellulaire du stratum corneum. *International Journal of Cosmetic Science*, **14**, 47-63.
- Frantz, S.W., Ballantyne, B., Beskitt, J.L., Tallant, M.J. and Greco, R.J. (1995) Pharmacokinetics of 2-ethyl-1,3-hexanediol. III. In vitro skin penetration comparisons using the excised skin of humans, rats, and rabbits. *Fundam Appl Toxicol*, **28**, 1-8.
- Franz, T.J. (1975) Percutaneous absorption on the relevance of in vitro data. *J Invest Dermatol*, **64**, 190-195.
- Freeman, A.E., Igel, H.J., Herrman, B.J. and Kleinfeld, K.L. (1976) Growth and characterization of human skin epithelial cell cultures. *In Vitro*, **12**, 352-362.
- Gandarillas, A., Goldsmith, L.A., Gschmeissner, S., Leigh, I.M. and Watt, F.M. (1999) Evidence that apoptosis and terminal differentiation of epidermal keratinocytes are distinct processes. *Exp Dermatol*, **8**, 71-79.
- Gawkrodger, D.J. (1995) *Mikroanatomie der Haut*. Churchill Livingstone, Stuttgart.
- Gazel, A., Ramphal, P., Rosdy, M., De Wever, B., Tornier, C., Hosein, N., Lee, B., Tomic-Canic, M. and Blumenberg, M. (2003) Transcriptional profiling of epidermal keratinocytes: comparison of genes expressed in skin, cultured keratinocytes, and reconstituted epidermis, using large DNA microarrays. *J Invest Dermatol*, **121**, 1459-1468.
- Georgia, K.A., Samanidou, V.F. and Papadoyannis, I.N. (2001) Use of novel solid-phase extraction sorbent materials for high-performance liquid chromatography quantitation of caffeine metabolism products methylxanthines and methyluric acids in samples of biological origin. *J Chromatogr B Biomed Sci Appl*, **759**, 209-218.
- Gibbs, S., Boelsma, E., Kempenaar, J. and Ponec, M. (1998) Temperature-sensitive regulation of epidermal morphogenesis and the expression of cornified envelope precursors by EGF and TGF alpha. *Cell Tissue Res*, **292**, 107-114.
- Gibbs, S., Vicanova, J., Bouwstra, J., Valstar, D., Kempenaar, J. and Ponec, M. (1997) Culture of reconstructed epidermis in a defined medium at 33 degrees C shows a delayed epidermal maturation, prolonged lifespan and improved stratum corneum. *Arch Dermatol Res*, **289**, 585-595.
- Gilchrest, B.A., Nemore, R.E. and Maciag, T. (1980) Growth of human keratinocytes on fibronectin -coated plates. *Cell Biol Int Rep*, **4**, 1009-1016.
- Gniadecki, R., Gajkowska, B., Bartosik, J., Hansen, M. and Wulf, H.C. (1998) Variable expression of apoptotic phenotype in keratinocytes treated with ultraviolet radiation, ceramide, or suspended in semisolid methylcellulose. *Acta Derm Venereol*, **78**, 248-257.
- Gorodischer, R., Zmora, E., Ben-Zvi, Z., Warszwaski, D., Yaari, A., Sofer, S. and Arnaud, M.J. (1986) Urinary metabolites of caffeine in the premature infant. *Eur J Clin Pharmacol*, **31**, 497-499.
- Goto, S., Uchida, T., Lee, C.K., Yasutake, T. and Zhang, J.B. (1993) Effect of various vehicles on ketoprofen permeation across excised hairless mouse skin. *J Pharm Sci*, **82**, 959-963.
- Gysler, A., Kleuser, B., Sippl, W., Lange, K., Korting, H.C., Holtje, H.D. and Korting, H.C. (1999) Skin penetration and metabolism of topical glucocorticoids in

- reconstructed epidermis and in excised human skin. *Pharm Res*, **16**, 1386-1391.
- Gysler, A., Lange, K., Korting, H.C. and Schafer-Korting, M. (1997) Prednicarbate biotransformation in human foreskin keratinocytes and fibroblasts. *Pharm Res*, **14**, 793-797.
- Ha, H.R., Chen, J., Freiburghaus, A.U. and Follath, F. (1995) Metabolism of theophylline by cDNA-expressed human cytochromes P-450. *Br J Clin Pharmacol*, **39**, 321-326.
- Haberland, A., Santos Maia, C., Jores, K., Durrfeld, M., Mehnert, W., Schimke, I., Christ, B. and Schafer-Korting, M. (2003) Albumin effects on drug absorption and metabolism in reconstructed epidermis and excised pig skin. *Altex*, **20**, 3-9.
- Hadgraft, J. (2001) Modulation of the barrier function of the skin. *Skin Pharmacol Appl Skin Physiol*, **14 Suppl 1**, 72-81.
- Hadgraft, J., Walters, K.A. and Guy, R.H. (1992) Epidermal lipids and topical drug delivery. *Semin Dermatol*, **11**, 139-144.
- Han, Y.T., Han, Z.W., Yu, G.Y., Wang, Y.J., Cui, R.Y. and Wang, C.B. (2004) Inhibitory effect of polypeptide from Chlamys farreri on ultraviolet A-induced oxidative damage on human skin fibroblasts in vitro. *Pharmacol Res*, **49**, 265-274.
- Harris, I.R., Siefken, W., Beck-Oldach, K., Brandt, M., Wittern, K.P. and Pollet, D. (2002a) Comparison of activities dependent on glutathione S-transferase and cytochrome P-450 IA1 in cultured keratinocytes and reconstructed epidermal models. *Skin Pharmacol Appl Skin Physiol*, **15 Suppl 1**, 59-67.
- Harris, I.R., Siefken, W., Beck-Oldach, K., Wittern, K.P. and Pollet, D. (2002b) NAD(P)H:quinone reductase activity in human epidermal keratinocytes and reconstructed epidermal models. *Skin Pharmacol Appl Skin Physiol*, **15 Suppl 1**, 68-73.
- Harrison, S.M., Barry, B.W. and Dugard, P.H. (1984) Effects of freezing on human skin permeability. *J Pharm Pharmacol*, **36**, 261-262.
- Hartung, T., Bremer, S., Casati, S., Coecke, S., Corvi, R., Fortaner, S., Gribaldo, L., Halder, M., Hoffmann, S., Roi, A.J., Prieto, P., Sabbioni, E., Scott, L., Worth, A. and Zuang, V. (2004) A modular approach to the ECVAM principles on test validity. *Altern Lab Anim*, **32**, 467-472.
- Henry, G., Li, W., Garner, W. and Woodley, D.T. (2003) Migration of human keratinocytes in plasma and serum and wound re-epithelialisation. *Lancet*, **361**, 574-576.
- Henseleit, U., Rosenbach, T. and Kolde, G. (1996) Induction of apoptosis in human HaCaT keratinocytes. *Arch Dermatol Res*, **288**, 676-683.
- Hewitt, P.G., Perkins, J. and Hotchkiss, S.A. (2000) Metabolism of fluroxypyrr, fluroxypyrr methyl ester, and the herbicide fluroxypyrr methylheptyl ester. I: during percutaneous absorption through fresh rat and human skin in vitro. *Drug Metab Dispos*, **28**, 748-754.
- Heylings, J.R., Van de Sandt, J.J., Gilde, A.J. and Ward, R.J. (1998) Evaluation of SkinEthic human reconstituted epidermis for in vitro percutaneous absorption testing. *Poster presentation: ICT*.
- HHS. (1999) Guidance for Industry: Validation of Analytical Procedures: Methodology. *U.S. Department of Health and Human Services*.
- Hirel, B., Chesne, C., Pailheret, J.P. and Guillouzo, A. (1995) In Vitro expression of drug metabolizing enzyme activities in human adult keratinocytes under

- various culture conditions and their response to inducers. *Toxicology in vitro*, **9**, 49-56.
- Hirvonen, J., Rytting, J.H., Paronen, P. and Urtti, A. (1991) Dodecyl N,N-dimethylamino acetate and azone enhance drug penetration across human, snake, and rabbit skin. *Pharm Res*, **8**, 933-937.
- Hoeller, D., Huppertz, B., Roos, T.C., Poblete Gutierrez, P., Merk, H.F., Frank, J. and Jugert, F.K. (2001) An improved and rapid method to construct skin equivalents from human hair follicles and fibroblasts. *Exp Dermatol*, **10**, 264-271.
- Hofer, T., Gerner, I., Gundert-Remy, U., Liebsch, M., Schulte, A., Spielmann, H., Vogel, R. and Wettig, K. (2004) Animal testing and alternative approaches for the human health risk assessment under the proposed new European chemicals regulation. *Arch Toxicol*.
- Hornstein, O.P. and Nürnberg, E. (1985) *Die Haut als Aufnahme- und Wirkungsort der externen Therapie*. Thieme, Stuttgart.
- Howes, D., Guy, R., Hadgraft, J., Heylings, J.R., Hoeck, U., Kemper, F., Maibach, H., Marty, J.P., Merck, H., Parra, J., Rekkas, D., Rondilli, I., Schäfer, H., Tauber, U. and Verbiese, N. (1996) Methods for assessing percutaneous absorption. The report and recommendations of ECVAM Workshop 13. *ATLA*, **24**, 81-106.
- Hueber, F., Besnard, M., Schaefer, H. and Wepierre, J. (1994) Percutaneous absorption of estradiol and progesterone in normal and appendage-free skin of the hairless rat: lack of importance of nutritional blood flow. *Skin Pharmacol*, **7**, 245-256.
- Huggins, J. (2003) Alternatives to animal testing: research, trends, validation, regulatory acceptance. *Altex*, **20**, 3-61.
- Indans, I. (2002) The use and interpretation of in vitro data in regulatory toxicology: cosmetics, toiletries and household products. *Toxicol Lett*, **127**, 177-182.
- Isseroff, R.R., Martinez, D.T. and Ziboh, V.A. (1985) Alterations in fatty acid composition of murine keratinocytes with in vitro cultivation. *J Invest Dermatol*, **85**, 131-134.
- Jaattela, M. and Tschopp, J. (2003) Caspase-independent cell death in T lymphocytes. *Nat Immunol*, **4**, 416-423.
- Jaeckle, E., Schaefer, U.F. and Loth, H. (2003) Comparison of effects of different ointment bases on the penetration of ketoprofen through heat-separated human epidermis and artificial lipid barriers. *J Pharm Sci*, **92**, 1396-1406.
- Jansen, B.J., van Ruissen, F., Cerneus, S., Cloin, W., Bergers, M., van Erp, P.E. and Schalkwijk, J. (2003) Tumor necrosis factor related apoptosis inducing ligand triggers apoptosis in dividing but not in differentiating human epidermal keratinocytes. *J Invest Dermatol*, **121**, 1433-1439.
- Jugert, F.K., Agarwal, R., Kuhn, A., Bickers, D.R., Merk, H.F. and Mukhtar, H. (1994) Multiple cytochrome P450 isozymes in murine skin: induction of P450 1A, 2B, 2E, and 3A by dexamethasone. *J Invest Dermatol*, **102**, 970-975.
- Kandarova, H., Liebsch, M., Genschow, E., Gerner, I., Traue, D., Slawik, B. and Spielmann, H. (2004) Optimisation of the EpiDerm test protocol for the upcoming ECVAM validation study on in vitro skin irritation tests. *Altex*, **21**, 107-114.
- Kao, J. and Carver, M.P. (1991) *Skin Metabolism*. Hemisphere Publication Corporation, USA.
- Keeney, D.S., Skinner, C., Wei, S., Friedberg, T. and Waterman, M.R. (1998) A keratinocyte-specific epoxygenase, CYP2B12, metabolizes arachidonic acid

- with unusual selectivity, producing a single major epoxyeicosatrienoic acid. *J Biol Chem*, **273**, 9279-9284.
- Kellner, H.M., Eckert, H.G., Fehlhaber, H.W., Hornke, I. and Oekonomopoulos, R. (1986) [Pharmacokinetics and biotransformation following topical use of the local corticoid prednicarbate]. *Z Hautkr*, **61 Suppl 1**, 18-40.
- Kerr, J.F., Wyllie, A.H. and Currie, A.R. (1972) Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br J Cancer*, **26**, 239-257.
- Khan, I.U., Bickers, D.R., Haqqi, T.M. and Mukhtar, H. (1992) Induction of CYP1A1 mRNA in rat epidermis and cultured human epidermal keratinocytes by benz(a)anthracene and beta-naphthoflavone. *Drug Metab Dispos*, **20**, 620-624.
- Kietzmann, M., Loscher, W., Arens, D., Maass, P. and Lubach, D. (1993) The isolated perfused bovine udder as an in vitro model of percutaneous drug absorption. Skin viability and percutaneous absorption of dexamethasone, benzoyl peroxide, and etofenamate. *J Pharmacol Toxicol Methods*, **30**, 75-84.
- Kitagawa, S. and Li, H. (1999) Effects of removal of stratum corneum, delipidization and addition of enhancers, ethanol and l-menthol, on skin permeation of benzoic acid and its 4-n-alkyl substituents in excised guinea pig dorsal skin. *Chem Pharm Bull (Tokyo)*, **47**, 44-47.
- Kitano, Y., Okada, N. and Sasai, S. (1990) Growth of human keratinocytes in a defined medium supplemented with growth factor of serum. *Dermatologica*, **180**, 236-239.
- Kizu, J., Arakawa, M., Arakawa, Y., Komoda, F., Takamizawa, M., Iwata, T., Hayakawa, T. and Imai, K. (1999a) Enhanced theophylline metabolism in patients with bronchial asthma at age 4 and under. *Biomed Chromatogr*, **13**, 462-464.
- Kizu, J., Watanabe, S., Yasuno, N., Arakawa, Y., Uzu, S., Kanda, S., Komoda, F., Iwata, T., Hayakawa, H., Hayakawa, T. and Imai, K. (1999b) Development and clinical application of high performance liquid chromatography for the simultaneous determination of plasma levels of theophylline and its metabolites without interference from caffeine. *Biomed Chromatogr*, **13**, 15-23.
- Kratz, G. (1998) Modeling of wound healing processes in human skin using tissue culture. *Microsc Res Tech*, **42**, 345-350.
- Krishnaiah, Y.S., Satyanarayana, V. and Karthikeyan, R.S. (2002) Effect of the solvent system on the in vitro permeability of nicardipine hydrochloride through excised rat epidermis. *J Pharm Pharm Sci*, **5**, 123-130.
- Krul, C. and Hageman, G. (1998) Analysis of urinary caffeine metabolites to assess biotransformation enzyme activities by reversed-phase high-performance liquid chromatography. *J Chromatogr B Biomed Sci Appl*, **709**, 27-34.
- Kubota, K., Ademola, J. and Maibach, H.I. (1994) Metabolism and degradation of betamethasone 17-valerate in homogenized living skin equivalent. *Dermatology*, **188**, 13-17.
- Lenoir, M.C. and Bernard, B.A. (1990) Architecture of reconstructed epidermis on collagen lattices varies according to the method used: a comparative study. *Skin Pharmacol*, **3**, 97-106.
- Levang, A.K., Zhao, K. and Singh, J. (1999) Effect of ethanol/propylene glycol on the in vitro percutaneous absorption of aspirin, biophysical changes and macroscopic barrier properties of the skin. *Int J Pharm*, **181**, 255-263.
- Li, P., Nijhawan, D., Budihardjo, I., Srinivasula, S.M., Ahmad, M., Alnemri, E.S. and Wang, X. (1997) Cytochrome c and dATP-dependent formation of Apaf-

- 1/caspase-9 complex initiates an apoptotic protease cascade. *Cell*, **91**, 479-489.
- Liebsch, M., Traue, D., Barrabas, C., Spielmann, H., Uphill, P., Wilkins, S., McPherson, J.P., Wiemann, C., Kaufmann, T., Remmeli, M., Holzhütter, H.G. (2000) The ECVAM prevalidation study on the use of EpiDerm for skin corrosivity testing. *Alternatives to Laboratory Animals*, **28**, 371-401.
- Lindner, W. and Wainer, I.W. (1998) Requirements for initial assay validation and publication in J. Chromatography B. *J Chromatogr B Biomed Sci Appl*, **707**, 1-2.
- Lippens, S., Kockx, M., Knaapen, M., Mortier, L., Polakowska, R., Verheyen, A., Garmyn, M., Zwijsen, A., Formstecher, P., Huylebroeck, D., Vandebaele, P. and Declercq, W. (2000) Epidermal differentiation does not involve the pro-apoptotic executioner caspases, but is associated with caspase-14 induction and processing. *Cell Death Differ*, **7**, 1218-1224.
- Liu, Z.G., Baskaran, R., Lea-Chou, E.T., Wood, L.D., Chen, Y., Karin, M. and Wang, J.Y. (1996) Three distinct signalling responses by murine fibroblasts to genotoxic stress. *Nature*, **384**, 273-276.
- Lockshin, R.A. and Zakeri, Z. (2004) Caspase-independent cell death? *Oncogene*, **23**, 2766-2773.
- Lotte, C., Patouillet, C., Zanini, M., Messager, A. and Roguet, R. (2002) Permeation and skin absorption: reproducibility of various industrial reconstructed human skin models. *Skin Pharmacol Appl Skin Physiol*, **15 Suppl 1**, 18-30.
- Luft, J.H. (1961) Improvements in epoxy resin embedding methods. *J Biophys Biochem Cytol*, **9**, 409-414.
- Lundehn, J.R. (1992) Uniform principles for safeguarding the health of applicators of plant protection products (uniform principles for operator protection). *Mitteilungen aus der Biologischen Bundesanstalt, Heft 277*.
- Maciag, T., Nemore, R.E., Weinstein, R. and Gilchrest, B.A. (1981) An endocrine approach to the control of epidermal growth: serum-free cultivation of human keratinocytes. *Science*, **211**, 1452-1454.
- Mackenzie, I.C. and Fusenig, N.E. (1983) Regeneration of organized epithelial structure. *J Invest Dermatol*, **81**, 189s-194s.
- Mahmoud, A., Haberland, A., Durrfeld, M., Heydeck, D., Wagner, S. and Schafer-Korting, M. (2005) Cutaneous estradiol permeation, penetration and metabolism in pig and man. *Skin Pharmacol Physiol*, **18**, 27-35.
- Man, M.Q., Feingold, K.R. and Elias, P.M. (1993) Exogenous lipids influence permeability barrier recovery in acetone-treated murine skin. *Arch Dermatol*, **129**, 728-738.
- Marjukka Suhonen, T., Pasonen-Seppanen, S., Kirjavainen, M., Tammi, M., Tammi, R. and Urtti, A. (2003) Epidermal cell culture model derived from rat keratinocytes with permeability characteristics comparable to human cadaver skin. *Eur J Pharm Sci*, **20**, 107-113.
- Märkl, H.H. (1979) *Plasmaproteins*. Ciba-Geigy Ltd., Basel.
- Mattie, D.R., Grabau, J.H. and McDougal, J.N. (1994) Significance of the dermal route of exposure to risk assessment. *Risk Anal*, **14**, 277-284.
- Menon, G.K. and Elias, P.M. (1997) Morphologic basis for a pore-pathway in mammalian stratum corneum. *Skin Pharmacol*, **10**, 235-246.
- Merk, H.F. (1989) [Geriatric dermatopharmacology]. *Z Hautkr*, **64**, 1015-1018.
- Merk, H.F. (1998) *Skin Metabolism*. Springer Verlag, Berlin.
- Meyer, W. and Neurand, K. (1976) The distribution of enzymes in the skin of the domestic pig. *Lab Anim*, **10**, 237-247.

- Micali, G., Bhatt, R.H., Distefano, G., Caltabiano, L., Cook, B., Fischer, J.H., Solomon, L.M. and West, D.P. (1993) Evaluation of transdermal theophylline pharmacokinetics in neonates. *Pharmacotherapy*, **13**, 386-390.
- Moser, K., Kriwet, K., Naik, A., Kalia, Y.N. and Guy, R.H. (2001) Passive skin penetration enhancement and its quantification in vitro. *Eur J Pharm Biopharm*, **52**, 103-112.
- Mosmann, T. (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods*, **65**, 55-63.
- Moss, G.P. and Cronin, M.T. (2002) Quantitative structure-permeability relationships for percutaneous absorption: re-analysis of steroid data. *Int J Pharm*, **238**, 105-109.
- Munster, U., Nakamura, C., Haberland, A., Jores, K., Mehnert, W., Rummel, S., Schaller, M., Korting, H.C., Zouboulis Ch, C., Blume-Peytavi, U. and Schafer-Korting, M. (2005) RU 58841-myristate--prodrug development for topical treatment of acne and androgenetic alopecia. *Pharmazie*, **60**, 8-12.
- Nakazawa, K., Nakazawa, H., Collombel, C. and Damour, O. (1995) Keratinocyte extracellular matrix-mediated regulation of normal human melanocyte functions. *Pigment Cell Res*, **8**, 10-18.
- Naline, E., Flouvat, B., Advenier, C. and Pays, M. (1987) Determination of theophylline and its metabolites in plasma and urine by reversed-phase liquid chromatography using an amine modifier. *J Chromatogr*, **419**, 177-189.
- Naughton, G., Jacob, L. and Naughton, B. (1989) *A physiological skin model for in vitro toxicity studies*. Mary Ann Liebert, New York.
- Netzlaff, F., Kostka, K.H., Lehr, C.M. and Schaefer, U.F. (2005a) TEWL measurements as a routine method for evaluating the integrity of epidermis sheets in static Franz type diffusion cells in vitro. Limitations shown by transport data testing. *Eur J Pharm Biopharm*.
- Netzlaff, F., Lehr, C.M., Wertz, P.W. and Schaefer, U.F. (2005b) The human epidermis models EpiSkin, SkinEthic and EpiDerm: an evaluation of morphology and their suitability for testing phototoxicity, irritancy, corrosivity, and substance transport. *Eur J Pharm Biopharm*, **60**, 167-178.
- Nicoli, S., Colombo, P. and Santi, P. (2005) Release and permeation kinetics of caffeine from bioadhesive transdermal films. *Aaps J*, **7**, E218-223.
- Nokhodchi, A., Shokri, J., Dashbolaghi, A., Hassan-Zadeh, D., Ghafourian, T. and Barzegar-Jalali, M. (2003) The enhancement effect of surfactants on the penetration of lorazepam through rat skin. *Int J Pharm*, **250**, 359-369.
- OECD. (2003) Guidance Document No 28 for the conduct of skin absorption studies. *Adopted at 35<sup>th</sup> Joint Meeting August 2003*.
- OECD. (2004a) Test Guideline 427: Skin Absorption: in vivo Method. *Adopted on 13th April 2004*.
- OECD. (2004b) Test Guideline 428: Skin Absorption: in vitro Method. *Adopted on 13th April 2004*.
- OECD. (2004c) Test Guideline 431: In vitro skin corrosion: Human Skin Model Test. *Adopted on 13th April 2004*.
- OECD. (2004d) Test Guideline 432: In vitro 3T3 NRU phototoxicity test. *Adopted on 13th April 2004*.
- Otberg, N., Richter, H., Schaefer, H., Blume-Peytavi, U., Sterry, W. and Lademann, J. (2004) Variations of hair follicle size and distribution in different body sites. *J Invest Dermatol*, **122**, 14-19.

- Pan, G., O'Rourke, K. and Dixit, V.M. (1998) Caspase-9, Bcl-XL, and Apaf-1 form a ternary complex. *J Biol Chem*, **273**, 5841-5845.
- Pendlington, R.U., Williams, D.L., Naik, J.T. and Sharma, R.K. (1994) Distribution of xenobiotic metabolising enzymes in skin. *Toxicology in vitro*, **8**, 525-527.
- Pfundt, R., van Vlijmen-Willems, I., Bergers, M., Wingens, M., Cloin, W. and Schalkwijk, J. (2001) In situ demonstration of phosphorylated c-jun and p38 MAP kinase in epidermal keratinocytes following ultraviolet B irradiation of human skin. *J Pathol*, **193**, 248-255.
- Pham, M.A., Magdalou, J., Siest, G., Lenoir, M.C., Bernard, B.A., Jamoulle, J.C. and Shroot, B. (1990) Reconstituted epidermis: a novel model for the study of drug metabolism in human epidermis. *J Invest Dermatol*, **94**, 749-752.
- Pisansarakit, P., du Cros, D. and Moore, G.P. (1990) Cultivation of keratinocytes derived from epidermal explants of sheep skin and the roles of growth factors in the regulation of proliferation. *Arch Dermatol Res*, **281**, 530-535.
- Pistrutto, G., Jost, M., Srinivasula, S.M., Baffa, R., Poyet, J.L., Kari, C., Lazebnik, Y., Rodeck, U. and Alnemri, E.S. (2002) Expression and transcriptional regulation of caspase-14 in simple and complex epithelia. *Cell Death Differ*, **9**, 995-1006.
- Pittermann, W., Kietzmann, M. and Jackwerth, B. (1995) [The isolated perfused Bovine Udder Skin]. *Altex*, **12**, 196-200.
- Plewig, G., Jansen, T. and Schurer, N.Y. (1997) [Stratum corneum]. *Hautarzt*, **48**, 510-521; quiz 520.
- Poet, T.S. and McDougal, J.N. (2002) Skin absorption and human risk assessment. *Chem Biol Interact*, **140**, 19-34.
- Ponec, M. (1991) Reconstruction of human epidermis on de-epidermized dermis: Expression of differentiation-specific protein markers and lipid composition. *Toxicology in vitro*, **5**, 597-606.
- Ponec, M., Boelsma, E., Gibbs, S. and Mommaas, M. (2002) Characterization of reconstructed skin models. *Skin Pharmacol Appl Skin Physiol*, **15 Suppl 1**, 4-17.
- Ponec, M., Boelsma, E., Weerheim, A., Mulder, A., Bouwstra, J. and Mommaas, M. (2000) Lipid and ultrastructural characterization of reconstructed skin models. *Int J Pharm*, **203**, 211-225.
- Ponec, M. and Weerheim, A. (1990) *Retinoids and lipid changes in keratinocytes*. Academic Press, San Diego.
- Ponec, M., Weerheim, A., Kempenaar, J., Mommaas, A.M. and Nugteren, D.H. (1988) Lipid composition of cultured human keratinocytes in relation to their differentiation. *J Lipid Res*, **29**, 949-961.
- Ponec, M., Weerheim, A., Kempenaar, J., Mulder, A., Gooris, G.S., Bouwstra, J. and Mommaas, A.M. (1997) The formation of competent barrier lipids in reconstructed human epidermis requires the presence of vitamin C. *J Invest Dermatol*, **109**, 348-355.
- Pont, A.R., Charron, A.R. and Brand, R.M. (2004) Active ingredients in sunscreens act as topical penetration enhancers for the herbicide 2,4-dichlorophenoxyacetic acid. *Toxicol Appl Pharmacol*, **195**, 348-354.
- Popa, C., Dicker, A.J., Dahler, A.L. and Saunders, N.A. (1999) Cytochrome P450, CYP26A1, is expressed at low levels in human epidermal keratinocytes and is not retinoic acid-inducible. *Br J Dermatol*, **141**, 460-468.
- Potard, G., Laugel, C., Baillet, A., Schaefer, H. and Marty, J.P. (1999) Quantitative HPLC analysis of sunscreens and caffeine during in vitro percutaneous penetration studies. *Int J Pharm*, **189**, 249-260.

- Prenosil, J.E. and Kino-oka, M. (1999) Computer controlled bioreactor for large-scale production of cultured skin grafts. *Ann N Y Acad Sci*, **875**, 386-397.
- Puviani, M., Marconi, A., Cozzani, E. and Pincelli, C. (2003) Fas ligand in pemphigus sera induces keratinocyte apoptosis through the activation of caspase-8. *J Invest Dermatol*, **120**, 164-167.
- Qiao, G.L. and Riviere, J.E. (2002) Systemic uptake and cutaneous disposition of pentachlorophenol in a sequential exposure scenario: effects of skin preexposure to benzo[a]pyrene. *J Toxicol Environ Health A*, **65**, 1307-1331.
- Ramsey, J.D., Woollen, B.H., Auton, T.R. and Scott, R.C. (1994) The predictive accuracy of in vitro measurements for the dermal absorption of a lipophilic penetrant (fluazifop-butyl) through rat and human skin. *Fundam Appl Toxicol*, **23**, 230-236.
- Raza, H. and Mukhtar, H. (1993) Differences in inducibility of cytochrome P-4501A1, monooxygenases and glutathione S-transferase in cutaneous and extracutaneous tissues after topical and parenteral administration of beta-naphthoflavone to rats. *Int J Biochem*, **25**, 1511-1516.
- Regnier, M., Asselineau, D. and Lenoir, M.C. (1990) Human epidermis reconstructed on dermal substrates in vitro: an alternative to animals in skin pharmacology. *Skin Pharmacol*, **3**, 70-85.
- Regnier, M., Schweizer, J., Michel, S., Bailly, C. and Prunieras, M. (1986) Expression of high molecular weight (67K) keratin in human keratinocytes cultured on dead de-epidermized dermis. *Exp Cell Res*, **165**, 63-72.
- Reifenrath, W.G., Hawkins, G.S. and Kurtz, M.S. (1989) Evaporation and skin penetration characteristics of mosquito repellent formulations. *J Am Mosq Control Assoc*, **5**, 45-51.
- Rendl, M., Ban, J., Mrass, P., Mayer, C., Lengauer, B., Eckhart, L., Declerq, W. and Tschachler, E. (2002) Caspase-14 expression by epidermal keratinocytes is regulated by retinoids in a differentiation-associated manner. *J Invest Dermatol*, **119**, 1150-1155.
- Riviere, J.E., Baynes, R.E., Brooks, J.D., Yeatts, J.L. and Monteiro-Riviere, N.A. (2003) Percutaneous absorption of topical N,N-diethyl-m-toluamide (DEET): effects of exposure variables and coadministered toxicants. *J Toxicol Environ Health A*, **66**, 133-151.
- Robert, M., Dusser, I., Muriel, M.P., Noel-Hudson, M.S., Aubery, M. and Wepierre, J. (1997) Barrier function of reconstructed epidermis at the air-liquid interface: influence of dermal cells and extracellular components. *Skin Pharmacol*, **10**, 247-260.
- Rosdy, M. and Clauss, L.C. (1990) Terminal epidermal differentiation of human keratinocytes grown in chemically defined medium on inert filter substrates at the air-liquid interface. *J Invest Dermatol*, **95**, 409-414.
- Roskos, K.V., Maibach, H.I. and Guy, R.H. (1989) The effect of aging on percutaneous absorption in man. *J Pharmacokinet Biopharm*, **17**, 617-630.
- Rougier, A., Lotte, C. and Maibach, H.I. (1987) In vivo percutaneous penetration of some organic compounds related to anatomic site in humans: predictive assessment by the stripping method. *J Pharm Sci*, **76**, 451-454.
- Roy, S.D. and Flynn, G.L. (1990) Transdermal delivery of narcotic analgesics: pH, anatomical, and subject influences on cutaneous permeability of fentanyl and sufentanil. *Pharm Res*, **7**, 842-847.
- Saeki, M., Saito, Y., Nagano, M., Teshima, R., Ozawa, S. and Sawada, J. (2002) mRNA expression of multiple cytochrome p450 isozymes in four types of cultured skin cells. *Int Arch Allergy Immunol*, **127**, 333-336.

- Sainio, E.L., Rantanen, T. and Kanerva, L. (2000) Ingredients and safety of cellulite creams. *Eur J Dermatol*, **10**, 596-603.
- Santos Maia, C., Mehnert, W., Schaller, M., Korting, H.C., Gysler, A., Haberland, A. and Schafer-Korting, M. (2002) Drug targeting by solid lipid nanoparticles for dermal use. *J Drug Target*, **10**, 489-495.
- Sarkar, M.A., Hunt, C., Guzelian, P.S. and Karnes, H.T. (1992) Characterization of human liver cytochromes P-450 involved in theophylline metabolism. *Drug Metab Dispos*, **20**, 31-37.
- SCCNFP. (2003) The SCCNFP's Notes of Guidance for the testing of cosmetic ingredients and their safety evaluation. *Adopted at 25<sup>th</sup> plenary meeting October 2003*.
- Schaefer, H. and Redelmeier, T.E. (1996) *Structure and dynamics of the skin barrier*. Karger, Basel.
- Schulze-Osthoff, K., Ferrari, D., Los, M., Wesselborg, S. and Peter, M.E. (1998) Apoptosis signaling by death receptors. *Eur J Biochem*, **254**, 439-459.
- Scott, R.C., Batten, P.L., Clowes, H.M., Jones, B.K. and Ramsey, J.D. (1992) Further validation of an in vitro method to reduce the need for in vivo studies for measuring the absorption of chemicals through rat skin. *Fundam Appl Toxicol*, **19**, 484-492.
- Seko, N., Bando, H., Lim, C.W., Yamashita, F. and Hashida, M. (1999) Theoretical analysis of the effect of cutaneous metabolism on skin permeation of parabens based on a two-layer skin diffusion/metabolism model. *Biol Pharm Bull*, **22**, 281-287.
- Sheikh, M.S., Antinore, M.J., Huang, Y. and Fornace, A.J., Jr. (1998) Ultraviolet-irradiation-induced apoptosis is mediated via ligand independent activation of tumor necrosis factor receptor 1. *Oncogene*, **17**, 2555-2563.
- Simon, G.A. and Maibach, H.I. (2000) The pig as an experimental animal model of percutaneous permeation in man: qualitative and quantitative observations--an overview. *Skin Pharmacol Appl Skin Physiol*, **13**, 229-234.
- Sivaramakrishnan, R., Nakamura, C., Mehnert, W., Korting, H.C., Kramer, K.D. and Schafer-Korting, M. (2004) Glucocorticoid entrapment into lipid carriers--characterisation by parelectric spectroscopy and influence on dermal uptake. *J Control Release*, **97**, 493-502.
- Slivka, S.R., Landeen, L.K., Zeigler, F., Zimber, M.P. and Bartel, R.L. (1993) Characterization, barrier function, and drug metabolism of an in vitro skin model. *J Invest Dermatol*, **100**, 40-46.
- Smith, D.A. (1994) Design of drugs through a consideration of drug metabolism and pharmacokinetics. *Eur J Drug Metab Pharmacokinet*, **19**, 193-199.
- Spielmann, H., Muller, L., Averbeck, D., Balls, M., Brendler-Schwaab, S., Castell, J.V., Curren, R., de Silva, O., Gibbs, N.K., Liebsch, M., Lovell, W.W., Merk, H.F., Nash, J.F., Neumann, N.J., Pape, W.J., Ulrich, P. and Vohr, H.W. (2000) The second ECVAM workshop on phototoxicity testing. The report and recommendations of ECVAM workshop 42. *Altern Lab Anim*, **28**, 777-814.
- Sultana, S. and Saleem, M. (2004) Salix caprea inhibits skin carcinogenesis in murine skin: inhibition of oxidative stress, ornithine decarboxylase activity and DNA synthesis. *J Ethnopharmacol*, **91**, 267-276.
- Suzuki, Y., Nomura, J., Hori, J., Koyama, J., Takahashi, M. and Horii, I. (1993) Detection and characterization of endogenous protease associated with desquamation of stratum corneum. *Arch Dermatol Res*, **285**, 372-377.
- Takahashi, A., Suzuki, S., Kawasaki, N., Kubo, W., Miyazaki, S., Loebenberg, R., Bachynsky, J. and Attwood, D. (2002) Percutaneous absorption of non-

- steroidal anti-inflammatory drugs from in situ gelling xyloglucan formulations in rats. *Int J Pharm*, **246**, 179-186.
- Takahashi, K., Sawasaki, Y., Hata, J., Mukai, K. and Goto, T. (1990) Spontaneous transformation and immortalization of human endothelial cells. *In Vitro Cell Dev Biol*, **26**, 265-274.
- Takahashi, T., Ogo, M. and Hibino, T. (1998) Partial purification and characterization of two distinct types of caspases from human epidermis. *J Invest Dermatol*, **111**, 367-372.
- Tinois, E., Tiollier, J., Gaucherand, M., Dumas, H., Tardy, M. and Thivolet, J. (1991) In vitro and post-transplantation differentiation of human keratinocytes grown on the human type IV collagen film of a bilayered dermal substitute. *Exp Cell Res*, **193**, 310-319.
- Todd, C., Hewitt, S.D., Kempenaar, J., Noz, K., Thody, A.J. and Ponec, M. (1993) Co-culture of human melanocytes and keratinocytes in a skin equivalent model: effect of ultraviolet radiation. *Arch Dermatol Res*, **285**, 455-459.
- Treffel, P., Muret, P., Muret-D'Aniello, P., Coumes-Marquet, S. and Agache, P. (1992) Effect of occlusion on in vitro percutaneous absorption of two compounds with different physicochemical properties. *Skin Pharmacol*, **5**, 108-113.
- Tsai, J.C., Lin, C.Y., Sheu, H.M., Lo, Y.L. and Huang, Y.H. (2003) Noninvasive characterization of regional variation in drug transport into human stratum corneum in vivo. *Pharm Res*, **20**, 632-638.
- Tu, A.S., Robertson, R.S., Callahan, M.M. and Thayer, P.S. (1982) Caffeine-metabolizing capability of primary rat hepatocytes, Chinese hamster V79 cells and normal human fibroblasts in culture. *Mutat Res*, **95**, 475-492.
- Ueda, O., Kitamura, S., Ohashi, K., Sugihara, K. and Ohta, S. (2003) Xanthine oxidase-catalyzed metabolism of 2-nitrofluorene, a carcinogenic air pollutant, in rat skin. *Drug Metab Dispos*, **31**, 367-372.
- Van de Craen, M., Van Loo, G., Pype, S., Van Criekinge, W., Van den brande, I., Moelmans, F., Fiers, W., Declercq, W. and Vandenameele, P. (1998) Identification of a new caspase homologue: caspase-14. *Cell Death Differ*, **5**, 838-846.
- van de Sandt, J.J., van Burgsteden, J.A., Cage, S., Carmichael, P.L., Dick, I., Kenyon, S., Korinth, G., Larese, F., Limasset, J.C., Maas, W.J., Montomoli, L., Nielsen, J.B., Payan, J.P., Robinson, E., Sartorelli, P., Schaller, K.H., Wilkinson, S.C. and Williams, F.M. (2004) In vitro predictions of skin absorption of caffeine, testosterone, and benzoic acid: a multi-centre comparison study. *Regul Toxicol Pharmacol*, **39**, 271-281.
- van Ravenzwaay, B. and Leibold, E. (2004) The significance of in vitro rat skin absorption studies to human risk assessment. *Toxicol In Vitro*, **18**, 219-225.
- Vecchini, F., Mace, K., Magdalou, J., Mahe, Y., Bernard, B.A. and Shroot, B. (1995) Constitutive and inducible expression of drug metabolizing enzymes in cultured human keratinocytes. *Br J Dermatol*, **132**, 14-21.
- Vermes, I., Haanen, C., Steffens-Nakken, H. and Reutelingsperger, C. (1995) A novel assay for apoptosis. Flow cytometric detection of phosphatidylserine expression on early apoptotic cells using fluorescein labelled Annexin V. *J Immunol Methods*, **184**, 39-51.
- Viard, I., Wehrli, P., Bullani, R., Schneider, P., Holler, N., Salomon, D., Hunziker, T., Saurat, J.H., Tschopp, J. and French, L.E. (1998) Inhibition of toxic epidermal necrolysis by blockade of CD95 with human intravenous immunoglobulin. *Science*, **282**, 490-493.

- Wagner, H., Kostka, K.H., Lehr, C.M. and Schaefer, U.F. (2000) Drug distribution in human skin using two different in vitro test systems: comparison with in vivo data. *Pharm Res*, **17**, 1475-1481.
- Wagner, S.M., Nogueira, A.C., Paul, M., Heydeck, D., Klug, S. and Christ, B. (2003) The isolated normothermic hemoperfused porcine forelimb as a test system for transdermal absorption studies. *J Artif Organs*, **6**, 183-191.
- Walker, N.I., Harmon, B.V., Gobe, G.C. and Kerr, J.F. (1988) Patterns of cell death. *Methods Achiev Exp Pathol*, **13**, 18-54.
- Wallach, D., Boldin, M., Varfolomeev, E., Beyaert, R., Vandenabeele, P. and Fiers, W. (1997) Cell death induction by receptors of the TNF family: towards a molecular understanding. *FEBS Lett*, **410**, 96-106.
- Welss, T., Basketter, D.A. and Schroder, K.R. (2004) In vitro skin irritation: facts and future. State of the art review of mechanisms and models. *Toxicol In Vitro*, **18**, 231-243.
- Wertz, P.W. and Downing, D.T. (1989) *Stratum corneum: biological and biochemical considerations*. Marcel Dekker, New York.
- Wester, R.C. and Maibach, H.I. (1976) Relationship of topical dose and percutaneous absorption in rhesus monkey and man. *J Invest Dermatol*, **67**, 518-520.
- Wille, J.J., Jr., Pittelkow, M.R., Shipley, G.D. and Scott, R.E. (1984) Integrated control of growth and differentiation of normal human prokeratinocytes cultured in serum-free medium: clonal analyses, growth kinetics, and cell cycle studies. *J Cell Physiol*, **121**, 31-44.
- Yagi, S., Nakayama, K., Kurosaki, Y., Higaki, K. and Kimura, T. (1998) Factors determining drug residence in skin during transdermal absorption: studies on beta-blocking agents. *Biol Pharm Bull*, **21**, 1195-1201.
- Zhang, M.C., Liu, H.P., Demchik, L.L., Zhai, Y.F. and Yang da, J. (2004) LIGHT sensitizes IFN-gamma-mediated apoptosis of HT-29 human carcinoma cells through both death receptor and mitochondria pathways. *Cell Res*, **14**, 117-124.
- Zoellner, H., Hofler, M., Beckmann, R., Hufnagl, P., Vanyek, E., Bielek, E., Wojta, J., Fabry, A., Lockie, S. and Binder, B.R. (1996) Serum albumin is a specific inhibitor of apoptosis in human endothelial cells. *J Cell Sci*, **109 (Pt 10)**, 2571-2580.