

6 REFERENCES

- Akeson EC, Lambert JP, Narayanswami S, Gardiner K, Bechtel LJ, et al. (2001). Ts65Dn -- localization of the translocation breakpoint and trisomic gene content in a mouse model for Down syndrome. *Cytogenet Cell Genet* 93: 270-276.
- Amano K, Sago H, Uchikawa C, Suzuki T, Kotliarova SE, et al. (2004). Dosage-dependent overexpression of genes in the trisomic region of Ts1Cje mouse model for Down syndrome. *Hum Mol Genet* 13: 1333-1340.
- Antonarakis SE (1991). Parental origin of the extra chromosome in trisomy 21 as indicated by analysis of DNA polymorphisms. *Down Syndrome Collaborative Group. N Engl J Med* 324: 872-876.
- Antonarakis SE (1992). Third International Workshop on Human Chromosome 21. *Genomics* 14: 1126-1132.
- Antonarakis SE (1993). Human chromosome 21: genome mapping and exploration, circa 1993. *Trends Genet* 9: 142-148.
- Antonarakis SE (2001). Chromosome 21: from sequence to applications. *Curr Opin Genet Dev* 11: 241-246.
- Antonarakis SE, Adelsberger PA, Petersen MB, Binkert F, Schinzel AA (1990). Analysis of DNA polymorphisms suggests that most de novo dup(21q) chromosomes in patients with Down syndrome are isochromosomes and not translocations. *Am J Hum Genet* 47: 968-972.
- Antonarakis SE, Lyle R, Dermitzakis ET, Reymond A, Deutsch S (2004). Chromosome 21 and down syndrome: from genomics to pathophysiology. *Nat Rev Genet* 5: 725-738.
- Aylward EH, Habbak R, Warren AC, Pulsifer MB, Barta PE, et al. (1997). Cerebellar volume in adults with Down syndrome. *Arch Neurol* 54: 209-212.
- Bahn S, Mimmack M, Ryan M, Caldwell MA, Jauniaux E, et al. (2002). Neuronal target genes of the neuron-restrictive silencer factor in neurospheres derived from fetuses with Down's syndrome: a gene expression study. *Lancet* 359: 310-315.
- Balkan W, Martin RH (1983). Segregation of chromosomes into the spermatozoa of a man heterozygous for a 14;21 Robertsonian translocation. *Am J Med Genet* 16: 169-172.
- Baxter LL, Moran TH, Richtsmeier JT, Troncoso J, Reeves RH (2000). Discovery and genetic localization of Down syndrome cerebellar phenotypes using the Ts65Dn mouse. *Hum Mol Genet* 9: 195-202.
- Becker L, Mito T, Takashima S, Onodera K (1991). Growth and development of the brain in Down syndrome. *Prog Clin Biol Res* 373: 133-152.
- Berger R (1997). Acute lymphoblastic leukemia and chromosome 21. *Cancer Genet Cytogenet* 94: 8-12.
- Birchler JA, Hiebert JC, Paigen K (1990). Analysis of autosomal dosage compensation involving the alcohol dehydrogenase locus in *Drosophila melanogaster*. *Genetics* 124: 679-686.
- Borota J, Milosevic-Tasic M, Katanic D (1997). [Homocystinuria--biochemical, clinical and genetic aspects]. *Med Pregl* 50: 187-193.
- Bovicelli L, Orsini LF, Rizzo N, Montacuti V, Bacchetta M (1982). Reproduction in Down syndrome. *Obstet Gynecol* 59: 13S-17S.
- Brenner S (1974). The genetics of *Caenorhabditis elegans*. *Genetics* 77: 71-94.
- Brown WT (1987). Premature aging syndromes. *Curr Probl Dermatol* 17: 152-165.

- Byrne JL, Ward K (1994). Genetic factors in recurrent abortion. *Clin Obstet Gynecol* 37: 693-704.
- Cairns NJ (1999). Neuropathology. *J Neural Transm Suppl* 57: 61-74.
- Chalfie M, White J (1988) The nervous system. In: Wood BW, editor. *The nematode Caenorhabditis elegans*. New York: Cold Spring Harbor Laboratory Press. pp. 337-392.
- Chomczynski P, Sacchi N (1987). Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162: 156-159.
- Chrast R, Scott HS, Papasavvas MP, Rossier C, Antonarakis ES, et al. (2000). The mouse brain transcriptome by SAGE: differences in gene expression between P30 brains of the partial trisomy 16 mouse model of Down syndrome (Ts65Dn) and normals. *Genome Res* 10: 2006-2021.
- Citron M, Oltersdorf T, Haass C, McConlogue L, Hung AY, et al. (1992). Mutation of the beta-amyloid precursor protein in familial Alzheimer's disease increases beta-protein production. *Nature* 360: 672-674.
- Clamp M, Cuff J, Searle SM, Barton GJ (2004). The Jalview Java alignment editor. *Bioinformatics* 20: 426-427.
- Clark SG, Lu X, Horvitz HR (1994). The *Caenorhabditis elegans* locus lin-15, a negative regulator of a tyrosine kinase signaling pathway, encodes two different proteins. *Genetics* 137: 987-997.
- Colon EY (1972). The structure of the cerebral cortex in Down's syndrome: quantitative analysis. *Neuropediatrics* 3: 362-376.
- Crome L, Cowie V, Slater E (1966). Statistical note on cerebellar and brain-stem weight in mongolism. *Journal of Mental Deficiency Research* 10: 69-72.
- Dauphinot L, Lyle R, Rivals I, Dang MT, Moldrich RX, et al. (2005). The cerebellar transcriptome during postnatal development of the Ts1Cje mouse, a segmental trisomy model for Down syndrome. *Hum Mol Genet* 14: 373-384.
- Davidoff LM (1928). Brain in mongolian idiocy: report of 10 cases. *Archives of Neurology and Psychiatry* 20: 1229-1257.
- Davisson MT, Akeson EC (1987). An improved method for preparing G-banded chromosomes from mouse peripheral blood. *Cytogenet Cell Genet* 45: 70-74.
- Davisson MT, Schmidt C, Akeson EC (1990). Segmental trisomy of murine chromosome 16: a new model system for studying Down syndrome. *Prog Clin Biol Res* 360: 263-280.
- Delabar JM, Chettouh Z, Rahmani Z, Theophile D, Blouin JL, et al. (1992). Gene dosage mapping of 30 DNA markers on chromosome 21. *Genomics* 13: 887-889.
- Delabar JM, Theophile D, Rahmani Z, Chettouh Z, Blouin JL, et al. (1993). Molecular mapping of twenty-four features of Down syndrome on chromosome 21. *Eur J Hum Genet* 1: 114-124.
- Demas GE, Nelson RJ, Krueger BK, Yarowsky PJ (1996). Spatial memory deficits in segmental trisomic Ts65Dn mice. *Behav Brain Res* 82: 85-92.
- Deutsch S, Lyle R, Dermitzakis ET, Attar H, Subrahmanyam L, et al. (2005). Gene expression variation and expression quantitative trait mapping of human chromosome 21 genes. *Hum Mol Genet* 14: 3741-3749.
- Devlin RH, Holm DG, Grigliatti TA (1982). Autosomal dosage compensation *Drosophila melanogaster* strains trisomic for the left arm of chromosome 2. *Proc Natl Acad Sci U S A* 79: 1200-1204.

- Devlin RH, Holm DG, Grigliatti TA (1988). The influence of whole-arm trisomy on gene expression in *Drosophila*. *Genetics* 118: 87-101.
- Dinani S, Carpenter S (1990). Down's syndrome and thyroid disorder. *J Ment Defic Res* 34 (Pt 2): 187-193.
- Down JL (1995). Observations on an ethnic classification of idiots. 1866. *Ment Retard* 33: 54-56.
- Driever W, Nusslein-Volhard C (1988). The bicoid protein determines position in the *Drosophila* embryo in a concentration-dependent manner. *Cell* 54: 95-104.
- Egeo A, Di Lisi R, Sandri C, Mazzocco M, Lapide M, et al. (2000). Developmental expression of the SH3BGR gene, mapping to the Down syndrome heart critical region. *Mech Dev* 90: 313-316.
- Elbashir SM, Harborth J, Lendeckel W, Yalcin A, Weber K, et al. (2001). Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature* 411: 494-498.
- Epstein CJ (1986) The consequences of chromosome imbalance Principles, mechanisms, and models; Barlow P, Green PB, Wyllie CC, editors. Cambridge: Cambridge University Press.
- Epstein CJ (1990). The consequences of chromosome imbalance. *Am J Med Genet Suppl* 7: 31-37.
- Epstein CJ (2001) Down syndrome (trisomy 21); Scriver CR, Beaudet AL, Sly WS, Valle D, editors. New York: McGraw Hill.
- Epstein CJ, Cox DR, Epstein LB (1985). Mouse trisomy 16: an animal model of human trisomy 21 (Down syndrome). *Ann N Y Acad Sci* 450: 157-168.
- Fernandez AG, Gunsalus KC, Huang J, Chuang LS, Ying N, et al. (2005). New genes with roles in the *C. elegans* embryo revealed using RNAi of ovary-enriched ORFeome clones. *Genome Res* 15: 250-259.
- Finnish German APECED Consortium (1997). An autoimmune disease, APECED, caused by mutations in a novel gene featuring two PHD-type zinc-finger domains. The Finnish-German APECED Consortium. Autoimmune Polyendocrinopathy-Candidiasis-Ectodermal Dystrophy. *Nat Genet* 17: 399-403.
- Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, et al. (1998). Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391: 806-811.
- FitzPatrick DR, Ramsay J, McGill NI, Shade M, Carothers AD, et al. (2002). Transcriptome analysis of human autosomal trisomy. *Hum Mol Genet* 11: 3249-3256.
- Fong CT, Brodeur GM (1987). Down's syndrome and leukemia: epidemiology, genetics, cytogenetics and mechanisms of leukemogenesis. *Cancer Genet Cytogenet* 28: 55-76.
- Ford CE, Hamerton JL (1956). The chromosomes of man. *Acta Genet Stat Med* 6: 264-266.
- Fraser AG, Kamath RS, Zipperlen P, Martinez-Campos M, Sohrmann M, et al. (2000). Functional genomic analysis of *C. elegans* chromosome I by systematic RNA interference. *Nature* 408: 325-330.
- Friedman DL, Kastner T, Pond WS, O'Brien DR (1989). Thyroid dysfunction in individuals with Down syndrome. *Arch Intern Med* 149: 1990-1993.
- Gallant S, Gilkeson G (2006). ETS transcription factors and regulation of immunity. *Arch Immunol Ther Exp (Warsz)* 54: 149-163.

- Gardiner K, Fortna A, Bechtel L, Davisson MT (2003). Mouse models of Down syndrome: how useful can they be? Comparison of the gene content of human chromosome 21 with orthologous mouse genomic regions. *Gene* 318: 137-147.
- Giannone S, Strippoli P, Vitale L, Casadei R, Canaider S, et al. (2004). Gene expression profile analysis in human T lymphocytes from patients with Down Syndrome. *Ann Hum Genet* 68: 546-554.
- Gilchrest BA (1981). Premature aging syndromes affecting the skin. *Birth Defects Orig Artic Ser* 17: 227-241.
- Gitton Y, Dahmane N, Baik S, Ruiz i Altaba A, Neidhardt L, et al. (2002). A gene expression map of human chromosome 21 orthologues in the mouse. *Nature* 420: 586-590.
- Goate A, Chartier-Harlin MC, Mullan M, Brown J, Crawford F, et al. (1991). Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature* 349: 704-706.
- Grasso M, Giovannucci Uzielli ML, Pierluigi M, Tavellini F, Perroni L, et al. (1989). Isochromosome not translocation in trisomy 21q21q. *Hum Genet* 84: 63-65.
- Gregor P, Reeves RH, Jabs EW, Yang X, Dackowski W, et al. (1993). Chromosomal localization of glutamate receptor genes: relationship to familial amyotrophic lateral sclerosis and other neurological disorders of mice and humans. *Proc Natl Acad Sci U S A* 90: 3053-3057.
- Gross SJ, Ferreira JC, Morrow B, Dar P, Funke B, et al. (2002). Gene expression profile of trisomy 21 placentas: a potential approach for designing noninvasive techniques of prenatal diagnosis. *Am J Obstet Gynecol* 187: 457-462.
- Guipponi M, Vuagniaux G, Wattenhofer M, Shibuya K, Vazquez M, et al. (2002). The transmembrane serine protease (TMPRSS3) mutated in deafness DFNB8/10 activates the epithelial sodium channel (ENaC) in vitro. *Hum Mol Genet* 11: 2829-2836.
- Hall H, Hunt P, Hassold T (2006). Meiosis and sex chromosome aneuploidy: how meiotic errors cause aneuploidy; how aneuploidy causes meiotic errors. *Curr Opin Genet Dev* 16: 323-329.
- Hassold T, Hunt P (2001). To err (meiotically) is human: the genesis of human aneuploidy. *Nat Rev Genet* 2: 280-291.
- Hattori M, Fujiyama A, Taylor TD, Watanabe H, Yada T, et al. (2000). The DNA sequence of human chromosome 21. *Nature* 405: 311-319.
- Hess EJ (1996). Identification of the weaver mouse mutation: the end of the beginning. *Neuron* 16: 1073-1076.
- Heurgue-Hamard V, Champ S, Engstrom A, Ehrenberg M, Buckingham RH (2002). The hemK gene in Escherichia coli encodes the N(5)-glutamine methyltransferase that modifies peptide release factors. *Embo J* 21: 769-778.
- Hill DP, Blake JA, Richardson JE, Ringwald M (2002). Extension and integration of the gene ontology (GO): combining GO vocabularies with external vocabularies. *Genome Res* 12: 1982-1991.
- Hogg N, Bates PA (2000). Genetic analysis of integrin function in man: LAD-1 and other syndromes. *Matrix Biol* 19: 211-222.
- Hook EB (1981). Unbalanced Robertsonian translocations associated with Down's syndrome or Patau's syndrome: chromosome subtype, proportion inherited, mutation rates, and sex ratio. *Hum Genet* 59: 235-239.

- Horvitz HR, Brenner S, Hodgkin J, Herman RK (1979). A uniform genetic nomenclature for the nematode *Caenorhabditis elegans*. *Mol Gen Genet* 175: 129-133.
- Huret JL, Delabar JM, Marlhens F, Aurias A, Nicole A, et al. (1987). Down syndrome with duplication of a region of chromosome 21 containing the CuZn superoxide dismutase gene without detectable karyotypic abnormality. *Hum Genet* 75: 251-257.
- Hutton M, Hardy J (1997). The presenilins and Alzheimer's disease. *Hum Mol Genet* 6: 1639-1646.
- Inoue T, Shinohara T, Takehara S, Inoue J, Kamino H, et al. (2000). Specific impairment of cardiogenesis in mouse ES cells containing a human chromosome 21. *Biochem Biophys Res Commun* 273: 219-224.
- Ives JH, Dagna-Bricarelli F, Basso G, Antonarakis SE, Jee R, et al. (1998). Increased levels of a chromosome 21-encoded tumour invasion and metastasis factor (TIAM1) mRNA in bone marrow of Down syndrome children during the acute phase of AML(M7). *Genes Chromosomes Cancer* 23: 61-66.
- Jackson JF, North ER, 3rd, Thomas JG (1976). Clinical diagnosis of Down's syndrome. *Clin Genet* 9: 483-487.
- Jacobs PA, Baikie AG, Court Brown WM, Strong JA (1959). The somatic chromosomes in mongolism. *Lancet* 1: 710.
- Jernigan TL, Bellugi U (1990). Anomalous brain morphology on magnetic resonance images in Williams syndrome and Down syndrome. *Arch Neurol* 47: 529-533.
- Kahlem P, Sultan M, Herwig R, Steinfath M, Balzereit D, et al. (2004). Transcript level alterations reflect gene dosage effects across multiple tissues in a mouse model of down syndrome. *Genome Res* 14: 1258-1267.
- Kamath RS, Ahringer J (2003). Genome-wide RNAi screening in *Caenorhabditis elegans*. *Methods* 30: 313-321.
- Kamath RS, Fraser AG, Dong Y, Poulin G, Durbin R, et al. (2003). Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature* 421: 231-237.
- Kamath RS, Martinez-Campos M, Zipperlen P, Fraser AG, Ahringer J (2001). Effectiveness of specific RNA-mediated interference through ingested double-stranded RNA in *Caenorhabditis elegans*. *Genome Biol* 2: RESEARCH0002.
- Karlsson B, Gustafsson J, Hedov G, Ivarsson SA, Anneren G (1998). Thyroid dysfunction in Down's syndrome: relation to age and thyroid autoimmunity. *Arch Dis Child* 79: 242-245.
- Keating DJ, Chen C, Pritchard MA (2006). Alzheimer's disease and endocytic dysfunction: Clues from the Down syndrome-related proteins, DSCR1 and ITSN1. *Ageing Res Rev*.
- Kennedy RL, Jones TH, Cuckle HS (1992). Down's syndrome and the thyroid. *Clin Endocrinol (Oxf)* 37: 471-476.
- Kennerdell JR, Carthew RW (1998). Use of dsRNA-mediated genetic interference to demonstrate that frizzled and frizzled 2 act in the wingless pathway. *Cell* 95: 1017-1026.
- Kim JK, Gabel HW, Kamath RS, Tewari M, Pasquinelli A, et al. (2005). Functional genomic analysis of RNA interference in *C. elegans*. *Science* 308: 1164-1167.

- King RA, Rotter JI, Motulsky AJ (2002) The genetic basis of common diseases: Oxford University Press.
- Klipp E, Herwig R, Kowald A, Wierling C, Lehrach H (2005) Systems Biology in Practice. Concepts, Implementation and Application. Weinheim: WILEY-VCH Verlag GmbH.
- Korenberg JR, Chen XN, Schipper R, Sun Z, Gonsky R, et al. (1994). Down syndrome phenotypes: the consequences of chromosomal imbalance. *Proc Natl Acad Sci U S A* 91: 4997-5001.
- Kraus J, Packman S, Fowler B, Rosenberg LE (1978). Purification and properties of cystathionine beta-synthase from human liver. Evidence for identical subunits. *J Biol Chem* 253: 6523-6528.
- Kravit W, Good RA (1957). Simultaneous occurrence of mongolism and leukemia; report of a nationwide survey. *AMA J Dis Child* 94: 289-293.
- Kwong YL, Ching LM, Liu HW, Lee CP, Pollock A, et al. (1993). 8;21 translocation and multilineage involvement. *Am J Hematol* 43: 212-216.
- Lafon-Cazal M, Pietri S, Culcasi M, Bockaert J (1993). NMDA-dependent superoxide production and neurotoxicity. *Nature* 364: 535-537.
- Lagos-Quintana M, Rauhut R, Yalcin A, Meyer J, Lendeckel W, et al. (2002). Identification of tissue-specific microRNAs from mouse. *Curr Biol* 12: 735-739.
- Lamb BT, Sisodia SS, Lawler AM, Slunt HH, Kitt CA, et al. (1993). Introduction and expression of the 400 kilobase amyloid precursor protein gene in transgenic mice [corrected]. *Nat Genet* 5: 22-30.
- Lander ES (1999). Array of hope. *Nat Genet* 21: 3-4.
- Larocca LM, Piantelli M, Valitutti S, Castellino F, Maggiano N, et al. (1988). Alterations in thymocyte subpopulations in Down's syndrome (trisomy 21). *Clin Immunol Immunopathol* 49: 175-186.
- Lau LL, Spain LM (2000). Altered aging-related thymic involution in T cell receptor transgenic, MHC-deficient, and CD4-deficient mice. *Mech Ageing Dev* 114: 101-121.
- Leeuwen FN, Kain HE, Kammen RA, Michiels F, Kranenburg OW, et al. (1997). The guanine nucleotide exchange factor Tiam1 affects neuronal morphology; opposing roles for the small GTPases Rac and Rho. *J Cell Biol* 139: 797-807.
- Lejeune J, Turpin R, Gautier M (1959). [Chromosomal diagnosis of mongolism.]. *Arch Fr Pediatr* 16: 962-963.
- Lie SO, Jonmundsson G, Mellander L, Siimes MA, Yssing M, et al. (1996). A population-based study of 272 children with acute myeloid leukaemia treated on two consecutive protocols with different intensity: best outcome in girls, infants, and children with Down's syndrome. Nordic Society of Paediatric Haematology and Oncology (NOPHO). *Br J Haematol* 94: 82-88.
- Lietard J, Theret N, Rehn M, Musso O, Dargere D, et al. (2000). The promoter of the long variant of collagen XVIII, the precursor of endostatin, contains liver-specific regulatory elements. *Hepatology* 32: 1377-1385.
- Livingstone CD, Barton GJ (1993). Protein sequence alignments: a strategy for the hierarchical analysis of residue conservation. *Comput Appl Biosci* 9: 745-756.
- Lockhart DJ, Winzeler EA (2000). Genomics, gene expression and DNA arrays. *Nature* 405: 827-836.

- Lott IT (1982). Down's syndrome, aging, and Alzheimer's disease: a clinical review. *Ann N Y Acad Sci* 396: 15-27.
- Lu QR, Sun T, Zhu Z, Ma N, Garcia M, et al. (2002). Common developmental requirement for Olig function indicates a motor neuron/oligodendrocyte connection. *Cell* 109: 75-86.
- Lyle R, Gehrig C, Neergaard-Henrichsen C, Deutsch S, Antonarakis SE (2004). Gene expression from the aneuploid chromosome in a trisomy mouse model of down syndrome. *Genome Res* 14: 1268-1274.
- Mann DM (1988). Alzheimer's disease and Down's syndrome. *Histopathology* 13: 125-137.
- Mann DM, Jones D, Prinja D, Purkiss MS (1990). The prevalence of amyloid (A4) protein deposits within the cerebral and cerebellar cortex in Down's syndrome and Alzheimer's disease. *Acta Neuropathol (Berl)* 80: 318-327.
- Mao R, Wang X, Spitznagel EL, Jr., Frelin LP, Ting JC, et al. (2005). Primary and secondary transcriptional effects in the developing human Down syndrome brain and heart. *Genome Biol* 6: R107.
- Mao R, Zielke CL, Ronald Zielke H, Pevsner J (2003). Global up-regulation of chromosome 21 gene expression in the developing down syndrome brain. *Genomics* 81: 457-467.
- Martinez-Frias ML (2005). The real earliest historical evidence of Down syndrome. *Am J Med Genet A* 132: 231.
- Matalon R, Supple P, Wyandt H, Rosenthal IM (1990). Transmission of ring 14 chromosome from mother to two sons. *Am J Med Genet* 36: 381-385.
- McClure HM, Belden KH, Pieper WA, Jacobson CB (1969). Autosomal trisomy in a chimpanzee: resemblance to Down's syndrome. *Science* 165: 1010-1012.
- McCormick MK, Schinzel A, Petersen MB, Stetten G, Driscoll DJ, et al. (1989). Molecular genetic approach to the characterization of the "Down syndrome region" of chromosome 21. *Genomics* 5: 325-331.
- McKay SJ, Johnsen R, Khattra J, Asano J, Baillie DL, et al. (2003). Gene expression profiling of cells, tissues, and developmental stages of the nematode *C. elegans*. *Cold Spring Harb Symp Quant Biol* 68: 159-169.
- Melcher T, Maas S, Herb A, Sprengel R, Higuchi M, et al. (1996). RED2, a brain-specific member of the RNA-specific adenosine deaminase family. *J Biol Chem* 271: 31795-31798.
- Mello CC, Kramer JM, Stinchcomb D, Ambros V (1991). Efficient gene transfer in *C.elegans*: extrachromosomal maintenance and integration of transforming sequences. *Embo J* 10: 3959-3970.
- Miki R, Kadota K, Bono H, Mizuno Y, Tomaru Y, et al. (2001). Delineating developmental and metabolic pathways in vivo by expression profiling using the RIKEN set of 18,816 full-length enriched mouse cDNA arrays. *Proc Natl Acad Sci U S A* 98: 2199-2204.
- Mikkelsen M (1977). Down syndrome: cytogenetical epidemiology. *Hereditas* 86: 45-50.
- Moore CS, Lee JS, Birren B, Stetten G, Baxter LL, et al. (1999). Integration of cytogenetic with recombinational and physical maps of mouse chromosome 16. *Genomics* 59: 1-5.
- Morley M, Molony CM, Weber TM, Devlin JL, Ewens KG, et al. (2004). Genetic analysis of genome-wide variation in human gene expression. *Nature* 430: 743-747.

- Muragaki Y, Timmons S, Griffith CM, Oh SP, Fadel B, et al. (1995). Mouse Col18a1 is expressed in a tissue-specific manner as three alternative variants and is localized in basement membrane zones. *Proc Natl Acad Sci U S A* 92: 8763-8767.
- Murgolo NJ, Brown JE, Bayne ML, Strader CD (1996). Presenilin mutations in Alzheimer's disease: molecular models suggest a potential functional locus. *Trends Pharmacol Sci* 17: 389-393.
- Nagamine K, Peterson P, Scott HS, Kudoh J, Minoshima S, et al. (1997). Positional cloning of the APECED gene. *Nat Genet* 17: 393-398.
- Nobrega MA, Zhu Y, Plajzer-Frick I, Afzal V, Rubin EM (2004). Megabase deletions of gene deserts result in viable mice. *Nature* 431: 988-993.
- O'Doherty A, Ruf S, Mulligan C, Hildreth V, Errington ML, et al. (2005). An aneuploid mouse strain carrying human chromosome 21 with Down syndrome phenotypes. *Science* 309: 2033-2037.
- Olson LE, Richtsmeier JT, Leszl J, Reeves RH (2004). A chromosome 21 critical region does not cause specific Down syndrome phenotypes. *Science* 306: 687-690.
- Olson LE, Roper RJ, Baxter LL, Carlson EJ, Epstein CJ, et al. (2004). Down syndrome mouse models Ts65Dn, Ts1Cje, and Ms1Cje/Ts65Dn exhibit variable severity of cerebellar phenotypes. *Dev Dyn* 230: 581-589.
- Oster J, Mikkelsen M, Nielsen A (1975). Mortality and life-table in Down's syndrome. *Acta Paediatr Scand* 64: 322-326.
- Patel P, Goldberg D, Moss S (1993). Psychiatric morbidity in older people with moderate and severe learning disability. II: The prevalence study. *Br J Psychiatry* 163: 481-491.
- Petersen MB, Adelsberger PA, Schinzel AA, Binkert F, Hinkel GK, et al. (1991). Down syndrome due to de novo Robertsonian translocation t(14q;21q): DNA polymorphism analysis suggests that the origin of the extra 21q is maternal. *Am J Hum Genet* 49: 529-536.
- Pierpont ME, Markwald RR, Lin AE (2000). Genetic aspects of atrioventricular septal defects. *Am J Med Genet* 97: 289-296.
- Polani PE, Briggs JH, Ford CE, Clarke CM, Berg JM (1960). A Mongol girl with 46 chromosomes. *Lancet* 1: 721-724.
- Pritchard MA, Kola I (1999). The "gene dosage effect" hypothesis versus the "amplified developmental instability" hypothesis in Down syndrome. *J Neural Transm Suppl* 57: 293-303.
- Pueschel SM, Jackson IM, Giesswein P, Dean MK, Pezzullo JC (1991). Thyroid function in Down syndrome. *Res Dev Disabil* 12: 287-296.
- Pueschel SM, Pezzullo JC (1985). Thyroid dysfunction in Down syndrome. *Am J Dis Child* 139: 636-639.
- Rabinowe SL, Rubin IL, George KL, Adri MN, Eisenbarth GS (1989). Trisomy 21 (Down's syndrome): autoimmunity, aging and monoclonal antibody-defined T-cell abnormalities. *J Autoimmun* 2: 25-30.
- Rahmani Z, Blouin JL, Creau-Goldberg N, Watkins PC, Mattei JF, et al. (1989). Critical role of the D21S55 region on chromosome 21 in the pathogenesis of Down syndrome. *Proc Natl Acad Sci U S A* 86: 5958-5962.
- Ramirez-Solis R, Liu P, Bradley A (1995). Chromosome engineering in mice. *Nature* 378: 720-724.

- Raz N, Torres IJ, Briggs SD, Spencer WD, Thornton AE, *et al.* (1995). Selective neuroanatomic abnormalities in Down's syndrome and their cognitive correlates: evidence from MRI morphometry. *Neurology* 45: 356-366.
- Reeves RH, Baxter LL, Richtsmeier JT (2001). Too much of a good thing: mechanisms of gene action in Down syndrome. *Trends Genet* 17: 83-88.
- Reeves RH, Irving NG, Moran TH, Wohn A, Kitt C, *et al.* (1995). A mouse model for Down syndrome exhibits learning and behavior deficits. *Nat Genet* 11: 177-184.
- Reeves RH, Yao J, Crowley MR, Buck S, Zhang X, *et al.* (1994). Astrocytosis and axonal proliferation in the hippocampus of S100b transgenic mice. *Proc Natl Acad Sci U S A* 91: 5359-5363.
- Rex AP, Preus M (1982). A diagnostic index for Down syndrome. *J Pediatr* 100: 903-906.
- Reymond A, Marigo V, Yaylaoglu MB, Leoni A, Uclat C, *et al.* (2002). Human chromosome 21 gene expression atlas in the mouse. *Nature* 420: 582-586.
- Richtsmeier JT, Zumwalt A, Carlson EJ, Epstein CJ, Reeves RH (2002). Craniofacial phenotypes in segmentally trisomic mouse models for Down syndrome. *Am J Med Genet* 107: 317-324.
- Robert K, Vialard F, Thiery E, Toyama K, Sinet PM, *et al.* (2003). Expression of the cystathionine beta synthase (CBS) gene during mouse development and immunolocalization in adult brain. *J Histochem Cytochem* 51: 363-371.
- Robertson WRB (1916). Chromosome studies. I. Taxonomic relationships shown in the chromosomes of Tettigidae and Acrididae. V-shaped chromosomes and their significance in Acrididae, Locustidae and Gryllidae: chromosome and variation. *Journal of Morphology* 27: 179-331.
- Rockman-Greenberg C, Ray M, Evans JA, Canning N, Hamerton JL (1982). Homozygous Robertsonian translocations in a fetus with 44 chromosomes. *Hum Genet* 61: 181-184.
- Rogaev EI, Sherrington R, Rogaeva EA, Levesque G, Ikeda M, *et al.* (1995). Familial Alzheimer's disease in kindreds with missense mutations in a gene on chromosome 1 related to the Alzheimer's disease type 3 gene. *Nature* 376: 775-778.
- Roper RJ, Baxter LL, Saran NG, Klinedinst DK, Beachy PA, *et al.* (2006). Defective cerebellar response to mitogenic Hedgehog signaling in Down [corrected] syndrome mice. *Proc Natl Acad Sci U S A* 103: 1452-1456.
- Roth GM, Sun B, Greensite FS, Lott IT, Dietrich RB (1996). Premature aging in persons with Down syndrome: MR findings. *AJNR Am J Neuroradiol* 17: 1283-1289.
- Rual JF, Ceron J, Koreth J, Hao T, Nicot AS, *et al.* (2004). Toward improving *Caenorhabditis elegans* phenotype mapping with an ORFeome-based RNAi library. *Genome Res* 14: 2162-2168.
- Ryan ET, Ecker JL, Christakis NA, Folkman J (1992). Hirschsprung's disease: associated abnormalities and demography. *J Pediatr Surg* 27: 76-81.
- Sago H, Carlson EJ, Smith DJ, Kilbridge J, Rubin EM, *et al.* (1998). Ts1Cje, a partial trisomy 16 mouse model for Down syndrome, exhibits learning and behavioral abnormalities. *Proc Natl Acad Sci U S A* 95: 6256-6261.
- Sago H, Carlson EJ, Smith DJ, Rubin EM, Crnic LS, *et al.* (2000). Genetic dissection of region associated with behavioral abnormalities in mouse models for Down syndrome. *Pediatr Res* 48: 606-613.

- Saito T, Guan F, Papolos DF, Lau S, Klein M, et al. (2001). Mutation analysis of SYNJ1: a possible candidate gene for chromosome 21q22-linked bipolar disorder. *Mol Psychiatry* 6: 387-395.
- Salehi A, Delcroix JD, Belichenko PV, Zhan K, Wu C, et al. (2006). Increased App expression in a mouse model of Down's syndrome disrupts NGF transport and causes cholinergic neuron degeneration. *Neuron* 51: 29-42.
- Saran NG, Pletcher MT, Natale JE, Cheng Y, Reeves RH (2003). Global disruption of the cerebellar transcriptome in a Down syndrome mouse model. *Hum Mol Genet* 12: 2013-2019.
- Satge D, Sasco AJ, Geneix A, Malet P (1998). Another reason to look for tumor suppressor genes on chromosome 21. *Genes Chromosomes Cancer* 21: 1.
- Scartezzini P, Egeo A, Colella S, Fumagalli P, Arrigo P, et al. (1997). Cloning a new human gene from chromosome 21q22.3 encoding a glutamic acid-rich protein expressed in heart and skeletal muscle. *Hum Genet* 99: 387-392.
- Schmucker D, Clemens JC, Shu H, Worby CA, Xiao J, et al. (2000). Drosophila Dscam is an axon guidance receptor exhibiting extraordinary molecular diversity. *Cell* 101: 671-684.
- Schuchter K, Hafner E, Stangl G, Ogris E, Philipp K (2001). Sequential screening for trisomy 21 by nuchal translucency measurement in the first trimester and maternal serum biochemistry in the second trimester in a low-risk population. *Ultrasound Obstet Gynecol* 18: 23-25.
- Seger R, Buchinger G, Stroder J (1977). On the influence of age on immunity in Down's syndrome. *Eur J Pediatr* 124: 77-87.
- Selkoe DJ (1997). Alzheimer's disease: genotypes, phenotypes, and treatments. *Science* 275: 630-631.
- Shaffer LG, Jackson-Cook CK, Stasiowski BA, Spence JE, Brown JA (1992). Parental origin determination in thirty de novo Robertsonian translocations. *Am J Med Genet* 43: 957-963.
- Shapiro BL (1999). The Down syndrome critical region. *J Neural Transm Suppl* 57: 41-60.
- Sheridan R, Llerena J, Jr., Watkins S, Debenham P, Cawood A, et al. (1989). Fertility in a male with trisomy 21. *J Med Genet* 26: 294-298.
- Sherman S (1992). Correction of the evaluation of recombination in meiosis I and II nondisjunction in trisomy 21. *Am J Hum Genet* 50: 1137-1138.
- Sherman SL, Takaesu N, Freeman SB, Grantham M, Phillips C, et al. (1991). Trisomy 21: association between reduced recombination and nondisjunction. *Am J Hum Genet* 49: 608-620.
- Sherzer AL (1992) The adolescent. In: Rogers PT, Coleman M, editors. *Medical care in Down syndrome: a preventive medicine approach*. New York: Marcel Dekker Inc. pp. 115-144.
- Shinohara T, Tomizuka K, Miyabara S, Takehara S, Kazuki Y, et al. (2001). Mice containing a human chromosome 21 model behavioral impairment and cardiac anomalies of Down's syndrome. *Hum Mol Genet* 10: 1163-1175.
- Simmer F, Moorman C, van der Linden AM, Kuijk E, van den Berghe PV, et al. (2003). Genome-wide RNAi of *C. elegans* using the hypersensitive rrf-3 strain reveals novel gene functions. *PLoS Biol* 1: E12.
- Simmer F, Tijsterman M, Parrish S, Koushika SP, Nonet ML, et al. (2002). Loss of the putative RNA-directed RNA polymerase RRF-3 makes *C. elegans* hypersensitive to RNAi. *Curr Biol* 12: 1317-1319.

- Slordahl SH, Smeland EB, Holte H, Grønn M, Lie SO, et al. (1993). Leukemic blasts with markers of four cell lineages in Down's syndrome ("megakaryoblastic leukemia"). *Med Pediatr Oncol* 21: 254-258.
- Snijders RJ, Noble P, Sebire N, Souka A, Nicolaides KH (1998). UK multicentre project on assessment of risk of trisomy 21 by maternal age and fetal nuchal-translucency thickness at 10-14 weeks of gestation. *Fetal Medicine Foundation First Trimester Screening Group. Lancet* 352: 343-346.
- Sonnichsen B, Koski LB, Walsh A, Marschall P, Neumann B, et al. (2005). Full-genome RNAi profiling of early embryogenesis in *Caenorhabditis elegans*. *Nature* 434: 462-469.
- Steinfath M, Wruck W, Seidel H, Lehrach H, Radelof U, et al. (2001). Automated image analysis for array hybridization experiments. *Bioinformatics* 17: 634-641.
- Sulston J, Hodgkin J (1988) Methods. In: Wood WB, editor. *The Nematode Caenorhabditis elegans*. New York: Cold Spring Harbor Laboratory Press. pp. 587-576.
- Tabara H, Grishok A, Mello CC (1998). RNAi in *C. elegans*: soaking in the genome sequence. *Science* 282: 430-431.
- Taketo MM, Araki Y, Matsunaga A, Yokoi A, Tsuchida J, et al. (1997). Mapping of eight testis-specific genes to mouse chromosomes. *Genomics* 46: 138-142.
- Teipel SJ, Alexander GE, Schapiro MB, Moller HJ, Rapoport SI, et al. (2004). Age-related cortical grey matter reductions in non-demented Down's syndrome adults determined by MRI with voxel-based morphometry. *Brain* 127: 811-824.
- The *C. elegans* Sequencing Consortium (1998). Genome sequence of the nematode *C. elegans*: a platform for investigating biology. *Science* 282: 2012-2018.
- Timmons L, Fire A (1998). Specific interference by ingested dsRNA. *Nature* 395: 854.
- Tinel N, Diochot S, Borsotto M, Lazdunski M, Barhanin J (2000). KCNE2 confers background current characteristics to the cardiac KCNQ1 potassium channel. *Embo J* 19: 6326-6330.
- Towbin JA, Vatta M (2001). Molecular biology and the prolonged QT syndromes. *Am J Med* 110: 385-398.
- Toyoda A, Noguchi H, Taylor TD, Ito T, Pletcher MT, et al. (2002). Comparative genomic sequence analysis of the human chromosome 21 Down syndrome critical region. *Genome Res* 12: 1323-1332.
- Vacik T, Forejt J (2003). Quantification of expression and methylation of the Igf2r imprinted gene in segmental trisomic mouse model. *Genomics* 82: 261-268.
- Vogel F, Motulsky AJ (1986) Human genetics Problems and approaches: Springer Verlag Berlin Heidelberg New York Tokyo.
- White NS, Alkire MT, Haier RJ (2003). A voxel-based morphometric study of nondemented adults with Down Syndrome. *Neuroimage* 20: 393-403.
- Wilson L, Curtis A, Korenberg JR, Schipper RD, Allan L, et al. (1993). A large, dominant pedigree of atrioventricular septal defect (AVSD): exclusion from the Down syndrome critical region on chromosome 21. *Am J Hum Genet* 53: 1262-1268.
- Wisniewski HM, Rabe A (1986). Discrepancy between Alzheimer-type neuropathology and dementia in persons with Down's syndrome. *Ann N Y Acad Sci* 477: 247-260.

- Wisniewski KE (1990). Down syndrome children often have brain with maturation delay, retardation of growth, and cortical dysgenesis. *Am J Med Genet Suppl* 7: 274-281.
- Wisniewski KE, Dalton AJ, McLachlan C, Wen GY, Wisniewski HM (1985). Alzheimer's disease in Down's syndrome: clinicopathologic studies. *Neurology* 35: 957-961.
- Wisniewski KE, Laure-Kamionowska M, Wisniewski HM (1984). Evidence of arrest of neurogenesis and synaptogenesis in brains of patients with Down's syndrome. *N Engl J Med* 311: 1187-1188.
- Yamakawa K, Huot YK, Haendelt MA, Hubert R, Chen XN, et al. (1998). DSCAM: a novel member of the immunoglobulin superfamily maps in a Down syndrome region and is involved in the development of the nervous system. *Hum Mol Genet* 7: 227-237.
- Yang Q, Rasmussen SA, Friedman JM (2002). Mortality associated with Down's syndrome in the USA from 1983 to 1997: a population-based study. *Lancet* 359: 1019-1025.
- Yang X, Aoki Y, Li X, Sakamoto O, Hiratsuka M, et al. (2001). Structure of human holocarboxylase synthetase gene and mutation spectrum of holocarboxylase synthetase deficiency. *Hum Genet* 109: 526-534.
- Zambrowicz BP, Friedrich GA (1998). Comprehensive mammalian genetics: history and future prospects of gene trapping in the mouse. *Int J Dev Biol* 42: 1025-1036.
- Zhang G, Brokx S, Weiner JH (2006). Extracellular accumulation of recombinant proteins fused to the carrier protein YebF in *Escherichia coli*. *Nat Biotechnol* 24: 100-104.
- Zigman W, Schupf N, Haveman M, Silverman W (1997). The epidemiology of Alzheimer disease in intellectual disability: results and recommendations from an international conference. *J Intellect Disabil Res* 41 (Pt 1): 76-80.
- Zipursky A, Peeters M, Poon A (1987). Megakaryoblastic leukemia and Down's syndrome--a review. *Prog Clin Biol Res* 246: 33-56.