

H. LITERATURVERZEICHNIS

- AKIYAMA, M., I. MATSUO, H. SHIMUZU (2002):
Formation of cornified cell envelope in human hair follicle development.
British Journal of Dermatology 146, 968-976
- ANDREADIS, ST., K.E. HAMOEN, M.L. YARMUSH, J.R. MORGAN (2001):
Keratinocyte growth factor induces hyperproliferation and differentiation in a skin equivalent model system.
FASEB J. 15 (6), 898-906
- ANDRIANI, F., J. GARFIELD, N. E. FUSENIG, J. A. GARLICK (2004):
Basement membrane proteins promote progression of intraepithelial neoplasia in 3-dimensional models of human stratified epithelium.
Int J Cancer 108 (3), 348-357
- ANTHAUER, K., CH. MÜLLING, K. D. BUDRAS (2005):
Membrane-coating granules and the intercellular cementing substance (membrane-coating material) in the epidermis in different regions of the equine hoof.
Anat Histol Embryol. 34 (5), 298-306
- ASHKENAS, J., J. MUSCHLER, M. J BISSELL (1996):
The extracellular matrix in epithelial biology: shared molecules and common themes in distant phyla.
Dev Biol 180 (2), 433-444
- BADEN, H. P., L. D. LEE, J. KUBILUS (1976):
The fibrous proteins of stratum corneum.
J Invest Dermatol 67 (5), 573-576
- BADEN, H. P., J. KUBILUS (1984):
A comparative study of the immunologic properties of hoof and nail fibrous proteins.
J Invest Dermatol 83 (8), 327-331
- BADEN, H. P. AND J. C. KVEDAR (1993):
Epithelial cornified envelope precursors are in the hair follicle and nail.
J Invest Dermatol 101 (1 Suppl), 72-74
- BANG, F. B. (1977):
History of tissue culture at Johns Hopkins.
Bull Hist Med 51, 516-537
- BANNO, T., A. GAZEL, M. BLUMENBERG (2004):
Effects of tumor necrosis factor-alpha (TNF alpha) in epidermal keratinocytes revealed using global transcriptional profiling.
J Biol Chem 279 (31), 32633-32642
- BERG, J. M. T., JOHN L.; STRYER, LUBERT (2003):
Biochemie.
Heidelberg, Berlin, New York, Spectrum.

BICKENBACH, J. R., E. CHISM (1998):

Selection and extended growth of murine epidermal stem cells in culture.
Exp Cell Res 244 (1), 184-195

BILLINGHAM, R. E., J. REYNOLDS (1952):

Transplantation studies on sheets of pure epidermal epithelium and on epidermal cell suspensions.
Br J Plast Surg 5, 25-36

BIRKNER, S., S. WEBER, J. KUHLMANN, O. MÜLLER, W. FÖLLMANN (1998):

Development of a new molecular biological in vitro assay for toxicological studies using cultured bovine urinary bladder epithelial cells.
Nauyn-Schmiedeberg's Arch. Pharmacol.

BLACK, R. A., C. T. RAUCH, C. J. KOZLOSKY, J. J. PESCHON, J. L. SLACK, M. F. WOLFSON, B. J. CASTNER, K. L. STOCKING, P. REDDY, S. SRINIVASAN, N. NELSON, N. BOIANI, K.A. SCHOOLEY, M. GERHART, R. DAVIS, J. N. FITZNER, R. S. JOHNSON, R. J. PAXTON, C. J. MARCH, D. P. CERRETTI (1997):

A metalloproteinase disintegrin that releases tumor-necrosis-factor- α from cells.
Nature 385, 729-733

BLUNK, T., A. L. SIEMINSKI, K. J. GOOCH, D. L. COURTER, A. P. HOLLANDER, A. M. NAHIR, R. LANGE, G. VUNJAK-NOVAKOVIC, L. E. FREED (2002):

Differential effects of growth factors on tissue-engineered cartilage.
Tissue Eng 8 (1), 73-84

BORAS, V. V., N. CIKES, J. LUKAC, A. CEKIC-ARAMBASIN, M. VIRAG, A. BOSNJAK (2004):

The significance of salivary and serum interleukin 6 and basic fibroblast growth factor levels in patients with Sjogrens syndrome.
Coll Antropol., 28 Suppl 2, 305-309

BOWDEN, P. E., H. J. STARK, D. BREITKREUTZ, N. E. FUSENIG (1987):

Expression and modification of keratins during terminal differentiation of mammalian epidermis.
Curr Top Dev Biol 22, 35-68

BRAGULLA, H. (1991):

The deciduous hoof capsule (*Capsula ungulae decidua*) of the equine fetus and newborn foal
Anat Histol Embryol 20 (1), 66-74

BRAGULLA, H., CH. MÜLLING (1992):

Light- and ultramicroscopic studies of hoof horn keratinization with special reference to aseptic inflammation of the bovine and equine digit.
Proc 7th International Symposium on Disorders of the Ruminant Digit.
21.-25.06.1992, Rebild/DK, S8

- BRAGULLA, H., CH. MÜLLING (1994):
Veränderungen der Architektur der Hornzellen und des Hornzellverbandes bei Klauenerkrankungen.
Proc 20th Congress of the European Association of Veterinary Anatomists.
15.-19.08.1994, Zürich/CH, S 43-44
- BRAGULLA, H., K. D. BUDRAS, J. D. REILLY (1998):
Fetal development of the white line (Zona alba) of the equine hoof.
Equine Vet J Suppl (26), 22-6
- BRAUN, S., U. AUF DEM KELLER, H. STEILING, S. WERNER (2004):
Factors in epithelial repair and cytoprotection.
Philos Trans R Soc Lond B Biol Sci; 359(1445) 753-757
- BUCHER (1997):
Cytologie, Histologie und mikroskopische Anatomie des Menschen.
Hans Huber Verlag
- BUDRAS, K. D., A. WÜNSCHE (2002):
Atlas der Anatomie des Rindes.
Hannover, Schlütersche
- BUDRAS, K. D., R. L. HULLINGER, W. O. SACK (1989):
Light and electron microscopy of keratinization in the laminar epidermis of the equine hoof with reference to laminitis.
Am J Vet Res 50 (7), 1150-1160
- BUDRAS, K. D., M. SEIDEL (1992):
The segmental organization and horny structure of the claw of the dog.
Anat Histol Embryol 21 (4), 348-363
- BUDRAS, K. D., CH. MÜLLING, A. HOROWITZ (1996):
Rate of keratinization of the wall segment of the hoof and its relation to width and structure of the zona alba (white line) with respect to claw disease in cattle.
Am J Vet Res 57 (4), 444-455
- BUDRAS, K. D., B. HUSKAMP (1995):
Die Hornqualität des Pferdehufes und deren Verbesserung nach einer orthopädischen Behandlung der Hufrehe.
In: KNEZEVIC, P.F. (Hrsg.): Orthopädie bei Huf- und Klauentieren.
Stuttgart, Schattauer Verlag, 252-268
- BYRNE, C., M. HARDMAN, K. NIELD (2003):
Covering the limb-formation of the integument.
J Anat 202 (1), 113-123
- CASASCO, A., M. CASASCO, N. ZERBINATI, A. ICAROCORNAGLIA, A. CALLIGARO (2001):
Cell proliferation and differentiation in a model of human skin equivalent
The anatomical record 264, 261-272

- CHEN, G., D. V. GOEDDEL (2002):
TNF-R1 signaling: a beautiful pathway.
Science 296 (5573), 1634-1635
- COHEN, S. (1962):
Isolation of a mouse submaxillary gland protein accelerating incisor eruption and eyelid opening in the new-born animal.
J Biol Chem 237, 1555–1562
- COHEN, S. (1965):
The stimulation of epidermal proliferation by a specific protein (EGF).
Develop Biol 12, 394–399
- COOPER, D., T.-T. SUN (1986):
Monoclonal antibody analysis of bovine epithelial keratins.
J Biol Chem 261 (10), 4646-4654
- DINARELLO, C. A. (1994):
The biological properties of interleukin-1.
Eur Cytokine Netw 5 (6), 517-531
- DIRKS, C. (1985):
Makroskopische, licht- und elektronenmikroskopische Untersuchungen über den Rückenteil der Rinderklaue.
Berlin: FUB, FB Vet Med, Diss
- DUNNWALD, M., A. TOMANEK-CHALKLEY, D. ALEXANDRUNAS, J. FISHBAUGH, J. R. BICKENBACH (2001):
Isolating a pure population of epidermal stem cells for use in tissue engineering.
Exp Dermatol 10 (1), 45-54
- ECKHART, L., C. REINISCHINOUE, P. MESSNER, M. DOCKAL, C. MAYER, E. TSCHACHLER (2003):
A basement membrane-like matrix formed by cell-released proteins at the medium/air interface supports growth of keratinocytes.
Eur J Cell Biol 82 (11), 549-555
- ECKFALCK, A., B. JONES, N. OBEL (1991):
Possible importance for laminitis research of recent studies on substances influencing the differentiation of cultured keratinocytes.
J Vet Med A Physiol Pathol Clin Med 38, 721-727
- EGGERS, T. (2001):
Die Wundheilung des Rusterholzschen Klauengeschwüres beim Rind: Licht- und transmissionselektronenmikroskopische Auswertung einer kontrollierten Studie zur Wundheilung und zum Einfluss von Biotin auf den Heilungsverlauf.
Berlin: FUB, FB Vet Med, Diss
- EL GHALBZOURI, A., E. LAMME, M. PONEC (2002):
Crucial role of fibroblasts in regulating epidermal morphogenesis.
Cell Tissue Res 310 (2), 189-199

ENGEL, M. (1994):

Paradigmenwechsel und Exodus.
Zellbiologie, Zellchemie und Biochemie in Berlin.
Fischer Verlag

FALEIROS, R. R., A. M. STOKES, S. C. EADES, D. Y. KIM, D. B. PAULSEN, R. M. MOORE (2004):

Assessment of apoptosis in epidermal lamellar cells in clinically normal horses and those with laminitis.
Am J Vet Res 65 (5), 578-585

FINCH, P. W., J. S. RUBIN (2004):

Keratinocyte growth factor/fibroblast growth factor 7, a homeostatic factor with therapeutic potential for epithelial protection and repair.
Adv Cancer Res. 91, 69-136

FRÖDE, D. (1998):

Untersuchungen zum Einfluss freier Fettsäuren auf in vitro kultivierte humane Keratinozyten.
Dissertation, Math.-Nat.-Tech. Fak., Martin-Luther-Universität Halle-Wittenberg

FROHNES, A. K. (1999):

Struktur, Verhornung und Hornqualität im Sohlen- und Ballen-Strahlsegment des Pferdehufes.
Berlin: FUB, FB Vet Med, Diss

FÜRST, A. (1992):

Makroskopische und mikroskopische Anatomie der Rinderklaue.
Zürich: Univ., Vet Med Fak, Diss.

FUCHS, E., A. L. TYNER, G. J. GIUDICE, D. MARCHUK, A. R. CHAUDHURY, M. ROSENBERG (1987):

The human keratin genes and their differential expression.
Curr Top Dev Biol 22, 5-34

GALLE, J., M. LOEFFLER, D. DRASDO (2005):

Modelling the effect of deregulated proliferation and apoptosis on the growth dynamics of epithelial cell populations in vitro.
Biophys J. Vol. 88 (1), 62-75

GAN, S. Q., O. W. MCBRIDE, W. W. IDLER, N. MARKOVA, P. M. STEINERT (1991):

Organization, structure, and polymorphisms of the human profilaggrin gene.
Biochemistry 30 (23), 5814

GEORGE K., K. GEORGE (2001):

TNF and receptors in organ-specific autoimmune disease: multi-layered functioning mirrored in animal models.

J Clin Invest, Volume 107, Number 12, 1507-1508

GREEN, K. J., J. C. JONES (1996):

Desmosomes and hemidesmosomes: structure and function of molecular components.
Faseb J 10 (8), 871-881

GRONE, A. (2002):

Keratinocytes and cytokines.
Vet Immunol Immunopathol 88 (1-2), 1-12

HABERMEHL, K. H. (1984):

Haut und Hautorgane. In: Nickel, Schummer et Seiferle (Hrsg.)
Lehrbuch der Anatomie der Haustiere 3.Aufl. Bd. III. Berlin, Parey Verlag

HAFTEK, M., G. SERRE, V. MILS, J. THIVOLET (1991):

Immunocytochemical evidence for a possible role of cross-linked keratinocyte envelopes in stratum corneum cohesion.
J Histochem Cytochem 39 (11), 1531-1538

HASHIMOTO, K. (1971):

Ultrastructure of the human toenail. II. Keratinization and formation of the marginal band. Ultrastruct Res 36 (3), 391-410

HAYWARD, A. F. (1979):

Membrane Coating Granules.
Int. Rev. Cytol. 59, 97-127

HENDRY, K. A., A. J. MACCALLUM, C. H. KNIGHT, C. J. WILDE (2001):

Synthesis and distribution of cytokeratins in healthy and ulcerated bovine claw epidermis.

J Dairy Res 68 (4), 525-537

HENDRY, K. A., C. H. KNIGHT, H. GALBRAITH, C. J. WILDE (2003):

Basement membrane integrity and keratinisation in healthy and ulcerated bovine hoof tissue.

J Dairy Res 70, 19-27

HINES, M. D., H. C. JIN, M. J. WHEELOCK, P. J. JENSEN (1999):

Inhibition of cadherin function differentially affects markers of terminal differentiation in cultured human keratinocytes.

J Cell Sci 112 (Pt 24), 4569-4579

HINTERHUBER, G., Y. MARQUARDT, E. DIEM, K. RAPPERSBERGER, K. WOLFF, D. FOEDINGER (2002):

Organotypic keratinocyte coculture using normal human serum:
An immunomorphological study at light and electron microscopic levels.
Exp Dermatol 123, 503-515

- HINTNER, H., G. R. NEISES, T. J. LAWLEY (1985):
Immunologic properties of enzymatically degraded human keratin intermediate filaments.
J Invest Dermatol 84 (2), 108-113
- HIRSCHBERG, R. M., CH. MÜLLING, H. BRAGULLA (1999):
Microvasculature of the bovine claw demonstrated by improved micro-corrosion-casting technique.
Microsc Res Tech 45 (3), 184-197
- HOCHSTETTER, T. (1998):
Die Hornqualität der Rinderklaue unter Einfluss einer Biotinsupplementierung.
Berlin: FUB, FB Vet Med, Diss
- HOFFMANN, D., U. NEBEL, CH: MÜLLING, K. D. BUDRAS (2004):
Long-term cultivation of bovine hoof cells in a novel perfusion chamber system.
25th Congress of the European Association of Veterinary Anatomists.
Norwegian School of Veterinary Science 2004, 94
- HOHL, D., B. RUF OLANO, P.A. DE VIRAGH, M. HUBER, C. J. DETRISAC, U. W. SCHNYDER, D. R. ROOP (1993):
Expression patterns of loricrin in various species and tissues.
Differentiation 54 (1), 25-34
- HOHL, D. (2005):
Formation of the cornified envelope.
Exp Dermatol. Oct; 14 (10), 770-780
- HOHMANN, H. (1902):
Untersuchungen über die Klauenleiderhaut des Rindes.
Monatsh Prakt Tierheilkd 13, 49-96
- HUANG, J., T. ASAWA, T. TAKATO, R. SAKAI (2003):
Cooperative roles of Fyn and cortactin in cell migration of metastatic murine melanoma.
J Biol Chem 278 (48), 48367-48376
- HUBER, M., O. DISTL, F. GRAF, H. KRÄUSLICH (1984):
Die Entwicklung der Klauenform von Jungbüffeln im Alter von 6-12 Monaten
Zbl. Vet. Med. A 31, 499-507
- IIJIMA M., T. HASHIMOTO, Y. MATSUDA, T. NAGAI, Y. YAMANO, T. ICHI, T. OSAKI, S. KAWABATA (2005):
Comprehensive sequence analysis of horseshoe crab cuticular proteins and their involvement in transglutaminase-dependent cross-linking.
Febs J. 272 (18), 4774-4786

ISHIDA-YAMAMOTO, A., H. TAKAHASHI, R. B. PRESLAND, B. A. DALE, H. IIZUKA (1998):

Translocation of profilaggrin N-terminal domain into keratinocyte nuclei with fragmented DNA in normal human skin and loricrin keratoderma.
Lab Invest 78 (10), 1245-1253

ISHIDA-YAMAMOTO, A., H. TANAKA, H. NAKANE, H. TAKAHASHI, Y. HASHIMOTO, H. IIZUKA (1999):

Programmed cell death in normal epidermis and loricrin keratoderma. Multiple functions of profilaggrin in keratinization.
J Investig Dermatol Symp Proc 4 (2), 145-149

JACOBSON, M. D., M. WEIL, M. C. RAFF (1997):

Programmed Cell Death.
Animal Development Cell 88, 374-354

JAHAVERIAN, A., M. VACCARIELO, N. E. FUSENIG, J.A. GARLICK (1998):

Normal keratinocytes suppress early stages of neoplastic progression in stratified epithelium.
Cancer Res 58, 2200-2208

JARNIK, M., M. N. SIMON, A. C. STEVEN (1998):

Cornified cell envelope assembly: a model based on electron microscopic determinations of thickness and projected density.
J Cell Sci 111 (Pt 8), 1051-1060

JESSEN, H., O. BEHNKE (1986):

Selective binding of colloidal gold-protein conjugates to epidermal phosphorus-rich keratoxyaline granules and cornified cells.
J Invest Dermatol 87 (6), 737-740

KAMMANN, U., M. LACORN, G. PIECHOTTA, W. WOSNIOK, T. J. SIMAT, T. LANG, W. E. G. MÜLLER, H. C. SCHRÖDER, H.-S. JENKE, H. STEINHART (2001):

Annual cycles of apoptosis, heat shock proteins, strand scission factor, and metallothionein isoforms in the marine flatfish dab (*Limanda limanda*): Influences of natural factors and consequences for biomonitoring.
Biomarkers 6 (2), 108–126

KITAHARA, T., H. OGAWA (1994):

Variation of differentiation in nail and bovine hoof cells
J Invest Dermatol 102, 725-729

KOPPELSTÄTTER, CH. (2000):

Zellkultur unter kontinuierlicher Nährstoff-Perfusion als neuer Weg bei in vitro Toxizitätstest. Eine Alternative zu Tierversuchen
Universität Innsbruck Dr. med. univ. Diss.

KOWALEWSKI, C., A. KOZLOWSKA, M. ZAWADZKA, K. WOZNIAK, M.
BLASZCZYK, S. JABLONSKA (2004):

Alterations of Basement Membrane Zone in Bullous and Non-Bullous Variants of
Extragenital Lichen Sclerosus.
Am J Dermatopathol 26 (2), 96-101

KUBILUS, J., I. SCOTT, C. R. HARDING, J. YENDLE, J. KVEDAR, H. P. BADEN (1985):
The occurrence of profilaggrin and its processing in cultured keratinocytes.
J Invest Dermatol 85 (6), 513-517

KÜNZEL, E. (1990):

Haut (Integumentum commune).

In: Mosimann, W., T. Kohler: Zytologie, Histologie und mikroskopische Anatomie
der Haussäugetiere, Kap 16 Berlin, Parey Verlag, 259-287

KVEDAR, J. C., J. KUBILUS, H.P. BADEN (1986):

Cytokeratins of the bovine hoof: classification and studies on expression.
Biochim Biophys Acta 884 (3), 462-473

LAKARTIDN, S. (1960):

Carl August Ljunggren-an international tissue culture pioneer.
Swedish Article 57, 1707-1709

LANDMANN, L. (1988):

The permeability barrier.
Anat. Embryol. 178, 1-13

LANDMANN, L. (1991):

The permeability barrier of the skin.
Pharm Unserer Zeit 20 (4), 155-163

LANGER, R. (2000):

Tissue Engineering.
Mol. Therapy., vol. 1, number 1, 12-15

LARSSON, B., N. OBEL, B. ABERG (1956):

On the biochemistry of keratinization in the matrix of horses hoof in normal conditions
and in laminitis.
Nord vet Med 8, 761-776

LEACH, D. H. (1993):

Structural changes in intercellular junctions during keratinization of the stratum
medium of the equine hoof wall.
Acta Anat (Basel) 147 (1), 45-55

LEE, L. D., H. P. BADEN (1976):

Organisation of the polypeptide chains in mammalian keratin.
Nature 264 (5584), 377-379

LEE, L. D., H. P. BADEN, J. KUNILUS, B. F. FLEMING (1976):
Immunology of epidermal fibrous proteins.
J Invest Dermatol 67 (4), 521-525

LEE, C. H., C. L. YU, W. T. LIAO, Y. H. KAO, C. Y. CHAI, G. S. CHEN, H. S. YU (2004):
Effects and interactions of low doses of arsenic and UVB on keratinocyte apoptosis.
Chem Res Toxicol 17 (9), 1199-205

LEHMANN, P., S. KLOTH, J. AIGNER, R. DAMMER, W.W. MINUTH (1997):
Lebende Langzeitkonservierung von humaner Gingiva in der Perfusionskultur.
Mund Kiefer Gesichts Chir 1, 26-30

LOCKSLEY, R. M., N. KILLEEN, M. J. LENARDO (2001):
The TNF and TNF receptor superfamilies: integrating mammalian biology.
Cell 104 (4), 487-501

MAAS-SZABOWSKI, N., A. SHIMOTOYODOME, N. E. FUSENIG (1999):
Keratinocyte growth regulation in fibroblast cocultures via a double paracrine mechanism.
J Cell Sci 112 (Pt 12), 1843-1853

MAAS-SZABOWSKI, N., A. SZABOWSKI, H.-J. STARK, S. ANDRECHT, A. KOLBUS,
M. SCHROPP-KISTNER, P. ANGEL, N. E. FUSENIG (2001):
Organotypic cocultures with genetically modified mouse fibroblasts as a tool to dissect
molecular mechanisms regulating keratinocyte growth and differentiation.
J Invest Dermatol 116, 816-820

MAKELA, M., T. SALO, H. LARJAVA (1998):
MMP-9 from TNF alpha-stimulated keratinocytes binds to cell membranes and type I
collagen: a cause for extended matrix degradation in inflammation?
Biochem Biophys Res Commun, 253 (2), 325-335

MARKOVA, N. G., L. N. MAREKOV, C. C. CHIPEV, S. Q. GAN, W. W. IDLER, P. M.
STEINERT (1993):
Profilaggrin is a major epidermal calcium-binding protein.
Mol Cell Biol 13 (1), 613-625

MATOLTSY, A. G., P. F. PARRAKAL (1965):
Membrane-coating granules of keratinizing epithelia.
J. Cell Biol. 24, 297-307

MATOLTSY, A. G. (1966):
Membrane-coating granules of the epidermis.
J. Ultrastruc. Res. 15, 510-515

MATOLTSY, A. G. (1976):
Keratinization.
J Invest Dermatol 67 (1), 20-25

MCCALL, C. A., J. J. COHEN (1991):

Programmed cell death in terminally differentiating keratinocytes: role of endogenous endonuclease.

J Invest Dermatol 97 (1), 111-114

MCMILLAN, J. R., H. SHIMIZU (2001):

Desmosomes: structure and function in normal and diseased epidermis.

J Dermatol 28 (6), 291-298

MEHREL, T., D. HOHL, J. A. ROTHNAGEL, M. A. LONGLEY, D. BUNDMAN, C.

CHENG, U. LICHTI, M. E. BISHER, A. C. STEVEN, P. M. STEINERT (1990):

Identification of a major keratinocyte cell envelope protein, loricrin.

Cell 61 (6), 1103-1112

MERKER, H. J. (1994):

Morphology of the basement membrane.

Microsc Res Tech 28 (2), 95-124

MILSTONE, L. M., J. MCGUIRE (1981):

Different polypeptides form the intermediate filaments in bovine hoof and esophageal epithelium and in aortic endothelium.

J Cell Biol 88 (2), 312-6

MINUTH, W. W., K. SCHUMACHER, R. STREHL (2005):

Renal epithelia in long term gradient culture for biomaterial testing and tissue engineering.

Biomed Mater Eng 15 (1-2), 51-63

MINUTH, W. W., K. SCHUMACHER, R. STREHL, S. KLOTH (2000):

Physiological and cell biological aspects of perfusion culture technique employed to generate differentiated tissues for long term biomaterial testing and tissue engineering.

J Biomater Sci Polym Ed 11 (5), 495-522

MINUTH, W. W., P. STEINER, R. STREHL, K. SCHUMACHER, U. DE VRIES, S.

KLOTH (1999):

Modulation of cell differentiation in perfusion culture.

Exp Nephrol 7 (5-6), 394-406

MINUTH, W. W., R. STREHL, K. SCHUMACHER (2004):

Tissue factory: conceptual design of a modular system for the in vitro generation of functional tissues.

Tissue Eng 10 (1-2), 285-294

MINUTH, W. W., R. STREHL, K. SCHUMACHER, U. DE VRIES (2001):

Long term culture of epithelia in a continuous fluid gradient for biomaterial testing and tissue engineering.

J Biomater Sci Polym Ed 12 (3), 353-365

MOCHIZUKI, R., M. KAMIYAMA, K. Y. ARAI, K. ARAI, K. UEHARA (2001):

Expression of desmosomal proteins in rat keratinocytes during in vitro differentiation.

J Vet Med Sci 64 (2), 123-127

- MOLL, R., W. W. FRANKE, B. VOLC-PLATZER, R. KREPLER (1982):
Different keratin polypeptides in epidermis and other epithelia of human skin: a specific cytokeratin of molecular weight 46,000 in epithelia of the pilosebaceous tract and basal cell epitheliomas.
J Cell Biol 95 (1), 285-295
- MOSSNER, R., I. BECKMANN, C. HALLERMANN, C. NEUMANN, K. REICH (2004):
Granulocyte colony-stimulating-factor-induced psoriasisform dermatitis resembles psoriasis with regard to abnormal cytokine expression and epidermal activation.
Exp. Dermatol. 13 (6), 340-346
- MÜLLING, CH. (1993):
Struktur, Verhornung und Hornqualität in Ballen, Sohle und weißer Linie der Rinderklaue und ihre Bedeutung für Klauenerkrankungen.
Berlin: FUB, FB Vet Med, Diss
- MÜLLING, CH., H. BRAGULLA, K. D. BUDRAS, S. REESE (1994):
Structural factors influencing the horn quality and predilection sites for diseases at the bottom surface of the bovine hoof.
Schweiz Arch Tierheilkd 136 (2), 49-57
- MÜLLING, CH., K. D. BUDRAS (1998):
Subsurface crypts and interstitial cells in the ovary of the seal (*Phoca vitulina vitulina*): light and electron microscopic findings.
Ital J Anat Embryol 103 (4 Suppl 1), 167-181
- MÜLLING, CH., CH. LISCHER (2002):
Neue Aspekte zur Ätiologie und Pathogenese der Klauenrehe.
Vet-Med Report, Sonderausgabe V4, 12-13
- MÜLLING, CH., K. D. BUDRAS (2002):
Morphology of the dermo-epidermal junction of the bovine hoof with reference to its biological function.
Wien. Tierärztl. Mschr. 89, 188-196
- NEBEL, U. (2005):
In vitro Modelle proliferierender und differenzierender epidermaler Keratinozyten der Rinderklaue.
Berlin: FUB, FB Vet Med, Diss
- NEUß-STEIN, S. (2004):
Untersuchungen zur Bedeutung mesenchymaler Stammzellen in der Gewebe-regeneration.
TH Aachen, Diss
- PAPINI S., D. CECCHETTI, D. CAMPANI, W. FITZGERALD, J. C. GRIVEL, S. CHEN, L. MARGOLIS, R.P. REVOLTELLA (2003):
Isolation and clonal analysis of human epidermal keratinocyte stem cells in long-term culture.
Stem cells 21, 481-494

- PARSHLEY, M. S., H. S. SIMMS (1950):
Cultivation of adult skin epithelial cells (chicken and human) in vitro.
Am J Anat 86, 163–189
- POTTEN, C. S., R. J. MORRIS (1988):
Epithelial stem cells in vivo.
J Cell Sci Suppl 10, 45-62
- POUMAY, Y., M. LECLERCQ-SMEKENS (1998):
In Vitro models of epidermal differentiation.
Folia Med (Plovdiv) 40 (2), 5-12
- REILLY, J. D., D. F. COTTRELL, R. J. MARTIN AND D. J. CUDDEFORD (1998):
Effect of supplemental dietary biotin on hoof growth and hoof growth rate in ponies: a controlled trial.
Equine Vet J. 26(Suppl.), 51-57
- RENNER, U., M. PAEZ-PEREDA, E. ARZT, G. K. STALLA (2004):
Growth factors and cytokines: function and molecular regulation in pituitary adenomas.
Front Horm Res 32, 96-109
- RHEINWALD, J. G., H. GREEN (1975):
Serial cultivation of strains of human epidermal keratinocytes: the formation of keratinizing colonies from single cells.
Cell 6, 331–344
- RHEINWALD, J. G., H. GREEN (1977):
Epidermal growth factor and the multiplication of cultured human epidermal keratinocytes.
Nature 265, 421–424
- ROMEIS, B. (1989):
Mikroskopische Technik, 17. Aufl.
München, Urban u. Schwarzenberg
- RUCKERT, R., G. LINDNER, S. BULFONE-PAUS, R. PAUS (2000):
High-dose proinflammatory cytokines induce apoptosis of hair bulb keratinocytes in vivo.
Br J Dermatol 143 (5), 1036-1039
- RUHRBERG, C., M. A. HAJIBAGHERI, D. A. PARRY, F. M. WATT, F. M (1997):
Periplakin, a novel component of cornified envelopes and desmosomes that belongs to the plakin family and forms complexes with envoplakin.
J Cell Biol 139 (7), 1835-1849
- RYO, Y., T. YOSHIHIRO (2004):
Development of tissue engineering skin based on acellular dermal matrix.
Wound Repair Regen 12 (1), A11

SCHELFHOUT, V. R., E. D. COENE, B. DELAEY, A. A. WAEYTENS, L. DE RYCKE, M. DELEU, C. R. DE POTTER (2002):

The role of heregulin-alpha as a motility factor and amphiregulin as a growth factor in wound healing.

J Pathol 198 (4), 523-533

SCHILLER, D. L., W. W. FRANKE, B. GEIGER (1982):

A subfamily of relatively large and basic cytokeratin polypeptides as defined by mapping is represented by one or several polypeptides in epithelial cells.

EMBO J 1 (6), 761-769

SCHUMACHER, K., R. STREHL, U. DE VRIES, W. W. MINUTH (2002):

Advanced technique for long term culture of epithelia in a continuous luminal-basal medium gradient.

Biomaterials 23 (3) 805-815

SCOTT, I. R., S. RICHARDS, C. HARDING, J. E. LIDDELL, C. G. CURTIS (1988):

Does catabolism of stratum corneum proteins yield functionally active molecules?

Ann N Y Acad Sci 548, 125-136

SHIRAKATA, Y., T. KOMURASAKI, H. TOYODA, Y. HANAKAWA, K. YAMASAKI, S. TOKUMARU, K. SAYAMA, K. HASHIMOTO (2000):

Epiregulin, a novel member of the epidermal growth factor family, is an autocrine growth factor in normal human keratinocytes.

J Biol Chem 275 (8), 5748-5753

SMOLA, H., M. REINKE, P. SHEPHARD, T. KRIEG, S. HESS (1999):

Autologous patient serum for the culture of keratinocyte transplants reduces risk of transmittable disease.

Lancet 353 (9153), 641-642

STEINERT, P. M., W. W. IDLER (1975):

The polypeptide composition of bovine epidermal alpha-keratin.

Biochem J 151(3), 603-614.

STEINERT, P. M., L. N. MAREKOV (1997):

Direct evidence that involucrin is a major early isopeptide cross-linked component of the keratinocyte cornified cell envelope.

J Biol Chem 272 (3), 2021-2030

STENN, K. S., L. A. FERNANDEZ, S. J. TIREL (1988):

The angiogenic properties of the rat vibrissa hair follicle associate with the bulb.

J Invest Dermatol 90 (3), 409-411

STEVEN, A. C., P. M. STEINERT (1994):

Protein composition of cornified cell envelopes of epidermal keratinocytes.

J Cell Sci 107 (Pt 2), 693-700

STREHL, R., K. SCHUMACHER, U. DE VRIES, W. W. MINUTH (2002):

Proliferating cells versus differentiated cells in tissue engineering.

Tissue Eng 8 (1), 37-42

- STREHL, R., K. SCHUMACHER, W. W. MINUTH (2004):
Controlled respiratory gas delivery to embryonic renal epithelial explants in perfusion culture.
Tissue Eng 10 (7-8), 1196-1203
- SULTAN, K. R., H. P. HAAGSMAN (2001):
Species-specific primary cell culture: a research tool in veterinary science.
Veterinary Sciences Tomorrow 1
www.vetscite.org/issue1/reviews/sultan_0800.pdf
- SUN, T. T., R. EICHNER (1983):
Keratin classes: molecular markers for different types of epithelial differentiation.
J Invest Dermatol 81 (1 Suppl), 109-115
- TAKEDA, H. (2004):
cis-Dimer formation of E-cadherin is independent of cell-cell adhesion assembly in vivo.
Biochem Biophys Res Commun 316 (3), 822-826
- TANG P., M.-C. HUNG, J. KLOSTERGAARD (1996):
Human pro-tumor necrosis factor is a homotrimer.
Biochemistry 35, 8216-8222
- TARLTON, J. F., D. E. HOLAH, K. M. EVANS, S. JONES, G. R. PEARSON, A. J. WEBSTER (2002):
Biomechanical and histopathological changes in the support structures of bovine hooves around the time of first calving.
Vet J. 163 (2), 196-204
- TOMIC-CANIC M., M. KOMINE, I.M. FREEDBERG, M. BLUMENBERG (1998):
Epidermal signal transduction and transcription factor activation in activated keratinocytes.
J Dermatol Sci 17 (3), 167-181
- UCHI, H., H. TERAO, T. KOGA, M. FURUE (2000):
Cytokines and chemokines in the epidermis.
J Dermatol Sci 24 Suppl 1, 29-38
- VERMUNT, J. J., P. R. GREENOUGH (1995):
Structural characteristics of the bovine claw: horn growth and wear, horn hardness and claw conformation.
Br Vet J 151 (2), 157-180
- WASCHKE, J., D. DRENCKHAHN, R. H. ADAMSON, F. E. CURRY (2004):
The role of adhesion and contraction in Rac 1-regulated endothelial barrier function in vivo and in vitro.
Am J Physiol Heart Circ Physiol 287, 704-711
- WERNER, S., H. SMOLA (2001):
Paracrine regulation of keratinocyte proliferation and differentiation.
Trends Cell Biol 11 (4), 143-146

WERTZ, P. W., D. C. SWARTZENDRUBER, D. J. KITKO, K. C. MADISON, D. T. DOWNING (1989):

The role of the corneocyte lipid envelopes in cohesion of the stratum corneum.
J Invest Dermatol 93 (1), 169-172

WEYRAUCH, K. D., A. SMOLLICH (1998):

Histologie-Kurs für Veterinärmediziner.
Enke Verlag

WILKENS, H. (1963):

Zur makroskopischen und mikroskopischen Morphologie der Rinderklaue mit einem Vergleich der Architektur von Klauen- und Hufröhrchen.
Hannover: tierärztl. Hochsch., Habil.-Schr.

WOHLRAB, F., S. SCHMIDT, P. CAFFIER (1973):

The distribution of the LDH-isoenzymes in isolated glomerula of kidneys of normal and streptozotocin-diabetic rats (author's transl).
Zentralbl Allg Pathol 117 (5), 512-523

WUNN, D., K. J. WARDROP, K. MEYERS, J. KRAMER, C. RAGLE (1999):

Culture and characterization of equine terminal arch endothelial cells and hoof keratinocytes.
Am J Vet Res 60 (1), 128-132

YANG Y, X. FU, J. LI (2002):

Effect of keratinocyte growth factor-2 on proliferation of human adult keratinocytes.
Chin J Traumatol. Dec;5(6), 342-345

ZELLMER, S., F. GAUNITZ, J. SALVETTER, A. SUROVOY, D. REISSIG, R. GEBHARDT (2001):

Long-term expression of foreign genes in normal human epidermal keratinocytes after transfection with lipid/DNA complexes.
Histochem Cell Biol 115 (1), 41-47

ZIETZSCHMANN, O. (1918):

Das Zehenendorgan der rezenten Säugetiere: Kralle, Nagel, Huf.
Schweiz Arch Tierheilkd 60 (6), 241-272

ZHANG, D., S. KARUNARATNE (2002):

Characterization of mouse profilaggrin: evidence for nuclear engulfment and translocation of the profilaggrin B-domain during epidermal differentiation.
J Invest Dermatol 119 (4), 905-912

ZHOU, J., L. ZHOU, X. JIN (2004):

Expression of the integrin alpha 6 beta 4 and effects of integrin alpha 6 beta 4 on adhesive and invasive behaviors of human laryngeal carcinoma cell lines.
Lin Chuang Er Bi Yan Hou Ke Za Zhi 18 (3), 164-1