

7 Literaturverzeichnis

- Abelson, J. L. und Curtis, G. C. (1996): Hypothalamic-pituitary-adrenal axis activity in panic disorder. 24-hour secretion of corticotropin and cortisol, *Arch Gen Psychiatry* 53 [4], Seite 323-31.
- Andersch, S.; Rosenberg, N. K.; Kullingsjo, H.; Ottosson, J. O.; Bech, P.; Bruun-Hansen, J.; Hanson, L.; Lorentzen, K.; Møllergaard, M.; Rasmussen, S. und et al. (1991): Efficacy and safety of alprazolam, imipramine and placebo in treating panic disorder. A Scandinavian multicenter study, *Acta Psychiatr Scand Suppl* 365, Seite 18-27.
- APA (1994): *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition*, Washington D.C.
- Arnou, B.; Taylor, C.; Agras, W. und Telch, M. (1985): Enhancing agoraphobia treatment outcome by changing couple communication patterns, *Behav Ther* [16], Seite 452-267.
- Aronson, T.A. und Logue, C.M. (1988): Phenomenology of panic attacks: a descriptive study of panic disorder patients' self-reports, *J Clin Psychol* [49], Seite 8-13.
- Babjak, M.; Blumenthal, J. A.; Herman, S.; Khatri, P.; Doraiswamy, M.; Moore, K.; Craighead, W. E.; Baldewicz, T. T. und Krishnan, K. R. (2000): Exercise treatment for major depression: maintenance of therapeutic benefit at 10 months, *Psychosom Med* 62 [5], Seite 633-8.
- Bailey, S. P.; Davis, J. M. und Ahlborn, E. N. (1992): Effect of increased brain serotonergic activity on endurance performance in the rat, *Acta Physiol Scand* 145 [1], Seite 75-6.
- Bakker, A.; van Balkom, A.J. und Spinhoven, P. (2002): SSRIs vs. TCAs in the treatment of panic disorder: a meta-analysis, *Acta Psychiatr Scand.* [106(3)], Seite 163-7.
- Bandelow, B. (2001): *Panik und Agoraphobie - Diagnose, Ursachen, Behandlung*, Springer Verlag, Wien.
- Bandelow, B. (2003): *Angst- und Panikerkrankungen. Ätiologie - Diagnostik - Therapie*, UNI-MED Verlag, Bremen.
- Bandelow, B.; Alvarez Tichauer, G.; Spath, C.; Broocks, A.; Hajak, G.; Bleich, S. und Ruther, E. (2001): Separation anxiety and actual separation experiences during childhood in patients with panic disorder, *Can J Psychiatry* 46 [10], Seite 948-52.
- Bandelow, B.; Spath, C.; Tichauer, G. A.; Broocks, A.; Hajak, G. und Ruther, E. (2002a): Early traumatic life events, parental attitudes, family history, and birth risk factors in patients with panic disorder, *Compr Psychiatry* 43 [4], Seite 269-78.
- Bandelow, B.; Zohar, J.; Hollander, E.; Kasper, S. und Møller, H. J. (2002b): World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for the pharmacological treatment of anxiety, obsessive-compulsive and posttraumatic stress disorders, *World J Biol Psychiatry* 3 [4], Seite 171-99.
- Barlow, D.H. und Craske, M.G. (1994): *Mastery of Your Anxiety and Panic*, Graywind Publications, Albany, NY.
- Beck, A (1988): *Cognitive approaches to panic disorder - Theory and therapy*, Lawrence Erlbaum, Hillsdale.
- Bidzseranova, A.; Gueron, J.; Toth, G.; Penke, B.; Varga, J. und Telegdy, G. (1992): Behavioral effects of atrial and brain natriuretic peptides in rats, *Neuroreport* 3 [3], Seite 283-5.

- Blumenthal, J. A.; Babyak, M. A.; Moore, K. A.; Craighead, W. E.; Herman, S.; Khatri, P.; Waugh, R.; Napolitano, M. A.; Forman, L. M.; Appelbaum, M.; Doraiswamy, P. M. und Krishnan, K. R. (1999): Effects of exercise training on older patients with major depression, *Arch Intern Med* 159 [19], Seite 2349-56.
- Boulenger, J. P.; Patel, J. und Marangos, P. J. (1982): Effects of caffeine and theophylline on adenosine and benzodiazepine receptors in human brain, *Neurosci Lett* 30 [2], Seite 161-6.
- Bradwejn, J. und Koszycki, D. (1991a): Comparison of the panicogenic effect of cholecystokinin 30-33 and carbon dioxide in panic disorder, *Prog Neuropsychopharmacol Biol Psychiatry* 15 [2], Seite 237-9.
- Bradwejn, J.; Koszycki, D. und Meterissian, G. (1990): Cholecystokinin-tetrapeptide induces panic attacks in patients with panic disorder, *Can J Psychiatry* 35 [1], Seite 83-5.
- Bradwejn, J.; Koszycki, D. und Shriqui, C. (1991b): Enhanced sensitivity to cholecystokinin tetrapeptide in panic disorder. Clinical and behavioral findings, *Arch Gen Psychiatry* 48 [7], Seite 603-10.
- Brambilla, F.; Bellodi, L.; Perna, G.; Battaglia, M.; Sciuto, G.; Diaferia, G.; Petraglia, F.; Panerai, A. und Sacerdote, P. (1992): Psychoimmunoendocrine aspects of panic disorder, *Neuropsychobiology* 26 [1-2], Seite 12-22.
- Broocks, A. und Bandelow, B. (2003): Panikstörung und Agoraphobie, *Neurologie, Psychiatrie und Sport*, Thieme Verlag, Stuttgart.
- Broocks, A.; Bandelow, B.; George, A.; Jestrabeck, C.; Opitz, M.; Bartmann, U.; Gleiter, C. H.; Meineke, I.; Roed, I. S.; Ruther, E. und Hajak, G. (2000): Increased psychological responses and divergent neuroendocrine responses to m-CPP and ipsapirone in patients with panic disorder, *Int Clin Psychopharmacol* 15 [3], Seite 153-61.
- Broocks, A.; Bandelow, B.; Pekrun, G.; George, A.; Meyer, T.; Bartmann, U.; Hillmer-Vogel, U. und Ruther, E. (1998): Comparison of aerobic exercise, clomipramine, and placebo in the treatment of panic disorder, *Am J Psychiatry* 155 [5], Seite 603-9.
- Broocks, A.; Meyer, T. F.; Bandelow, B.; George, A.; Bartmann, U.; Ruther, E. und Hillmer-Vogel, U. (1997): Exercise avoidance and impaired endurance capacity in patients with panic disorder, *Neuropsychobiology* 36 [4], Seite 182-7.
- Broocks, A.; Meyer, T.; George, A.; Hillmer-Vogel, U.; Meyer, D.; Bandelow, B.; Hajak, G.; Bartmann, U.; Gleiter, C. H. und Ruther, E. (1999): Decreased neuroendocrine responses to meta-chlorophenylpiperazine (m-CPP) but normal responses to ipsapirone in marathon runners, *Neuropsychopharmacology* 20 [2], Seite 150-61.
- Broocks, A.; Meyer, T.; Opitz, M.; Bartmann, U.; Hillmer-Vogel, U.; George, A.; Pekrun, G.; Wedekind, D.; Ruther, E. und Bandelow, B. (2003): 5-HT_{1A} responsivity in patients with panic disorder before and after treatment with aerobic exercise, clomipramine or placebo, *Eur Neuropsychopharmacol* 13 [3], Seite 153-64.
- Broocks, A.; Schweiger, U. und Pirke, K. M. (1991): The influence of semistarvation-induced hyperactivity on hypothalamic serotonin metabolism, *Physiol Behav* 50 [2], Seite 385-8.
- Cameron, O. G. und Hudson, C. J. (1986): Influence of exercise on anxiety level in patients with anxiety disorders, *Psychosomatics* 27 [10], Seite 720-3.
- Carr, D. B.; Bullen, B. A.; Skrinar, G. S.; Arnold, M. A.; Rosenblatt, M.; Beitins, I. Z.; Martin, J. B. und McArthur, J. W. (1981): Physical conditioning facilitates the

- exercise-induced secretion of beta-endorphin and beta-lipotropin in women, *N Engl J Med* 305 [10], Seite 560-3.
- Chaouloff, F.; Laude, D. und Elghozi, J. L. (1989): PHysical exercise: evidence for differential consequences of tryptophan on 5-HT synthesis and metabolism in central serotonergic cell bodies and terminals, *J Neural Transm* 78 [2], Seite 121-30.
- Charney, D. S.; Heninger, G. R. und Breier, A. (1984): Noradrenergic function in panic anxiety. Effects of yohimbine in healthy subjects and patients with agoraphobia and panic disorder, *Arch Gen Psychiatry* 41 [8], Seite 751-63.
- Charney, D. S.; Woods, S. W.; Krystal, J. H.; Nagy, L. M. und Heninger, G. R. (1992): Noradrenergic neuronal dysregulation in panic disorder: the effects of intravenous yohimbine and clonidine in panic disorder patients, *Acta Psychiatr Scand* 86 [4], Seite 273-82.
- Cicardo, V. H.; Carbone, S. E.; de Rondina, D. C. und Mastronardi, I. O. (1986): Stress by forced swimming in the rat: effects of mianserin and moclobemide on GABAergic-monoaminergic systems in the brain, *Comp Biochem Physiol C* 83 [1], Seite 133-5.
- CNCPS (1992): Drug treatment of panic disorder. Comparative efficacy of alprazolam, imipramine, and placebo. Cross-National Collaborative Panic Study, Second Phase Investigators, *Br J Psychiatry* 160, Seite 191-202; discussion 202-5.
- Cohen, A.S.; Barlow, D.H. und Blanchard, E.B. (1985): Psychophysiology of relaxation-associated panic attacks, *J Abnorm Psychol* [94], Seite 96-101.
- Coplan, J. D.; Andrews, M. W.; Rosenblum, L. A.; Owens, M. J.; Friedman, S.; Gorman, J. M. und Nemeroff, C. B. (1996): Persistent elevations of cerebrospinal fluid concentrations of corticotropin-releasing factor in adult nonhuman primates exposed to early-life stressors: implications for the pathophysiology of mood and anxiety disorders, *Proc Natl Acad Sci U S A* 93 [4], Seite 1619-23.
- Coplan, J. D.; Liebowitz, M. R.; Gorman, J. M.; Fyer, A. J.; Dillon, D. J.; Campeas, R. B.; Davies, S. O.; Martinez, J. und Klein, D. F. (1992): Noradrenergic function in panic disorder. Effects of intravenous clonidine pretreatment on lactate induced panic, *Biol Psychiatry* 31 [2], Seite 135-46.
- Cox, B. J.; Endler, N. S.; Lee, P. S. und Swinson, R. P. (1992): A meta-analysis of treatments for panic disorder with agoraphobia: imipramine, alprazolam, and in vivo exposure, *J Behav Ther Exp Psychiatry* 23 [3], Seite 175-82.
- Cramer, S. R.; Nieman, D. C. und Lee, J. W. (1991): The effects of moderate exercise training on psychological well-being and mood state in women, *J Psychosom Res* 35 [4-5], Seite 437-49.
- Davis, M. (1992): The role of the amygdala in fear and anxiety, *Annu Rev Neurosci* 15, Seite 353-75.
- de Bold, A. J. (1985): Atrial natriuretic factor: a hormone produced by the heart, *Science* 230 [4727], Seite 767-70.
- de Bold, A. J.; Borenstein, H. B.; Veress, A. T. und Sonnenberg, H. (1981): A rapid and potent natriuretic response to intravenous injection of atrial myocardial extract in rats, *Life Sci* 28 [1], Seite 89-94.
- de Montigny, C. (1989): Cholecystokinin tetrapeptide induces panic-like attacks in healthy volunteers. Preliminary findings, *Arch Gen Psychiatry* 46 [6], Seite 511-7.

- Deacon, B.J. und Abramowitz, J.S. (2004): Cognitive and behavioral treatments for anxiety disorders: a review of meta-analytic findings, *J Clin Psychol* [60(4)], Seite 429-41.
- Dengler, W. und Selbmann, H. (2000): Leitlinien zur Diagnostik und Therapie von Angsterkrankungen. Ergebnis einer Konsensuskonferenz, Steinkopff, Darmstadt.
- Dey, S.; Singh, R. H. und Dey, P. K. (1992): Exercise training: significance of regional alterations in serotonin metabolism of rat brain in relation to antidepressant effect of exercise, *Physiol Behav* 52 [6], Seite 1095-9.
- Dillon, D. J.; Gorman, J. M.; Liebowitz, M. R.; Fyer, A. J. und Klein, D. F. (1987): Measurement of lactate-induced panic and anxiety, *Psychiatry Res* 20 [2], Seite 97-105.
- Dimeo, F.; Bauer, M.; Varahram, I.; Proest, G. und Halter, U. (2001): Benefits from aerobic exercise in patients with major depression: a pilot study, *Br J Sports Med* 35 [2], Seite 114-7.
- Doyme, E. J.; Ossip-Klein, D. J.; Bowman, E. D.; Osborn, K. M.; McDougall-Wilson, I. B. und Neimeyer, R. A. (1987): Running versus weight lifting in the treatment of depression, *J Consult Clin Psychol* 55 [5], Seite 748-54.
- Dratcu, L. (2000): Panic, hyperventilation and perpetuation of anxiety, *Prog Neuropsychopharmacol Biol Psychiatry* 24 [7], Seite 1069-89.
- Dratcu, L. (2001): Physical exercise: an adjunctive treatment for panic disorder? *Eur Psychiatry* 16 [6], Seite 372-4.
- Dunn, A. J. und Berridge, C. W. (1990): Physiological and behavioral responses to corticotropin-releasing factor administration: is CRF a mediator of anxiety or stress responses? *Brain Res Brain Res Rev* 15 [2], Seite 71-100.
- Dunn, A. L.; Trivedi, M. H.; Kampert, J. B.; Clark, C. G. und Chambless, H. O. (2005): Exercise treatment for depression: efficacy and dose response, *Am J Prev Med* 28 [1], Seite 1-8.
- Esquivel, G.; Schruers, K.; Kuipers, H. und Griez, E. (2002): The effects of acute exercise and high lactate levels on 35% CO₂ challenge in healthy volunteers, *Acta Psychiatr Scand* 106 [5], Seite 394-7.
- Farmer, M. E.; Locke, B. Z.; Moscicki, E. K.; Dannenberg, A. L.; Larson, D. B. und Radloff, L. S. (1988): Physical activity and depressive symptoms: the NHANES I Epidemiologic Follow-up Study, *Am J Epidemiol* 128 [6], Seite 1340-51.
- Farrell, P. A.; Gates, W. K.; Maksud, M. G. und Morgan, W. P. (1982): Increases in plasma beta-endorphin/beta-lipotropin immunoreactivity after treadmill running in humans, *J Appl Physiol* 52 [5], Seite 1245-9.
- Geese, R. und Reimers, C.D. (2003): Grundlagen des Trainings, Reimers, C.D. und Broocks, A., *Neurologie, Psychiatrie und Sport* Seite 8-45, Thieme Verlag, Stuttgart.
- Goddard, A. W.; Brouette, T.; Almai, A.; Jetty, P.; Woods, S. W. und Charney, D. (2001): Early coadministration of clonazepam with sertraline for panic disorder, *Arch Gen Psychiatry* 58 [7], Seite 681-6.
- Goetz, R. R.; Klein, D. F. und Gorman, J. M. (1996): Symptoms essential to the experience of sodium lactate-induced panic, *Neuropsychopharmacology* 14 [5], Seite 355-66.
- Goldfarb, A. H. und Jamurtas, A. Z. (1997): Beta-endorphin response to exercise. An update, *Sports Med* 24 [1], Seite 8-16.

- Goldstein, A.J.; De Beurs, E.; Chambless, D.L. und Wilson, K.A. (2000): EMDR for panic disorder with agoraphobia: comparison with waiting list and credible attention-placebo control conditions, *J Consult Clin Psychol* [68], Seite 947-56.
- Gorman, J. M.; Fyer, M. R.; Goetz, R.; Askanazi, J.; Liebowitz, M. R.; Fyer, A. J.; Kinney, J. und Klein, D. F. (1988): Ventilatory physiology of patients with panic disorder, *Arch Gen Psychiatry* 45 [1], Seite 31-9.
- Gorman, J. M.; Kent, J. M.; Sullivan, G. M. und Coplan, J. D. (2000): Neuroanatomical hypothesis of panic disorder, revised, *Am J Psychiatry* 157 [4], Seite 493-505.
- Gorman, J. M.; Liebowitz, M. R.; Fyer, A. J. und Stein, J. (1989): A neuroanatomical hypothesis for panic disorder, *Am J Psychiatry* 146 [2], Seite 148-61.
- Grawe, K.; Donati, R. und Bernauer, F. (1994): Psychotherapie im Wandel. Von der Konfession zur Profession, Hogrefe, Göttingen.
- Guszkowska, M. (2004): [Effects of exercise on anxiety, depression and mood], *Psychiatr Pol* 38 [4], Seite 611-20.
- Gutkowska, J.; Antunes-Rodrigues, J. und McCann, S. M. (1997): Atrial natriuretic peptide in brain and pituitary gland, *Physiol Rev* 77 [2], Seite 465-515.
- Herrmann-Lingen, C.; Binder, L.; Klinge, M.; Sander, J.; Schenker, W.; Beyermann, B.; von Lewinski, D. und Pieske, B. (2003): High plasma levels of N-terminal pro-atrial natriuretic peptide associated with low anxiety in severe heart failure, *Psychosom Med* 65 [4], Seite 517-22.
- Hoehn-Saric, R.; McLeod, D. R. und Zimmerli, W. D. (1991): Psychophysiological response patterns in panic disorder, *Acta Psychiatr Scand* 83 [1], Seite 4-11.
- Hofmann, A. (2005): EMDR in der Therapie psychotraumatischer Belastungssyndrome, Thieme, Stuttgart.
- Horwath, E.; Adams, P.; Wickramaratne, P.; Pine, D. und Weissman, M. M. (1997): Panic disorder with smothering symptoms: evidence for increased risk in first-degree relatives, *Depress Anxiety* 6 [4], Seite 147-53.
- Janal, M. N.; Colt, E. W.; Clark, W. C. und Glusman, M. (1984): Pain sensitivity, mood and plasma endocrine levels in man following long-distance running: effects of naloxone, *Pain* 19 [1], Seite 13-25.
- Kahn, R. S.; Wetzler, S.; van Praag, H. M.; Asnis, G. M. und Strauman, T. (1988): Behavioral indications for serotonin receptor hypersensitivity in panic disorder, *Psychiatry Res* 25 [1], Seite 101-4.
- Kaiya, H.; Umekage, T.; Harada, S.; Okazaki, Y. und Sasaki, T. (2005): Factors associated with the development of panic attack and panic disorder: survey in the Japanese population, *Psychiatry Clin Neurosci* 59 [2], Seite 177-82.
- Kasper, S. und Resinger, E. (2001): Panic disorder: the place of benzodiazepines and selective serotonin reuptake inhibitors, *Eur Neuropsychopharmacol* 11 [4], Seite 307-21.
- Katzman, M. (2004): Venlafaxine in the treatment of anxiety disorders, *Expert Rev Neurother* [4], Seite 371-81.
- Keck, P. E., Jr.; McElroy, S. L.; Tugrul, K. C.; Bennett, J. A. und Smith, J. M. (1993b): Antiepileptic drugs for the treatment of panic disorder, *Neuropsychobiology* 27 [3], Seite 150-3.
- Keck, P. E., Jr.; Taylor, V. E.; Tugrul, K. C.; McElroy, S. L. und Bennett, J. A. (1993a): Valproate treatment of panic disorder and lactate-induced panic attacks, *Biol Psychiatry* 33 [7], Seite 542-6.
- Kellner, M.; Herzog, L.; Yassouridis, A.; Holsboer, F. und Wiedemann, K. (1995): Possible role of atrial natriuretic hormone in pituitary-adrenocortical

- unresponsiveness in lactate-induced panic, *Am J Psychiatry* 152 [9], Seite 1365-7.
- Kellner, M.; Wiedemann, K. und Holsboer, F. (1992): Atrial natriuretic factor inhibits the CRH-stimulated secretion of ACTH and cortisol in man, *Life Sci* 50 [24], Seite 1835-42.
- Kessler, R.C.; McGonagle, K.A.; S., Zhao.; Nelson, C.B.; Hughes, M.; Eshleman, S.; Wittchen, H.U. und Kendler, K.S. (1994): Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey, *Arch Gen Psychiatry* [51(1)], Seite 8-19.
- King, A. C.; Taylor, C. B.; Haskell, W. L. und DeBusk, R. F. (1989): Influence of regular aerobic exercise on psychological health: a randomized, controlled trial of healthy middle-aged adults, *Health Psychol* 8 [3], Seite 305-24.
- Klein, D. F. (1993): False suffocation alarms, spontaneous panics, and related conditions. An integrative hypothesis, *Arch Gen Psychiatry* 50 [4], Seite 306-17.
- Kraemer, W. J.; Hakkinen, K.; Newton, R. U.; Nindl, B. C.; Volek, J. S.; McCormick, M.; Gotshalk, L. A.; Gordon, S. E.; Fleck, S. J.; Campbell, W. W.; Putukian, M. und Evans, W. J. (1999): Effects of heavy-resistance training on hormonal response patterns in younger vs. older men, *J Appl Physiol* 87 [3], Seite 982-92.
- Krüger, M.B. und Dahl, A.A. (1999): The efficacy and safety of moclobemide compared to clomipramine in the treatment of panic disorder, *European Archives Of Psychiatry And Clinical Neuroscience* [249], Seite S19-24.
- Kujala, U. M.; Kaprio, J.; Sarna, S. und Koskenvuo, M. (1998): Relationship of leisure-time physical activity and mortality: the Finnish twin cohort, *Jama* 279 [6], Seite 440-4.
- Lawlor, D. A. und Hopker, S. W. (2001): The effectiveness of exercise as an intervention in the management of depression: systematic review and meta-regression analysis of randomised controlled trials, *Bmj* 322 [7289], Seite 763-7.
- Lecrubier, Y.; Bakker, A.; Dunbar, G. und Judge, R. (1997): A comparison of paroxetine, clomipramine and placebo in panic disorder, *Acta Psychiatr Scand* [95], Seite 145-52.
- Lennox, S. S.; Bedell, J. R. und Stone, A. A. (1990): The effect of exercise on normal mood, *J Psychosom Res* 34 [6], Seite 629-36.
- Leon, A. S.; Connett, J.; Jacobs, D. R., Jr. und Rauramaa, R. (1987): Leisure-time physical activity levels and risk of coronary heart disease and death. The Multiple Risk Factor Intervention Trial, *Jama* 258 [17], Seite 2388-95.
- Loerch, B.; Graf-Morgenstern, M.; Hautzinger, M.; Schlegel, S.; Hain, C.; Sandmann, J. und Benkert, O. (1999): Randomised placebo-controlled trial of moclobemide, cognitive-behavioural therapy and their combination in panic disorder with agoraphobia, *Br J Psychiatry* 174, Seite 205-12.
- Löffler, G. (1998): Elektronentransport und oxidative Phosphorylierung, *Biochemie und Pathobiochemie* Seite 495-518, Springer-Verlag, Berlin Heidelberg New York.
- Long, B.C. und van Stavel, R.V. (1995): Effects of exercise training on anxiety: a meta analysis, *J Appl Sport Psychol* [7], Seite 167-89.
- Lydiard, R.B. (1995): Drug treatment of panic disorder. Hypnotics and Anxiolytics, *Baillieres Clin Psychiatry* [1], Seite 427-46.

- Magee, W.J.; Eaton, W.W.; Wittchen, H.U.; McGonagle, K.A. und Kessler, R.C. (1996): Agoraphobia, simple phobia, and social phobia in the National Comorbidity Survey, *Arch Gen Psychiatry* [53], Seite 159-68.
- Mandroukas, K.; Zakas, A.; Aggelopoulou, N.; Christoulas, K.; Abatzides, G. und Karamouzis, M. (1995): Atrial natriuretic factor responses to submaximal and maximal exercise, *Br J Sports Med* 29 [4], Seite 248-51.
- Margraf und Ruhmland (2001).
- Marks, I.M. (1987): *Fears, phobias and rituals*, Oxford University Press, New York, Oxford.
- McDaniel, W. W. (1988): Panic disorder and exercise, *Am J Psychiatry* 145 [2], Seite 269.
- Meeusen, R. und De Meirleir, K. (1995): Exercise and brain neurotransmission, *Sports Med* 20 [3], Seite 160-88.
- Moses, J.; Steptoe, A.; Mathews, A. und Edwards, S. (1989): The effects of exercise training on mental well-being in the normal population: a controlled trial, *J Psychosom Res* 33 [1], Seite 47-61.
- Muller, B. und Armstrong, H.E. (1975): A further note on the running treatment for anxiety, *Psychother Theory Res Pract* [12], Seite 385-87.
- Munjack, D. J.; Crocker, B.; Cabe, D.; Brown, R.; Usigli, R.; Zulueta, A.; McManus, M.; McDowell, D.; Palmer, R. und Leonard, M. (1989): Alprazolam, propranolol, and placebo in the treatment of panic disorder and agoraphobia with panic attacks, *J Clin Psychopharmacol* 9 [1], Seite 22-7.
- NIH (1991): *Treatment of Panic Disorder*, National Institutes of Health Consensus Development Conference Statement September 25-27 [1-26].
- North, T. C.; McCullagh, P. und Tran, Z. V. (1990): Effect of exercise on depression, *Exerc Sport Sci Rev* 18, Seite 379-415.
- Orwin, A. (1973): 'The running treatment': a preliminary communication on a new use for an old therapy (physical activity) in the agoraphobic syndrome, *Br J Psychiatry* 122 [567], Seite 175-9.
- Orwin, A. (1974): Treatment of a situational phobia--a case for running, *Br J Psychiatry* 125 [0], Seite 95-8.
- Öst, L.G. und Westling, B.E. (1995): Applied relaxation vs cognitive behavior therapy in the treatment of panic disorder, *Behav Res Ther* [33], Seite 145-58.
- Öst, L.G.; Westling, B.E. und Hellstrom, K. (1993): Applied relaxation, exposure in vivo and cognitive methods in the treatment of panic disorder with agoraphobia, *Behav Res Ther* [31], Seite 383-94.
- Paluska, S. A. und Schwenk, T. L. (2000): Physical activity and mental health: current concepts, *Sports Med* 29 [3], Seite 167-80.
- Perna, G.; Bertani, A.; Caldirola, D. und Bellodi, L. (1996): Family history of panic disorder and hypersensitivity to CO₂ in patients with panic disorder, *Am J Psychiatry* 153 [8], Seite 1060-4.
- Petruzzello, S. J.; Landers, D. M.; Hatfield, B. D.; Kubitz, K. A. und Salazar, W. (1991): A meta-analysis on the anxiety-reducing effects of acute and chronic exercise. Outcomes and mechanisms, *Sports Med* 11 [3], Seite 143-82.
- Pitts, F. N., Jr. und McClure, J. N., Jr. (1967): Lactate metabolism in anxiety neurosis, *N Engl J Med* 277 [25], Seite 1329-36.
- Regier, D.A.; Rae, D.S.; Narrow, W.E.; Kaelber, C.T. und Schatzberg, A.F. (1998): Prevalence of anxiety disorders and their comorbidity with mood and addictive disorders, *Br J Psychiatry* [173], Seite 24-28.

- Reiman, E. M.; Raichle, M. E.; Robins, E.; Butler, F. K.; Herscovitch, P.; Fox, P. und Perlmutter, J. (1986): The application of positron emission tomography to the study of panic disorder, *Am J Psychiatry* 143 [4], Seite 469-77.
- Riemann, F. (1961): Grundformen der Angst: eine tiefenpsychologische Studie, Ernst Reinhardt, GmbH & Co, München.
- Rogers, C.R. (1951): *Client-centred Psychotherapy*, Houghton Mifflin, Boston.
- Rogers, P. J.; Tyce, G. M.; Bailey, K. R. und Bove, A. A. (1991): Exercise-induced increases in atrial natriuretic factor are attenuated by endurance training, *J Am Coll Cardiol* 18 [5], Seite 1236-41.
- Rosenblum, L. A.; Coplan, J. D.; Friedman, S.; Bassoff, T.; Gorman, J. M. und Andrews, M. W. (1994): Adverse early experiences affect noradrenergic and serotonergic functioning in adult primates, *Biol Psychiatry* 35 [4], Seite 221-7.
- Roy-Byrne, P. P.; Geraci, M. und Uhde, T. W. (1986a): Life events and the onset of panic disorder, *Am J Psychiatry* 143 [11], Seite 1424-7.
- Roy-Byrne, P. P.; Uhde, T. W.; Post, R. M.; Gallucci, W.; Chrousos, G. P. und Gold, P. W. (1986b): The corticotropin-releasing hormone stimulation test in patients with panic disorder, *Am J Psychiatry* 143 [7], Seite 896-9.
- Salmon, P. (2001): Effects of physical exercise on anxiety, depression, and sensitivity to stress: a unifying theory, *Clin Psychol Rev* 21 [1], Seite 33-61.
- Schnabel, G. (1998): *Trainingswissenschaft*, SVB Sportverlag, Berlin.
- Schreiber, W.; Lauer, C. J.; Krumrey, K.; Holsboer, F. und Krieg, J. C. (1996): Dysregulation of the hypothalamic-pituitary-adrenocortical system in panic disorder, *Neuropsychopharmacology* 15 [1], Seite 7-15.
- Schruers, K.; Pols, H.; Overbeek, T. und Griez, E. (2000): 10 microg CCK-4 premedication and 35% CO₂ challenge in healthy volunteers, *Prog Neuropsychopharmacol Biol Psychiatry* 24 [3], Seite 409-18.
- Scully, D.; Kremer, J.; Meade, M. M.; Graham, R. und Dudgeon, K. (1998): Physical exercise and psychological well being: a critical review, *Br J Sports Med* 32 [2], Seite 111-20.
- Sexton, H.; Maere, A. und Dahl, N. H. (1989): Exercise intensity and reduction in neurotic symptoms. A controlled follow-up study, *Acta Psychiatr Scand* 80 [3], Seite 231-5.
- Sheehan, D. V.; Raj, A. B.; Harnett-Sheehan, K.; Soto, S. und Knapp, E. (1993): The relative efficacy of high-dose buspirone and alprazolam in the treatment of panic disorder: a double-blind placebo-controlled study, *Acta Psychiatr Scand* 88 [1], Seite 1-11.
- Siegel, R.E. (1968): *Galen's System of Physiology and Medicine*, Karger, New York.
- Skre, I.; Onstad, S.; Torgersen, S.; Lygren, S. und Kringlen, E. (1993): A twin study of DSM-III-R anxiety disorders, *Acta Psychiatr Scand* 88 [2], Seite 85-92.
- Smits, J. A. und Zvolensky, M. J. (2006): Emotional vulnerability as a function of physical activity among individuals with panic disorder, *Depress Anxiety* 23 [2], Seite 102-6.
- Spielberger, C.D.; Gorsuch, R.L.; Lushene, R.; Vagg, P.R. und Jacobs, G.A. (1983): *Manual for the State-Trait Anxiety Inventory*.
- Stephoe, A.; Edwards, S.; Moses, J. und Mathews, A. (1989): The effects of exercise training on mood and perceived coping ability in anxious adults from the general population, *J Psychosom Res* 33 [5], Seite 537-47.
- Strauß, B. und Burgmeier-Lohse, M. (1994): Evaluation einer stationären Langzeitgruppenpsychotherapie. Ein Beitrag zur differentiellen Psychotherapieforschung im stationären Feld, *Psychosom Med Psychol* [44], Seite 184-92.

- Ströhle, A. (2002): Neurobiologie von Angst: Natriuretische Peptide und neuroaktive Steroide, unveröffentlichte Habilitationsschrift, München.
- Ströhle, A. (2003a): [The neuroendocrinology of stress and the pathophysiology and therapy of depression and anxiety], *Nervenarzt* 74 [3], Seite 279-91; quiz 292.
- Ströhle, A. (2003b): Die experimentelle Provokation von Panikattacken als humanexperimentelles Angstmodell, *Nervenarzt* 74, Seite 733-93.
- Ströhle, A.; Holsboer, F. und Rupprecht, R. (2000): Increased ACTH concentrations associated with cholecystinin tetrapeptide-induced panic attacks in patients with panic disorder, *Neuropsychopharmacology* 22 [3], Seite 251-6.
- Ströhle, A.; Jahn, H.; Montkowski, A.; Liebsch, G.; Boll, E.; Landgraf, R.; Holsboer, F. und Wiedemann, K. (1997): Central and peripheral administration of atriopeptin is anxiolytic in rats, *Neuroendocrinology* 65 [3], Seite 210-5.
- Ströhle, A.; Kellner, M.; Holsboer, F. und Wiedemann, K. (1998): Atrial natriuretic hormone decreases endocrine response to a combined dexamethasone-corticotropin-releasing hormone test, *Biol Psychiatry* 43 [5], Seite 371-5.
- Ströhle, A.; Kellner, M.; Holsboer, F. und Wiedemann, K. (2001): Anxiolytic activity of atrial natriuretic peptide in patients with panic disorder, *Am J Psychiatry* 158 [9], Seite 1514-6.
- Ströhle, A.; Romeo, E.; di Michele, F.; Pasini, A.; Hermann, B.; Gajewsky, G.; Holsboer, F. und Rupprecht, R. (2003): Induced panic attacks shift gamma-aminobutyric acid type A receptor modulatory neuroactive steroid composition in patients with panic disorder: preliminary results, *Arch Gen Psychiatry* 60 [2], Seite 161-8.
- Sullivan, G. M.; Coplan, J. D.; Kent, J. M. und Gorman, J. M. (1999): The noradrenergic system in pathological anxiety: a focus on panic with relevance to generalized anxiety and phobias, *Biol Psychiatry* 46 [9], Seite 1205-18.
- Suter, E.; Marti, B.; Tschopp, A. und Wanner, H. U. (1991): [Effects of jogging on mental well-being and seasonal mood variations: a randomized study with healthy women and men], *Schweiz Med Wochenschr* 121 [35], Seite 1254-63.
- Tanaka, H.; Shindo, M.; Gutkowska, J.; Kinoshita, A.; Urata, H.; Ikeda, M. und Arakawa, K. (1986): Effect of acute exercise on plasma immunoreactive-atrial natriuretic factor, *Life Sci* 39 [18], Seite 1685-93.
- Targum, S. D. und Marshall, L. E. (1989): Fenfluramine provocation of anxiety in patients with panic disorder, *Psychiatry Res* 28 [3], Seite 295-306.
- Tate, A. K. und Petruzzello, S. J. (1995): Varying the intensity of acute exercise: implications for changes in affect, *J Sports Med Phys Fitness* 35 [4], Seite 295-302.
- Teusch, L.; Böhme, H. und Gastpar, M. (1997): The benefit of an insight-oriented and experiential approach on panic and agoraphobia symptoms. Results of a controlled comparison of client-centered therapy alone and in combination with behavioral exposure, *Psychother Psychosom* [66(6)], Seite 293-301.
- Tiller, J.W.; Bouwer, C. und Behnke, K. (1999): Moclobemide and fluoxetine for panic disorder. International Panic Disorder Study Group., *European Archives Of Psychiatry And Clinical Neuroscience* [249], Seite S7-10.
- Tweed, J. L.; Schoenbach, V. J.; George, L. K. und Blazer, D. G. (1989): The effects of childhood parental death and divorce on six-month history of anxiety disorders, *Br J Psychiatry* 154, Seite 823-8.
- Uhde, T. W.; Stein, M. B. und Post, R. M. (1988): Lack of efficacy of carbamazepine in the treatment of panic disorder, *Am J Psychiatry* 145 [9], Seite 1104-9.
- van Balkom, A.J.; Bakker, A.; Spinhoven, P.; Blauw, B.M.; Smeenk, S. und Ruesink, B. (1997): A meta-analysis of the treatment of panic disorder with or without

- agoraphobia: a comparison of psychopharmacological, cognitive-behavioral, and combination treatments, *J Nerv Ment Dis* [185(8)], Seite 510-6.
- Versiani, M.; Cassano, G.; Perugi, G.; Benedetti, A.; Mastalli, L.; Nardi, A. und Savino, M. (2002): Reboxetine, a selective norepinephrine reuptake inhibitor, is an effective and well-tolerated treatment for panic disorder, *J Clin Psychiatry* 63 [1], Seite 31-7.
- Wade, A.G.; Lepola, U.; Koponen, H.J.; Pedersen, V. und Pedersen, T. (1997): The effect of citalopram in panic disorder, *Br J Psychiatry* [170], Seite 549-553.
- Wedekind, D.; Bandelow, B.; Broocks, A.; Hajak, G. und Ruther, E. (2000): Salivary, total plasma and plasma free cortisol in panic disorder, *J Neural Transm* 107 [7], Seite 831-7.
- Weissman, M. M.; Fyer, A. J.; Haghghi, F.; Heiman, G.; Deng, Z.; Hen, R.; Hodge, S. E. und Knowles, J. A. (2000): Potential panic disorder syndrome: clinical and genetic linkage evidence, *Am J Med Genet* 96 [1], Seite 24-35.
- WHO (1994a): Weltgesundheitsorganisation - Internationale Klassifikation psychischer Störungen. ICD-10 Kapitel V(F). Forschungskriterien, Verlag Hans Huber, Bern/Göttingen.
- WHO (1994b): Erklärung von Weltgesundheitsorganisation und Weltsporthärzteverband in Köln, *Dtsch Ztschr Sportmed* [45], Seite 170.
- Wiborg, I.M. und Dahl, A.A. (1996): Does brief dynamic psychotherapy reduce the relapse rate of panic disorder? *Arch Gen Psychiatry* [53(8)], Seite 689-94.
- Youngstedt, S. D.; O'Connor, P. J.; Crabbe, J. B. und Dishman, R. K. (1998): Acute exercise reduces caffeine-induced anxiogenesis, *Med Sci Sports Exerc* 30 [5], Seite 740-5.
- Zwanzger, P.; Eser, D.; Padberg, F.; Baghai, T. C.; Schule, C.; Rupprecht, R.; di Michele, F.; Romeo, E.; Pasini, A. und Ströhle, A. (2004): Neuroactive steroids are not affected by panic induction with 50 microg cholecystokinin-tetrapeptide (CCK-4) in healthy volunteers, *J Psychiatr Res* 38 [2], Seite 215-7.