

6 Literaturverzeichnis

Abelló P. & Frankland R.J. 1997. Population characteristics of the neustonic isopod *Idotea metallica* (Crustacea, Isopoda, Idoteidae) in the western Mediterranean (June 1993). *Sci. Mar.* 61: 409-414.

Abrams P.A. 2000. Character shifts of prey species that share predators. *Am. Nat.* 156 (suppl.): 45-61.

Adema J.P.H.M. & Huwae P.H.M. 1982. New and supplementary records of marine Isopoda for the Netherlands and the southern North Sea since 1956, with a note on *Peltogaster paguri* (Crustacea, Cirripedia). *Zoologische Bijdragen* 28: 33-57.

Arnaud F., Arnaud P.M., Intès A. & Le Loeuff P. 1976. Transport d'invertébrés benthiques entre l'Afrique du Sud et Sainte Hélène par les laminaires (Phaeophyceae). *Bull. Mus. Natl. Hist. Nat. Paris Sér. 3* 384: 49-55.

Baan S.M. van der & Holthuis L.B. 1969. On the occurrence of Isopoda in the surface plankton in the North Sea near the lightship "Texel". *Neth. J. Sea Res.* 4: 354-363.

Backus R.H., Flierl G.R., Kester D.R., Olson D.B., Richardson P.L., Vastano A.C., Wiebe P.H. & Wormuth J.H. 1981. Gulf stream cold-core rings: their physics, chemistry, and biology. *Science* 212: 1091-1100.

Barnes D.K.A. 2002. Invasions by marine life on plastic debris. *Nature* 416: 808-809.

Barstow S.F. 1983. The ecology of Langmuir circulation: A review. *Mar. Environ. Res.* 9 : 211-236.

Biesiot P.M., Wang S.Y., Perry H.M. & Trigg, C., 1999. Organic reserves in the midgut gland and fat body of the giant deep-sea isopod *Bathynomus giganteus*. *J. Crust. Biol.* 19: 450-458.

Borowsky B., 1987. Laboratory studies of the pattern of reproduction of the isopod crustacean *Idotea baltica*. Fish. Bull. 85: 377-380.

Butler J.N. 1975. Pelagic tar. Scient. Amer. 232: 90-97.

Cărăușu A. 1954. Contributii la studiul isopodelor (Crustacea Malacostraca) marii negre (litoralul rominesc si regiunile invecinate) III. Familia Idoteidae. Anal. Stiint. Univ. 'Al. I. Cuza' Din Iasi, Sect. II. T. I, fasc. 1-2: 137-216.

Carlson L. 1991. Seasonal variation in growth, reproduction and nitrogen content of *Fucus vesiculosus* L in the Oeresund, southern Sweden. Bot. Mar. 34: 447-453.

Carlton J. T. & Geller J.B. 1993. Ecological roulette: the global transport of non-indigenous marine organisms. Science 261: 78-82.

Caswell H. 1982. Stable population structure and reproductive value for populations with complex life cycles. Ecology 63: 1223-1231.

Caswell H. 1989. Analysis of life table response experiments. I. Decomposition of effects on population growth rate. Ecol. Model. 46: 221-237.

Caswell H. 2001. Matrix population models, 2nd edition. Sinauer Associates, Inc., Sunderland.

Caughley G. 1994. Directions in conservation biology. J. Anim. Ecol. 63: 215-244.

Collin R. 2001. The effects of mode of development on phylogeography and population structure of North Atlantic *Crepidula* (Gastropoda: Calyptraeidae). Mol. Ecol. 10: 2249-2262.

Coston-Clements L., Settle L.R., Hoss D.E. & Cross F.A. 1991. Utilization of the *Sargassum* habitat by marine invertebrates and vertebrates – a review. NOAA Technical Memorandum NMFS-SEFSC-296.

Dellinger T., Davenport J. & Wortz P. 1997. Comparisons of social structure of Columbus crabs living on Loggerhaed sea turtles and inanimate flotsam. J. mar. biol. Ass. U.K. 77: 185-194.

Deudero S. & Moralis-Nin, B. 2000. Occurrence of *Polyprion americanus* under floating objects in western Mediterranean oceanic waters, inference from stomach contents analysis. J. mar. biol. Ass. U.K. 80: 751-752.

Dietrich G., Kalle K., Krauss W. & Siedler G. 1980. General Oceanography, an Introduction. Wiley & Sons, New York.

Dixon & Dixon 1983. Marine litter distribution and composition in the North Sea. Mar. Poll. Bull. 14: 145-148.

Dooley J.K. 1972. Fishes associated with the pelagic *Sargassum* complex, with a discussion of the *Sargassum* community. Contr. mar. Sci. 16: 1-32.

Dow T.G. & Menzies R.J. 1958. The pelagic isopod *Idotea metallica* in the Mediterranean. Pubbl. Staz. Zool. Napoli 30: 330-336.

Ebbesmeyer C.C. & Ingraham J.W. Jr. 1992. Shoe Spill in the North Pacific. EOS, Transactions, American Geophysical Union 73: 361-368.

Edgar G.J. 1987. Dispersal of faunal and floral propagules associated with drifting *Macrocystis pyrifera* plants. Mar. Biol. 95: 599-610.

Ehrlich P.R. & Murphy D.D. 1981. The population biology of butterflies (Euphydryas). Biologisches Zentralblatt 100: 613-629.

Engkvist R., Malm T. & Tobiasson S. 2000. Density dependent grazing effects of the isopod *Idotea baltica* Pallas on *Fucus vesiculosus* L in the Baltic Sea. Aquat. Ecol. 34: 253-360.

Fava G., Zangaglia A. & Cervelli M. 1992. Ecology of *Idotea baltica* (Pallas) populations in the lagoon of Venice. *Oceanologica Acta* 15: 651-660.

Fisher R.A. 1930. The genetical theory of natural selection. Clarendon Press, Oxford.

Franke H.-D. (im Druck). The mechanisms of competition, coexistence and ecological diversification in marine congeners.

Franke H.-D., Gutow L. & Janke M. 1999. The recent arrival of the oceanic isopod *Idotea metallica* Bosc off Helgoland (German Bight, North Sea): an indication of a warming trend in the North Sea? *Helgoländer Meeresunters.* 52: 347-357.

Franke H.-D. & Janke M. 1998. Mechanisms and consequences of intra- and interspecific interference competition in *Idotea baltica* (Pallas) and *Idotea emarginata* (Fabricius) (Crustacea: Isopoda): A laboratory study of possible proximate causes of habitat segregation. *J. exp. mar. Biol. Ecol.* 227: 1-21.

Furness R.W. & Todd C.M. 1984. Diets and feeding of Fulmars *Fulmarus glacialis* during the breeding season: a comparison between St Kilda and Shetland colonies. *Ibis* 126: 379-387.

Gaylord B. & Gaines S.D. 2000. Temperature or transport? Range limits in marine species mediated solely by flow. *Am. Nat.* 155: 769-789.

Gerlach S.A. 1977. Means of meiofaunal dispersal. *Mikrofauna Meeresbod.* 61: 89-103.

Gorelova T.A. & Fedoryako B.I. 1986. Topic and trophic relationships of fishes associated with drifting *Sargassum* vegetation. *J. Ichtyol.* 26: 63-72.

Gregory M.R. 1978. Accumulation and distribution of virgin plastic granules on New Zealand beaches. *N.Z. J. Mar. Freshwat. Res.* 12: 399-414.

Greve W. 1994. The 1989 German Bight invasion of *Muggiaea atlantica*. ICES J. mar. Sci. 51: 355-358.

Guarino S.M., Gambardella C. & de Nicola M. 1993. Biology and population dynamics of *Idotea baltica* (Crustacea, Isopoda) in the Gulf of Naples, the Tyrrhenian sea. Vie Milieu 43: 125-136.

Gutow L. & Franke H.-D. 2001. On the current and possible future status of the neustonic isopod *Idotea metallica* Bosc in the North Sea: a laboratory study. J. Sea Res. 45: 37-44.

Hammrich A. 2003. Vergleichende Untersuchungen zur ökologischen Diversifikation der Helgoländer *Idotea*-Arten. Diplomarbeit, Universität Hamburg.

Hamner W.M. & Schneider D. 1986. Regularly spaced rows of medusae in the Bering Sea: Role of Langmuir circulation. Limnol. Oceanogr. 31: 171-177.

Hanski I. 1997. Metapopulation dynamics: From concepts and observations to predictive models. In: Hanski, I. & Gilpin, M.E. (Herausg.). Metapopulation biology: ecology, genetics, and evolution. Academic Press, London, S. 69-91.

Hanski I. & Simberloff D. 1997. The metapopulation approach, its history, conceptual domain, and application to conservation. In: Hanski, I. & Gilpin, M.E. (Herausg.). Metapopulation biology: ecology, genetics, and evolution. Academic Press, London, S. 5-26.

Harms J. 1999. The neozoan *Elminius modestus* Darwin (Crustacea, Cirripedia): possible explanations for its successful invasion in European water. Helgoländer Meeresunters. 52: 337-346.

Harms J. & Anger K. 1989. Settlement of the barnacle *Elminius modestus* Darwin on test panels at Helgoland (North Sea): A ten year study. Sci. Mar. 53: 417-421.

Hartmann J. 1976. Der Jahresgang von Neuston im Golf von Neapel. Int. Revue ges. Hydrobiol. 61: 825-840.

Helmuth B.S., Veit R.R. & Holberton R. 1994. Long-distance dispersal of a Subantarctic brooding bivalve (*Gaimardia trapesina*) by kelp rafting. Mar. Biol. 120: 421-426.

Hemmi A. & Jormalainen, V. 2002. Nutrient enhancement increases performance of a marine herbivore via quality of its food alga. Ecology 83: 1052-1064.

Hentschel E. 1922. Über den Bewuchs auf treibenden Tangen der Sargassosee. Mitt. Zool. Staatsinst. Zool. Mus. Hamburg 38: 1-26.

Herring P.J. 1969. Pigmentation and carotinoid metabolism of the marine isopod *Idotea metallica*. J. mar. biol. Ass. U.K. 49: 766-779.

Hervant F. & Renault D. 2002. Long-term fasting and realimentation in hypogean and epigean isopods: a proposed adaptive strategy for groundwater organisms. J. exp. Biol. 205: 2079-2087.

Highsmith R.C. 1985. Floating and algal rafting as potential dispersal mechanisms in brooding invertebrates. Mar. Ecol. Prog. Ser. 25: 169-179.

Hobday A.J. 2000. Abundance and dispersal of drifting kelp *Macrocystis pyrifera* rafts in the Southern California Bight. Mar. Ecol. Prog. Ser. 195: 101-116.

Holdway P. & Maddock L. 1983. A comparative survey of neuston: geographical and temporal distribution patterns. Mar. Biol. 76: 263-270.

Horsman P.V. 1982. The amount of garbage pollution from merchant ships. Mar. Poll. Bull. 13: 167-169.

Howard K.L. & Menzies R.J. 1969. Distribution and production of *Sargassum* in the waters off the Carolina coast. Bot. Mar. 12: 244-254.

Huey R.B. & Berrigan D. 2001. Temperature, demography, and ectotherm fitness. *Am. Nat.* 158: 204-210.

Hurrell J.W., Kushnir Y., Ottersen G. & Visbeck M. 2003. An overview of the North Atlantic Oscillation. In: Hurrell J.W., Kushnir Y., Ottersen G. & Visbeck M. (Herausg.). *Geophysical Monograph* 134, S. 1-35.

Huysman M. & Huwae P.H.M. 1978. De eerste vondst van *Idotea metallica* Bosc 1801-1802 (Crustacea Isopoda) aan de Nederlandse kust. *Het Zeepard* 38: 121-122.

Ingólfsson A. 1995. Floating clumps of seaweed around Iceland: Natural microcosms and a means of dispersal for shore fauna. *Mar. Biol.* 122: 13-21.

Ingólfsson A. 1998. Dynamics of macrofaunal communities of floating seaweed clumps off western Iceland: a study of patches on the surface of the sea. *J. exp. mar. Biol. Ecol.* 231: 119-137.

Ingólfsson A. 2000. Colonization of floating seaweed by pelagic and subtidal benthic animals in southwestern Iceland. *Hydrobiologia* 44: 181-189.

Johannesson K. 1988. The paradox of Rockall: Why is a brooding gastropod (*Littorina saxatilis*) more widespread than one having a planktonic larval dispersal stage (*L. littorea*)? *Mar. Biol.* 99: 507-513.

Johnson D.L. & Richardson P.L. 1977. On the wind-induced sinking of *Sargassum*. *J. exp. mar. Biol. Ecol.* 28: 255-267.

Jormalainen V., Honkanen T. & Heikkilä N. 2001. Feeding preferences and performance of a marine isopod on seaweed hosts: cost of habitat specialisation. *Mar. Ecol. Prog. Ser.* 220: 219-230.

Kangas P., Autio H., Hällfors G., Luthre H., Niemi Å & Salemaa H. 1982. A general model of the decline of *Fucus vesiculosus* at Tvärminne, south coast of Finland in 1977-81. Acta Bot. Fennica 118: 1-27.

Kingsford 1995. Drift algae: a contribution to near-shore habitat complexity in the pelagic environment and an attractant for fish. Mar. Ecol. Prog. Ser. 116: 297-301.

Korczynski R.E. 1989. Biochemical composition of the isopod *Mesidotea entomon* (Linnaeus) from the western Arctic. Polar Biol. 9: 391-395.

Kouwenberg J. & Pinkster S. 1984. Population dynamics of three brackish water isopod species (Crustacea) in the lagoon system of Bages-Sigean (France) I. General aspects and distribution. Vie Milieu 34: 229-240.

Kouwenberg J., Tan S., Snoek B. & Pinkster S. 1987. Population dynamics of brackish isopod species (Crustacea) in the lagoon system of Bages-Sigean (France) III. Reproductive rates of *Idotea baltica* and *Idotea granulosa*. Vie Milieu 37: 105-114.

Levin L., Caswell H., Bridges T., DiBacco C., Cabrera D. & Plaia, G. 1996. Demographic responses of estuarine polychaetes to pollutants: life table response experiments. Ecol. Appl. 6: 1295-1313.

Levins R. 1969. Some demographic and genetic consequences of environmental heterogeneity for biological control. Bull. Entomol. Soc. Am. 15: 237-240.

Locke A. & Corey S. 1989. Amphipods, isopods and surface currents: a case for passive dispersal in the Bay of Fundy, Canada. Journal of Plankt. Res. 11: 419-430.

Ludwig D. 1996. The distribution of population survival times. Am. Nat. 147: 506-526.

Ludyanskiy M.L., McDonald D. & MacNeill, D. 1993. Impact of the Zebra Mussel, a Bivalve Invader. BioScience 43: 533-544.

McGrath D. 1980. A record of the marine isopod *Idotea metallica* Bosc. off the west coast of Ireland. Ir. Nat. J. 20: 126.

Meinesz A. & Hesse B. 1991. Introduction et invasion de l'algue tropicale *Caulerpa taxifolia* en Méditerranée nord-occidentale. Oceanologica Acta 14: 415-426.

Miller M.A. 1968. Isopoda and Tanaidacea from buoys in coastal waters of the continental United States, Hawaii, and the Bahamas (Crustacea). Proc. U.S. Nat. Mus. 125: 1-53.

Mitchell C.T. & Hunter J.R. 1970. Fishes associated with drifting kelp, *Macrocystis pyrifera*, off the coast of southern California and northern Baja California. Calif. Fish. Game 56: 288-297.

Moore C.J., Moore S.L. Leecaster M.K. & Weisberg S.B. 2001. A comparison of plastic and plankton in the North Pacific central gyre. Mar. Poll. Bull. 42: 1297-1300.

Moreira P.S. 1972. Species of marine Isopoda (Crustacea, Peracarida) from Southern Brazil. Bolm. Inst. Oceanogr., S Paulo 21: 163-179.

Morris R.J. 1980. Floating plastic debris in the Mediterranean. Mar. Poll. Bull. 11: 125.

Müller H.G. 1989. Marine Isopoda from the Levantine Basin of the Mediterranean Sea. Senckenbergiana marit. 20: 237-240.

Naylor E. 1955. The diet and feeding mechanism of *Idotea*. J. mar. biol. Ass. U.K. 34: 347-355.

Naylor E. 1957. The occurrence of *Idotea metallica* Bosc in british waters. J. mar. biol. Ass. U.K. 36: 599-602.

Naylor E. 1972. British Marine Isopods. Linn. Soc. Synopsis of the British Fauna (new series) No. 3, Academic Press, London.

Olivar M.P., Sabates A., Abelló P. & Garcia, M. 1998. Transitory hydrographic structures and distribution of fish larvae and neustonic crustaceans in the north-western Mediterranean. *Oceanologica Acta* 21: 95-104.

Ott J. 1996. Meereskunde. Ulmer, Stuttgart.

Park T., Mertz D.B., Grodzinski W. & Prus T. 1965. Cannibalistic predation in populations of flour beetles. *Physiol. Zool.* 38: 289-321.

Parr A.E. 1939. Quantitative observations on the pelagic *Sargassum* vegetation of the western North Atlantic. *Bull. Bingham Oceanogr. Coll.* 6: 1-94.

Parson T.R., Takahashi M. & Hargrave B. 1984. Biological oceanographic processes. 3rd edition. Pergamon Press, Oxford.

Pavia H., Cervin G., Lindgren A. & Åberg P. 1997. Effects of UV-B radiation and simulated herbivory on phlorotannins in the brown alga *Ascophyllum nodosum*. *Mar. Ecol. Prog. Ser.* 157: 139-146.

Pickard G.L. & Emery W.J. 1990. Descriptive Physical Oceanography. An introduction. 5th enlarged edition. Pergamon Press, Oxford.

Reich M. & Grimm V. 1996. Das Metapopulationskonzept in Ökologie und Naturschutz: Eine kritische Bestandsaufnahme. *Z. Ökologie u. Naturschutz* 5: 123-139.

Reid P.C., Planque B. & Edwards M. 1998. Is observed variability in the long-term results of the Continuous Plankton Recorder survey a response to climate change? *Fish. Oceanogr.* 7: 282-288.

Romestand B. & Trilles J.-P. 1976. Les protéines, les glucides et les lipides dans l'hémolymphe et l'hépatopancréas de quelques isopodes cymothoidae, et leurs variations au cours du jeûne expérimental. *Bull. Soc. Zool. France* 101: 445-455.

Roy K., Jablonski D., Valentine J.W. & Rosenberg G. 1998. Marine latitudinal diversity gradients: Tests of causal hypotheses. *Proc. Natl. Acad. Sci. USA* 95: 3699-3702.

Ruiz G.M., Carlton J.T., Grosholz E.D. & Hines A.H. 1997. Global invasions of marine estuarine habitats by non-indigenous species: mechanisms, extent, and consequences. *Amer. Zool.* 37: 621-632.

Salemaa H. 1979. Ecology of *Idotea* spp. (Isopoda) in the northern Baltic. *Ophelia* 18: 133-150.

Salemaa H. 1987. Herbivory and microhabitat preferences of *Idotea* spp. (Isopoda) in the northern Baltic Sea. *Ophelia* 27: 1-15.

Schoener A. & Rowe G.T. 1970. Pelagic *Sargassum* and its presence among the deep sea benthos. *Deep Sea Res.* 17: 923-925.

Seymour R.J., Tegner M.J., Dayton P.K. & Parnell P.E. 1989. Storm wave induced mortality of giant kelp, *Macrocystis pyrifera*, in southern California. *Estuar. Coast. Shelf Sci.* 28: 277-292.

Steeves H.R. III 1969. Lipid content of the hepatopancreas of the isopod *Bathynomus giganteus* A. Milne Edwards, 1879. *Crustaceana* 16: 135-138.

Tegner M.J., Dayton P.K., Edwards P.B. & Riser K.L. 1995. Sea urchin cavitation of giant kelp (*Macrocystis pyrifera* C. Agardh) holdfasts and its effects on kelp mortality across a large California forest. *J. exp. mar. Biol. Ecol.* 191: 83-99.

Thiel M., Hinojosa N., Vásquez N. & Macaya E. 2003. Floating marine debris in coastal waters of the SE-Pacific (Chile). *Mar. Poll. Bull.* 46: 224-231.

Thurman H.V. 1997. Introductory oceanography, 8th ed. Prentice Hall, Upper Saddle River.

Timmermann G. 1932. Biogeographische Untersuchungen über die Lebensgemeinschaft des treibenden Golfkrautes. Z. Morph. Ökol. Tiere 25: 288-335.

Travis J. 1993. Invader threatens Black, Azov Seas. Science 262: 1366-1367.

Tugwell S. & Branch G.M. 1992. Effects of herbivore gut surfactants on kelp polyphenol defenses. Ecology 73: 205-215.

Tully O. & C  idigh P.  . 1986. The ecology of *Idotea* species (Isopoda) and *Gammarus locusta* (Amphipoda) on surface driftweed in Galway Bay (West of Ireland). J. mar. biol. Ass. U.K. 66: 931-942.

Tully O. & McGrath D. 1987. The status of *Idotea metallica* Bosc (Isopoda) in Irish waters. Ir. Nat. J. 22: 190-192.

Tuomi J., Jormalainen V. & Iivessalo H. 1988. Does the aquatic isopod *Idotea baltica* minimize the survival costs of reproduction? Oikos 52: 245-249.

Vasquez J.A. 1993. Effects on the animal community of dislodgement of holdfasts of *Macrocystis pyrifera*. Pac. Sci. 47: 180-184.

Wares J.P. 2001. Intraspecific variation and geographic isolation in *Idotea balthica* (Isopoda: Valvifera). J. Crust. Biol. 21: 1007-1013.

Westheide W., Ha  -Cordes E., Krabusch M. & M  ller M.C.M. 2003. *Ctenodrilus serratus* (Polychaeta: Ctenodrilidae) is a truly ampho-Atlantic meiofauna species – evidence from molecular data. Mar. Biol. 142: 637-642.

Wiebe P.H. 1991. Ringe am Rande des Golfstroms. In: Biologie der Meere. Spektrum Akad. Verl., Heidelberg, S. 46-57.

Wiencke C., G  mez I., Pakker H., Flores-Moya A., Altamirano M., Hanelt D., Bischof K. & Figueroa F.L. 2000. Impact of UV-radiation on viability, photosynthetic

characteristics and DNA of brown algal zoospores: implications for depth zonation. Mar. Ecol. Prog. Ser. 197: 217-229.

Winston J.E. 1982. Drift plastic – an expanding niche for a marine invertebrate? Mar. Poll. Bull. 13: 348-351.

Wolanski E. & Hamner W.M. 1988. Topographically controlled fronts in the ocean and their biological influence. Science 241: 177-181.

Wong C.S., Green D.R. & Cretney W.J. 1974. Quantitative tar and plastic waste distributions in the Pacific Ocean. Nature 247: 30-32.

Zöllner N. & Kirsch K. 1962. Über die quantitative Bestimmung von Lipoiden (Mikromethode) mittels der vielen natürlichen Lipoiden (allen bekannten Plasmalipoiden) gemeinsamen Sulphosphovanillin-Reaktion. Z. ges. Exp. Med. 135: 545-561.