

7 Literatur

- [1] Allison, T.; McCarthy, G.; Wood, C. C.; Darcey, T. M.; Spencer, D. D. und Williamson, P. D. (1989): Human cortical potentials evoked by stimulation of the median nerve. II. Cytoarchitectonic areas generating short-latency activity., J.Neurophysiol. (Band 62), Nr. 3, Seite 694-710.
- [2] Allison, T.; McCarthy, G.; Wood, C. C.; Williamson, P. D.; Spencer, D. D. und Darcey, T. M. (1989): Human cortical potentials evoked by stimulation of the median nerve. II. Cytoarchitectonic areas generating long-latency activity., J.Neurophysiol. (Band 62), Nr. 3, Seite 711-722.
- [3] Allison, T.; Wood, C. C.; McCarthy, G. und Spencer, D. D. (1991): Cortical somatosensory evoked potentials. II. Effects of excision of somatosensory or motor cortex in humans and monkeys., J.Neurophysiol. (Band 66), Nr. 1, Seite 64-82.
- [4] Alper, J. (1993): EEG + MRI: a sum greater than the parts., Science (Band 261), Nr. 5121, Seite 559.
- [5] Bandettini, P. A.:MRI studies of brain activation: Temporal characteristics. Berkely, CA.
- [6] Bandettini, P. A.; Wong, E. C.; Hinks, R. S.; Tikofsky, R. S. und Hyde, J. S. (1992): Time course EPI of human brain function during task activation., Magn.Reson.Med. (Band 25), Nr. 2, Seite 390-397.
- [7] Baumgartner, C.; Barth, D. S.; Levesque, M. F. und Sutherling, W. W. (1992): Human hand and lip sensorimotor cortex as studied on electrocorticography., Electroencephalogr.Clin.Neurophysiol. (Band 84), Nr. 2, Seite 115-126.
- [8] Baumgartner, C.; Doppelbauer, A.; Sutherling, W. W.; Zeitlhofer, J.; Lindinger, G.; Lind, C.; Deecke, L. und Barth, D. S. (1991): Human somatosensory cortical finger representation as studied by combined neuromagnetic and neuroelectric measurements.
Neuromagnetic investigation of somatotopy of human hand somatosensory cortex., Neurosci.Lett. (Band 134), Nr. 1, Seite 103-108.
- [9] Beisteiner, R.; Gomiscek, G.; Erdler, M.; Teichtmeister, C. ; Moser, E. und Deecke, L. (1995): Comparing localization of conventional functional magnetic resonance imaging and magnetoencephalography., Eur.J.Neurosci. (Band 7), Nr. 5, Seite 1121-1124.
- [10] Belliveau, J. W.; Kennedy, D. N. Jr; McKinstry, R. C.; Buchbinder, B. R.; Weisskoff, R. M.; Cohen, M. S.; Vevea, J. M.; Brady, T. J. und Rosen, B. R. (1991): Functional mapping of the human visual cortex by magnetic resonance imaging., Science (Band 254), Nr. 5032, Seite 716-719.

- [11] Belliveau, J. W.; Rosen, B. R.; Kantor, H. L.; Rzedzian, R. R.; Kennedy, D. N.; McKinstry, R. C.; Vevea, J. M.; Cohen, M. S.; Pykett, I. L. und Brady, T. J. (1990): Functional cerebral imaging by susceptibility-contrast NMR., Magn.Reson.Med. (Band 14), Nr. 3, Seite 538-546.
- [12] Biermann, K.; Schmitz, F.; Witte, O. W.; Konczak, J.; Freund, H. J. und Schnitzler, A. (1998): Interaction of finger representation in the human first somatosensory cortex: a neuromagnetic study., Neurosci.Lett. (Band 251), Nr. 1, Seite 13-16.
- [13] Binder, J. R.; Rao, S. M.; Hammeke, T. A.; Frost, J. A.; Bandettini, P. A. und Hyde, J. S. (1994): Effects of stimulus rate on signal response during functional magnetic resonance imaging of auditory cortex., Brain Res.Cogn.Brain Res. (Band 2), Nr. 1, Seite 31-38.
- [14] Blamire, A. M.; Ogawa, S.; Ugurbil, K.; Rothman, D.; McCarthy, G.; Ellermann, J. M.; Hyder, F.; Rattner, Z. und Shulman, R. G. (1992): Dynamic mapping of the human visual cortex by high-speed magnetic resonance imaging., Proc.Natl.Acad.Sci.U.S.A. (Band 89), Nr. 22, Seite 11069-11073.
- [15] Blankenberg, F. G. und Strauss, H. W. (2002): Nuclear medicine applications in molecular imaging., J.Magn.Reson.Imaging (Band 16), Nr. 4, Seite 352-361.
- [16] Blankenburg, F.; Ruben, J.; Meyer, R.; Schwemann, J. und Villringer, A. (2003): Evidence for a rostral-to-caudal somatotopic organization in human primary somatosensory cortex with mirror-reversal in areas 3b and 1, Cereb.Cortex (Band 13), Nr. 9, Seite 987-993.
- [17] Blankenburg, F.; Taskin, B.; Ruben, J.; Moosmann, M.; Ritter, P.; Curio, G. und Villringer, A. (2003): Imperceptible stimuli and sensory processing impediment, Science (Band 299), Nr. 5614, Seite 1864.
- [18] Blood, A. J. und Toga, A. W. (1998): Optical intrinsic signal imaging responses are modulated in rodent somatosensory cortex during simultaneous whisker and forelimb stimulation., J.Cereb.Blood Flow Metab. (Band 18), Nr. 9, Seite 968-977.
- [19] Bodegard; Ledberg, A.; Geyer, S.; Naito, E.; Zilles, K. und Roland, P. E. (2000): T object shape differences reflected by somatosensory cortical activation [In Process Citation], J.Neurosci. (Band 20), Nr. 1, Seite RC51.
- [20] Bodegard, A.; Geyer, S.; Naito, E.; Zilles, K. und Roland, P. E. (2000): Somatosensory areas in man activated by moving stimuli: cytoarchitectonic mapping and PET., Neuroreport (Band 11), Nr. 1, Seite 187-191.
- [21] Brodmann, K. (1909): Vergleichende Lokalisationslehre der Grosshirnrinde in ihre Prinzipien dargestellt auf Grund des Zellenbaus, J. A. Barth, Leipzig.
- [22] Brooks, R. A.; Battocletti, J. H.; Sances, A. Jr; Larson, S. J.; Bowman, R. L. und Kudravcev, V. (1975): Nuclear magnetic relaxation in blood., IEEE Trans.Biomed.Eng. (Band 22), Nr. 1, Seite 12-18.

- [23] Buchner, H.; Adams, L. ; Muller, A.; Ludwig, I.; Knepper, A.; Thron, A.; Niemann, K. und Scherg, M. (1995): Somatotopy of human hand somatosensory cortex revealed by dipole source analysis of early somatosensory evoked potentials and 3D-NMR tomography., *Electroencephalogr.Clin.Neurophysiol.* (Band 96), Nr. 2, Seite 121-134.
- [24] Buchner, H.; Fuchs, M. ; Wischmann, H. A.; Dossel, O.; Ludwig, I.; Knepper, A. und Berg, P. (1994): Source analysis of median nerve and finger stimulated somatosensory evoked potentials: multichannel simultaneous recording of electric and magnetic fields combined with 3D-MR tomography., *Brain Topogr.* (Band 6), Nr. 4, Seite 299-310.
- [25] Burton, H.; MacLeod, A. M.; Videen, T. O. und Raichle, M. E. (1997): Multiple foci in parietal and frontal cortex activated by rubbing embossed grating patterns across fingerpads: a positron emission tomography study in humans., *Cereb.Cortex.* (Band 7), Nr. 1, Seite 3-17.
- [26] Cannestra, A. F.; Black, K. L.; Martin, N. A.; Cloughesy, T.; Burton, J. S.; Rubinstein, E. ; Woods, R. P. und Toga, A. W. (1998): Topographical and temporal specificity of human intraoperative optical intrinsic signals., *Neuroreport* (Band 9), Nr. 11, Seite 2557-2563.
- [27] Cannestra, A. F.; Blood, A. J.; Black, K. L. und Toga, A. W. (1996): The evolution of optical signals in human and rodent cortex., *Neuroimage* (Band 3), Nr. 3 Pt 1, Seite 202- 208.
- [28] Cannestra, A. F.; Pouratian, N.; Bookheimer, S. Y.; Martin, N. A.; Beckerand, D. P. und Toga, A. W. (2001): Temporal spatial differences observed by functional MRI and human intraoperative optical imaging., *Cereb.Cortex.* (Band 11), Nr. 8, Seite 773-782.
- [29] Cho, Z. H.; Ro, Y. M. und Lim, T. H. (1992): NMR venography using the susceptibility effect produced by deoxyhemoglobin., *Magn.Reson.Med.* (Band 28), Nr. 1, Seite 25-38.
- [30] Davis, T. (1997): St. Louis
- [31] Dreyer, D. A.; Loe, P. R.; Metz, C. B. und Whitsel, B. L. (1975): Representation of head and face in postcentral gyrus of the macaque., *J.Neurophysiol.* (Band 38), Nr. 3, Seite 714-733.
- [32] Duus, P. (1990): *Neurologischtopische Diagnostik*, 5. Auflage , Thieme.
- [33] Economo, v. C. und Koskinas, G. N. (1925): *Die Cytoarchitektonik der Hirnrinde des erwachsenen Menschen.*, Springer.
- [34] Engel, S. A. (1996): Looking into the black box: new directions in neuroimaging., *Neuron* (Band 17), Nr. 3, Seite 375-378.
- [35] Favorov, O. V. und Diamond, M. E. (1990): Demonstration of discrete place-defined columns--segregates--in the cat SI., *J.Comp.Neurol.* (Band 298), Nr. 1, Seite 97-112.

- [36] Ferrington, D. G. und Rowe, M. (1980): Differential contributions to coding of cutaneous vibratory information by cortical somatosensory areas I and II., J.Neurophysiol. (Band 43), Nr. 2, Seite 310-331.
- [37] Forss, N.; Salmelin, R. und Hari, R. (1994): Comparison of somatosensory evoked fields to airpuff and electric stimuli., Electroencephalogr.Clin.Neurophysiol. (Band 92), Nr. 6, Seite 510-517.
- [38] Fox, P. T.; Burton, H. und Raichle, M. E. (1987): Mapping human somatosensory cortex with positron emission tomography., J.Neurosurg. (Band 67), Nr. 1, Seite 34-43.
- [39] Fox, P. T.; Mintun, M. A.; Raichle, M. E.; Miezin, F. M. ; Allman, J. M. und Van, Essen DC (1986): Mapping human visual cortex with positron emission tomography., Nature (Band 323), Nr. 6091, Seite 806-809.
- [40] Fox, P. T. und Raichle, M. E. (1984): Stimulus rate dependence of regional cerebral blood flow in human striate cortex, demonstrated by positron emission tomography., J.Neurophysiol. (Band 51), Nr. 5, Seite 1109-1120.
- [41] Fox, P. T.; Raichle, M. E.; Mintun, M. A. und Dence, C. (1988): Nonoxidative glucose consumption during focal physiologic neural activity., Science (Band 241), Nr. 4864, Seite 462-464.
- [42] Frackowiak, R. S.; Friston, K. J.; Frith, C. D.; Dolan, R. J. und Mazziotta, J. C. (1997): Human Brain Function, 1. Auflage , Academic Press, San Diego, London, Boston, New York, Sydney, Tokyo, Toronto, ISBN: 0-12-264840-4.
- [43] Frackowiak, R. S.; Lenzi, G. L.; Jones, T. und Heather, J. D. (1980): Quantitative measurement of regional cerebral blood flow and oxygen metabolism in man using ^{15}O and positron emission tomography: theory, procedure, and normal values., J.Comput.Assist.Tomogr. (Band 4), Nr. 6, Seite 727-736.
- [44] Frahm, J.; Bruhn, H.; Merboldt, K. D. und Hanicke, W. (1992): Dynamic MR imaging of human brain oxygenation during rest and photic stimulation., J.Magn.Reson.Imaging (Band 2), Nr. 5, Seite 501-505.
- [45] Frahm, J.; Merboldt, K. D. und Hanicke, W. (1993): Functional MRI of human brain activation at high spatial resolution., Magn.Reson.Med. (Band 29), Nr. 1, Seite 139-144.
- [46] Frahm, J.; Merboldt, K. D.; Hanicke, W.; Kleinschmidt, A. und Boecker, H. (1994): Brain or vein--oxygenation or flow? On signal physiology in functional MRI of human brain activation., NMR.Biomed. (Band 7), Nr. 1-2, Seite 45-53.
- [47] Francis, S. T.; Kelly, E. F.; Bowtell, R.; Dunseath, W. J.; Folger, S. E. und McGlone, F. (2000): fMRI of the Responses to Vibratory Stimulation of Digit Tips., Neuroimage (Band 11), Nr. 3, Seite 188-202.
- [48] Fransson, P.; Krüger, G.; Merboldt, K. D.; and Frahm, J. (1998):

- [49] Friston, K. J.; Williams, S.; Howard, R.; Frackowiak, R. S. und Turner, R. (1996): Movement-related effects in fMRI time-series., Magn.Reson.Med. (Band 35), Nr. 3, Seite 346-355.
- [50] Frostig, R. D.; Lieke, E. E.; Ts und Grinvald, A. (1990): Cortical functional architecture and local coupling between neuronal activity and the microcirculation revealed by in vivo high-resolution optical imaging of intrinsic signals., Proc.Natl.Acad.Sci.U.S.A. (Band 87), Nr. 16, Seite 6082-6086.
- [51] Gelnar, P. A.; Krauss, B. R.; Szeverenyi, N. M. und Apkarian, A. V. (1998): Fingertip Representation in the Human Somatosensory Cortex: An fMRI Study., Neuroimage. (Band 7), Nr. 4, Seite 261-283.
- [52] Geyer, S.; Ledberg, A. ; Schleicher, A.; Kinomura, S.; Schormann, T.; Burgel, U.; Klingberg, T.; Larsson, J.; Zilles, K. und Roland, P. E. (1996): Two different areas within the primary motor cortex of man., Nature (Band 382), Nr. 6594, Seite 805-807.
- [53] Geyer, S.; Schleicher, A. und Zilles, K. (1997): The somatosensory cortex of human: cytoarchitecture and regional distributions of receptor-binding sites., Neuroimage. (Band 6), Nr. 1, Seite 27-45.
- [54] Geyer, S.; Schleicher, A. und Zilles, K. (1999): Areas 3a, 3b, and 1 of human primary somatosensory cortex., Neuroimage. (Band 10), Nr. 1, Seite 63-83.
- [55] Godde, B.; Hilger, T.; von Seelen W.; Berkefeld, T. und Dinse, H. R. (1995): Optical imaging of rat somatosensory cortex reveals representational overlap as topographic principle., Neuroreport (Band 7), Nr. 1, Seite 24-28 .
- [56] Grimm, C.; Schreiber, M.; Kristeva-Feige, R.; Mergner, T.; Hennig, J. und Lucking, C. H. (1998): A comparison between electric source localisation and fMRI during somatosensory stimulation., Electroencephalogr.Clin.Neurophysiol. (Band 106), Nr. 1, Seite 22-29.
- [57] Grinvald, A.; Lieke, E.; Frostig, R. D.; Gilbert, C. D. und Wiesel, T. N. (1986): Functional architecture of cortex revealed by optical imaging of intrinsic signals., Nature (Band 324), Nr. 6095, Seite 361-364.
- [58] Grubb, R. L. Jr; Raichle, M. E.; Eichling, J. O. und Ter-Pogossian, M. M. (1974): The effects of changes in PaCO₂ on cerebral blood volume, blood flow, and vascular mean transit time., Stroke (Band 5), Nr. 5, Seite 630-639.
- [59] Gyngell, M. L.; Bock, C.; Schmitz, B.; Hoehn-Berlage, M. und Hossmann, K. A. (1996): Variation of functional MRI signal in response to frequency of somatosensory stimulation in alpha-chloralose anesthetized rats, Magn Reson.Med. (Band 36), Nr. 1, Seite 13-15.
- [60] Haase, A.; Frahm, J.; Matthaei, D.; Hänicke, W. und Merboldt, K. D. (1986): FLASH imaging. Rapid NMR imaging using low flip-angle pulses., J.Magn.Reson.Imaging (Band 67), Seite 258-258.

- [61] Hamalainen, M. S. (1992): Magnetoencephalography: a tool for functional brain imaging., Brain Topogr. (Band 5), Nr. 2, Seite 95-102.
- [62] Hansson, T. und Brismar, T. (1999): Tactile stimulation of the hand causes bilateral cortical activation: a functional magnetic resonance study in humans., Neurosci.Lett. (Band 271), Nr. 1, Seite 29-32.
- [63] Hari, R.; Karhu, J.; Hamalainen, M.; Knuutila, J.; Salonen, O.; Sams, M. und Vilkman, V. (1993): Functional organization of the human first and second somatosensory cortices: a neuromagnetic study., Eur.J.Neurosci. (Band 5), Nr. 6, Seite 724-734.
- [64] Hari, R. und Kaukoranta, E. (1985): Neuromagnetic studies of somatosensory system: principles and examples., Prog.Neurobiol. (Band 24), Nr. 3, Seite 233-256.
- [65] Harrington, G. S. und Hunter, Downs J. (2001): FMRI mapping of the somatosensory cortex with vibratory stimuli. Is there a dependency on stimulus frequency?, Eur.J.Neurosci. (Band 897), Nr. 1-2, Seite 188-192.
- [66] Heeger, D. J.; Huk, A. C.; Geisler, W. S. und Albrecht, D. G. (2000): Spikes versus BOLD: what does neuroimaging tell us about neuronal activity?, Nat.Neurosci. (Band 3), Nr. 7, Seite 631-633.
- [67] Heilman, K. und Watson, S. (1977): The Neglect Syndrome - A Unilateral Defect of the Orienting Response., Harnad, S.; Doty, R.; Goldstein, L.; Jaynes, J. und Krauthamer, G., Lateralization in the Nervous System. Academic Press, New York.
- [68] Horwitz, B. und Poeppel, D. (2002): How can EEG/MEG and fMRI/PET data be combined?, Hum.Brain Mapp. (Band 17), Nr. 1, Seite 1-3 .
- [69] Hoshi, Y. und Tamura, M. (1993): Detection of dynamic changes in cerebral oxygenation coupled to neuronal function during mental work in man, Neuroscience Letters (Band 150), Seite 5-8.
- [70] Hoshiyama, M.; Kakigi, R.; Koyama, S.; Kitamura, Y.; Shimojo, M. und Watanabe, S. (1996): Somatosensory evoked magnetic fields following stimulation of the lip in humans., Electroencephalogr.Clin.Neurophysiol. (Band 100), Nr. 2, Seite 96-104.
- [71] Hyvarinen, J. und Poranen, A. (1978): Receptive field integration and submodality convergence in the hand area of the post-central gyrus of the alert monkey, J.Physiol (Band 283), Seite 539-556.
- [72] Ibanez, V.; Deiber, M. P.; Sadato, N.; Toro, C.; Grissom, J.; Woods, R. P.; Mazziotta, J. C. und Hallett, M. (1995): Effects of stimulus rate on regional cerebral blood flow after median nerve stimulation., Brain (Band 118), Nr. Pt 5, Seite 1339-1351.
- [73] Iwamura, Y.; Iriki, A. und Tanaka, M. (1994): Bilateral hand representation in the postcentral somatosensory cortex., Nature (Band 369), Nr. 6481, Seite 554-556.

- [74] Iwamura, Y. und Tanaka, M. (1978): Functional organization of receptive fields in the cat somatosensory cortex. II: Second representation of the forepaw in the ansate region., Eur.J.Neurosci. (Band 151), Nr. 1, Seite 61-72.
- [75] Iwamura, Y.; Tanaka, M. und Hikosaka, O. (1980): Overlapping representation of fingers in the somatosensory cortex (area 2) of the conscious monkey., Eur.J.Neurosci. (Band 197), Nr. 2, Seite 516- 520.
- [76] Iwamura, Y.; Tanaka, M.; Sakamoto, M. und Hikosaka, O. (1983): Converging patterns of finger representation and complex response properties of neurons in area 1 of the first somatosensory cortex of the conscious monkey., Exp.Brain Res. (Band 51), Seite 327-337.
- [77] Johansson, R. S.; Trulsson, M.; Olsson, K. A. und Westberg, K. G. (1988): Mechanoreceptor activity from the human face and oral mucosa., Exp.Brain Res. (Band 72), Nr. 1, Seite 204-208.
- [78] Jones, T.; Chesler, D. A. und Ter-Pogossian, M. M. (1976): The continuous inhalation of oxygen-15 for assessing regional oxygen extraction in the brain of man., Br.J.Radiol. (Band 49), Nr. 580, Seite 339-343.
- [79] Kaas, J. H. (1983): What, if anything, is SI? Organization of first somatosensory area of cortex., Physiol.Rev. (Band 63), Nr. 1, Seite 206-231.
- [80] Kaas, J. H.; Nelson, R. J.; Sur, M.; Lin, C. S. und Merzenich, M. M. (1979): Multiple representations of the body within the primary somatosensory cortex of primates., Science (Band 204), Nr. 4392, Seite 521-523.
- [81] Kampe, K. K.; Jones, R. A. und Auer, D. P. (2000): Frequency dependence of the functional MRI response after electrical median nerve stimulation., Hum.Brain Mapp. (Band 9), Nr. 2, Seite 106-114.
- [82] Kandel, E. und Jessell, T. M. (1991): Touch, Kandel, E.; Schwartz, J. und Jessell, T. M., Principles of Neural Science , Seite 367-384, Appleton and Lange, Norwalk.
- [83] Kandel, E.; Schwartz, J. und Jessel, T. (1996): Neurowissenschaften, 1. Auflage , Spektrum Akademischer Verlag, Heidelberg, Berlin, Oxford, ISBN: 3-86025-391-3.
- [84] Kandel, E.; Schwartz, J. und Jessel, T. (2000): Principles of Neural Science, 4. Auflage , McGraw-Hill, New York, ISBN: 0-07-112000-9.
- [85] Kato, T.; Kamei, A.; Takashima, S. und Ozaki, T. (1993): Human visual cortecal function during photic stimulation monitoring by means of near-infrared spectroscopy, J.Cereb.Blood Flow Metab. (Band 13), Seite 516-520.

- [86] Kaukoranta, E.; Hamalainen, M.; Sarvas, J. und Hari, R. (1986): Mixed and sensory nerve stimulations activate different cytoarchitectonic areas in the human primary somatosensory cortex SI. Neuromagnetic recordings and statistical considerations., Exp.Brain Res. (Band 63), Nr. 1, Seite 60-66.
- [87] Kelly, E. F. und Folger, S. E. (1999): EEG evidence of stimulus-directed response dynamics in human somatosensory cortex., Eur.J.Neurosci. (Band 815), Nr. 2, Seite 326-336.
- [88] Kischka, U.; Wallesch, C.-W. und Wolf, G. (1997): Methoden der Hirnforschung, Spektrum, Akademischer Verlag, Heidelberg, Berlin, ISBN: 3-8274-0108-9 .
- [89] Korvenoja, A.; Wikstrom, H.; Huttunen, J.; Virtanen, J.; Laine, P.; Aronen, H. J.; Seppäläinen, A. M. und Ilmoniemi, R. J. (1995): Activation of ipsilateral primary sensorimotor cortex by median nerve stimulation., Neuroreport (Band 6), Nr. 18, Seite 2589-2593.
- [90] Krause, T.; Kurth, R.; Ruben, J.; Schwiemann, J.; Villringer, K.; Deuchert, M.; Moosmann, M.; Brandt, S.; Wolf, K.; Curio, G. und Villringer, A. (2001): Representational overlap of adjacent fingers in multiple areas of human primary somatosensory cortex depends on electrical stimulus intensity: an fMRI study., Eur.J.Neurosci. (Band 899), Nr. 1-2, Seite 36-46.
- [91] Kurth, R.; Villringer, K.; Curio, G.; Wolf, K. J.; Krause, T.; Repenthin, J.; Schwiemann, J.; Deuchert, M. und Villringer, A. (2000): fMRI shows multiple somatotopic digit representations in human primary somatosensory cortex, Neuroreport (Band 11), Nr. 7, Seite 1487-1491.
- [92] Kurth, R.; Villringer, K.; Mackert, B. M.; Schwiemann, J.; Braun, J.; Curio, G.; Villringer, A. und Wolf, K. J. (1998): fMRI assessment of somatotopy in human Brodmann area 3b by electrical finger stimulation., Neuroreport (Band 9), Nr. 2, Seite 207 - 212.
- [93] Kwong, K. K.; Belliveau, J. W.; Chesler, D. A.; Goldberg, I. E.; Weisskoff, R. M.; Poncelet, B. P.; Kennedy, D. N.; Hoppel, B. E. ; Cohen, M. S. und Turner, R. (1992): Dynamic magnetic resonance imaging of human brain activity during primary sensory stimulation., Proc.Natl.Acad.Sci.U.S.A. (Band 89), Nr. 12, Seite 5675-5679.
- [94] Lai, S.; Hopkins, A. L.; Haacke, E. M.; Li, D.; Wasserman, B. A.; Buckley, P.; Friedman, L.; Meltzer, H.; Hedera, P. und Friedland, R. (1993): Identification of vascular structures as a major source of signal contrast in high resolution 2D and 3D functional activation imaging of the motor cortex at 1.5T: preliminary results., Magn.Reson.Med. (Band 30), Nr. 3, Seite 387-392.
- [95] Lauritzen, M. (2001): Relationship of spikes, synaptic activity, and local changes of cerebral blood flow, J.Cereb.Blood Flow Metab (Band 21), Nr. 12, Seite 1367-1383.
- [96] Lauritzen, M. und Gold, L. (2003): Brain function and neurophysiological correlates of signals used in functional neuroimaging, J.Neurosci. (Band 23), Nr. 10, Seite 3972-3980.

- [97] Le, Bihan D.; Mangin, J. F.; Poupon, C.; Clark, C. A.; Pappata, S.; Molko, N. und Chabriat, H. (2001): Diffusion tensor imaging: concepts and applications., *J.Magn.Reson.Imaging* (Band 13), Nr. 4, Seite 534-546.
- [98] Lee (1992): Induction of high frequency activity in the somatosensory thalamus of rats in vivo results in long-term potentiation of responses in SI cortex, *Experimental Brain Research*, Seite 253-261.
- [99] Leniger-Follert, E. und Hossmann, K. A. (1979): Simultaneous measurements of microflow and evoked potentials in the somatomotor cortex of the cat brain during specific sensory activation., *Pflugers Arch.* (Band 380), Nr. 1, Seite 85-89.
- [100] Lin, W.; Kuppusamy, K.; Haacke, E. M. und Burton, H. (1996): Functional MRI in human somatosensory cortex activated by touching textured surfaces., *J.Magn.Reson.Imaging* (Band 6), Nr. 4, Seite 565-572.
- [101] Logothetis, N. K.; Pauls, J.; Augath, M.; Trinath, T. und Oeltermann, A. (2001): Neurophysiological investigation of the basis of the fMRI signal, *Nature* (Band 412), Nr. 6843, Seite 150-157.
- [102] Lotze, M.; Flor, H.; Grodd, W.; Larbig, W. und Birbaumer, N. (2001): Phantom movements and pain. An fMRI study in upper limb amputees, *Brain* (Band 124), Nr. Pt 11, Seite 2268-2277.
- [103] Maldjian, J. A.; Gottschalk, A.; Patel, R. S.; Detre, J. A. und Alsop, D. C. (1999): The sensory somatotopic map of the human hand demonstrated at 4 Tesla., *Neuroimage* (Band 10), Nr. 1 , Seite 55-62.
- [104] Manger, P. R.; Woods, T. M. und Jones, E. G. (1995): Representation of the face and intraoral structures in area 3b of the squirrel monkey (*Saimiri sciureus*) somatosensory cortex, with special reference to the ipsilateral representation., *J.Comp.Neurol.* (Band 362), Nr. 4, Seite 597-607.
- [105] Manger, P. R.; Woods, T. M. und Jones, E. G. (1996): Representation of face and intra-oral structures in area 3b of macaque monkey somatosensory cortex., *J.Comp.Neurol.* (Band 371), Nr. 4, Seite 513-521.
- [106] Mansfield, P. und Maudsley, A. A. (1977): Medical imaging by NMR., *Br.J.Radiol.* (Band 50), Nr. 591, Seite 188-194.
- [107] Martin, J. H. und Jessell, T. M. (1991): Anatomy of the Somatic Sensory System, Kandel, E.; Schwartz, J. und Jessell, T. M., *Principles of Neural Science* , Seite 353-366, Appleton and Lange, Norwalk, CT.
- [108] McCarthy, G. und Allison, T. (1995): Trigeminal evoked potentials in somatosensory cortex of the Macaca mulatta., *J.Neurosurg.* (Band 82), Nr. 6, Seite 1015-1020.
- [109] McCarthy, G.; Allison, T. und Spencer, D. D. (1993): Localization of the face area of human sensorimotor cortex by intracranial recording of somatosensory evoked potentials., *J.Neurosurg.* (Band 79), Nr. 6, Seite 874-884.

- [110] McCarthy, G.; Wood, C. C. und Allison, T. (1991): Cortical somatosensory evoked potentials. I. Recordings in the monkey *Macaca fascicularis.*, *J.Neurophysiol.* (Band 66), Nr. 1, Seite 53-63.
- [111] Menon, R. S.; Gati, J. S.; Goodyear, B. G.; Luknowsky, D. C. und Thomas, C. G. (1998): Spatial and temporal resolution of functional magnetic resonance imaging [In Process Citation], *Biochem.Cell Biol.* (Band 76), Nr. 2-3, Seite 560-571.
- [112] Mogilner, A.; Grossman, J. A.; Ribary, U.; Joliot, M.; Volkmann, J.; Rapaport, D.; Beasley, R. W. und Llinas, R. R. (1993): Somatosensory cortical plasticity in adult humans revealed by magnetoencephalography., *Proc.Natl.Acad.Sci.U.S.A.* (Band 90), Nr. 8, Seite 3593-3597.
- [113] Mogilner, A.; Nomura, M.; Ribary, U.; Jagow, R.; Lado, F.; Rusinek, H. und Llinas, R. (1994): Neuromagnetic studies of the lip area of primary somatosensory cortex in humans: evidence for an oscillotopic organization., *Exp.Brain Res.* (Band 99), Nr. 1, Seite 137-147.
- [114] Moore, C. I.; Stern, C. E.; Corkin, S.; Fischl, B.; Gray, A. C.; Rosen, B. R. und Dale, A. M. (2000): Segregation of somatosensory activation in the human rolandic cortex using fMRI., *J.Neurophysiol.* (Band 84), Nr. 1, Seite 558-569.
- [115] Mountcastle, V. B. (1984): Central nervous mechanisms in mechanoreceptive sensibility., Darian-Smith, *Handbook of Physiology* , Seite 789-878, American Physiological Society, Bethesda.
- [116] Mountcastle, V. B. und POWELL, T. P. (1959): Neural mechanisms subserving cutaneous sensibility, with special reference to the role of afferent inhibition in sensory perception and discrimination, *Bull.Johns.Hopkins.Hosp.* (Band 105), Seite 201-232.
- [117] Nakamura, A.; Yamada, T.; Goto, A.; Kato, T.; Ito, K.; Abe, Y.; Kachi, T. und Kakigi, R. (1998): Somatosensory Homunculus as Drawn by MEG., *Neuroimage.* (Band 7), Nr. 4, Seite 377-386.
- [118] Nelson, R. J.; Sur, M.; Felleman, D. J. und Kaas, J. H. (1980): Representations of the body surface in postcentral parietal cortex of *Macaca fascicularis.*, *J.Comp.Neurol.* (Band 192), Nr. 4, Seite 611-643.
- [119] Ngai, A. C.; Jolley, M. A.; D'Ambrosio, R.; Meno, J. R. und Winn, H. R. (1999): Frequency-dependent changes in cerebral blood flow and evoked potentials during somatosensory stimulation in the rat [In Process Citation], *Eur.J.Neurosci.* (Band 837), Nr. 1-2, Seite 221-228.
- [120] Nielsen, A. und Lauritzen, M. (2001): Coupling and uncoupling of activity-dependent increases of neuronal activity and blood flow in rat somatosensory cortex., *J.Physiol.* (Band 533), Nr. Pt 3, Seite 773-785.
- [121] Ogawa, S. und Lee, T. M. (1990): Magnetic resonance imaging of blood vessels at high fields: in vivo and in vitro measurements and image simulation., *Magn.Reson.Med.* (Band 16), Nr. 1, Seite 9-18.

- [122] Ogawa, S.; Lee, T. M. und Barrere, B. (1993): The sensitivity of magnetic resonance image signals of a rat brain to changes in the cerebral venous blood oxygenation., Magn.Reson.Med. (Band 29), Nr. 2, Seite 205-210.
- [123] Ogawa, S.; Lee, T. M. ; Kay, A. R. und Tank, D. W. (1990): Brain magnetic resonance imaging with contrast dependent on blood oxygenation., Proc.Natl.Acad.Sci.U.S.A. (Band 87), Nr. 24, Seite 9868-9872.
- [124] Ogawa, S.; Lee, T. M. ; Nayak, A. S. und Glynn, P. (1990): Oxygenation-sensitive contrast in magnetic resonance image of rodent brain at high magnetic fields., Magn.Reson.Med. (Band 14), Nr. 1, Seite 68-78.
- [125] Ogawa, S.; Tank, D. W.; Menon, R.; Ellermann, J. M.; Kim, S. G.; Merkle, H. und Ugurbil, K. (1992): Intrinsic signal changes accompanying sensory stimulation: functional brain mapping with magnetic resonance imaging., Proc.Natl.Acad.Sci.U.S.A. (Band 89), Nr. 13, Seite 5951-5955.
- [126] Okada, Y. C.; Tanenbaum, R.; Williamson, S. J. und Kaufman, L. (1984): Somatotopic organization of the human somatosensory cortex revealed by neuromagnetic measurements., Exp.Brain Res. (Band 56), Nr. 2, Seite 197-205.
- [127] Pauling, L. und Coryell, C. (1936): The magnetic properties and structure of hemoglobin and carbonmonoxyhemoglobin., Proc.Natl.Acad.Sci.U.S.A. (Band 22), Seite 210-216.
- [128] Penfield, W. und Boldrey, E. (1937): Somatic motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation., Brain (Band 60), Seite 389-443.
- [129] Pons, T. P.; Garraghty, P. E.; Cusick, C. G. und Kaas, J. H. (1985): The somatotopic organization of area 2 in macaque monkeys.
Connections of area 2 of somatosensory cortex with the anterior pulvinar and subdivisions of the ventroposterior complex in macaque monkeys. , J.Comp.Neurol. (Band 241), Nr. 4, Seite 445-466.
- [130] Pratt, H.; Politoske, D. und Starr, A. (1980): Mechanically and electrically evoked somatosensory potentials in humans: effects of stimulus presentation rate, Electroencephalogr.Clin.Neurophysiol. (Band 49), Nr. 3-4, Seite 240-249.
- [131] Preibisch, C. und Haase, A. (1999): Functional MR imaging of the human brain using FLASH: influence of various imaging parameters, J.Magn Reson. (Band 140), Nr. 1, Seite 162-171.
- [132] Price, C.; Wise, R.; Ramsay, S.; Friston, K.; Howard, D.; Patterson, K. und Frackowiak, R. (1992): Regional response differences within the human auditory cortex when listening to words., Neurosci.Lett. (Band 146), Nr. 2, Seite 179-182.
- [133] Rao, S. M.; Bandettini, P. A.; Binder, J. R.; Bobholz, J. A.; Hammeke, T. A.; Stein, E. A. und Hyde, J. S. (1996): Relationship between finger movement rate and functional magnetic resonance signal change in human primary motor cortex., J.Cereb.Blood Flow Metab. (Band 16), Nr. 6, Seite 1250-1254.

- [134] Rao, S. M.; Binder, J. R.; Hammeke, T. A.; Bandettini, P. A.; Bobholz, J. A.; Frost, J. A.; Myklebust, B. M.; Jacobson, R. D. und Hyde, J. S. (1995): Somatotopic mapping of the human primary motor cortex with functional magnetic resonance imaging [see comments], Neurology (Band 45), Nr. 5, Seite 919- 924.
- [135] Rees, G.; Friston, K. und Koch, C. (2000): A direct quantitative relationship between the functional properties of human and macaque V5, Nat.Neurosci. (Band 3), Nr. 7, Seite 716-723.
- [136] Rees, G.; Howseman, A.; Josephs, O.; Frith, C. D.; Friston, K. J.; Frackowiak, R. S. und Turner, R. (1997): Characterizing the relationship between BOLD contrast and regional cerebral blood flow measurements by varying the stimulus presentation rate., Neuroimage. (Band 6), Nr. 4, Seite 270-278.
- [137] Regan, D. (1993): Human Brain Electrophysiology, Elsevier, London.
- [138] Rohlfing, T.; West, J. B.; Beier, J.; Liebig, T.; Taschner, C. A. und Thomale, U. W. (2000): Registration of functional and anatomical MRI: accuracy assessment and application in navigated neurosurgery., Comput.Aided.Surg. (Band 5), Nr. 6, Seite 414-425.
- [139] Roland, P. E. und Zilles, K. (1994): Brain atlases--a new research tool., Trends.Neurosci. (Band 17), Nr. 11, Seite 458-467.
- [140] Rosen, B. R.; Belliveau, J. W.; Aronen, H. J.; Kennedy, D.; Buchbinder, B. R.; Fischman, A.; Gruber, M.; Glas, J.; Weisskoff, R. M. und Cohen, M. S. (1991): Susceptibility contrast imaging of cerebral blood volume: human experience., Magn.Reson.Med. (Band 22), Nr. 2, Seite 293-299.
- [141] Rosen, B. R.; Belliveau, J. W.; Vevea, J. M. und Brady, T. J. (1990): Perfusion imaging with NMR contrast agents., Magn.Reson.Med. (Band 14), Nr. 2, Seite 249-265.
- [142] Ruben, J.; Schwiemann, J.; Deuchert, M.; Meyer, R.; Krause, T.; Curio, G.; Villringer, K.; Kurth, R. und Villringer, A. (2001): Somatotopic Organization of Human Secondary Somatosensory Cortex., Cereb.Cortex. (Band 11), Nr. 5, Seite 463-473.
- [143] Sadato; Ibanez und Hallett (1997): Frequency-Dependent changes of regional cerebral blood flow during finger movements: functional MRI compared to PET, Journal of cerebral blood flow and metabolism (Band 17), Nr. 1997, Seite 670-679.
- [144] Sadato, N.; Campbell, G.; Ibanez, V.; Deiber, M. und Hallett, M. (1996): Complexity affects regional cerebral blood flow change during sequential finger movements., J.Neurosci. (Band 16), Nr. 8, Seite 2691-2700.
- [145] Sadato, N.; Ibanez, V.; Deiber, M. P.; Campbell, G.; Leonardo, M. und Hallett, M. (1996): Frequency-dependent changes of regional cerebral blood flow during finger movements., J.Cereb.Blood Flow Metab. (Band 16), Nr. 1, Seite 23-33.

- [146] Schleicher, A.; Amunts, K.; Geyer, S.; Morosan, P. und Zilles, K. (1999): Observer-independent method for microstructural parcellation of cerebral cortex: A quantitative approach to cytoarchitectonics., Neuroimage. (Band 9), Nr. 1, Seite 165 -177.
- [147] Schmidt, R. F. (1995): Neuro- und Sinnesphysiologie, 2. Auflage .
- [148] Shoham, D. und Grinvald, A. (2001): The cortical representation of the hand in macaque and human area S-I: high resolution optical imaging., J.Neurosci. (Band 21), Nr. 17, Seite 6820-6835.
- [149] Sicard, K.; Shen, Q.; Brevard, M. E.; Sullivan, R.; Ferris, C. F.; King, J. A. und Duong, T. Q. (2003): Regional cerebral blood flow and BOLD responses in conscious and anesthetized rats under basal and hypercapnic conditions: implications for functional MRI studies, J.Cereb.Blood Flow Metab (Band 23), Nr. 4, Seite 472-481.
- [150] Singh, K. D. (1995): Functional imaging of the brain using superconducting magnetometry., Endeavour. (Band 19), Nr. 1, Seite 39-44.
- [151] Singh, M.; Khosla, D. ; and Patel, P. (1996): Correlation of fMRI and EEG imaging during visual stimulation, Proc.4th Int.Conference of the International Society for Magnetic Resonance in Medicine (Abstract Band), Seite 1851.
- [152] Sloan, T. B. (1998): Anesthetic effects on electrophysiologic recordings., J.Clin.Neurophysiol. (Band 15), Nr. 3, Seite 217-226.
- [153] Sobol, WT. (1986): Dedicated Coils in Magnetic Resonance Imaging, Rev.Magn.Res.Med. (Band 1(2)), Seite 181-224.
- [154] Sobotta, J. (1993): Obere Extremität, Putz, R. und Pabst, R., Atlas der Anatomie des Menschen , 20. Auflage, Seite 160-250, Urban&Schwarzenberg, München, Wien, Baltimore.
- [155] Stehling, M. K.; Turner, R. und Mansfield, P. (1991): Echo-planar imaging: magnetic resonance imaging in a fraction of a second., Science (Band 254), Nr. 5028, Seite 43-50.
- [156] Stippich, C.; Freitag, P.; Kassubek, J.; Soros, P.; Kamada, K.; Kober, H.; Scheffler, K.; Hopfengartner, R.; Bilecen, D.; Radu, E. W. und Vieth, J. B. (1998): Motor, somatosensory and auditory cortex localization by fMRI and MEG., Neuroreport (Band 9), Nr. 9, Seite 1953-1957.
- [157] Sur, M.; Merzenich, M. M. und Kaas, J. H. (1980): Magnification, receptive-field area, and "hypercolumn" size in areas 3b and 1 of somatosensory cortex in owl monkeys, J.Neurophysiol. (Band 44), Nr. 2, Seite 295-311.
- [158] Sur, M.; Nelson, R. J. und Kaas, J. H. (1982): Representations of the body surface in cortical areas 3b and 1 of squirrel monkeys: comparisons with other primates., J.Comp.Neurol. (Band 211), Nr. 2, Seite 177-192.

- [159] Sutherling, W. W.; Levesque, M. F. und Baumgartner, C. (1992): Cortical sensory representation of the human hand: size of finger regions and nonoverlapping digit somatotopy., Neurology (Band 42), Nr. 5, Seite 1020-1028.
- [160] Thees, S.; Blankenburg, F.; Taskin, B.; Curio, G. und Villringer, A. (2003): Dipole source localization and fMRI of simultaneously recorded data applied to somatosensory categorization, Neuroimage. (Band 18), Nr. 3, Seite 707-719.
- [161] Thulborn, K. R.; Waterton, J. C.; Matthews, P. M. und Radda, G. K. (1982): Oxygenation dependence of the transverse relaxation time of water protons in whole blood at high field., Biochim.Biophys.Acta (Band 714), Nr. 2, Seite 265-270.
- [162] Tommerdahl, M.; Delemos, K. A.; Whitsel, B. L.; Favorov, O. V. und Metz, C. B. (1999): Response of anterior parietal cortex to cutaneous flutter versus vibration., J.Neurophysiol. (Band 82), Nr. 1, Seite 16-33.
- [163] Tommerdahl, M.; Whitsel, B. L.; Favorov, O. V.; Metz, C. B. und O'Quinn, B. L. (1999): Responses of contralateral SI and SII in cat to same-site cutaneous flutter versus vibration., J.Neurophysiol. (Band 82), Nr. 4, Seite 1982-1992.
- [164] Turner, R.; Jezzard, P.; Wen, H.; Kwong, K. K.; Le, Bihan D.; Zeffiro, T. und Balaban, R. S. (1993): Functional mapping of the human visual cortex at 4 and 1,5 Tesla using deoxygenation contrast EPI, Magn.Reson.Imaging, Seite 277-279.
- [165] Turner, R.; Le, Bihan D.; Moonen, C. T.; Despres, D. und Frank, J. (1991): Echo-planar time course MRI of cat brain oxygenation changes., Magn.Reson.Med. (Band 22), Nr. 1, Seite 159-166.
- [166] van Westen, D.; Fransson, P.; Olsrud, J.; Rosen, B.; Lundborg, G. und Larsson, E. M. (2004): Fingersomatotopy in area 3b: an fMRI-study, BMC.Neurosci. (Band 5), Nr. 1, Seite 28.
- [167] Villringer, A. und Dirnagl, U. (1995): Coupling of brain activity and cerebral blood flow: basis of functional neuroimaging., Cerebrovasc.Brain Metab.Rev. (Band 7), Nr. 3, Seite 240-276.
- [168] Villringer, A.; Them, A.; Lindauer, U.; Einhaupl, K. und Dirnagl, U. (1994): Capillary perfusion of the rat brain cortex. An in vivo confocal microscopy study., Circ.Res. (Band 75), Nr. 1, Seite 55-62.
- [169] Vogt, O. und Vogt, C. (1919): Allgemeine Ergebnisse unserer Hirnforschung, J.Psych.Neurol. (Band 25), Seite 279-461.
- [170] Weisskoff, R. M.; Chesler, D.; Boxerman, J. L. und Rosen, B. R. (1993): Pitfalls in MR measurement of tissue blood flow with intravascular tracers: which mean transit time?, Magn.Reson.Med. (Band 29), Nr. 4, Seite 553-558.
- [171] Weisskoff, R. M. und Kihne, S. (1992): MRI susceptometry: image-based measurement of absolute susceptibility of MR contrast agents and human blood., Magn.Reson.Med. (Band 24), Nr. 2, Seite 375-383.

[172] Whitsel, B. L.; Favorov, O.; Delemos, K. A.; Lee, C.; Tommerdahl, M.; Essick, G. K. und Nakhle, B. (1999): SI neuron response variability is stimulus tuned and NMDA receptor dependent., J.Neurophysiol. (Band 81), Nr. 6, Seite 2988-3006.

[173] Wustenberg, T.; Jordan, K.; Giesel, F. L. und Villringer, A. (2003): [Physiological and technical limitations of functional magnetic resonance imaging (fMRI)--consequences for clinical use]
Physiologische und technische Grenzen der funktionellen Magnetresonanztomographie und die damit verbundenen Konsequenzen fur die klinische Anwendung. DEP - 20030624, Radiologe (Band 43), Nr. 7, Seite 552-557.

[174] Yoo, S. S.; Talos, I. F.; Golby, A. J.; Black, P. M. und Panych, L. P. (2004): Evaluating requirements for spatial resolution of fMRI for neurosurgical planning, Hum.Brain Mapp. (Band 21), Nr. 1, Seite 34-43.

[175] Yousry, T. A.; Schmid, U. D.; Alkadhi, H.; Schmidt, D.; Peraud, A.; Buettner, A. und Winkler, P. (1997): Localization of the motor hand area to a knob on the precentral gyrus. A new landmark., Brain (Band 120), Nr. Pt 1, Seite 141-157.

[176] Zilles, K. und Rehkämper, G. (1998): Funktionelle Neuroanatomie, 3 th. Auflage , Springer, 1998, Berlin; Heidelberg; New York; Barcelona; Budapest; Hongkong; London; Mailand; Paris; Singapur; Tokio; ISBN: 3-540-64293-5.