

8 Literaturverzeichnis

- Abi-Dargham A., Gil R., Krystal J. et al. (1998) Increase striatal dopamine transmission in schizophrenia: confirmation in a second cohort. *Am. J. Psychiatry* 155, 761-767
- Aleman A., Kahn R.S. (2005) Strange feelings: Do amygdala abnormalities dysregulate the emotional brain in schizophrenia? *Progress in Neurobiology* 77, 283-298
- Altshuler L.L., Bartzokis G., Grieder, T. et al. (2000) An MRI study of temporal lobe structures in men with bipolar disorder or schizophrenia. *Biol. Psychiatry* 48, 147-162
- Andreasen N.C., Roy M.A., Flaum F. (1987) Positive and negative symptoms. In Hirsch S.R., Weinberger D.R. (1995) *Schizophrenia*. Blackwell Science, Oxford, 28,45
- Andreasen N.C., Flashman L., Flaum M. (1994) Regional brain abnormalities in schizophrenia-measured with magnetic resonance imaging. *JAMA* 272, 1763-1769
- Bechdorf A., Ruhrmann S., Lansen B. et al. (2004) Früherkennung und –intervention bei Personen mit erhöhtem Psychoserisiko. *psychoneuro* 30, 606-614
- Benes F.M., Sorensen I., Bird E.D. (1991) Deduced neuronal size in posterior hippocampus of schizophrenic patients. *Schizophr. Bull.* 17, 597-608
- Berger M. (2000) *Psychiatrie und Psychotherapie*. Urban & Fischer, München - Jena
- Bigler E.D., Tate D.F. (2001) Brain volume, intracranial volume and dementia. *Invest Radiol* 36, 539-546
- Bleuler E. (1911) *Dementia praecox oder Gruppe der Schizophrenien*. Deuticke, Leipzig-Wien
- Bleuler M. (1972) *Die schizophrenen Geistesstörungen im Lichte langjähriger Kranken- und Familiengeschichten*. Thieme, Stuttgart-New York
- Bogerts B., Ashtari, M., Degreef, G. et al. (1990) Reduced temporal limbic structure volumes on magnetic resonance images in first episode schizophrenia. *Psychiatry Res.* 35, 1-13

-
- Bonilha L., Kobayashi E., Cendes F., Min Li L. (2004) Protocol for volumetric segmentation of medial temporal structures using high-resolution 3-D magnetic resonance imaging. *Hum Brain Mapp.* 22(2), 145-154
- Breier A., Buchanan R.W., Elkashef A. et al. (1992) Brain morphology and schizophrenia. A magnetic resonance imaging study of, prefrontal cortex and caudate structures. *Arch. Gen. Psychiatry* 49, 921-926
- Brown A.S., Begg M.D., Gravenstein S. (2004) Serologic evidence of prenatal influenza in the etiology of schizophrenia. *Arch Gen Psychiatry* 61(8),774-80
- Byrne M., Hodges A., Grant E. et al. (1999) Neuropsychological assessment of young people at high genetic risk for developing schizophrenia compared with controls: preliminary findings of Edinburgh High Risk Study. *Psychological Medicine* 29, 1161-1173
- Chance S.A., Esiri M.M., Crow T.J. (2002) Amygdala volume in schizophrenia: post-mortem study and review of magnetic resonance imaging findings. *British J. of Psychiatry* 180, 331-338
- Chua S. E., Sharma T., Takei N. et al. (2000) A magnetic resonance imaging study of corpus callosum size in familial schizophrenic subjects, their relatives and normal controls. *Schizophrenia Res.* 41, 397-403
- Ciompi L., Müller C. (1976) Lebensweg und Alter der Schizophrenen. Eine katamnestiche Langzeitstudie bis ins Senium. Springer, Berlin-Heidelberg-New York
- Convit A., McHugh P., Wolf O.T., de Leon M.J., Bobinski M., De Santi S., Roche A., Tsui W. (1999) MRI volume of the amygdala : a reliable method allowing separation from the hippocampal formation. *Psychiatry Research: Neuroimaging Section* 90, 113-123
- Copolov D., Velakoulis D., McGorry P., et al. (2000) Neurobiological findings in early phase schizophrenia. *Brain Res.* 31, 157-165

- Corey-Bloom J, Jernigan T., Archibald S. et al. (1995) Quantitative magnetic resonance imaging of the brain in late-life schizophrenia. *Am. J. Psychiatry* 152, 447-449
- Crow T.J. (1980) The molecular pathology of schizophrenia. More than one disease process. *Br. Med. J.* 280, 66-68
- Csernansky J.G., Wang L., Jones D. et al. (2002) Hippocampal deformities in schizophrenia characterized by high dimensional brain mapping. *Am. J. Psychiatry* 159, 2000-2006
- Danos P., Baumann B., Kramer A. et al. (2003) Volumes of association thalamic nuclei in schizophrenia: a postmortem study. *Schizophrenia Res.* 60, 141-155
- Degreef G., Ashtari M., Bogerts B. et al. (1992) Volumes of ventricular system subdivisions measured from magnetic resonance imaging in first-episode schizophrenic patients. *Arch. Gen. Psychiatry* 49, 531-537
- DeLisi L.E., Sakuma M., Ge S. et al. (1998) Association of brain structural change with the heterogeneous course of schizophrenia from early childhood through five years subsequent to a first hospitalization. *Psychiatry Res.* 84, 75-88
- Dickey C.C., McCarley R.W., Voglmaier M.M. et al. (1999) Schizotypal personality disorder and MRI abnormalities of temporal lobe gray matter. *Biol. Psychiatry* 45, 1393-1402
- Duvernoy H. (2004) *The Human Hippocampus*. Springer, Berlin
- Eaton W.W. et al. (1995) Prodromes and precursors: epidemiologic data for primary prevention of disorders with slow onset. *Am J Psychiatry* 152, 967-972
- Erlenmeyer-Kimling L., Rock D., Roberts S.A. et al. (2000) Attention, memory and motor skills as childhood predictors of schizophrenia-related psychoses: The New York high-risk project. *J. of Psychiatry* 157, 1416-1422
- Exner C., Boucsein K., Degner D. et al. (2004) Impaired emotional learning and reduced amygdala size in schizophrenia: a 3-month follow-up. *Schizophrenia Research* 71, 493-503

-
- Falkai P., Maier W. (2006) Fortschritte in der neurobiologischen Erforschung der Schizophrenie. *Der Nervenarzt (Suppl 2) 77*, S65-S76
- Free S.L., Bergin P.S., Fish D.R. et al. (1995) Methods of normalisation of hippocampal volumes measured with MR. *Am. J. Neuroradiol.* 16, 637-643
- Frederikse M., Lu A., Aylward E. et al. (2000) Sex differences in inferior parietal lobule volume in schizophrenia. *Am. J. Psychiatry* 157, 422-427
- Gallinat J., Mulert C., Bajbouj M. et al. (2002): Frontal and temporal dysfunction of auditory stimulus processing in schizophrenia. *NeuroImage* 17: 110-127
- Geuze E., Vermetten E. Bremner J.D. (2004) MR-based in vivo hippocampal volumetrics: 1. Review of methodologies currently employed. *Molecular Psychiatry*, 1-13
- Goldstein J.M., Goodman J.M., Seidmann L.J. et al. (1999) Cortical abnormalities in schizophrenia identified by structural magnetic resonance imaging. *Arch. Gen. Psychiatry* 56, 537-547
- Gross G., Huber G., Klosterkötter J., Linz M. (1987) Bonner Skala für die Beurteilung von Basis-symptomen (BSABS, Bonn Scale for the Assessment of Basic Symptoms). Berlin, Heidelberg, New York: Springer
- Gur R.E., Turetsky B.I., Cowell P.E. et al. (2000) Temporolimbic volume reductions in schizophrenia. *Arch. Gen. Psychiatry* 57, 769-775
- Halberstadt A.L. (1995) The phencyclidine-glutamate model of schizophrenia. *Clin Neuropharmacol.* 18, 237-49
- Hecker E. (1871) Die Hebephrenie. *Archiv für pathologische Anatomie und Physiologie und für klinische Medizin*; 52: 394-429.
- Heckers S., Heinsen H., Heinsen Y.C. et al. (1990) Limbic structures and lateral ventricle in schizophrenia. a quantitative postmortem study. *Archives of General Psychiatry* 47, 1016-1022

- Hietala J., Syvalahti E., Vuorio K. et al. (1995) Presynaptic dopamine function in striatum of neuroleptic naive schizophrenic patients. *Lancet* 346, 1130-1131
- Hackert V.H., den Heijer T., Ouskerk M. et al. (2002) Hippocampal head size associated with verbal memory performance in nondemented elderly. *NeuroImage* 17, 1365-1372
- Häfner H., Riecher-Rössler A., Maurer K. et al. (1992) First onset and early symptomatology of schizophrenia. A chapter of epidemiological and neurubiological research into age and sex differences. *Europ. Arch. Psychiat. Clin. Neurosci.* 242, 109-118
- Häfner H., Maurer K., Löffler W. et al. (1995) Onset and early course of schizophrenia. In: Häfner H., Gattaz H. *Search for the courses of schizophrenia*, Berlin, Springer, 43-66
- Hirayasu Y., McCarley R.W., Salisbury D.F. et al. (2000) Planum temporale and Hesch gyrus volume reduction in schizophrenia: A magnetic resonance imaging study of first-episode patients. *Arch. Gen. Psychiatry* 57, 692-699
- Ho B. C., Alicata D., Mola C. et al. (2005) Hippocampus volume and treatment delays in first-episode schizophrenia. *Am. J. Psychiatry* 162, 1527-1529
- Honeycutt N. A., Smith P.D., Aylward E., Li Q., Chan M., Barta P.E., Pearlson G.D. (1998) Mesial temporal lobe measurements on magnetic resonance imaging scans. *Psychiatry Research: Neuroimaging Section* 83, 85-94
- Hooker C., Park S. (2002) Emotion processing and its relationship to social functioning in schizophrenia patients. *Psychiatry Res.* 112, 41-50
- Huber G., Gross G., Schüttler R. (1979) *Schizophrenie. Eine verlaufs- und sozialpsychiatrische Langzeitstudie.* Springer, Berlin-Heidelberg-New York
- Jablensky A. (1995) Schizophrenia: The epidemiological horizon. in Hirsch S.R., Weinberger D.R. *Schizophrenia.* Blackwell Science, Oxford, 206-252

-
- Job D.E., Whalley H.C., McConnell S. et al. (2003) voxel-based morphometry of grey matter densities in subjects at high risk of schizophrenia. *Schizophrenia Research* 64, 1-13
- Jacobsen L.K., Giedd J.N., Vaituzis A.C. (1996) Temporal lobe morphology in childhood-onset schizophrenia. *Am. J. Psychiatry* 153, 355-361
- Johnstone EC, Crow TJ, Frith CD, Husband J, Kreel L. (1976) Cerebral ventricular size and cognitive impairment in chronic schizophrenia. *Lancet*; 2: 924-926.
- Joyal C.C., Laakso M.P., Tiihonen J. et al. (2003) The amygdala and schizophrenia: a volumetric magnetic resonance imaging study in first-episode, neuroleptic-naive patients. *Biol. Psychiatry* 54, 1302-1304
- Juckel G., Schultze-Luther F., Rurmann S. (2004) Früherkennung beginnender schizophrener Erkrankungen. *psychoneuro* 30, 153-159
- Kahle W., Leonardt H., Platzer, W: (1991) Taschenatlas der Anatomie, Band 3: Nervensystem und Sinnesorgane. Thieme, Stuttgart
- Kalus P., Slotboom J., Gallinat J. et al. (2005) The amygdala in schizophrenia: a trimodal magnetic resonance imaging study. *Neuroscience Letters* 375, 151-156
- Kawasaki Y., Maeda Y., Urata K. et al. (1993) A quantitative magnetic resonance imaging study of patients with schizophrenia. *Eur. Arch. Psychiatry Clin. Neurosci.* 242, 268-272
- Keshavan M.S., Haas G.L., Kahn C.E. et al. (1998) Superior temporal gyrus and the course of early schizophrenia: progressive, static or reversible? *J. Psychiatry Res.* 32, 161-167
- Keshavan M.S., Dick E., Mankowski I. et al. (2002) Decrease left amygdala and hippocampal volumes in young offspring at risk for schizophrenia. *Schizophrenia Res.* 58, 173-183
- Keshavan M.S., Berger G. Zipurski R.B. et al. (2005) Neurobiology of early psychosis. *British J. of Psychiatry* 187, 8-18

- Kirch D.G. (1993) Infection and autoimmunity as etiologic factors in schizophrenia: A review and reappraisal. *Schizophr. Bull.* 19, 355-370
- Klosterkötter J., Hellmich M., Steinmeyer E.M. (2001) Diagnosing schizophrenia in the initial prodromal phase. *Arch. Gen. Psychiatry* 58, 158-164
- Kraepelin E. (1893) *Psychiatrie*. 4. Auflage Abel (Meixner), Leipzig
- Kwon J.S., McCarley R.W., Hirayasu Y. et al. (1999) Left planum temporale volume reduction in schizophrenia. *Arch. Gen. Psychiatry* 56, 142-148
- Kwon J.S., Shin Y-W., Kim C-W., Kim Y.I., Youn T., Han M.H. Chang K-H., Kim J-J. (2003) Similarity and disparity of obsessive-compulsive disorder and schizophrenia in MR volumetric abnormalities of the hippocampus-amygdala complex. *Journal of Neurology Neurosurgery and Psychiatry* 74, 962-964
- Laakso M.P., Tiihonen J., Syvälahti E. Et al. (2001) A morphometric MRI study of the hippocampus in first-episode, neuroleptic-naive schizophrenia. *Schizophrenia Res.* 50, 3-7
- Lawrie S.M., Whalley H.C., Kestelmann J.N. et al. (1999) Magnetic resonance imaging of brain in people at high risk of developing schizophrenia. *Lancet* 353, 30-33
- Lawrie S.M., Whalley H.C., Abukmeil S.S. et al. (2001) Brain structure genetic liability and psychotic symptoms in subjects at high risk of developing schizophrenia. *Biol. Psychiatry* 49, 811-823
- Lawrie S.M., Whalley H.C., Abukmeil S.S. et al. (2002) Temporal lobe volume changes in people at high risk of schizophrenia with psychotic symptoms. *British J. of Psychiatry* 181, 138-143
- Lawrie S.M., Whalley H.C., Job D.E. et al. (2003) Structural and functional abnormalities of the amygdala in schizophrenia. *Ann. N.Y. Acad. Scii.* 985, 445-460
- LeDoux J.E. (2000) Emotion circuits in the brain. *Annu. Rev. Neuroscience* 23, 155-184

-
- Lehrl S.(1978) Mehrfachwahl-Wortschatz-Intelligenztest MWT-B. Erlangen: Dr. med. Straube Verlag
- Levitt J.J., McCarley R.W., Nesto P.G. (1999) Quantitative volumetric MRI study of the cerebellum and vermis in schizophrenia: clinical and cognitive correlates. *Am. J. Psychiatry* 156, 1105-1107
- Maier W., Rietschel M., Lichtermann D. et al. (1999) Family and genetic studies on the relationship of schizophrenia to affective disorders. *Eur. Arch. Psychiatry clin. Neurosci.* 249, 57-61
- McGlashan T., Miller T.J., Woods S.W. (2001) A scale for the assessment of prodromal symptoms and states. In: Miller T., Mednick S.A., McGlashan T.H., Libiger J., Johannessen J.O., Early Intervention in Psychotic Disorders. Dordrecht, Netherlands: Kluwer Academic, 135-150.
- McGorry P.D., Singh B.S. (1995) Schizophrenia: risk and possibility of prevention. In: Handbook of studies on preventive psychiatry. Raphael B., Burrows G.D. (eds), New York, Elsevier, 491-514
- Mc Neil T.F. (1991) Review article: Obstetric complication in schizophrenic parents. *Schizophr. Research* 5, 89-101
- McNeil T.F., CantorGraae E., Weinberger D.R. (2000) Relationship of obstetric complications and differences in size of brain structures in monozygotic twin pairs discordant for schizophrenia. *Am. J. Psychiatry* 157, 203-212
- Miyahira Y., Yu J., Hiramatsu K., Shimazaki Y., Takeda Y. (2004) Brain volumetric MRI study in healthy elderly persons using statistical parametric mapping. *Seishin Shinkeigaku Zasshi* 106(2), 138-151
- Miller, E. (1984). Verbal fluency as a function of a measure of verbal intelligence and in relation to different types of pathology. *British Journal of Clinical Psychology*, 23, 53-57.

- Murray C.J.L., Lopez A.D. (1997) Global mortality, disability and the contribution of risk factors: Global burden of disease study. *Lancet* 349, 1436-1442
- Myin-Germeys I., Krabbendam L., Delespaul P.A. (2004) Sex differences in emotional reactivity to daily life stress in psychosis. *J. Clin. Psychiatrie* 65, 805-809
- Narr K. L., Thompson P. M., Sharma T. (2000) Mapping morphology of the corpus callosum in schizophrenia. *Cereb. Cortex* 10, 40-49
- Narr K. L., Thompson P. M., Szeszko P. et al. (2004) Regional specificity of hippocampal volume reductions in first- episode schizophrenia. *NeuroImage* 21,1563-1575
- Niemann K., Hammers A., Coenen V:A., et al. (2000) Evidence of a smaller left hippocampus and left temporal horn in both patients with first episode schizophrenia and normal control subjects. *Psychiatry Res.* 99, 93-110
- Niznikiewicz M., Donnino R., McCarley R. W. Et al. (2000) Abnormal angular gyrus asymmetry in schizophrenia. *Am. J. Psychiatry* 157, 428-437
- Norman R.N.G., Malla A.K. (1993) Stressful life events and schizophrenia. A review of the research. *Brit. J. Psychiatry* 162, 161-166
- Nuechterlein K.H., Dawson M.E., Gitlin M. (1992) Developmental Processes in Schizophrenic Disorders: longitudinal studies of vulnerability and stress. *Schizophr. Bull.* 18(3), 387-425
- O'Driscoll G.A., Florencio P.S., Gagnon D. et al. (2001) Amygdala-hippocampal volume and verbal memory in first-degree relatives of schizophrenic patients. *Psychiatry Res.* 107, 75-85
- Oldfield R.C. (1971) Assessment and Analysis of Handedness - Edinburgh Inventory. *Neuropsychologia* 9, 97-113.
- Pakkenberg B. (1990) Pronounced reduction of total neuron number in mediodorsal thalamic nucleus and nucleus accumbens in schizophrenics. *Arch. of General Psychiatry* 47, 1023-1028

-
- Pantel J., O'Leary D.S.; Cretsingher K., Bockholt H.J., Keefe H., Magnotta V.A., Andreasen N.C. (2000) A new method for the in vivo volumetric measurement of the human hippocampus with high neuroanatomical accuracy. *Hippocampus* 10, 752-758
- Pantelis C., Velakoulis D., McGorry P.D. et al. (2003) Neuroanatomical abnormalities before and after onset of psychosis: a cross-sectional and longitudinal MRI comparison. *The Lancet* 361, 281-288
- Panfield W., Roberts L. (1959) *Speech and Brain Mechanisms*. Princeton University Press, Princeton, NJ
- Paradiso S., Andreasen N.C., Crespino-Facorro et al. (2003) Emotions in unmedicated patients with schizophrenia during evaluation with positron emission tomography. *Am. J. Psychiatry* 160, 1775-1783
- Pegues M. P., Rogers L. J., Amend D. et al. (2003) Anterior hippocampal volume reduction in male patients with schizophrenia. *Schizophrenia Research* 60, 105-115
- Phillips L. J., Velakoulis D., Pantelis C. et al. (2002) Non-reduction in hippocampal volume is associated with higher risk of psychosis. *Schizophrenia Res.* 58, 145-158
- Pruessner J.C., Li L.M., Serles W. et al. (2000) Volumetry of hippocampus and amygdala with high resolution MRI and three-dimensional analysis software : minimizing the discrepancies between laboratories. *Cerebral Cortex* 10, 433-442
- Rajarethinam R., DeQuardo J.R., Miedler J. et al. (2001) Hippocampus and amygdala in schizophrenia: assessment of the relationship of neuroanatomy to psychopathology. *Psychiatry Res.* 108, 79-87
- Resch F. (2003) Schizophrenie. In Herpertz-Dahlmann, Resch, Schulte-Markwort, Warnke (2003) *Entwicklungspsychiatrie*. Stuttgart, Schattauer-Verlag
- Roth B.L., Hanizavareh S.M., Blum A.E. (2004) Serotonin receptors represent highly favorable molecular targets for cognitive enhancement in schizophrenia and other disorders. *Psychopharmacology* 174, 17-24.

- Roy P.D., Zipurski R.B., SaintCyr J.A., et al. (1998) Temporal horn enlargement is present in schizophrenia and bipolar disorder. *Biol. Psychiatry* 44, 418-422
- Sachdev P., Brodaty H., Cheang D. et al. (2005) Hippocampus and amygdalavolumes in elderly schizophrenic patients as assessed by magnetic resonance imaging. *Psychiatry Clin. Neuroscience* 54, 105-112
- Sanderson T. L., Best J.J., Doody G.A. et al. (1999) Neuroanatomy of comorbid schiziphrenia and learning disability: a controlled study. *Lancet* 354, 1867-1871
- Sanfilipo M., Lafargue T., Rusinek H. (2000) Volumetric measure of the frontal and temporal lobe regions in schizophrenia: relationship to negative symptoms. *Arch. Gen. Psychiatry* 57,471-480
- Schneider K. (1992) *Klinische Psychopathologie*. 14. Auflage, Thieme, Stuttgart-New York
- Schneider F., Gur R.C., Gur R.E. et al. (1995) Emotional processing in schizophrenia: neurobehavioral probes in relation to psychopathology. *Schizophr. Res.* 17, 67-75
- Schulze K., McDonald C. Frangou S. et al. (2003) Hippocampal Volume in Familial and Non-familial Schizophrenic Probands and Their unaffected Relatives. *Biol. Psychiatry* 53, 562-570
- Seidman L.J., Faraone S.V., Goldstein J.M. et al. (1999) Thalamic and amygdalahippocampal volume reduction in firstdegree relatives of patients with schizophrenia: an MRIbased morphometric analysis. *Biol. Psychiatry* 46, 941-954
- Seidman L.J., Goldstein J.M., Makris N.N. et al. (2000) Subcortical brain abnormalities in patients with schizophrenia: an MRI morphometric study. *Biol. Psychiatry* 47 (8s), 24s
- Seidmann L.J., Faraone S.V., Goldstein J.M. et al. (2002) Left Hippocampal Volume as a Vulnerability Indicator for Schizophrenia. *Arch. Gen. Psychiatry* 59, 839-849
- Shapleske J., Rossell S.L., Woodruff P.W. et al. (1999) The planum temporale: a systematic, quantitative review of its structural, functional and clinical significance. *Brain Res.* 29, 26-49

-
- Shaw P., Lawrence E.J., Radbourne C. et al. (2004) The impact of early and late damage to the human amygdala on “theory of mind” reasoning. *Brain* 127, 1535-1548
- Sheehan D.V., Lecrubier Y., Sheehan K.H., Amorim P., Janavs J., Weiller E., Hergueta T., Baker R., Dunbar G.C. (1998) The Mini-International Neuropsychiatric Interview (MINI): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry*. 59, 22-33.
- Shenton M.E., Kikinis R., Jolesz F. A. et al. (1992) Abnormalities of the left temporal lobe and thought disorder in schizophrenia. A quantitative magnetic resonance imaging study. *N. Engl. J. Med.* 327, 604-612
- Shenton M.E., Wible C.G., McCarley R.W, et al. (1997) A review of magnetic resonance imaging studies of brain abnormalities in schizophrenia. In Krishnan K., Doraiswamy P.(Hrg) *Brain Imaging in Clinical Psychiatrie*. Marcel Decker, New York, 297-380
- Shenton M.E., Dickey C.C., Frumin M., McCarley R.W. (2001) A review of MRI findings in schizophrenia. *Schizophrenia. Research* 49, 1-52
- Shenton M.E., Gerig G., McCarley R.W. et al. (2002) Amygdala-hippocampal shape differences in schizophrenia: the application of 3D shape models to volumetric MR data. *Psychiatry Res.* 115, 15-35
- Sim K., DeWitt I., Ditman T. (2006) Hippocampal and Parahippocampal volumes in schizophrenia: a structural MRI study. *Schizophr. Bull.* 32, 332-340
- Smith G.S., Schloesser R., Brodie J.D. et al. (1998) Glutamate modulation of dopamine measured in vivo with positron emission tomography (PET) and 11C-raclopride in normal human subjects. *Neuropsychopharmacology* 18, 18-25
- Smith G.N., Lang D.J., Kopala L.C. et al. (2003) Developmental abnormalities of the hippocampus in first-episode schizophrenia. *Biol. Psychiatry* 53, 555-561

- Smith D.M., Mizumori S.J. (2006) Hippocampal place cells, context, and episodic memory. *Hippocampus* 16, 716-729
- Staal W.G., Hulshoff Pol H.E., Schnack H.G. et al. (2000) Structural brain abnormalities in patients with schizophrenia and their healthy siblings. *A. J. of Psychiatry* 157, 416-421
- Stefanis N., Frangou S., Yakeley J. et al. (1999) Hippocampal volume reduction in schizophrenia: effects of genetic risk and pregnancy and birth complications. *Biol. Psychiatry* 46, 697-702
- Suddath R.L., Casanova M.F., Goldberg T.E. (1989) Temporal lobe pathology in schizophrenia: a quantitative magnetic resonance imaging study. *Am. J. Psychiatry* 146, 464-472
- Sullivan E.V., Mathalon D.H., Lim K.O. (1998) Patterns of regional cortical dysmorphology distinguishing schizophrenia and chronic alcoholism. *Biol. Psychiatry* 43, 118- 131
- Sumich A., Chitnis X.A., Fannon D.G. et al. (2002) Temporal lobe abnormalities in first-episode psychosis. *Am J. Psychiatry* 159, 1232-1234
- Supprian T., Ulmar G., Bauer M. et al. (2000) Cerebellar vermis area in schizophrenic patients - a post-mortem study. *Schizophrenia Res.* 42, 19-28
- Suzuki M., Zhou S.Y., Takahashi T. et al. (2005) Differential contribution of prefrontal and temporo- limbic pathology to mechanism of psychosis. *Brain* 128, 2109-2122
- Szendi I., Kiss M., Racsmany M. et al. (2006) Correlation between clinical symptoms, working memory functions and structural brain abnormalities in men with schizophrenia. *Psychiatry Research* 147(1), 47-55
- Szeszko P. R., Goldberg E., Gunduz-Bruce H. et al. (2003) Smaller anterior hippocampal formation volume in antipsychotic-naive patients with first-episode schizophrenia. *Am. J. Psychiatry* 160, 2190-2197

-
- Tanskanen P., Veijola J.M., Piippo U.K. et al. (2005) Hippocampus and amygdala volumes in schizophrenia and other psychoses in Northern Finland 1966 birth cohort. *Schizophrenia Research* 75, 283-294
- Tepst R., Wang L., Miller M. I. et al. (2003) Hippocampal deformities in unaffected siblings of schizophrenia subjects. *Biol. Psychiatry* 54, 1234-1240
- Theberge J., Bartha R., Drost D.J. et al. (2002) Glutamat and glutamine measured with 4.0 T proton MRS in never-treated patients with schizophrenia and healthy volunteers. *Am. J. Psychiatry* 159, 1944-1946
- Theberge J., Al-Semaan Y., Williamson P.C. et al. (2003) Glutamate and glutamine in the anterior cingulate and thalamus of medicated patients with chronic schizophrenia and healthy comparison subjects measured with 4.0 T proton MRS. *Am. J. Psychiatry* 160, 2231-2233
- Van Erp T.G.M., Saleh P.A., Huttunen M. (2004) Hippocampal volumes in schizophrenic twins. *Arch. Gen. Psychiatry* 61, 346-353
- Velakoulis D., Pantelis C., McGorry P.D. et al. (1999) Hippocampal volume in first episode psychoses and chronic schizophrenia: a high resolution magnetic resonance imaging study. *Arch. Gen. Psychiatry* 56, 133-141
- Velakoulis D., Stuart G.W., Wood S.J. et al. (2001) Selective bilaterale hippocampal volume loss in chronic schizophrenia. *Biol. Psychiatrie* 50, 531-539
- Velakoulis D., Wood S.J., Wong M.T. et al. (2006) Hippocampal and amygdala volumes according to psychosis stage and diagnosis: a magnetic resonance imaging study of chronic schizophrenia, first-episode psychosis and ultra-high-risk individuals. *Arch. Gen. Psychiatry* 63, 139-139
- Wassink T.H., Andreasen N.C., Nopoulos P. Et al. (1999) Cerebellar morphology as a predictor of symptom and psychosozial outcome in schizophrenia. *Biol. Psychiatry* 45, 41-48

- Weltgesundheitsorganisation (2000) Internationale Klassifikation psychischer Störungen. 4. Auflage, Bern, Göttingen, Toronto, Seattle: Verlag Hans Huber
- Weinberger D.R. (1999) Cell biology of the hippocampal formation in schizophrenia. *Biol. Psychiatry* 45, 395-402
- Whitworth A.B., Kemmler G., Honeder M. et al. (2005) Longitudinal volumetric MRI study in first- and multiple-episode schizophrenia patients. *Psychiatry Res.* 140, 225-237
- Wible C.G., Shenton M.E., Hokama H. Et al. (1995) Prefrontal cortex and schizophrenia. A quantitative magnetic resonance imaging study. *Arch. Gen. Psychiatry* 52, 279-288
- Winston J.S., Strange B.A., O'Doherty J. et al. (2002) Automatic and intentional brain response during evaluation of trustworthiness of faces. *Nat. Neuroscience* 5, 277-283
- Wittchen H.U., Semler G. (1991) Composite International Diagnostic Interview - CIDI. Interviewheft und Manual. Weinheim: Beltz.
- Wittchen H.U., Zaudig M., Frydreich T.(1997) Strukturiertes klinisches Interview für DSM- IV (Skid-I und Skid-II). Achse I: Psychische Störungen/Achse II: Persönlichkeitstörungen. Göttingen, Hogrefe
- Wood S.J., Velakoulis D., Smith D.J. et al. (2001) Alongitudnal study of hippocampal volume in first episode psychosis and chronic schizophrenia. *Schizophrenia Resaerch* 52, 37-46
- Wood S.J., Pantelis C., Proffitt, T. et al. (2003) Spatial working memory ability is a marker of risk for psychosis. *Psychological Medicine* 33, 1239-1247
- Wood S.J., Yücel M., Velakoulis D. et al (2005) Hippocampal and anterior cingulate morphology in subjects at ultra-high-risk for psychosis: the role of famoily history of psychotic illness. *Schizophrenia Research* 75, 295-301
- Yung A.R., McGorry P.D. (1996)The prodromal phase of first-episode psychosis, past and current conceptualizations. *Schizophr. Bull.* 22, 353-370

Yung A.R., Yung A.R., McGorry P.D. et al. (1998) A step toward indicated prevention of schizophrenia. *Br. J. Psychiatry* 172, 14-20

Yung A.R., Yung A.R., Yuen H.P. et al. (2003) 12-month follow up of a high-risk ("prodromal") group. *Schizophr. Research* 60, 21-32

Zuzuki M., Hagino H., Nohara S., Zhou S., Kawasaki Y. Takahashi T., Matsui M., Seto H., Ono T., Kurachi M. (2004) Male-specific volume expansion of the human hippocampus during adolescence. *Cerebral Cortex*