

## 6. Literatur

Abbott BD, Schmid JE, Pitt JA, Buckalew AR, Wood CR, Held GA, Diliberto JJ. Adverse reproductive outcomes in the transgenic Ah receptor-deficient mouse. *Toxicol Appl Pharmacol.* 1999; 155: 62-70.

Abraham K, Knoll A, Ende M. Intake, fecal excretion and body burden of polychlorinated dibenzo-p-dioxins and dibenzofurans in breast-fed and formula-fed infants. *Pediatric Res.* 1996; 40: 671-679.

Ahlborg UG, Brouwer A, Fingerhut MA, Jacobson JL, Jacobson SW, Kennedy SW, Kettrup AA, Koeman JH, Poiger H, Rappe C, et al. Impact of polychlorinated dibenzo-p-dioxins, dibenzofurans, and biphenyls on human environmental health, with special emphasis on application of the toxic equivalency factor concept. *Eur J Pharmacol.* 1992; 228: 179-199.

Alfred LJ, Wojdani A. Effects of methylcholanthrene and benzanthracene on blastogenesis and aryl hydrocarbon hydroxylase induction in splenic lymphocytes from three inbred strains of mice. *Int J Immunopharmacol.* 1983; 5: 123-129.

Allen JR, Barsotti DA, Van Miller JP, Abrahamson LJ, Lalich JJ. Morphological changes in monkeys consuming a diet containing low levels of 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Food Cosmet Toxicol.* 1977; 15: 401-410.

Anderson D. Findings of US EPA/Paper industrycooperative dioxin study. The 104 mill. Study, in *Dioxin 90 - EPRI Seminar* (O. Hutzinger, H. Fiedler, Hrsg.) Vol. 3, Ecoinforma Press, Bayreuth. 1990; 207-210.

Annunziato F, Romagnani P, Cosmi L, Beltrame C, Steiner BH, Lazzeri E, Raport CJ, Galli G, Manetti R, Mavilia C, Vanini V, Chantry D, Maggi E, Romagnani S. Macrophage-derived chemokine and EBI1-ligand chemokine attract human thymocytes in different stage of development and are produced by distinct subsets

of medullary epithelial cells: possible implications for negative selection. *J Immunol.* 2000; 165: 238-246.

Baggiolini M. Chemokines and leukocyte traffic. *Nature.* 1999; 392: 565-568.

Bannister R, Davis D, Zacharewski T, Tizard I, Safe S. Aroclor 1254 as a 2,3,7,8-tetrachlorodibenzo-p-dioxin antagonist: effects on enzyme induction and immunotoxicity. *Toxicology.* 1987; 46: 29-42.

Barnes D, Alford-Stevens A, Birnbaum L, Kutz FW, Wood W, Patton D. Toxicity equivalency factors for PCBs ?, *Qual Assur.* 1991; 1: 70-81.

Birnbaum LS. The role of structure in the disposition of halogenated aromatic xenobiotics. *Environ Health Perspect.* 1985; 61: 11-20.

Birnbaum LS, McDonald MM, Blair PC, Clark AM, Harris MW. Differential toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in C57BL/6J mice congenic at the Ah Locus. *Fundam Appl Toxicol.* 1990; 15: 186-200.

Blaylock BL, Holladay SD, Comment CE, Heindel JJ, Luster MI. Exposure to tetrachlorodibenzo-p-dioxin (TCDD) alters fetal thymocyte maturation. *Toxicol Appl Pharmacol.* 1992; 112: 207-213.

Brewster DW, Matsumura F, Akera T. Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin on guinea pig heart muscle. *Toxicol Appl Pharmacol.* 1987; 89: 408-417.

Brewster DW, Bombick DW, Matsumura F. Rabbit serum hypertriglyceridemia after administration of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *J Toxicol Environ Health.* 1988; 25: 495-507.

- Brewster DW, Banks YB, Clark AM. Comparative dermal absorption of 2,3,7,8-tetrachlorodibenzo-p-dioxin and three polychlorinated dibenzofurans. *Toxicol Appl Pharmacol.* 1989; 97: 156-166.
- Burbach KM, Poland A, Bradfield CA. Cloning of the Ah-receptor cDNA reveals a novel ligand-activated transcription factor. *Proc Natl Acad Sci USA.* 1992; 89: 8185-8189.
- Burleson GR, Lebrec H, Yang YG, Ibanes JD, Pennington KN, Birnbaum LS. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on influenza virus host resistance in mice. *Fundam Appl Toxicol.* 1996 ; 29: 40-47.
- Canga L, Levi R, Rifkind AB. Heart as a target organ in 2,3,7,8-tetrachlorodibenzo-p-dioxin toxicity: decreased beta-adrenergic responsiveness and evidence of increased intracellular calcium. *Proc Natl Acad Sci U S A.* 1988; 85: 905-909.
- Carlos TM, Harlan JM. Leukocyte-endothelial adhesion molecules. *Blood.* 1994; 84: 2068-2101.
- Chantry D, Romagnani P, Raport CJ, Wood CL, Epp A, Romagnani S, Gray PW. Macrophage-derived chemokine is localized to thymic medullary epithelial cells and is a chemoattractant for CD3(+), CD4(+), CD8(low) thymocytes. *Blood.* 1999; 94: 1890-1898.
- Crawford RB, Holsapple MP, Kaminski NE. Leukocyte activation induces aryl hydrocarbon receptor up-regulation, DNA binding, and increased Cyp1a1 expression in the absence of exogenous ligand. *Mol Pharmacol.* 1997; 52: 921-927.
- Cuthill S, Wilhelmsson A, Mason GGF, Gillner M, Poellinger L, Gustafsson JA. The dioxin receptor: a comparison with the glucocorticoid receptor. *J Steroid Biochem.* 1988; 30: 277-280.

- Dahl P, Lindstrom G, Wiberg K, Rappe C. Absorption of polychlorinated biphenyls, dibenzo-p-dioxins and dibenzofurans by breast-fed infants. *Chemosphere*. 1995; 30: 2297-2306.
- Davis D, Safe S. Immunosuppressive activities of polychlorinated dibenzofuran congeners: quantitative structure-activity relationships and interactive effects. *Toxicol Appl Pharmacol*. 1988; 94: 141-149.
- Davis D, Safe S. Dose-response immunotoxicities of commercial polychlorinated biphenyls (PCBs) and their interaction with 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Toxicol Lett*. 1989; 48: 35-43.
- Davis D, Safe S. Immunosuppressive activities of polychlorinated biphenyls in C57BL/6N mice: structure-activity relationships as Ah receptor agonists and partial antagonists. *Toxicology*. 1990; 63: 97-111.
- De Krey GK, Baecher-Steppan L, Deyo JA, Smith B, Kerkvliet NI. Polychlorinated biphenyl-induced immune suppression: castration, but not adrenalectomy or RU 38486 treatment, partially restores the suppressed cytotoxic T lymphocyte response to alloantigen. *J Pharmacol Exp Ther*. 1993; 267: 308-315.
- De Krey GK, Baecher-Steppan L, Fowles JR, Kerkvliet NI. Polychlorinated biphenyl-induced suppression of cytotoxic T lymphocyte activity: role of prostaglandin-E2. *Toxicol Lett*. 1994; 74: 211-220.
- De Krey GK, Kerkvliet NI. Suppression of cytotoxic T lymphocyte activity by 2,3,7,8-tetrachlorodibenzo-p-dioxin occurs in vivo, but not in vitro, and is independent of corticosterone elevation. *Toxicology*. 1995; 97: 105-112.

- DeVito MJ, Birnbaum LS, Farland WH, Gasiewicz TA. Comparisons of estimated human body burdens of dioxinlike chemicals and TCDD body burdens in experimentally exposed animals. *Environ Health Perspect.* 1995; 103: 820-831.
- DeVito MJ, Menache MG, Diliberto JJ, Ross DG, Birnbaum LS. Dose-response relationships for induction of CYP1A1 and CYP1A2 enzyme activity in liver, lung, and skin in female mice following subchronic exposure to polychlorinated biphenyls. *Toxicol Appl Pharmacol.* 2000; 167: 157-172.
- Diliberto JJ, Kedderis LB, Jackson JA, Birnbaum LS. Effects of dose and routes of exposure on the disposition of 2,3,7,8-<sup>3</sup>H-tetrabromodibenzo-p-dioxin (TBDD) in the rat. *Toxicol Appl Pharmacol.* 1993a; 120: 315-326.
- Diliberto JJ, Akubue PI, Jackson JA. Dose-dependent tissue distribution of 2,3,7,8-TCDD in mice. *Toxicologist.* 1993b; 13: 195.
- Eadon G, Kaminsky L, Silkworth J, Aldous K, Hilker D, O'Keefe P, Smith R, Gierthy J, Hawley J, Kim N, et al. Calculation of 2,3,7,8-TCDD equivalent concentrations of complex environmental contaminant mixtures. *Environ Health Perspect.* 1986; 70: 221-227.
- Fassler R, Georges-Labouesse E, Hirsch E. Genetic analyses of integrin function in mice. *Curr Opin Cell Biol.* 1996; 8: 641-646.
- Fine JS, Gasiewicz TA, Silverstone AE. Lymphocyte stem cell alterations following perinatal exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Mol Pharmacol.* 1989; 35: 18-25.
- Fine JS, Silverstone AE, Gasiewicz TA. Impairment of prothymocyte activity by 2,3,7,8-tetrachlorodibenzo-p-dioxin. *J Immunol.* 1990; 144: 1169-1176.

- Flesch-Janys D, Becher H, Gurn P, Jung D, Konietzko J, Manz A, Papke O. Elimination of polychlorinated dibenzo-p-dioxins and dibenzofurans in occupationally exposed persons. *J Toxicol Environ Health*. 1996; 47: 363-378.
- Gahmberg CG, Tolvanen M, Kotovuori P. Leukocyte adhesion-structure and function of human leukocyte beta2-integrins and their cellular ligands. *Eur J Biochem*. 1997; 245: 215-232.
- Gasiewicz TA, Geiger LE, Rucci G, Neal RA. Distribution, excretion, and metabolism of 2,3,7,8-tetrachlorodibenzo-p-dioxin in C57BL/6J, DBA/2J, and B6D2F1/J mice. *Drug Metab Dispos*. 1983; 11: 397-403.
- Gillner M, Bergman J, Cambillau C, Alexandersson M, Fernstrom B, Gustafsson JA. Interactions of indolo[3,2-b]carbazoles and related polycyclic aromatic hydrocarbons with specific binding sites for 2,3,7,8-tetrachlorodibenzo-p-dioxin in rat liver. *Mol Pharmacol*. 1993; 44: 336-345.
- Greenlee WF, Dold KM, Irons RD, Osborne R. Evidence for direct action of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on thymic epithelium. *Toxicol Appl Pharmacol*. 1985; 79: 112-120.
- Hahn ME, Karchner SI, Shapiro MA, Perera SA. Molecular evolution of two vertebrate aryl hydrocarbon (dioxin) receptors (AHR1 and AHR2) and the PAS family. *Proc Natl Acad Sci U S A*. 1997; 94: 13743-13748.
- Hayashi S, Okabe-Kado J, Honma Y, Kawajiri K. Expression of Ah receptor (TCDD receptor) during human monocytic differentiation. *Carcinogenesis*. 1995; 16: 1403-1409.
- Haynes BF. The human thymic microenvironment. *Adv Immunol*. 1984; 36: 87-142.

- Haynes BF, Scearce RM, Lobach DF, Hensley LL. Phenotypic characterization and ontogeny of mesodermal-derived and endocrine epithelial components of the human thymic microenvironment. *J Exp Med.* 1984; 159: 1149-1168.
- Henck JM, New MA, Kociba RJ, Rao KS. 2,3,7,8-tetrachlorodibenzo-p-dioxin: acute oral toxicity in hamsters. *Toxicol Appl Pharmacol.* 1981; 59: 405-407.
- Hemler ME. VLA proteins in the integrin family: structures, functions, and their role on leukocytes. *Annu Rev Immunol.* 1990; 8: 365-400.
- Henry EC, Kende AS, Rucci G, Totleben MJ, Willey JJ, Dertinger SD, Pollenz RS, Jones JP, Gasiewicz TA. Flavone antagonists bind competitively with 2,3,7, 8-tetrachlorodibenzo-p-dioxin (TCDD) to the aryl hydrocarbon receptor but inhibit nuclear uptake and transformation. *Mol Pharmacol.* 1999; 55: 716-725.
- Holladay SD, Lindstrom P, Blaylock BL, Comment CE, Germolec DR, Heindell JJ, Luster MI. Perinatal thymocyte antigen expression and postnatal immune development altered by gestational exposure to tetrachlorodibenzo-p-dioxin (TCDD). *Teratology.* 1991; 44: 385-393.
- Holsapple MP, Morris DL, Wood SC, Snyder NK. 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced changes in immunocompetence: possible mechanisms. *Annu Rev Pharmacol Toxicol.* 1991a; 31: 73-100.
- Holsapple MP, Snyder NK, Wood SC, Morris DL. A review of 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced changes in immunocompetence: 1991 update. *Toxicology.* 1991b; 69: 219-255.
- Holsapple MP. Immunotoxicity of halogenated aromatic hydrocarbons. *Experimental immunotoxicology.* CRC press, Boca Raton. 1995; 265-305.

Hornung MW, Zabel EW, Peterson RE. Toxic equivalency factors of polybrominated dibenzo-p-dioxin, dibenzofuran, biphenyl, and polyhalogenated diphenyl ether congeners based on rainbow trout early life stage mortality. *Toxicol Appl Pharmacol.* 1996; 140: 227-234.

Iida T, Hirakawa H, Matsueda T, Nagayama J, Nagata T. Polychlorinated dibenzo-p-dioxins and related compounds: correlations of levels in human tissues and in blood. *Chemosphere.* 1999; 38: 2767-2774.

Ikuta T, Eguchi H, Tachibana T, Yoneda Y, Kawajiri K. Nuclear localization and export signals of the human aryl hydrocarbon receptor. *J Biol Chem.* 1998; 273: 2895-2904.

Ivanov V and Sandell E. Characterization of polychlorinated biphenyl isomers in sovol and trichlordiphenyl formulations by high-resolution gas chromatography with electron capture detection and high-resolution gas chromatography - mass spectrometry techniques. *Environ Sci Technol.* 26; 1992; 2012-2017.

Kamath AB, Xu H, Nagarkatti PS, Nagarkatti M. Evidence for the induction of apoptosis in thymocytes by 2,3,7,8-tetrachlorodibenzo-p-dioxin in vivo. *Toxicol Appl Pharmacol.* 1997; 142: 367-377.

Kamath AB, Nagarkatti PS, Nagarkatti M. Characterization of phenotypic alterations induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin on thymocytes in vivo and its effect on apoptosis. *Toxicol Appl Pharmacol.* 1998; 150: 117-124.

Kamath AB, Camacho I, Nagarkatti PS, Nagarkatti M. Role of Fas-Fas ligand interactions in 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD)-induced immunotoxicity: increased resistance of thymocytes from Fas-deficient (*lpr*) and Fas ligand-defective (*gld*) mice to TCDD-induced toxicity. *Toxicol Appl Pharmacol.* 1999; 160: 141-155.



- Kelling CK, Menahan LA, Peterson RE. Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin treatment on mechanical function of the rat heart. *Toxicol Appl Pharmacol.* 1987; 91: 497-501.
- Kerkvliet NI. Halogenated aromatic hydrocarbons (HAH) as immunotoxicants. *Prog Clin Biol Res.* 1984; 161: 369-387.
- Kerkvliet NI, Brauner JA, Matlock JP. Humoral immunotoxicity of polychlorinated diphenyl ethers, phenoxyphenols, dioxins and furans present as contaminants of technical grade pentachlorophenol. *Toxicology.* 1985; 36: 307-324.
- Kerkvliet NI, Brauner JA. Flow cytometric analysis of lymphocyte subpopulations in the spleen and thymus of mice exposed to an acute immunosuppressive dose of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *Environ Res.* 1990; 52: 146-154.
- Kerkvliet NI, Steppan LB, Brauner JA, Deyo JA, Henderson MC, Tomar RS, Buhler DR. Influence of the Ah locus on the humoral immunotoxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin: evidence for Ah-receptor-dependent and Ah-receptor-independent mechanisms of immunosuppression. *Toxicol Appl Pharmacol.* 1990a; 105: 26-36.
- Kerkvliet NI, Burleson GR. Immunotoxicity of TCDD and related halogenated aromatic hydrocarbons. *Immunotoxicology and immunopharmacology.* Second Ed., New York, 1994; Raven press: 97-121.
- Kerkvliet NI. Recent advances in understanding the mechanisms of TCDD immunotoxicity. *Int Immunopharmacol.* 2002; 2: 277-291.
- Kimmig J und Schulz KH. Chlorierte aromatische zyklische Äther als Ursache der sogenannten Chlorakne. *Dermatologica.* 1957; 115: 540-546.

- Kremer J, Lai ZW, Esser C. Evidence for the promotion of positive selection of thymocytes by Ah receptor agonist 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Eur J Pharmacol.* 1995; 293: 413-427.
- Lang DS, Becker S, Clark GC, Devlin RB, Koren HS. Lack of direct immunosuppressive effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on human peripheral blood lymphocyte subsets in vitro. *Arch Toxicol.* 1994; 68: 296-302.
- Lang DS, Becker S, Devlin RB, Koren HS. 2,3,7,8-TCDD induces cytochrome P450 enzyme activity but not proliferation or phenotypical changes in human peripheral blood lymphocytes. *Toxicol Lett.* 1996; 88: 317-325.
- Larson RS, Springer TA. Structure and function of leukocyte integrins. *Immunol Rev.* 1990; 114: 181-217.
- Lawrence BP, Leid M, Kerkvliet NI. Distribution and behavior of the Ah receptor in murine T lymphocytes. *Toxicol Appl Pharmacol.* 1996; 138: 275-284.
- Lawrence BP, Kerkvliet NI. Role of altered arachidonic acid metabolism in 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced immune suppression in C57Bl/6 mice. *Toxicol Sci.* 1998; 42: 13-22.
- Lawrence BP, Meyer M, Reed DJ, Kerkvliet NI. Role of glutathione and reactive oxygen intermediates in 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced immune suppression in C57Bl/6 mice. *Toxicol Sci.* 1999; 52: 50-60.
- Leece B, Denomme MA, Towner R, Li SM, Safe S. Polychlorinated biphenyls: correlation between in vivo and in vitro quantitative structure-activity relationships (QSARs). *J Toxicol Environ Health.* 1985; 16: 379-388.

Li H, Dong L, Whitlock JP Jr. Transcriptional activation function of the mouse Ah receptor nuclear translocator. *J Biol Chem.* 1994; 269: 28098-28105.

Lobach DF, Searce RM, Haynes BF. The human thymic microenvironment. Phenotypic characterization of Hassall's bodies with the use of monoclonal antibodies. *J Immunol.* 1985; 134: 250-257.

Lu YF, Santostefano M, Cunningham BD, Threadgill MD, Safe S. Substituted flavones as aryl hydrocarbon (Ah) receptor agonists and antagonists. *Biochem Pharmacol.* 1996; 51: 1077-1087.

Lundberg K, Gronvik KO, Goldschmidt TJ, Klareskog L, Dencker L. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) alters intrathymic T-cell development in mice. *Chem Biol Interact.* 1990; 74: 179-193.

Luster MI, Germolec DR, Clark G, Wiegand G, Rosenthal GJ. Selective effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin and corticosteroid on in vitro lymphocyte maturation. *J Immunol.* 1988; 140: 928-935.

Luster AD. Chemokines--chemotactic cytokines that mediate inflammation. *N Engl J Med.* 1998; 338: 436-445.

Marcus RS, Holsapple MP, Kaminski NE. Lipopolysaccharide activation of murine splenocytes and splenic B cells increased the expression of aryl hydrocarbon receptor and aryl hydrocarbon receptor nuclear translocator. *J Pharmacol Exp Ther.* 1998; 287: 1113-1118.

Masten SA, Shiverick KT. Characterization of the aryl hydrocarbon receptor complex in human B lymphocytes: evidence for a distinct nuclear DNA-binding form. *Arch Biochem Biophys.* 1996; 336: 297-308.

- Max SR, Silbergeld EK. Skeletal muscle glucocorticoid receptor and glutamine synthetase activity in the wasting syndrome in rats treated with 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Toxicol Appl Pharmacol.* 1987; 87: 523-527.
- McConnell EE, Moore JA, Dalgard DW. Toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in rhesus monkeys (*Macaca mulatta*) following a single oral dose. *Toxicol Appl Pharmacol.* 1978; 43: 175-187.
- McLachlan MS. Digestive tract absorption of polychlorinated dibenzo-p-dioxins, dibenzofurans and biphenyls in a nursing infant. *Toxicol Appl Pharmacol.* 1993; 123: 68-72.
- Miniero R, De Felip E, Ferri F, di Domenico A. An overview of TCDD half-life in mammals and its correlation to body weight. *Chemosphere.* 2001; 43: 839-844.
- Mocarelli P. Seveso: a teaching story. *Chemosphere.* 2001; 43: 391-402.
- Morris DL, Holsapple MP. Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on humoral immunity: II. B cell activation. *Immunopharmacology.* 1991; 21: 171-181.
- Nagarkatti PS, Sweeney GD, Gauldie J, Clark DA. Sensitivity to suppression of cytotoxic T cell generation by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) is dependent on the Ah genotype of the murine host. *Toxicol Appl Pharmacol.* 1984; 72: 169-176.
- NATO/CCMS, Scientific basis for the development of the International Toxicity Equivalency Factor (I-TEF) method of risk assessment for complex mixtures of dioxins and related compounds. 1989; Report No. 178.
- Neal RA, Olson JR, Gasiewicz TA, Geiger LE. The toxicokinetics of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin in mammalian systems. *Drug Metab Rev.* 1982; 13: 355-385.

Nessel CS, Amoruso MA, Umbreit TH, Gallo MA. Hepatic aryl hydrocarbon hydroxylase and cytochrome P450 induction following the transpulmonary absorption of TCDD from intratracheally instilled particles. *Fundam Appl Toxicol.* 1990; 15: 500-509.

Nessel CS, Amoruso MA, Umbreit TH, Meeker RJ, Gallo MA. Pulmonary bioavailability and fine particle enrichment of 2,3,7,8-tetrachlorodibenzo-p-dioxin in respirable soil particles. *Fundam Appl Toxicol.* 1992; 19: 279-285.

Neubert R, Jacob-Müller U, Stahlmann R, Helge H, Neubert D. Polyhalogenated dibenzo-p-dioxins and dibenzofurans and the immune system. 1. Effects on peripheral lymphocyte subpopulations of a non-human primate (*Callithrix jacchus*) after treatment with 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *Arch Toxicol.* 1990; 64: 345-359.

Neubert R, Stahlmann R, Korte M, van Loveren H, Vos JG, Golor G, Webb JR, Helge H, Neubert D. Effects of small doses of dioxins on the immune system of marmosets and rats. *Ann N Y Acad Sci.* 1993; 685: 662-686.

Norback DH, Allen JR. Biological responses of the nonhuman primate, chicken, and rat to chlorinated dibenzo-p-dioxin ingestion. *Environ Health Perspect.* 1973; 5: 233-240.

Norris EH. The morphogenesis and histogenesis of the thymus gland in man: in which the origin of the Hassal's corpuscles of the human thymus is discovered. *Contrib Embryol Carnegie Inst.* 1938; 27: 193.

Olson JR, Holscher MA, Neal RA. Toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in the golden Syrian hamster. *Toxicol Appl Pharmacol.* 1980; 55: 67-78.

- Pazdernik TL, Rozman KK. Effect of thyroidectomy and thyroxine on 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced immunotoxicity. *Life Sci.* 1985; 36: 695-703.
- Peterson RE, Seefeld MD, Christian BJ. The wasting syndrome in 2,3,7,8-tetrachlorodibenzo-p-dioxin toxicity: basic features and their interpretation. In: *Banbury Report: biological mechanisms of dioxin action.* Poland A, Kimbrough R (Hrsg.). 1984; Vol. 18: 291-308.
- Poiger H, Schlatter C. Pharmacokinetics of 2,3,7,8-TCDD in man. *Chemosphere.* 1986; 15: 1489-1494.
- Poland A, Knutson JC. 2,3,7,8-Tetrachlorodibenzo-p-dioxin and related aromatic hydrocarbons: examination of the mechanism of toxicity. *Annu Rev Pharmacol Toxicol.* 1982; 22: 517-554.
- Poland A, Glover E, Taylor BA. The murine Ah locus: a new allele and mapping to chromosome 12. *Mol Pharmacol.* 1987; 32: 471-478.
- Poland A, Glover E. Characterization and strain distribution pattern of the murine Ah receptor specified by the Ahd and Ahb-3 alleles. *Mol Pharmacol.* 1990; 38: 306-312.
- Pollenz RS, Sattler CA, Poland A. The aryl hydrocarbon receptor and aryl hydrocarbon receptor nuclear translocator protein show distinct subcellular localizations in Hepa 1c1c7 cells by immunofluorescence microscopy. *Mol Pharmacol.* 1994; 45: 428-438.
- Pongratz I, Mason GG, Poellinger L. Dual roles of the 90-kDa heat shock protein hsp90 in modulating functional activities of the dioxin receptor. Evidence that the dioxin receptor functionally belongs to a subclass of nuclear receptors which require hsp90 both for ligand binding activity and repression of intrinsic DNA binding activity. *J Biol Chem.* 1992; 267: 13728-13734.

Prell RA, Oughton JA, Kerkvliet NI. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on anti-CD3-induced changes in T-cell subsets and cytokine production. *Int J Immunopharmacol.* 1995; 17: 951-961.

Przyrembel H, Heinrich-Hirsch B, Vieth B. Exposition to and health effects of residues in human milk. *Adv Exp Med Biol.* 2000; 478: 307-325.

Riecke K, Nogueira AC, Alexi-Meskishvili V, Stahlmann R. Cross-reactivity of antibodies on thymic epithelial cells from humans and marmosets by flow-cytometry. *J Med Primatol.* 2000; 29: 343-349.

Riecke K, Grimm D, Shakibaei M, Kossmehl P, Schulze-Tanzil G, Paul M, Stahlmann R. Low doses of 2,3,7,8-tetrachlorodibenzo-p-dioxin increase transforming growth factor beta and cause myocardial fibrosis in marmosets (*Callithrix jacchus*). *Arch Toxicol.* 2002; 76: 360-366.

Reisz-Porszasz S, Probst MR, Fukunaga BN, Hankinson O. Identification of functional domains of the aryl hydrocarbon receptor nuclear translocator protein (ARNT). *Mol Cell Biol.* 1994; 14: 6075-6086.

Rizzardini M, Romano M, Tursi F, Salmona M, Vecchi A, Sironi M, Gizzi F, Benfenati E, Garattini S, Fanelli R. Toxilogical evaluation of urban waste incinerator emissions. *Chemosphere.* 1983; 12: 559-564.

Romagnani P, Annunziato F, Manetti R, Mavilia C, Lasagni L, Manuelli C, Vannelli GB, Vanini V, Maggi E, Pupilli C, Romagnani S. High CD30 ligand expression by epithelial cells and Hassal's corpuscles in the medulla of human thymus. *Blood.* 1998; 91: 3323-3332.

- Rose JQ, Ramsey JC, Wentzler TH, Hummel RA. The fate of 2,3,7,8-tetrachlorodibenzo-p-dioxin following single and repeated oral doses to the rat. *Toxicol Appl Pharmacol.* 1976; 36: 209-226.
- Rowlands JC, Gustafsson JA. Aryl hydrocarbon receptor-mediated signal transduction. *Crit Rev Toxicol.* 1997; 27: 109-134.
- Safe SH. Comparative toxicology and mechanism of action of polychlorinated dibenzo-p-dioxins and dibenzofurans. *Annu Rev Pharmacol Toxicol.* 1986; 26: 371-399.
- Safe SH. Polychlorinated biphenyls (PCBs), dibenzo-p-dioxins (PCDDs), dibenzofurans (PCDFs), and related compounds: environmental and mechanistic considerations which support the development of toxic equivalency factors (TEFs). *Crit Rev Toxicol.* 1990; 21: 51-88.
- Safe SH. Polychlorinated biphenyls (PCBs): environmental impact, biochemical and toxic responses, and implications for risk assessment. *Crit Rev Toxicol.* 1994; 24: 87-149.
- Salomon DR, Crisa L, Mojciak CF, Ishii JK, Klier G, Shevach EM. Vascular cell adhesion molecule-1 is expressed by cortical thymic epithelial cells and mediates thymocyte adhesion. Implications for the function of alpha4beta1 (VLA4) integrin in T-cell development. *Blood.* 1997; 89: 2461-2471.
- Savino W, Villa-Verde DM, Lannes-Vieira J. Extracellular matrix proteins in intrathymic T-cell migration and differentiation? *Immunol Today.* 1993; 14: 158-161.
- Savino W, Dardenne M, Carnaud C. The conveyor belt model for intrathymic T-cell migration. *Immunol Today.* 1996; 17: 97-98.



- Sawada M, Nagamine J, Takeda K, Utsumi K, Kosugi A, Tatsumi Y, Hamaoka T, Miyake K, Nakajima K, Watanabe T, et al. Expression of VLA-4 on thymocytes. Maturation stage-associated transition and its correlation with their capacity to adhere to thymic stromal cells. *J Immunol.* 1992; 149: 3517-3524.
- Schmidt JV, Bradfield CA. Ah receptor signaling pathways. *Annu Rev Cell Dev Biol.* 1996; 12: 55-89.
- Schwetz BA, Norris JM, Sparschu GL, Rowe UK, Gehring PJ, Emerson JL, Gerbig CG. Toxicology of chlorinated dibenzo-p-dioxins. *Environ Health Perspect.* 1973; 5: 87-99.
- Shepherd DM, Dearstyne EA, Kerkvliet NI. The effects of TCDD on the activation of ovalbumin (OVA)-specific DO11.10 transgenic CD4(+) T cells in adoptively transferred mice. *Toxicol Sci.* 2000; 56: 340-350.
- Sligh JE Jr, Ballantyne CM, Rich SS, Hawkins HK, Smith CW, Bradley A, Beaudet AL. Inflammatory and immune responses are impaired in mice deficient in intercellular adhesion molecule 1. *Proc Natl Acad Sci U S A.* 1993; 90: 8529-8533.
- Staunton DE, Marlin SD, Stratowa C, Dustin ML, Springer TA. Primary structure of ICAM-1 demonstrates interaction between members of the immunoglobulin and integrin supergene families. *Cell.* 1988; 52: 925-933.
- Stephens RD, Petreas MX, Hayward DG. Biotransfer and bioaccumulation of dioxins and furans from soil: chickens as a model for foraging animals. *Sci Total Environ.* 1995; 175: 253-273.
- Tarkowski S. WHO Coordinated intercountry studies on levels of PCDDs and PCDFs in human milk, *Chemosphere* 19, 1996, 995-1000.

- Thurmond TS, Silverstone AE, Baggs RB, Quimby FW, Staples JE, Gasiewicz TA. A chimeric aryl hydrocarbon receptor knockout mouse model indicates that aryl hydrocarbon receptor activation in hematopoietic cells contributes to the hepatic lesions induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Toxicol Appl Pharmacol.* 1999; 158: 33-40.
- U.S. Environmental Protection Agency (U.S. EPA). Ambient water quality criteria for 2,3,7,8-tetrachlorodibenzo-p-dioxin. Office of Water Regulations and Standards Division, Washington, DC. 1984; EPA 440/5-84-007.
- U.S. EPA. Health assessment for polychlorinated dibenzo-p-dioxins. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH, for the Office of Emergency and Remedial Response, Washington, DC. 1985; EPA 600/8-84/0146.
- U.S. EPA. Interim procedures for estimating risks associated with exposures to mixtures of chlorinated dibenzo-p-dioxines and –dibenzofurans (CDDs and CDFs). 1987; EPA 625/3-87/012.
- U.S. EPA. Workshop report on toxicity equivalency factors for polychlorinated biphenyls congeners. 1991; EPA/625/3-91/020.
- Van den Berg M, De Jongh J, Poiger H, Olson JR. The toxicokinetics and metabolism of polychlorinated dibenzo-p-dioxins (PCDDs) and dibenzofurans (PCDFs) and their relevance for toxicity. *Crit Rev Toxicol.* 1994; 24: 1-74.
- Van den Berg M, Birnbaum L, Bosveld AT, Brunstrom B, Cook P, Feeley M, Giesy JP, Hanberg A, Hasegawa R, Kennedy SW, Kubiak T, Larsen JC, van Leeuwen FX, Liem AK, Nolt C, Peterson RE, Poellinger L, Safe S, Schrenk D, Tillitt D, Tysklind M, Younes M, Waern F, Zacharewski T. Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. *Environ Health Perspect.* 1998; 106: 775-792.

- Van den Heuvel JP, Clark GC, Tritscher A, Lucier GW. Accumulation of polychlorinated dibenzo-p-dioxins and dibenzofurans in liver of control laboratory rats. *Fundam Appl Toxicol.* 1994; 23: 465-469.
- Van der Vieren M, Le Trong H, Wood CL, Moore PF, St John T, Staunton DE, Gallatin WM. A novel leukointegrin, alpha d beta 2, binds preferentially to ICAM-3. *Immunity.* 1995; 3: 683-690.
- Villa-Verde DM, Calado TC, Ocampo JS, Silva-Monteiro E, Savino W. The conveyor belt hypothesis for thymocyte migration: participation of adhesion and de-adhesion molecules. *Braz J Med Biol Res.* 1999; 32: 569-572.
- Viluksela M, Stahl BU, Birnbaum LS, Schramm KW, Kettrup A, Rozman KK. Subchronic/chronic toxicity of 1,2,3,4,6,7,8-heptachlorodibenzo-p-dioxin (HpCDD) in rats. Part I. Design, general observations, hematology, and liver concentrations. *Toxicol Appl Pharmacol.* 1997a; 146: 207-216.
- Viluksela M, Stahl BU, Birnbaum LS, Rozman KK. Subchronic/chronic toxicity of 1,2,3,4,6,7,8-heptachlorodibenzo-p-dioxin (HpCDD) in rats. Part II. Biochemical effects. *Toxicol Appl Pharmacol.* 1997b; 146: 217-226.
- Viluksela M, Stahl BU, Birnbaum LS, Schramm KW, Kettrup A, Rozman KK. Subchronic/chronic toxicity of a mixture of four chlorinated dibenzo-p-dioxins in rats. I. Design, general observations, hematology, and liver concentrations. *Toxicol Appl Pharmacol.* 1998; 151: 57-69.
- Vogel C, Abel J. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on growth factor expression in the human breast cancer cell line MCF-7. *Arch Toxicol.* 1995; 69: 259-265.

- Vorderstrasse BA, Steppan LB, Silverstone AE, Kerkvliet NI. Aryl hydrocarbon receptor-deficient mice generate normal immune responses to model antigens and are resistant to TCDD-induced immune suppression. *Toxicol Appl Pharmacol.* 2001; 171: 157-164.
- Vos JG, Luster MI. Immune alterations. Halogenated biphenyls, terphenyles, naphthalenes, dibenzodioxins and related products. Elsevier Science Publishers, Amsterdam. 1989; 295-322.
- Waller CL, McKinney JD. Three-dimensional quantitative structure-activity relationships of dioxins and dioxin-like compounds: model validation and Ah receptor characteristics. *Chem Res Toxicol.* 1995; 8: 847-858.
- Weber LW, Zesch A, Rozman K. Penetration, distribution and kinetics of 2,3,7,8-tetrachlorodibenzo-p-dioxin in human skin in vitro. *Arch Toxicol.* 1991; 65: 421-428.
- Weller GL. Development of the thyroid, parathyroid and thymus glands in man. *Contrib Embryol Carnegie Inst.* 1933; 22: 95.
- Wester RC, Maibach HI, Sedik L, Melendres J, Wade M. Percutaneous absorption of PCBs from soil: in vivo rhesus monkey, in vitro human skin, and binding to powdered human stratum corneum. *J Toxicol Environ Health.* 1993a; 39: 375-382.
- Wester RC, Maibach HI, Sedik L, Melendres J, Wade M, DiZio S. Percutaneous absorption of pentachlorophenol from soil. *Fundam Appl Toxicol.* 1993b; 20: 68-71.
- Whitlock JP Jr, Cooper HL, Gelboin VH. Aryl hydrocarbon (benzopyrene) hydroxylase is stimulated in human lymphocytes by mitogens and benz(a)anthracene. *Science.* 1972; 177: 618-619.

Whitlock JP. Genetic and molecular aspects of 2,3,7,8-tetrachlorodibenzo-p-dioxin action. *Annu Rev Pharmacol Toxicol.* 1990; 30: 251-77.

Whitlock JP. Induction of cytochrome P4501A1. *Annu Rev Pharmacol Toxicol.* 1999; 39: 103-125.

World Health Organisation / International Programme on Chemical Safety (WHO/IPCS). Polychlorinated dibenzo-p-dioxins and dibenzofurans. *Environmental Health Criteria.* 1989; 88.

World Health Organization, International Agency for Research on Cancer, IARC monographs on the evaluation of carcinogenic risks to humans, Polychlorinated Dibenzo-para-dioxins and Polychlorinated Dibenzofurans, Lyon. 1997; 47.

Xu H, Tong IL, De Fougerolles AR, Springer TA. Isolation, characterization, and expression of mouse ICAM-2 complementary and genomic DNA. *J Immunol.* 1992; 149: 2650-2655.