

## 5. References

- Abu-Issa R, Smyth G, Smoak I, Yamamura K, Meyers EN.2002. Fgf8 is required for pharyngeal arch and cardiovascular development in the mouse. *Development*. 129(19):4613-25.
- Ahlgren SC, Bronner-Fraser M.1999. Inhibition of sonic hedgehog signaling in vivo results in craniofacial neural crest cell death. *Curr Biol*. 9(22):1304-14.
- Alcaniz S, Silva FJ. 1997. Phenylalanine hydroxylase participation in synthesis of serotonin and pteridines in drosophila melanogaster. *Comp. Bioch. Physiol*. 116(3):205-212.
- Alenina N, Bashammakh S, Bader M. 2006. Specification and differentiation of serotonergic neurons. *Stem Cell Rev*. 2:5-10.
- Amores A, Force A, Yan YL, Joly L, Amemiya C, Fritz A, Ho RK, Langeland J, Prince V, Wang YL, Westerfield M, Ekker M, Postlethwait JH. 1998. Zebrafish hox clusters and vertebrate genome evolution. *Science*. 1998. 282:1711-4.
- Anichtchik O, Sallinen V, Peitsaro N, Panula P.2006. Distinct structure and activity of monoamine oxidase in the brain of zebrafish (*Danio rerio*). *J Comp Neurol*. 2006.498(5):593-610.
- Appelbaum L, Vallone D, Anzulovich A, Ziv L, Tom M, Foulkes NS, Gothilf Y. 2006. Zebrafish arylalkylamine-N-acetyltransferase genes - targets for regulation of the circadian clock. *J Mol Endocrinol*. 36(2):337-47.
- Aragon MA, Ayala ME, Marin M, Aviles A, Damian-Matsumura P, Dominguez R.. 2005. Serotonergic system blockage in the prepubertal rat inhibits spermatogenesis development. *Reproduction*. 129(6):717-27.
- Azmitia EC. 2001. Modern views on an ancient chemical: serotonin effects on cell proliferation, maturation, and apoptosis. *Brain Res Bull*. 56(5):413-24.
- Azmitia EC. 2007. Serotonin and brain: evolution, neuroplasticity, and homeostasis. *Int Rev Neurobiol*. 77:31-56.
- Bagdy G. 1996. Role of the hypothalamic paraventricular nucleus in 5-HT1A, 5-HT2A and 5-HT2C receptor-mediated oxytocin, prolactin and ACTH/corticosterone responses. *Behav Brain Res*. 73(1-2):277-80.
- Bally-Cuif L, Goutel C, Wassef M, Wurst W, Rosa F. 2000. Coregulation of anterior and posterior mesendodermal development by a hairy-related transcriptional repressor. *Genes Dev*. 14(13):1664-77.

- Barnes NM and Sharp T. 1999. A review of central 5-HT receptors and their function. *Neuropharmacology*. 38:1083 - 1152.
- Begay V, Falcon J, Cahil GM, Klein DC, Coon SL. 1998. Transcripts encoding two melatonin synthesis enzymes in the teleost pineal organ: circadian regulation in pike and zebrafish, but not in trout. *Endocrinology*. 139(3):905-12.
- Begbie J, Graham A. 2001. Integration between the epibranchial placodes and the hindbrain. *Science*. 294:595-598.
- Bellipanni G, Rink E, Bally-Cuif L. 2002. Cloning of two tryptophan hydroxylase genes expressed in the diencephalon of the developing zebrafish brain. *Gene Expr Patterns*. 2(3-4):251-6
- Berry-Kravis EM, Zhou L, Rand CM, Weese-Mayer DE. 2006. Congenital Central Hypoventilation Syndrome: PHOX2B Mutations and Phenotype. *Am J Respir Crit Care Med*. 174(10):1139-44.
- Bhasin N, Kernick E, Luo X, Seidel HE, Weiss ER, Lauder JM. 2004a. Differential regulation of chondrogenic differentiation by the serotonin<sub>2B</sub> receptor and retinoic acid in the embryonic mouse hindlimb. *Dev Dyn*. 2:201-9.
- Bhasin N, LaMantia AS, Lauder JM. 2004b. Opposing regulation of cell proliferation by retinoic acid and the serotonin<sub>2B</sub> receptor in the mouse frontonasal mass. *Anat Embryol (Berl)*. 208(2):135-43.
- Blaugrund E, Pham TD, Tennyson V M, Lo L, Sommer L, Anderson DJ, and Gershon MD. 1996. Distinct subpopulations of enteric neuronal progenitors defined by time of development, sympathoadrenal lineage markers and Mash-1-dependence. *Development*. 122:309-320.
- Borday-Birraux V, Van der Heyden C, Debais-Thibaud M, Verreijdt L, Stock DW, Huisseune A, Sire JY. 2006. Expression of Dlx genes during the development of the zebrafish pharyngeal dentition: evolutionary implications. *Evol Dev*. 8(2):130-41.
- Bradley SR, Pieribone VA, Wang W, Severson CA, Jacobs RA, Richerson GB. 2002. Chemosensitive serotonergic neurons are closely associated with large medullary arteries. *Nat Neurosci*. 5(5):401-2.
- Briscoe J, Sussel L, Serup P, Hartigan-O'Connor D, Jessell TM, Rubenstein JL, and Ericson J. 1999. Homeobox gene Nkx2.2 and specification of neuronal identity by graded Sonic hedgehog signalling. *Nature*. 398:622-627.

- Brito JM, Teillet MA, Le Douarin NM. 2006. An early role for sonic hedgehog from foregut endoderm in jaw development: ensuring neural crest cell survival .Proc. Natl. Acad. Sci. U S A. 103(31):11607-12.
- Britsch S, Goerich DE, Riethmacher D, Peirano RI, Rossner M, Nave KA, Birchmeier C, Wegner M. 2001. The transcription factor Sox10 is a key regulator of peripheral glial development. *Genes Dev.* 15(1):66-78.
- Brodski C, Weisenhorn DM, Signore M, Sillaber I, Oesterheld M, Broccoli V, Acampora D, Simeone A, Wurst W. 2003. Location and size of dopaminergic and serotonergic cell populations are controlled by the position of the midbrain-hindbrain organizer. *J Neurosci.* 23(10):4199-207.
- Bronner-Fraser, M. & Fraser, S. (1988).Cell lineage analysis shows multipotentiality of some avian neural crest cells. *Nature* .335:161-164.
- Brunet, J. F. and Pattyn, A. 2002. Phox2 genes - from patterning to connectivity. *Curr. Opin. Genet. Dev.* 12:435-440.
- Buzikov GA, Shmuker YB, Lauder JM. 1996. From oocyte to neuron: do neurotransmitters function in the same way throughout development?. *Cell Mol Neurobiol.* 16(5): 537-59.
- Buznikov GA, Nikitina LA, Voronezhskaya EE, Bezuglov VV, Dennis Willows AO, Nezlin LP. 2003. Localization of serotonin and its possible role in early embryos of *Tritonia diomedea*(Mollusca: Nudibranchia). *Cell Tissue Res.* 311(2):259-66.
- Buznikov GA, Peterson RE, Nikitina LA, Bezuglov VV, Lauder JM. 2005. The pre-nervous serotonergic system of developing sea urchin embryos and larvae: pharmacologic and immunocytochemical evidence.*Neurochem Res.* 30(6-7):825-37.
- Byrd NA, Meyers EN. 2005. Loss of Gbx2 results in neural crest cell patterning and pharyngeal arch artery defects in the mouse embryo. *Dev Biol.* 284(1):233-45.
- Byrden HW and Lawrence IE Jr. 1973. Presence of biogenic amines in early rat development. *Am J Anat.* 136(2):251-7.
- Calogero AE, Bagdy G, Szemerdi K, Tartaglia ME, Gold PW, Chrousos GP. 1990. Mechanisms of serotonin receptor agonist-induced activation of the hypothalamic-pituitary-adrenal axis in the rat. *Endocrinology.* 126(4):1888-94.
- Cau E, Wilson SW. 2003. Ash1a and Neurogenin1 function downstream of Floating head to regulate epiphyseal neurogenesis. *Development.* 130(11):2455-66.

- Cheng L, Chen CL, Luo P, Tan M, Qiu M, Johnson R, Ma Q. 2003. *Lmx1b*, *Pet-1*, and *Nkx2.2* coordinately specify serotonergic neurotransmitter phenotype. *J. Neurosci.* 23:9961-9967.
- Choi DS, Ward SJ, Messaddeg N, Launay JM, Maroteaux L. 1997. 5-HT<sub>2B</sub> receptor-mediated serotonin morphogenetic functions in mouse cranial neural crest and myocardial cells. *Development.* 124(9):1745-55.
- Clark MS, Lanigan TM, Page NM, Russo AF. 1995. Induction of a serotonergic and neuronal phenotype in thyroid C-cells. *J Neurosci.* 15(9):6167-78.
- Colas JF, Launay JM, Maroteaux L. 1999a. Maternal and zygotic control of serotonin biosynthesis are both necessary for *Drosophila* germband extension. *Mech Dev.* 87(1-2):67-76.
- Colas JF, Launay JM, Vonesch JL, Hickel P, Maroteaux L. 1999b. Serotonin synchronises convergent extension of ectoderm with morphogenetic gastrulation movements in *Drosophila*. *Mech Dev.* 87(1-2):77-91.
- Consolazione, A., and Cuello, A. 1982. C. CNS serotonin pathways. In *The Biology of Serotonergic Transmission*. New York: John Wiley & Sons, Ltd., pp. 29–61
- Couly G, Creuzet S, Bennaceur S, Vincent C, Le Douarin NM. 2002. Interactions between Hox-negative cephalic neural crest cells and the foregut endoderm in patterning the facial skeleton in the vertebrate head. *Development.* 129(4):1061-73.
- Craven SE, Lim KC, Ye W, Engel JD, de Sauvage F, Rosenthal A. 2004. *Gata2* specifies serotonergic neurons downstream of sonic hedgehog. *Development.* 13:1165-1173.
- Dahlstrom A, and Fuxe K. 1964. Evidence for the existence of monamine-containing neurons in the central nervous system. I. Demonstration of monamines in the cell bodies of brainstem neurons. *Acta Physiol Scand.* 62(232):1-55.
- Dager S, Pattyn A, Lofaso F, Gaultier C, Goridis C, Gallego J, Brunet JF. 2003. *Phox2b* controls the development of peripheral chemoreceptors and afferent visceral pathways. *Development.* 130(26):6635-42.
- Ding YQ, Marklund U, Yuan W, Yin J, Wegman L, Ericson J, Deneris E, Johnson RL, and Chen ZF. 2003. *Lmx1b* is essential for the development of serotonergic neurons. *Nat. Neurosci.* 6:933-938.
- Dutton KA, Pauliny A, Lopes SS, Elworthy S, Carney TJ, Rauch J, Geisler R, Haffter P, Kelsh RN. 2001. Zebrafish *colourless* encodes *sox10* and specifies non-ectomesenchymal neural crest fates. *Development.* 128(21):4113-25.

- Ehret M, Cash CD, Hamon M, Maitre M. 1989. Formal demonstration of the phosphorylation of rat brain tryptophan hydroxylase by Ca<sup>2+</sup>/calmodulin-dependent protein kinase. *J. Neurochem.* 52:1886–1891.
- Erspamer V, and Asero B. 1952. Identification of enteramine, specific hormone of enterochromaffin cells, as 5-hydroxytryptamine. *Nature.* 169:800-801.
- Fitzpatrick PF. 1999. Tetrahydropterin-dependent amino acid hydroxylases. *Annu. Rev. Biochem.* 68:355–381.
- Fukumoto T, Kema IP, Levin M. 2005. Serotonin signaling is a very early step in patterning of the left-right axis in chick and frog embryos. *Curr Biol.* 15(9):794-803.
- Gal EM, Patterson K. 1973. Rapid nonisotopic assay of tryptophan-5-hydroxylase activity in tissues. *Anal Biochem.* 52(2):625-9.
- Galter D, Bottner M, Krieglstein K, Schomig E, Unsicker K. 1999. Differential regulation of distinct phenotypic features of serotonergic neurons by bone morphogenetic proteins. *Eur J Neurosci.* 11(7):2444-52.
- Gammill LS, Bronner-Fraser M. 2003. Neural crest specification: migrating into genomics. *Nat Rev Neurosci.* 10:795-805
- Gaspar P, Cases O, Maroteaux L. 2003. The developmental role of serotonin: news from mouse molecular genetics. *Nat Rev Neurosci.* 4(12):1002-12.
- George J. Siegel M.D (editor). 1998. *Basic Neurochemistry. Molecular, Cellular and Medical Aspects. Sixth Edition*
- Gershon MD, Sherman DL, Pintar JE. 1990. Type-specific localization of monoamine oxidase in the enteric nervous system: relationship to 5-hydroxytryptamine, neuropeptides, and sympathetic nerves. *J Comp Neurol.* 301(2):191-213.
- Gershon MD, Tack J. 2007. The serotonin signaling system: from basic understanding to drug development for functional GI disorders. *Gastroenterology.* 132(1):397-414.
- Glennon A, Malgorzata Dukat, Richard B. Westkaemper. *Serotonin Receptors Subtypes and Ligands. American College of Neuropsychopharmacology. The Fourth Generation of Progress.* www.acnp.org
- Goldstein AM, Brewer KC, Doyle AM, Nagy N, Roberts DJ. 2005. BMP signaling is necessary for neural crest cell migration and ganglion formation in the enteric nervous system. *Mech Dev.* 122(6):821-33.
- Goridis C, Rohrer H. 2002. Specification of catecholaminergic and serotonergic neurons. *Nat Rev Neurosci.* 3(7):531-41.

- Gothilf Y, Coon SL, Toyama R, Chitnis A, Namboodiri MA, Klein DC. 1999. Zebrafish serotonin N-acetyltransferase-2: marker for development of pineal photoreceptors and circadian clock function. *Endocrinology*. 140(10):4895-903.
- Graham A, Begbie J, McGonnell I. 2004. Significance of the cranial neural crest. *Dev Dyn*. 229:5-13.
- Graham A, Francis-West P, Brickell P, Lumsden A. 1994. The signalling molecule BMP4 mediates apoptosis in the rhombencephalic neural crest. *Nature*. 372: 684-686.
- Graham A, Smith A. 2001. Patterning the pharyngeal arches. *Bioessays* 23: 54-61.
- Graham A. 2003. Development of the pharyngeal arches. *Am J Med Genet* 119A: 251-256.
- Graham A, Okabe M, Quinlan R. 2005. The role of the endoderm in the development and evolution of the pharyngeal arches. *J. Anat.* 207(5):479-487.
- Grahame-Smith DG. 1964. Tryptophan hydroxylation in brain. *Biochem. Biophys. Res. Commun.* 16:586-592.
- Grailhe R, Grabtree GW, Hen R. 2001. Human 5-HT(5) receptors: the 5-HT(5A) receptor is functional but the 5-HT(5B) receptor was lost during mammalian evolution. *Eur J Pharmacol.* 418(3):157-67.
- Guo S, Yamaguchi Y, Schilbach S, Wada T, Lee J, Goddard A, French D, Handa H, Rosenthal A. (2000) A regulator of transcriptional elongation controls vertebrate neuronal development. *Nature*. 408(6810):366-9
- Gustafson T, Toneby M. 1970 On the role of serotonin and acetylcholine in sea urchin morphogenesis. *Exp Cell Res.* 62(1):102-17.
- Hamalainen M, Kohonen J. 1989. Studies on the effect of monoamine antagonists on the morphogenesis of the newt. *Int J Dev Biol.* 33(1):157-63.
- Hanaoka R, Ohmori Y, Uyemura K, Hosoya T, Hotta Y, Shirao T, Okamoto H. 2004; Zebrafish *gcmb* is required for pharyngeal cartilage formation. *Mech Dev.* 121(10):1235-47.
- Hendricks TJ, Fyodorov DV, Wegman LJ, Lelutiu NB, Pehek EA, Yamamoto B, Silver J, Weeber E J, Sweatt JD, Deneris ES. 2003. Pet-1 ETS gene plays a critical role in 5-HT neuron development and is required for normal anxiety-like and aggressive behavior. *Neuron*. 37:233-247.
- Hendricks T, Francis N, Fyodorov D, Deneris ES. 1999. The ETS domain factor Pet-1 is an early and precise marker of central serotonin neurons and interacts with a conserved element in serotonergic genes. *J. Neurosci.* 19:10348-10356.

- Hobson RJ, Hapiak VM, Xiao H, Buehrer KL, Komuniecki PR, Komuniecki RW. 2006. SER-7, a *Caenorhabditis elegans* 5-HT7-like receptor, is essential for the 5-HT stimulation of pharyngeal pumping and egg laying. *Genetics*. 172(1):159-69.
- Hogan BM, Hunter MP, Oates AC, Crowhurst MO, Hall NE, Heath JK, Prince VE, Lieschke GJ. 2004; Zebrafish *gcm2* is required for gill filament budding from pharyngeal ectoderm. *Dev Biol*. 276(2):508-2
- Hogan BM, Hunter MP, Oates AC, Crowhurst MO, Hall NE, Heath JK, Prince VE, Lieschke GJ. 2004. Zebrafish *gcm2* is required for gill filament budding from pharyngeal ectoderm. *Dev Biol*. 276(2):508-22
- Hoyer D, Hannon JP, Martin GR. 2002. Molecular, pharmacological and functional diversity of 5-HT receptors. *Pharmacol Biochem Behav*. 71(4):533-54. Review.
- Hoyer D, and Martin G. 1997. 5-HT receptor classification and nomenclature: towards a harmonization with the human genome. *Neuropharmacology*. 36:419 - 428.
- Il'kova G, Rehak P, Vesela J, Cikos S, Fabian D, Czikkova S, Koppel J. 2004. Serotonin localization and its functional significance during mouse preimplantation embryo development. *Zygote*. 12(3):205-13.
- Jacobs and Azmitia. 1992. Structure and function of the brain serotonin system. *Physiol. Rev*. 72(1):165-229.
- Johansen PA, Jennings I, Cotton RG, Kuhn DM. 1996. Phosphorylation and activation of tryptophan hydroxylase by exogenous protein kinase A. *J. Neurochem*. 66:817-823.
- Jonz MG, Nurse CA. 2006. Ontogenesis of oxygen chemoreception in aquatic vertebrates. *Respir Physiol Neurobiol*. 154(1-2):139-52.
- Kameda Y. 2005. *Mash1* is required for glomus cell formation in the mouse carotid body. *Dev Biol*. 283(1):128-39.
- Kaslin J, Panula P. 2001. Comparative anatomy of the histaminergic and other aminergic systems in zebrafish (*Danio rerio*). *J Comp Neurol*. 440(4):342-77.
- Kaufman S. 1966. Coenzymes and hydroxylases: ascorbate and dopamine-beta-hydroxylase; tetrahydropteridines and phenylalanine and tyrosine hydroxylases. *Pharmacol Rev*. 18(1):61-9.
- Kline D, Simoncini L, Mandel G, Maue RA, Kado RT, Jaffe LA. 1988. Fertilization events induced by neurotransmitters after injection of mRNA in *Xenopus* eggs. *Science*. 241(4864):464-7.

- Kopinke D, Sasine J, Swift J, Stephens WZ, Piotrowski T. 2006. Retinoic acid is required for endodermal pouch morphogenesis and not for pharyngeal endoderm specification. *Dev Dyn.* 235(10):2695-709
- Koza A, Wilhelm M, Hiripi L, Elekes K, Csoknya M. 2006. Embryogenesis of the serotonergic system in the earthworm *Eisenia fetida* (Annelida, Oligochaeta): immunohistochemical and biochemical studies. *J Comp Neurol.* 497(3):451-67.
- Kuhn DH, Arthur R. Jr., States JC, 1972. Phosphorylation and activation of brain tryptophan hydroxylase: identification of serine-58 as a substrate site for protein kinase A. *J. Neurochem.* 68:2220–2223.
- Lallier T, Leblanc G, Artinger KB, Bronner-Fraser M. 1992. Cranial and trunk neural crest cells use different mechanisms for attachment to extracellular matrices. *Development* 116:531–541.
- Lanfumey L, Hamon M. 2004. 5-HT1 receptors. *Curr Drug Targets CNS Neurol Disord.* 3(1):1-10.
- Lauder JM, Zimmerman EF. 1988. Sites of serotonin uptake in epithelia of the developing mouse palate, oral cavity, and face: possible role in morphogenesis. *J Craniofac Genet Dev Biol.* 8(3):265-76.
- Lebrand C, Cases O, Adelbrecht C, Doye A, Alvarez C, El Mestikawy S, Seif I, Gaspar P. 1996. Transient uptake and storage of serotonin in developing thalamic neurons. *Neuron.* (5):823-35.
- LeDouarin, N. & Kalcheim, C. *The Neural Crest* (eds. Bard, J., Barlow, P. & Kirk, D.) Cambridge Univ. Press. 1999.
- Leibowitz SF. and Alexander JT. 1998. Hypothalamic serotonin in control of eating behavior, meal size, and body weight. *Biol Psychiatry.* 44(9):851-64. Review.
- Li Y, Xiang J, Duan CJ. 2005. Insulin-like growth factor-binding protein-3 plays an important role in regulating pharyngeal skeleton and inner ear formation and differentiation. *J Biol Chem.* 280(5):3613-20.
- Li YX, Zdanowicz M, Young L, Kumiski D, Leatherbury L, Kirby ML. 2003. Cardiac neural crest in zebrafish embryos contributes to myocardial cell lineage and early heart function. *Dev Dyn.* 226(3):540-50.
- Lister JA, Cooper C, Nguyen K, Modrell M, Grant K, Raible DW. 2006. Zebrafish *Foxd3* is required for development of a subset of neural crest derivatives. *Dev Biol.* 290(1):92-104.



- Lovenberg W, Jequier E, Sjoerdsma A, 1976. Tryptophan hydroxylation: measurement in pineal gland, brainstem, and carcinoid tumor. *Science*. 155:217–219.
- McArthur AJ, Hunt AE, Gillette MU. 1997. Melatonin action and signal transduction in the rat suprachiasmatic circadian clock: activation of protein kinase C at dusk and dawn. *Endocrinology*. 138(2):627-34.
- McLean DL, Fetcho JR. 2004. Ontogeny and innervation patterns of dopaminergic, noradrenergic, and serotonergic neurons in larval zebrafish. *J Comp Neurol*. 480(1):3856.
- McNulty CL, Peres JN, Bardine N, van den Akker WM, Durston AJ. 2005. Knockdown of the complete Hox paralogous group 1 leads to dramatic hindbrain and neural crest defects. *Development*. 132(12):2861-71.
- Meizel S, Turner KO. 1983. Serotonin or its agonist 5-methoxytryptamine can stimulate hamster sperm acrosome reactions in a more direct manner than catecholamines. *J Exp Zool*. 226(1):171-4.
- Metcalfe WK, Myers PZ, Trevarrow B, Bass MB, Kimmel CB. 1990. Primary neurons that express the L2/HNK-1 carbohydrate during early development in the zebrafish. *Development*. 110(2):491-504.
- Miller CT, Schilling TF, Lee K, Parker J, Kimmel CB. 2000. sucker encodes a zebrafish Endothelin-1 required for ventral pharyngeal arch development. *Development*. 127(17):3815-28.
- Moiseiwitsch JR and Lauder JM. 1995. Serotonin regulates mouse cranial neural crest migration. *Proc. Natl. Acad. Sci. USA*. 92 (16):7182-7186.
- Moiseiwitsch JR. 2000. The role of serotonin and neurotransmitters during craniofacial development. *Crit Rev Oral Biol Med*. 2:230-9.
- Morikawa K, Tsuneki K, Ito K. 2001. Expression patterns of HNK-1 carbohydrate and serotonin in sea urchin, amphioxus, and lamprey, with reference to the possible evolutionary origin of the neural crest. *Zoology*. 104(2):81-90.
- Nebigil CG, Etienne N, Schaelinger B, Hickel P, Launay JM, Maroteaux L. 2001. Developmentally regulated serotonin 5-HT<sub>2B</sub> receptors. *Int J Dev Neurosci*. 19(4):365-72.
- Nebigil CG, Choi DS, Dierich A, Hickel P, Le Meur M, Messaddeq N, Launay JM, Maroteaux L. 2000. Serotonin 2B receptor is required for heart development. *Proc Natl Acad Sci U S A*. 97(17):9508-13.

- Neckameyer WS, White K. 1992. A single locus encodes both phenylalanine hydroxylase and tryptophan hydroxylase activities in *Drosophila*. *J Biol Chem.* 267(6):4199-206.
- Nelson DL. 2004. 5-HT<sub>5</sub> receptors. *Curr Drug Targets CNS Neurol Disord.* 3(1):53-8.
- Parisi E, De Prisco P, Capasso A, del Prete M. 1984. Serotonin and sperm motility. *Cell Biol Int Rep.* 8(2):95.
- Patient RK, and McGhee JD. 2002. The GATA family (vertebrates and invertebrates)., *Curr. Opin. Genet. Dev.* 12:416-422.
- Pattyn A, Morin X, Cremer H, Goridis C, Brunet JF. 1999. The homeobox gene *Phox2b* is essential for the development of autonomic neural crest derivatives. *Nature.* 399(6734):366-70.
- Pattyn A, Simplicio N, van Doorninck JH, Goridis C, Guillemot F, Brunet JF. 2004. *Ascl1/Mash1* is required for the development of central serotonergic neurons. *Nat. Neurosci.* 7:589-595.
- Pattyn A, Vallstedt A, Dias JM, Samad OA, Krumlauf R, Rijli F M, Brunet JF, Ericson J. 2003. Coordinated temporal and spatial control of motor neuron and serotonergic neuron generation from a common pool of CNS progenitors. *Genes Dev.* 17:729-737.
- Pino GD, Moessner R, Lesch KP, Lauder JM, Persico AM. 2004. Roles for Serotonin in Neurodevelopment: More than just Neural Transmission. *Current Neuropharmacology.* 4(2):403-417.
- Piotrowski T, and Nusslein-Volhard C. 2000. The endoderm plays an important role in patterning the segmented pharyngeal region in zebrafish (*Danio rerio*). *Dev. Biol.* 225:339 -356.
- Puglisi-Allegra S, Cabib S, Pascucci T, Ventura R, Cali F, Romano V. 2000. Dramatic brain aminergic deficit in a genetic mouse model of phenylketonuria. *Neuroreport.* 11(6):1361-4.
- Ranganathan R, Cannon SC, Horvitz HR. 2000. MOD-1 is a serotonin-gated chloride channel that modulates locomotory behaviour in *C. elegans*. *Nature.* 408(6811):470-5.
- Richerson GB. 2004. Serotonergic neurons as carbon dioxide sensors that maintain pH homeostasis. *Nat Rev Neurosci.* 5(6):449-61.
- Ruhin B, Creuzet S, Vincent C, Benouaiche L, Le Douarin NM, Couly G. 2003. Patterning of the hyoid cartilage depends upon signals arising from the ventral foregut endoderm. *Dev Dyn.* 228(2):239-46.

- Samad OA, Geisen MJ, Caronia G, Varlet I, Zappavigna V, Ericson J, Goridis C, Rijli FM. 2004. Integration of anteroposterior and dorsoventral regulation of Phox2b transcription in cranial motoneuron progenitors by homeodomain proteins. *Development*. 131(16):4071-83.
- Scott MM, Wylie CJ, Lerch JK, Murphy R, Lobur, K, Herlitze S, Jiang W, Conlon RA, Strowbridge BW, Deneris ES. 2005. A genetic approach to access serotonin neurons for in vivo and in vitro studies. *Proc. Natl. Acad. Sci. U S A*. 102(45):16472-16477.
- Setini A, Pierucci F, Senatori O, Nicotra A. 2005. Molecular characterization of monoamine oxidase in zebrafish (*Danio rerio*). *Comp Biochem Physiol B Biochem Mol Biol*. 2005.140(1):153-61.
- Sheng Y, Wang L, Liu XS, Montplaisir V, Tiberi M, Baltz JM, Liu XJ. 2005. A serotonin receptor antagonist induces oocyte maturation in both frogs and mice: evidence that the same G protein-coupled receptor is responsible for maintaining meiosis arrest in both species. *J Cell Physiol*. 202(3):777-86.
- Shilling F, Mandel G, Jaffe LA. 1990. Activation by serotonin of starfish eggs expressing the rat serotonin 1c receptor. *Cell Regul*.1(6):465-9.
- Shuey DL, Sadler TW, Tamir H, Lauder JM. 1993. Serotonin and morphogenesis. Transient expression of serotonin uptake and binding protein during craniofacial morphogenesis in the mouse. *Anat Embryol (Berl)*.187(1):75-85.
- Smidt MP, Asbreuk CH, Cox JJ, Chen H, Johnson RL, Burbach JP. 2000. A second independent pathway for development of mesencephalic dopaminergic neurons requires Lmx1b. *Nat. Neurosci*. 3:337-341.
- Smith A, Graham A. 2001. Restricting Bmp-4 mediated apoptosis in hindbrain neural crest. *Dev Dyn*. 220(3):276-83.
- Smith TJ. 1990. Phylogenetic distribution and function of arylalkylamine N-acetyltransferase. *Bioessays*. 12(1):30-3.
- Sodhi MS and Sanders-Bush E. 2004. Serotonin and brain development. *Int. Rev. Neurobiol*. 59:111-174
- Stewart RA, Arduini BL, Berghmans S, George RE, Kanki JP, Henion PD, Look AT. 2006. Zebrafish foxd3 is selectively required for neural crest specification, migration and survival. *Dev Biol*. 292(1):174-88.
- Stornetta RL, Moreira TS, Takakura AC, Kang BJ, Chang DA, West GH, Brunet JF, Mulkey DK, Bayliss DA, Guyenet PG. 2006. Expression of Phox2b by brainstem

- neurons involved in chemosensory integration in the adult rat. *J Neurosci.* 26(40):10305-14.
- Sze JY, Victor M, Loer C, Shi Y, Ruvkun G.. 2000. Food and metabolic signalling defects in a *Caenorhabditis elegans* serotonin-synthesis mutant. *Nature.* 403(6769):560-4.
  - Teraoka H, Russell C, Regan J, Chandrasekhar A, Concha ML, Yokoyama R, Higashi K, Take-Uchi M, Dong W, Hiraga T, Holder N, Wilson SW. 2004 . Hedgehog and Fgf signaling pathways regulate the development of tphR-expressing serotonergic raphe neurons in zebrafish embryos. *J Neurobiol.* 60(3):275-88.
  - Thomas DR, Hagen JJ. 2004. 5-HT<sub>7</sub> receptors. *Curr Drug Targets CNS Neurol Disord.* 3(1):81-90.
  - Thomas T, Kurihara H, Yamagishi H, Kurihara Y, Yazaki Y, Olson EN, Srivastava D. 1998. A signaling cascade involving endothelin-1, dHAND and msx1 regulates development of neural-crest-derived branchial arch mesenchyme. *Development.* 125(16):3005-14.
  - Trainor PA, and Krumlauf R. 2000. Patterning the cranial neural crest: hindbrain segmentation and Hox gene plasticity. *Nat Rev Neurosci.* 1(2):116-2
  - Twarog BM, and Page ICH. 1953. Serotonin content of some mammalian tissues and urine and a method for its determination. *Am J Physiol.* 175(1):157-61.
  - Unsicker K, Huber K, Schutz G, Kalcheim C. 2005. The chromaffin cell and its development. *Neurochem Res.* 30(6-7):921-5. Review.
  - van Doorninck JH, van Der WJ, Karis A, Goedknecht E, Engel JD, Coesmans M, Rutteman M, Grosveld F, De Zeeuw CI. 1999. GATA-3 is involved in the development of serotonergic neurons in the caudal raphe nuclei. *J. Neurosci.* 19. RC12.
  - Vanderlaan G, Tyurina OV, Karlstrom RO, Chandrasekhar A. 2005. Gli function is essential for motor neuron induction in zebrafish. *Dev Biol.* 282(2):550-70.
  - Veitch E, Begbie J, Schilling TF, Smith MM, Graham A. 1999. Pharyngeal arch patterning in the absence of neural crest. *Curr Biol.* 9:1481-1484.
  - Vesela J, Rehak P, Mihalik J, Czikkova S, Pokorny J, Koppel J. 2003. Expression of serotonin receptors in mouse oocytes and preimplantation embryos. *Physiol Res.* 52(2):223-8.

- Vitalis T, Alvarez C, Chen K, Shih JC, Gaspar P, Cases O. 2003. Developmental expression pattern of monoamine oxidases in sensory organs and neural crest derivatives. *J. Comp. Neurol.* 464(3):392-403.
- Walker MB, Miller CT, Swartz ME, Eberhart JK, Kimmel CB. 2007. phospholipase C, beta 3 is required for Endothelin1 regulation of pharyngeal arch patterning in zebrafish. *Dev Biol.* 304(1):194-207.
- Walther DJ, and Bader M. 2003. A unique central tryptophan hydroxylase isoform. *Biochem Pharmacol.* 66(9):1673-80.
- Walther DJ, Peter JU, Bader M. 2002. 7-Hydroxytryptophan, a novel, specific, cytotoxic agent for carcinoids and other serotonin-producing tumors. *Cancer.* 94(12):3135-40.
- Walther DJ, Peter JU, Bashammakh S, Hortnagl H, Voits M, Fink H, Bader M. (2003). Synthesis of serotonin by a second tryptophan hydroxylase isoform. *Science.* 299(5603):76.
- Wang GY, Lee CG, Lee EJ. 2004. Genetic variability of arylalkylamine-N-acetyltransferase (AA-NAT) gene and human sleep/wake pattern. *Chronobiol Int.* 21(2):229-37.
- Wang Y, Takai R, Yoshioka H, Shirabe K. 2006; Characterization and expression of serotonin transporter genes in zebrafish. *Tohoku J Exp Med.* 208(3):267-74.
- Westbroek I, van der Plas A, de Rooji KE, Klein-Nulend J, Nijweide PJ. 2001. Expression of serotonin receptors in bone. *J Biol Chem.* 276(31):28961-8
- Westerfield, M. (2000). *The zebrafish book. A guide for the laboratory use of zebrafish (Danio rerio).* 4th ed., Univ. of Oregon Press, Eugene
- Whitlock KE, Smith KM, Kim H, Harden MV. 2005. A role for foxd3 and sox10 in the differentiation of gonadotropin-releasing hormone (GnRH) cells in the zebrafish *Danio rerio*. *Development.* 132(24):5491-502.
- Woodruff RH, Tekki-Kessaris N, Stiles CD, Rowitch DH, Richardson WD. 2001. Oligodendrocyte development in the spinal cord and telencephalon: common themes and new perspectives. *Int J Dev Neurosci.* 19(4):379-85.
- Woolley ML, Marsden CA, Fone KC. 2004. 5-HT<sub>6</sub> receptors. *Curr Drug Targets CNS Neurol Disord.* 3(1):59-79.
- Xiao H, Hapiak VM, Smith KA, Lin L, Hobson RJ, Plenefisch J, Komuniecki R. 2006. SER-1, a *Caenorhabditis elegans* 5-HT<sub>2</sub>-like receptor, and a multi-PDZ domain

containing protein (MPZ-1) interact in vulval muscle to facilitate serotonin-stimulated egg-laying. *Dev Biol.* 298(2):379-91.

- Yamagishi C, Yamagishi H, Maeda J, Tsuchihashi T, Ivey K, Hu T, Srivastava D..2006. Sonic hedgehog is essential for first pharyngeal arch development. *Pediatr Res.* 59(3):349-54.
- Yavarone MS, Shuey DL, Tamir H, Sadler TW, Lauder JM. 1993. Serotonin and cardiac morphogenesis in the mouse embryo. *Teratology.* 47(6):573-84.
- Ye W, Bouchard M, Stone D, Liu X, Vella F, Lee J, Nakamura H, Ang SL, Busslinger M and Rosenthal A. 2001. Distinct regulators control the expression of the mid-hindbrain organizer signal FGF8. *Nat. Neurosci.* 4:1175-1181.
- Ye W, Shimamura K, Rubenstein JL, Hynes MA and Rosenthal A.1998. FGF and Shh signals control dopaminergic and serotonergic cell fate in the anterior neural plate. *Cell.* 93: 755-766.
- Yohrling GJ, Jiang GC, Mockus SM, Vrana KE, 2000. Intersubunit binding domains within tyrosine hydroxylase and tryptophan hydroxylase. *J. Neurosci. Res.* 61:313–320.
- Zacccone G, Mauceri A, Fasulo S, Ainis L, Lo Cascio P, Ricca MB. 1996. Localization of immunoreactive endothelin in the neuroendocrine cells of fish gill. *Neuropeptides.* 30(1):53-7.
- Zhou QY, Quaife CJ, Palmiter RD. 1995. Targeted disruption of the tyrosine hydroxylase gene reveals that catecholamines are required for mouse fetal development. *Nature.* 374:640–643.