

## 9. Literatur

- Anderson, T. J., M. Klugmann, et al. (1999). "Distinct phenotypes associated with increasing dosage of the PLP gene: implications for CMT1A due to PMP22 gene duplication." Ann N Y Acad Sci **883**: 234-46.
- Bass, B. L., K. Nishikura, et al. (1997). "A standardized nomenclature for adenosine deaminases that act on RNA." Rna **3**(9): 947-9.
- Bass, B. L. and H. Weintraub (1988). "An unwinding activity that covalently modifies its double-stranded RNA substrate." Cell **55**(6): 1089-98.
- Beaudoing, E., S. Freier, et al. (2000). "Patterns of variant polyadenylation signal usage in human genes." Genome Res **10**(7): 1001-10.
- Black, D. L. (2000). "Protein diversity from alternative splicing: a challenge for bioinformatics and post-genome biology." Cell **103**(3): 367-70.
- Black, D. L. (2003). "Mechanisms of Alternative Pre-Messenger RNA Splicing." Annu Rev Biochem.
- Bork, P., L. Holm, et al. (1994). "The immunoglobulin fold. Structural classification, sequence patterns and common core." J Mol Biol **242**(4): 309-20.
- Buchner, G., E. Montini, et al. (1999). "MID2, a homologue of the Opitz syndrome gene MID1: similarities in subcellular localization and differences in expression during development." Hum Mol Genet **8**(8): 1397-407.
- Celotto, A. M. and B. R. Graveley (2001). "Alternative splicing of the Drosophila Dscam pre-mRNA is both temporally and spatially regulated." Genetics **159**(2): 599-608.
- Cox, T. C., L. R. Allen, et al. (2000). "New mutations in MID1 provide support for loss of function as the cause of X-linked Opitz syndrome." Hum Mol Genet **9**(17): 2553-2562.
- Dal Zotto, L., N. A. Quaderi, et al. (1998). "The mouse Mid1 gene: implications for the pathogenesis of Opitz syndrome and the evolution of the mammalian pseudoautosomal region." Hum Mol Genet **7**(3): 489-99.
- Docherty, K., R. J. Carroll, et al. (1982). "Conversion of proinsulin to insulin: involvement of a 31,500 molecular weight thiol protease." Proc Natl Acad Sci U S A **79**(15): 4613-7.
- Dufour, E. (1988). "Sequence homologies, hydrophobic profiles and secondary structures of cathepsins B, H and L: comparison with papain and actinidin." Biochimie **70**(10): 1335-42.
- Erdmann, V. A., M. Z. Barciszewska, et al. (2001). "Regulatory RNAs." Cell Mol Life Sci **58**(7): 960-77.

- Frischmeyer, P. A. and H. C. Dietz (1999). "Nonsense-mediated mRNA decay in health and disease." Hum Mol Genet **8**(10): 1893-900.
- Gaudenz, K., E. Roessler, et al. (1998). "Opitz G/BBB syndrome in Xp22: mutations in the MID1 gene cluster in the carboxy-terminal domain." Am J Hum Genet **63**(3): 703-10.
- Gonzalez, C. I., M. J. Ruiz-Echevarria, et al. (2000). "The yeast hnRNP-like protein Hrp1/Nab4 marks a transcript for nonsense-mediated mRNA decay." Mol Cell **5**(3): 489-99.
- Graveley, B. R. (2001). "Alternative splicing: increasing diversity in the proteomic world." Trends Genet **17**(2): 100-7.
- Hentze, M. W. and A. E. Kulozik (1999). "A perfect message: RNA surveillance and nonsense-mediated decay." Cell **96**(3): 307-10.
- Jackson, I. J. (1991). "A reappraisal of non-consensus mRNA splice sites." Nucleic Acids Res **19**(14): 3795-8.
- Kalvakolanu, D. V. and E. C. Borden (1996). "An overview of the interferon system: signal transduction and mechanisms of action." Cancer Invest **14**(1): 25-53.
- Kerppola, T. K. and C. M. Kane (1988). "Intrinsic sites of transcription termination and pausing in the c-myc gene." Mol Cell Biol **8**(10): 4389-94.
- Koop, B. F. and J. H. Nadeau (1996). "Pufferfish and new paradigm for comparative genome analysis." Proc Natl Acad Sci U S A **93**(4): 1363-5.
- Kozak, M. (1996). "Interpreting cDNA sequences: some insights from studies on translation." Mamm Genome **7**(8): 563-74.
- Kozak, M. (2002). "Pushing the limits of the scanning mechanism for initiation of translation." Gene **299**(1-2): 1-34.
- Kriventseva, E. V., I. Koch, et al. (2003). "Increase of functional diversity by alternative splicing." Trends Genet **19**(3): 124-8.
- Krystal, G. W., B. C. Armstrong, et al. (1990). "N-myc mRNA forms an RNA-RNA duplex with endogenous antisense transcripts." Mol Cell Biol **10**(8): 4180-91.
- Kumar, M. and G. G. Carmichael (1998). "Antisense RNA: function and fate of duplex RNA in cells of higher eukaryotes." Microbiol Mol Biol Rev **62**(4): 1415-34.
- Ladd, A. N. and T. A. Cooper (2002). "Finding signals that regulate alternative splicing in the post-genomic era." Genome Biol **3**(11): reviews0008.
- Lamba, J. K., M. Adachi, et al. (2003). "Nonsense mediated decay downregulates conserved alternatively spliced ABCC4 transcripts bearing nonsense codons." Hum Mol Genet **12**(2): 99-109.

- Lander, E. S., L. M. Linton, et al. (2001). "Initial sequencing and analysis of the human genome." Nature **409**(6822): 860-921.
- Landry, J. R. and D. L. Mager (2002). "Widely spaced alternative promoters, conserved between human and rodent, control expression of the Opitz syndrome gene MID1." Genomics **80**(5): 499-508.
- Lee, R. C., R. L. Feinbaum, et al. (1993). "The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*." Cell **75**(5): 843-54.
- Lewis, B. P., R. E. Green, et al. (2003). "Evidence for the widespread coupling of alternative splicing and nonsense-mediated mRNA decay in humans." Proc Natl Acad Sci U S A **100**(1): 189-92.
- Liu, J., T. D. Prickett, et al. (2001). "Phosphorylation and microtubule association of the Opitz syndrome protein mid-1 is regulated by protein phosphatase 2A via binding to the regulatory subunit alpha 4." Proc Natl Acad Sci U S A **98**(12): 6650-5.
- Lupski, J. R., C. A. Garcia, et al. (1991). "Discordance of muscular dystrophy in monozygotic female twins: evidence supporting asymmetric splitting of the inner cell mass in a manifesting carrier of Duchenne dystrophy." Am J Med Genet **40**(3): 354-64.
- Lykke-Andersen, J., M. D. Shu, et al. (2000). "Human Upf proteins target an mRNA for nonsense-mediated decay when bound downstream of a termination codon." Cell **103**(7): 1121-31.
- Marcus, P. I. (1981). "The interferon inducer moiety of viruses: a single molecule of dsRNA." Tex Rep Biol Med **41**: 70-5.
- Mitrovich, Q. M. and P. Anderson (2000). "Unproductively spliced ribosomal protein mRNAs are natural targets of mRNA surveillance in *C. elegans*." Genes Dev **14**(17): 2173-84.
- Modrek, B. and C. Lee (2002). "A genomic view of alternative splicing." Nat Genet **30**(1): 13-9.
- Modrek, B., A. Resch, et al. (2001). "Genome-wide detection of alternative splicing in expressed sequences of human genes." Nucleic Acids Res **29**(13): 2850-9.
- Mount, S. M. (2000). "Genomic sequence, splicing, and gene annotation." Am J Hum Genet **67**(4): 788-92.
- Nepveu, A. and K. B. Marcu (1986). "Intragenic pausing and anti-sense transcription within the murine *c-myc* locus." Embo J **5**(11): 2859-65.
- Nishikura, K. (1992). "Modulation of double-stranded RNAs in vivo by RNA duplex unwindase." Ann N Y Acad Sci **660**: 240-50.
- Opitz (1969). Birth Defects Orig. Art. Ser. **V**(2): 86-94.

- Peng, H., I. Feldman, et al. (2002). "Hetero-oligomerization among the TIF family of RBCC/TRIM domain-containing nuclear cofactors: a potential mechanism for regulating the switch between coactivation and corepression." *J Mol Biol* **320**(3): 629-44.
- Perry, J., K. M. Short, et al. (1999). "FX/Y2/MID2, a gene related to the X-linked Opitz syndrome gene FX/Y/MID1, maps to Xq22 and encodes a FNIII domain-containing protein that associates with microtubules." *Genomics* **62**(3): 385-94.
- Pickart, C. M. (2001). "Mechanisms underlying ubiquitination." *Annu Rev Biochem* **70**: 503-33.
- Pousi, B., T. Hautala, et al. (1994). "Alu-Alu recombination results in a duplication of seven exons in the lysyl hydroxylase gene in a patient with the type VI variant of Ehlers-Danlos syndrome." *Am J Hum Genet* **55**(5): 899-906.
- Quaderi, N. A., S. Schweiger, et al. (1997). "Opitz G/BBB syndrome, a defect of midline development, is due to mutations in a new RING finger gene on Xp22." *Nat Genet* **17**(3): 285-91.
- Ratnasinghe, D., J. M. Phang, et al. (1998). "Differential expression and activity of phosphatases and protein kinases in adriamycin sensitive and resistant human breast cancer MCF-7 cells." *Int J Oncol* **13**(1): 79-84.
- Roberts, G. C. and C. W. Smith (2002). "Alternative splicing: combinatorial output from the genome." *Curr Opin Chem Biol* **6**(3): 375-83.
- Schmucker, D., J. C. Clemens, et al. (2000). "Drosophila Dscam is an axon guidance receptor exhibiting extraordinary molecular diversity." *Cell* **101**(6): 671-84.
- Schweiger, S., J. Foerster, et al. (1999). "The Opitz syndrome gene product, MID1, associates with microtubules." *Proc Natl Acad Sci U S A* **96**(6): 2794-9.
- Schweiger, S. and R. Schneider (2003). "The MID1/PP2A complex: a key to the pathogenesis of Opitz BBB/G syndrome." *Bioessays* **25**(4): 356-66.
- Senapathy, P., M. B. Shapiro, et al. (1990). "Splice junctions, branch point sites, and exons: sequence statistics, identification, and applications to genome project." *Methods Enzymol* **183**: 252-78.
- Shendure, J. and G. M. Church (2002). "Computational discovery of sense-antisense transcription in the human and mouse genomes." *Genome Biol* **3**(9): RESEARCH0044.
- Short, K. M., B. Hopwood, et al. (2002). "MID1 and MID2 homo- and heterodimerise to tether the rapamycin-sensitive PP2A regulatory subunit, alpha 4, to microtubules:

- implications for the clinical variability of X-linked Opitz GBBB syndrome and other developmental disorders." BMC Cell Biol **3**(1): 1.
- Siebenlist, U., P. Bressler, et al. (1988). "Two distinct mechanisms of transcriptional control operate on c-myc during differentiation of HL60 cells." Mol Cell Biol **8**(2): 867-74.
- Slavov, D. and K. Gardiner (2002). "Phylogenetic comparison of the pre-mRNA adenosine deaminase ADAR2 genes and transcripts: conservation and diversity in editing site sequence and alternative splicing patterns." Gene **299**(1-2): 83-94.
- Sleutels, F., R. Zwart, et al. (2002). "The non-coding Air RNA is required for silencing autosomal imprinted genes." Nature **415**(6873): 810-3.
- Sontag, E. (2001). "Protein phosphatase 2A: the Trojan Horse of cellular signaling." Cell Signal **13**(1): 7-16.
- Stamm, S. (2002). "Signals and their transduction pathways regulating alternative splicing: a new dimension of the human genome." Hum Mol Genet **11**(20): 2409-16.
- Tone, M., Y. Tone, et al. (2001). "Regulation of CD40 function by its isoforms generated through alternative splicing." Proc Natl Acad Sci U S A **98**(4): 1751-6.
- Trockenbacher, A., V. Suckow, et al. (2001). "MID1, mutated in Opitz syndrome, encodes an ubiquitin ligase that targets phosphatase 2A for degradation." Nat Genet **29**(3): 287-94.
- Veldhoen, N., S. Metcalfe, et al. (1999). "A novel exon within the mdm2 gene modulates translation initiation in vitro and disrupts the p53-binding domain of mdm2 protein." Oncogene **18**(50): 7026-33.
- Wagner, R. W., J. E. Smith, et al. (1989). "A double-stranded RNA unwinding activity introduces structural alterations by means of adenosine to inosine conversions in mammalian cells and *Xenopus* eggs." Proc Natl Acad Sci U S A **86**(8): 2647-51.
- Wightman, B., I. Ha, et al. (1993). "Posttranscriptional regulation of the heterochronic gene *lin-14* by *lin-4* mediates temporal pattern formation in *C. elegans*." Cell **75**(5): 855-62.
- Williams, T. and M. Fried (1986). "A mouse locus at which transcription from both DNA strands produces mRNAs complementary at their 3' ends." Nature **322**(6076): 275-9.
- Wolfsberg, T. G., J. McEntyre, et al. (2001). "Guide to the draft human genome." Nature **409**(6822): 824-6.