

## 6 Literaturverzeichnis

1. **Akça** O, Lenhardt R, Fleischmann E, Treschan T, Greif R, Fleischhackl R, Kimberger O, Kurz A, Sessler DI (2004). Nitrous oxide increases the incidence of bowel distension in patients undergoing elective colon resection. *Acta Anaesthesiol Scand.* Aug;48(7):894-8.
2. **Aldrete** JA., Kroulik D (1970): A postanesthetic recovery score. *Anesth Analg*, 49 (6): 924-934.
3. **Andrews** PL (1992). Physiology of nausea and vomiting. *Br J Anaesth.*;69(7 Suppl 1):2S-19S.
4. **Apfel** CC, Greim CA, Haubitz I, Grundt D, Goepfert C, Sefrin P, Roewer N (1998). The discriminating power of a risk score for postoperative vomiting in adults undergoing various types of surgery. *Acta Anaesthesiol Scand.* May;42(5):502-9.
5. **Apfel** CC, Korttila K, Abdalla M, Kerger H, Turan A, Vedder I, Zernak C, Danner K, Jokela R, Pocock SJ, Trenkler S, Kredel M, Biedler A, Sessler DI, Roewer N; IMPACT Investigators (2004). A factorial trial of six interventions for the prevention of postoperative nausea and vomiting. *N Engl J Med.* Jun 10;350(24):2441-51.
6. **Apfel** CC, Kranke P, Katz MH, Goepfert C, Papenfuss T, Rauch S, Heineck R, Greim CA, Roewer N (2002). Volatile anaesthetics may be the main cause of early but not delayed postoperative vomiting: a randomized controlled trial of factorial design. *Br J Anaesth.* May;88(5):659-68.
7. **Apfel** CC, Läärä E, Koivuranta M, Greim CA, Roewer N. (1999). A simplified risk score for predicting postoperative nausea and vomiting: conclusions from cross-validations between two centers. *Anesthesiology.* Sep;91(3):693-700.
8. **Apfel** CC, Roewer N, Korttila K.(2002). How to study postoperative nausea and vomiting. *Acta Anaesthesiol Scand.* Sep;46(8):921-8. Review.
9. **Arellano** RJ, Pole ML, Rafuse SE, Fletcher M, Saad YG, Friedlander M, Norris A, Chung FF (2000). Omission of nitrous oxide from a propofol-based anesthetic does not affect the recovery of women undergoing outpatient gynecologic surgery. *Anesthesiology.* Aug;93(2):332-9.
10. **Bardenheuer** HJ, Taut F (1997). [Incidence and pathophysiology of postoperative nausea and vomiting] *Anesthesiol Intensivmed Notfallmed Schmerzther.* Oct;32(10):617-9. Review.

11. **Bauer P**, Köhne K (1994). Evaluation of experiments with adaptive interim analyses. *Biometrics*. Dec;50(4):1029-41.
12. **Beattie WS**, Lindblad T, Buckley DN, Forrest JB. (1993). Menstruation increases the risk of nausea and vomiting after laparoscopy. A prospective randomized study. *Anesthesiology*. Feb;78(2):272-6.
13. **Bellville JW** (1961). Postanesthetic nausea and vomiting. *Anesthesiology*; 22: 773-780.
14. **Bellville JW**, Bross ID, Howland WS (1960). Postoperative nausea and vomiting. IV. Factors related to postoperative nausea and vomiting. *Anesthesiology*. Mar-Apr;21:186-93.
15. **Bhandari P**, Bingham S, Andrews PL (1992). The neuropharmacology of loperamide-induced emesis in the ferret: the role of the area postrema, vagus, opiate and 5-HT<sub>3</sub> receptors. *Neuropharmacology*. Aug; 31(8):735-42.
16. **Bonica JJ**, Crepps W, Monk B, Bennett B.(1958): Postanesthetic nausea, retching and vomiting; evaluation of cyclizine (marezine) suppositories for treatment. *Anesthesiology*. 1958 Jul-Aug;19(4):532-40.
17. **Boogaerts JG**, Vanacker E, Seidel L, Albert A, Bardiau FM (2000). Assessment of postoperative nausea using a visual analogue scale. *Acta Anaesthesiol Scand*. Apr;44(4):470-4
18. **Carroll NV**, Miederhoff PA, Cox FM, Hirsch JD (1994). Costs incurred by outpatient surgical centers in managing postoperative nausea and vomiting. *J Clin Anesth*. Sep-Oct;6(5):364-9.
19. **Chapman CR**, Benedetti C (1979). Nitrous oxide effects on cerebral evoked potential to pain: partial reversal with a narcotic antagonist. *Anesthesiology*. Aug;51(2):135-8.
20. **Cohen MM**, Duncan PG, DeBoer DP, Tweed WA (1994). The postoperative interview: assessing risk factors for nausea and vomiting. *Anesth Analg*. Jan;78(1):7-16.
21. **Divatia JV**, Vaidya JS, Badwe RA, Hawaldar RW. (1996). Omission of nitrous oxide during anesthesia reduces the incidence of postoperative nausea and vomiting. A meta-analysis. *Anesthesiology*. Nov;85(5):1055-62.
22. **Eberhart LH**, Morin AM, Georgieff M (2000). [The menstruation cycle in the postoperative phase. Its effect of the incidence of nausea and vomiting] *Anaesthesist*. Jun;49(6):532-5. German

23. **Eger** EI 2nd, Lampe GH, Wauk LZ, Whitendale P, Cahalan MK, Donegan JH. (1990). Clinical pharmacology of nitrous oxide: an argument for its continued use. *Anesth Analg.* Dec;71(6):575-85.
24. **Fang** F, Guo TZ, Davies MF, Maze M. (1997). Opiate receptors in the periaqueductal gray mediate analgesic effect of nitrous oxide in rats. *Eur J Pharmacol.* Oct 8;336(2-3):137-41.
25. **Farrar** JT, Young JP Jr, LaMoreaux L, Werth JL, Poole RM. (2001). Clinical importance of changes in chronic pain intensity measured on an 11-point numerical pain rating scale. *Pain.* Nov;94(2):149-58.
26. **Felts** JA, Poler SM, Spitznagel EL.(1990). Nitrous oxide, nausea, and vomiting after outpatient gynecologic surgery. *J Clin Anesth.* 1990 May-Jun;2(3):168-71.
27. **Finck** AD, Samaniego E, Ngai SH (1995). Nitrous oxide selectively releases Met5-enkephalin and Met5-enkephalin-Arg6-Phe7 into canine third ventricular cerebrospinal fluid. *Anesth Analg.* Apr;80(4):664-70.
28. **Forrest** JB, Beattie WS, Goldsmith CH (1990). Risk factors for nausea and vomiting after general anaesthesia. *Can J Anaesth.* May;37(4 Pt 2):S90.
29. **Forrest** JB, Cahalan MK, Rehder K, Goldsmith CH, Levy WJ, Strunin L, Bota W, Boucek CD, Cucchiara RF, Dhamee S et al (1990). Multicenter study of general anesthesia. II. Results. *Anesthesiology.* Feb;72(2):262-8.
30. **Gan** TJ (2002). Postoperative nausea and vomiting – Can it be eliminated? *JAMA* 287:1233-1236.
31. **Goepfert** C, Apfel CC, Kranke P, Rauch S, Papenfuß T, Sefrin P, Roewer N (1997). Hat der Menstruationszyklus wirklich einen Einfluss auf das postoperative Erbrechen? *AINS*; 33 (Suppl 3):166
32. **Gold** BS, Kitz DS, Lecky JH, Neuhaus JM (1969). Unanticipated admission to the hospital following ambulatory surgery. *JAMA.* 262:3008-3010
33. **Hapfelmeier** G, Haseneder R, Kochs E, Beyerle M, Zieglgansberger W. (2001). Coadministered nitrous oxide enhances the effect of isoflurane on GABAergic transmission by an increase in open-channel block. *J Pharmacol Exp Ther.* Jul;298(1):201-8.
34. **Hartung** J (1996). Twenty-four of twenty-seven studies show a greater incidence of emesis associated with nitrous oxide than with alternative

- anesthetics. *Anesth Analg.* Jul;83(1):114-6. Erratum in: *Anesth Analg* 1997 Nov;85(5):986. *Anesth Analg* 1998 Nov;87(5):1215.
35. **Hirsch J** (1994). Impact of postoperative nausea and vomiting in the surgical setting. *Anaesthesia.* Jan;49 Suppl:30-3. Review.
36. **Honkavaara P**, Lehtinen AM, Hovorka J, Korttila K (1991). Nausea and vomiting after gynaecological laparoscopy depends upon the phase of the menstrual cycle. *Can J Anaesth.* Oct;38(7):876-9.
37. **Jevtovic-Todorovic V**, Todorovic SM, Mennerick S, Powell S, Dikranian K, Benshoff N, Zorumski CF, Olney JW. (1998). Nitrous oxide (laughing gas) is an NMDA antagonist, neuroprotectant and neurotoxin. *Nat Med.* Apr;4(4):460-3.
38. **Jin F**, Norris A, Chung F, Ganeshram T (1998). Should adult patients drink fluids before discharge from ambulatory surgery? *Anesth Analg.* Aug;87(2):306-11.
39. **Kingery WS**, Sawamura S, Agashe GS, Davies MF, Clark JD, Zimmer A (2001). Enkephalin release and opioid receptor activation does not mediate the antinociceptive or sedative/hypnotic effects of nitrous oxide. *Eur J Pharmacol.* Sep 7;427(1):27-35.
40. **Koivuranta M**, Läärä E, Snare L, Alahuhta S (1997). A survey of postoperative nausea and vomiting. *Anaesthesia.* May;52(5):443-9.
41. **Korttila K**, Hovorka J, Erkola O (1987). Nitrous oxide does not increase the incidence of nausea and vomiting after isoflurane anesthesia. *Anesth Analg.* Aug;66(8):761-5.
42. **Kovac AL** (2000). Prevention and treatment of postoperative nausea and vomiting. *Drugs.* Feb;59(2):213-43. Review.
43. **Kranke P**, Apfel CC, Papenfuss T, Rauch S, Löbmann U, Rübsam B, Greim CA, Roewer N. (2001). An increased body mass index is no risk factor for postoperative nausea and vomiting. A systematic review and results of original data. *Acta Anaesthesiol Scand.* Feb;45(2):160-6.
44. **Kranke P**, Eberhart LH, Gan TJ, Roewer N, Tramer MR (2007). Algorithms for the prevention of postoperative nausea and vomiting: an efficacy and efficiency simulation. *Eur J Anaesthesiol.* Jul 3; 1-12.
45. **Lerman J** (1992). Surgical and patient factors involved in postoperative nausea and vomiting. *Br J Anaesth.*;69(7 Suppl 1):24S-32S.

46. **Lindgren** L, Koivusalo AM, Kellokumpu I. (1995). Conventional pneumoperitoneum compared with abdominal wall lift for laparoscopic cholecystectomy. *Br J Anaesth.* Nov;75(5):567-72.
47. **Lonie** DS, Harper JN (1986). Nitrous oxide anaesthesia and vomiting. The effect of nitrous oxide anaesthesia on the incidence of vomiting following gynaecological laparoscopy. *Anaesthesia.* 1986 Jul;41(7):703-7. Erratum in: *Anaesthesia* 1986 Oct;41(10):1083.
48. **Madej** TH, Simpson KH (1986). Comparison of the use of domperidone, droperidol and metoclopramide in the prevention of nausea and vomiting following gynaecological surgery in day cases. *Br J Anaesth.* Aug;58(8):879-83.
49. **Meeks** GR, Waller GA, Meydrech EF, Flautt FH Jr (1992). Unscheduled hospital admission following ambulatory gynecologic surgery. *Obstet Gynecol.* Sep;80(3 Pt 1):446-50.
50. **Mennerick** S, Jevtovic-Todorovic V, Todorovic SM, Shen W, Olney JW, Zorumski CF (1998). Effect of nitrous oxide on excitatory and inhibitory synaptic transmission in hippocampal cultures. *J Neurosci.* Dec 1;18(23):9716-26..
51. **Muir** JJ, Warner MA, Offord KP, Buck CF, Harper JV, Kunkel SE (1987). Role of nitrous oxide and other factors in postoperative nausea and vomiting: a randomized and blinded prospective study. *Anesthesiology.* Apr;66(4):513-8.
52. **Murakawa** M, Adachi T, Nakao S, Seo N, Shingu K, Mori K (1994). Activation of the cortical and medullary dopaminergic systems by nitrous oxide in rats: a possible neurochemical basis for psychotropic effects and postanesthetic nausea and vomiting. *Anesth Analg.* Feb;78(2):376-81.
53. **Myles** PS, Leslie K, Chan MT, Forbes A, Paech MJ, Peyton P, Silbert BS, Pascoe E; ENIGMA Trial Group (2007). Avoidance of nitrous oxide for patients undergoing major surgery: a randomized controlled trial. *Anesthesiology.* Aug;107(2):221-31.
54. **Nader** ND, Simpson G, Reedy RL (2004). Middle ear pressure changes after nitrous oxide anesthesia and its effect on postoperative nausea and vomiting. *Laryngoscope.* May;114(5):883-6
55. **Neuhäuser** M (2004). Einige neue Methoden aus der Biometrie. *TumorDiagn u Ther.* 25: 160-163

56. **Orhan-Sungur** M, Apfel C, Akca O (2005). Effects of nitrous oxide on intraoperative bowel distension. *Curr Opin Anaesthesiol.* Dec;18(6):620-4.
57. **Palazzo** M, Evans R (1993). Logistic regression analysis of fixed patient factors for postoperative sickness: a model for risk assessment. *Br J Anaesth.* Feb;70(2):135-40.
58. **Rung** GW, Claybon L, Hord A, Patel C, Kallgren M, Koppel J, Bennedetti C, Creed M Asgharian A, Bryson J (1997). Intravenous Ondansetron for postsurgical opioid-induced nausea and vomiting. *Anesth Analg* 84:832-838
59. **Russell** GB, Snider MT, Richard RB, Loomis JL (1990). Hyperbaric nitrous oxide as a sole anesthetic agent in humans. *Anesth Analg.* Mar;70(3):289-95.
60. **Sinclair** DR, Chung F, Mezei G (1999). Can postoperative nausea and vomiting be predicted? *Anesthesiology.* Jul;91(1):109-18.
61. **Sukhani** R, Lurie J, Jabamoni R (1994). Propofol for ambulatory gynecologic laparoscopy: does omission of nitrous oxide alter postoperative emetic sequelae and recovery? *Anesth Analg.* May;78(5):831-5.
62. **Taylor** E, Feinstein R, White PF, Soper N. (1992). Anesthesia for laparoscopic cholecystectomy. Is nitrous oxide contraindicated? *Anesthesiology.* Apr;76(4):541-3.
63. **Thompson** PI, Bingham S, Andrews PL, Patel N, Joel SP, Slevin ML (1992). Morphine 6-glucuronide: a metabolite of morphine with greater emetic potency than morphine in the ferret. *Br J Pharmacol.* May;106(1):3-8.
64. **Tramèr** MR (2007). Clinical pharmacology: Postoperative nausea and vomiting. *Anaesthesist.* Jul;56(7):679-685.
65. **Tramèr** MR, Fuchs-Buder T, Sansonetti A, Rifat K (1997). Low incidence of the oculocardiac reflex and postoperative nausea and vomiting in adults undergoing strabismus surgery. *Can J Anaesth.* Aug;44(8):830-5.
66. **Tramèr** MR, Moore A, McQuay H. (1996). Omitting nitrous oxide in general anaesthesia: meta-analysis of intraoperative awareness and postoperative emesis in randomized controlled trials. *Br J Anaesth.* Feb;76(2):186-93.
67. **Turk** DC, Rudy TE, Sorkin BA (1993). Neglected topics in chronic pain treatment outcome studies: determination of success. *Pain.* 53:3-16.
68. **van den Berg** AA, Halliday E, Lule EK, Baloch MS (1999). The effects of tramadol on postoperative nausea, vomiting and headache after ENT surgery.

- A placebo-controlled comparison with equipotent doses of nalbuphine and pethidine. *Acta Anaesthesiol Scand.* Jan;43(1):28-33
69. **Wang** SC, Glaviano VV (1954). Locus of emetic action of morphine and hydergine in dogs. *J Pharmacol Exp Ther.* Jul;111 (3):329-34.
70. **Watcha** MF, White PF (1995). Post-operative nausea and vomiting: do they matter? *Eur J Anaesthesiol Suppl.* May;10:18-23.
71. **Watcha** MF, White PF (1992). Postoperative nausea and vomiting. Its etiology, treatment, and prevention. *Anesthesiology.* Jul;77(1):162-84. Review.