

6 Literaturverzeichnis

Abele, R. und Tampe, R. (2006) Modulation of the antigen transport machinery TAP by friends and enemies, FEBS Lett. 580, 1156-1163.

Abes, S., Williams, D., Prevot, P., Thierry, A., Gait, M. J. und Lebleu, B. (2006) Endosome trapping limits the efficiency of splicing correction by PNA-oligolysine conjugates, J. Control Release 110, 595-604.

Albertshofer, K., Siwkowski, A. M., Wancewicz, E. V., Esau, C. C., Watanabe, T., Nishihara, K. C., Kinberger, G. A., Malik, L., Eldrup, A. B., Manoharan, M., Geary, R. S., Monia, B. P., Swayze, E. E., Griffey, R. H., Bennett, C. F. und Maier, M. A. (2005) Structure-activity relationship study on a simple cationic peptide motif for cellular delivery of antisense peptide nucleic acid, J. Med. Chem. 48, 6741-6749.

Aldrian-Herrada, G., Desarmenien, M. G., Orcel, H., Boissin-Agasse, L., Mery, J., Brugidou, J. und Rabie, A. (1998) A peptide nucleic acid (PNA) is more rapidly internalized in cultured neurons when coupled to a retro-inverso delivery peptide. The antisense activity depresses the target mRNA and protein in magnocellular oxytocin neurons, Nucleic Acids Res. 26, 4910-4916.

Amador, M., Gonzalez, M. E., Cordova, L. und Perez, N. (1982) Diagnosing and misdiagnosing malnutrition, Acta Paediatr. Acad. Sci. Hung. 23, 391-401.

Anderson, D. C., Nichols, E., Manger, R., Woodle, D., Barry, M. und Fritzberg, A. R. (1993) Tumor cell retention of antibody Fab fragments is enhanced by an attached HIV TAT protein-derived peptide, Biochem. Biophys. Res. Commun. 194, 876-884.

Andrieu, M., Loing, E., Desoutter, J. F., Connan, F., Choppin, J., Gras-Masse, H., Hanau, D., Dautry-Varsat, A., Guillet, J. G. und Hosmalin, A. (2000) Endocytosis of an HIV-derived lipopeptide into human dendritic cells followed by class I-restricted CD8(+) T lymphocyte activation, Eur. J. Immunol. 30, 3256-3265.

Ashkenazi, A. und Dixit, V. M. (1998) Death receptors: signaling and modulation, Science 281, 1305-1308.

Aslam, M. und Dent, A. (1998) The preparation of protein-protein conjugates, in *Bioconjugation*, Grove's Dictionaries Inc, New York, 222-223.

Astriab, F. A., Sergueev, D. S., Fisher, M., Shaw, B. R. und Juliano, R. L. (2000) Antisense inhibition of P-glycoprotein expression using peptide-oligonucleotide conjugates, Biochem. Pharmacol. 60, 83-90.

Astriab-Fisher, A., Sergueev, D., Fisher, M., Shaw, B. R. und Juliano, R. L. (2002) Conjugates of antisense oligonucleotides with the Tat and antennapedia cell-penetrating peptides: Effects on cellular uptake, binding to target sequences, and biologic actions, Pharmaceutical Research 19, 744-754.

Aussedat, B., Sagan, S., Chassaing, G., Bolbach, G. und Burlina, F. (2006) Quantification of the efficiency of cargo delivery by peptidic and pseudo-peptidic Trojan carriers using MALDI-TOF mass spectrometry, Biochim. Biophys. Acta, in Druck.

Barany-Wallje, E., Keller, S., Serowy, S., Geibel, S., Pohl, P., Bienert, M. und Dathe, M. (2005) A critical reassessment of penetratin translocation across lipid membranes, Biophys. J. 89, 2513-2521.

- Beck, G. F., Irwin, W. J., Nicklin, P. L. und Akhtar, S. (1996) Interactions of phosphodiester and phosphorothioate oligonucleotides with intestinal epithelial Caco-2 cells, *Pharm. Res.* 13, 1028-1037.
- Bellet-Amalric, E., Blaudez, D., Desbat, B., Graner, F., Gauthier, F. und Renault, A. (2000) Interaction of the third helix of Antennapedia homeodomain and a phospholipid monolayer, studied by ellipsometry and PM-IRRAS at the air-water interface, *Biochim. Biophys. Acta* 1467, 131-143.
- Belting, M. (2003) Heparan sulfate proteoglycan as a plasma membrane carrier, *Trends. Biochem. Sci.* 28, 145-151.
- Bendifallah N., Rasmussen F.W., Zachar V., Ebbesen P., Nielsen P.E. und Koppelhus U. (2006) Evaluation of Cell-Penetrating Peptides (CPPs) as Vehicles for Intracellular Delivery of Antisense Peptide Nucleic Acids (PNA), *Bioconj. Chem.*, in Druck.
- Bennett, C. F., Chiang, M. Y., Chan, H., Shoemaker, J. E. und Mirabelli, C. K. (1992) Cationic lipids enhance cellular uptake and activity of phosphorothioate antisense oligonucleotides, *Mol. Pharmacol.* 41, 1023-1033.
- Berger, H., Albrecht, E., Wallukat, G. und Bienert, M. (1999) Antagonism by acetyl-RYYRIK-NH₂ of G protein activation in rat brain preparations and of chronotropic effect on rat cardiomyocytes evoked by nociceptin/orphanin FQ, *Br. J. Pharmacol.* 126, 555-558.
- Best, M. D., Tobey, S. L. und Anslyn, E. V. (2003) Abiotic guanidinium containing receptors for anionic species, *Coordination Chemistry Reviews* 240, 3-15.
- Binder, H. und Lindblom, G. (2003) Charge-dependent translocation of the Trojan peptide penetratin across lipid membranes, *Biophys. J.* 85, 982-995.
- Boffa, L. C., Morris, P. L., Carpaneto, E. M., Louissaint, M. und Allfrey, V. G. (1996) Invasion of the CAG triplet repeats by a complementary peptide nucleic acid inhibits transcription of the androgen receptor and TATA-binding protein genes and correlates with refolding of an active nucleosome containing a unique AR gene sequence, *J. Biol. Chem.* 271, 13228-13233.
- Bonham, M. A., Brown, S., Boyd, A. L., Brown, P. H., Bruckenstein, D. A., Hanvey, J. C., Thomson, S. A., Pipe, A., Hassman, F., Bisi, J. E., Froehler, B. C., Matteucci, M. D., Wagner, R. W., Noble, S. A. und Babiss, L. E. (1995) An Assessment of the Antisense Properties of RNase H-Competent and Steric-Blocking Oligomers, *Nucleic Acids Research* 23, 1197-1203.
- Boussif, O., Lezoualc'h, F., Zanta, M. A., Mergny, M. D., Scherman, D., Demeneix, B. und Behr, J. P. (1995) A versatile vector for gene and oligonucleotide transfer into cells in culture and in vivo: polyethylenimine, *Proc. Natl. Acad. Sci. U. S. A* 92, 7297-7301.
- Braasch, D. A. (2002) Synthesis and Purification of Peptide Nucleic Acids, in *Current Protocols in Nucleic Acid Chemistry* (John Wiley & Sons, Ed.), Hoboken, New York, 4.11.1-4.11.18.
- Braasch, D. A. und Corey, D. R. (2001) Synthesis, analysis, purification, and intracellular delivery of peptide nucleic acids, *Methods* 23, 97-107.
- Braasch, D. A. und Corey, D. R. (2002) Novel antisense and peptide nucleic acid strategies for controlling gene expression, *Biochemistry* 41, 4503-4510.
- Bradford, M. M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding, *Anal. Biochem.* 72, 248-254.
- Braun, K., Peschke, P., Pipkorn, R., Lampel, S., Wachsmuth, M., Waldeck, W., Friedrich, E. und Debus, J. (2002) A biological transporter for the delivery of peptide nucleic acids (PNAs) to the nuclear compartment of living cells, *J. Mol. Biol.* 318, 237-243.

- Braun, V. (1975) Covalent lipoprotein from the outer membrane of Escherichia coli, *Biochim. Biophys. Acta* 415, 335-377.
- Brokx, R. D., Bisland, S. K. und Gariepy, J. (2002) Designing peptide-based scaffolds as drug delivery vehicles, *J. Control Release* 78, 115-123.
- Brown, D. A., Kang, S. H., Gryaznov, S. M., DeDionisio, L., Heidenreich, O., Sullivan, S., Xu, X. und Nerenberg, M. I. (1994) Effect of phosphorothioate modification of oligodeoxynucleotides on specific protein binding, *J. Biol. Chem.* 269, 26801-26805.
- Brown, D. A. und London, E. (2000) Structure and function of sphingolipid- and cholesterol-rich membrane rafts, *J. Biol. Chem.* 275, 17221-17224.
- Bullough, P. A., Hughson, F. M., Skehel, J. J. und Wiley, D. C. (1994) Structure of influenza haemagglutinin at the pH of membrane fusion, *Nature* 371, 37-43.
- Burlina, F., Sagan, S., Bolbach, G. und Chassaing, G. (2005) Quantification of the Cellular Uptake of Cell-Penetrating Peptides by MALDI-TOF Mass Spectrometry, *Angew. Chem. Int. Ed Engl.* 44, 4244-4247.
- Burton, P. S., Conradi, R. A., Hilgers A.R., Ho N.F.H. und Maggiora L.L. (1992) The relationship between peptide structure and transport across epithelial cell monolayers., *J. Control Release* 19, 87-98.
- Campbell, J. M., Bacon, T. A. und Wickstrom, E. (1990) Oligodeoxynucleoside phosphorothioate stability in subcellular extracts, culture media, sera and cerebrospinal fluid, *J. Biochem. Biophys. Methods* 20, 259-267.
- Capecchi, M. R. (1980) High efficiency transformation by direct microinjection of DNA into cultured mammalian cells, *Cell* 22, 479-488.
- Caron, N. J., Quenneville, S. P. und Tremblay, J. P. (2004) Endosome disruption enhances the functional nuclear delivery of Tat-fusion proteins, *Biochem. Biophys. Res. Commun.* 319, 12-20.
- Carpino, L. A. (1993) 1-Hydroxy-7-Azabenzotriazole - An Efficient Peptide Coupling Additive, *Journal Of The American Chemical Society* 115, 4397-4398.
- Carpino, L. A., Imazumi, H., El Faham, A., Ferrer, F. J., Zhang, C. W., Lee, Y. S., Foxman, B. M., Henklein, P., Hanay, C., Mugge, C., Wenschuh, H., Klose, K., Beyermann, M. und Bienert, M. (2002) The uronium/guanidinium peptide coupling reagents: Finally the true uronium salts, *Angewandte Chemie-International Edition* 41, 442-445.
- Cartegni, L., Chew, S. L. und Krainer, A. R. (2002) Listening to silence and understanding nonsense: Exonic mutations that affect splicing, *Nature Reviews Genetics* 3, 285-298.
- Cech, T. R., Zaug, A. J. und Grabowski, P. J. (1981) In vitro splicing of the ribosomal RNA precursor of Tetrahymena: involvement of a guanosine nucleotide in the excision of the intervening sequence, *Cell* 27, 487-496.
- Chaubey, B., Tripathi, S., Ganguly, S., Harris, D., Casale, R. A. und Pandey, V. N. (2005) A PNA-transportan conjugate targeted to the TAR region of the HIV-1 genome exhibits both antiviral and virucidal properties, *Virology* 331, 418-428.
- Chen, H. C., Brown, J. H., Morell, J. L. und Huang, C. M. (1988) Synthetic magainin analogues with improved antimicrobial activity, *FEBS Lett.* 236, 462-466.

- Chiu, Y. L., Ali, A., Chu, C. Y., Cao, H. und Rana, T. M. (2004) Visualizing a correlation between siRNA localization, cellular uptake, and RNAi in living cells, *Chem. Biol.* 11, 1165-1175.
- Christensen, L., Fitzpatrick, R., Gildea, B., Petersen, K. H., Hansen, H. F., Koch, T., Egholm, M., Buchardt, O., Nielsen, P. E., Coull, J. und et. a. (1995) Solid-phase synthesis of peptide nucleic acids, *J. Pept. Sci.* 1, 175-183.
- Conner, S. D. und Schmid, S. L. (2003) Regulated portals of entry into the cell, *Nature* 422, 37-44.
- Console, S., Marty, C., Garcia-Echeverria, C., Schwendener, R. und Ballmer-Hofer, K. (2003) Antennapedia and HIV transactivator of transcription (TAT) "protein transduction domains" promote endocytosis of high molecular weight cargo upon binding to cell surface glycosaminoglycans, *J. Biol. Chem.* 278, 35109-35114.
- Coste, J., Lenguyen, D. und Castro, B. (1990) Pybop - A New Peptide Coupling Reagent Devoid of Toxic By-Product, *Tetrahedron Letters* 31, 205-208.
- Crespo, L., Sanclimens, G., Montaner, B., Perez-Tomas, R., Royo, M., Pons, M., Albericio, F. und Giralt, E. (2002) Peptide dendrimers based on polyproline helices, *J. Am. Chem. Soc.* 124, 8876-8883.
- Cutrona, G., Carpaneto, E. M., Ulivi, M., Roncella, S., Landt, O., Ferrarini, M. und Boffa, L. C. (2000) Effects in live cells of a c-myc anti-gene PNA linked to a nuclear localization signal, *Nat. Biotechnol.* 18, 300-303.
- Czajlik, A., Mesko, E., Penke, B. und Perczel, A. (2002) Investigation of penetratin peptides. Part 1. The environment dependent conformational properties of penetratin and two of its derivatives, *J. Pept. Sci.* 8, 151-171.
- Dathe, M., Weprecht, T., Nikolenko, H., Handel, L., Maloy, W. L., MacDonald, D. L., Beyermann, M. und Bienert, M. (1997) Hydrophobicity, hydrophobic moment and angle subtended by charged residues modulate antibacterial and haemolytic activity of amphipathic helical peptides, *FEBS Lett.* 403, 208-212.
- De Clercq, E., Eckstein, E. und Merigan, T. C. (1969) Interferon induction increased through chemical modification of a synthetic polyribonucleotide, *Science* 165, 1137-1139.
- Demidov, V. V., Potaman, V. N., Frank-Kamenetskii, M. D., Egholm, M., Buchard, O., Sonnichsen, S. H. und Nielsen, P. E. (1994) Stability of peptide nucleic acids in human serum and cellular extracts, *Biochem. Pharmacol.* 48, 1310-1313.
- Deres, K., Schild, H., Wiesmuller, K. H., Jung, G. und Rammensee, H. G. (1989) In vivo priming of virus-specific cytotoxic T lymphocytes with synthetic lipopeptide vaccine, *Nature* 342, 561-564.
- Derossi, D., Calvet, S., Trembleau, A., Brunissen, A., Chassaing, G. und Prochiantz, A. (1996) Cell internalization of the third helix of the antennapedia homeodomain is receptor-independent, *J. Biol. Chem.* 271, 18188-18193.
- Derossi, D., Chassaing, G. und Prochiantz, A. (1998) Trojan peptides: the penetratin system for intracellular delivery, *Trends Cell Biol.* 8, 84-87.
- Derossi, D., Joliot, A. H., Chassaing, G. und Prochiantz, A. (1994) The third helix of the Antennapedia homeodomain translocates through biological membranes, *J. Biol. Chem.* 269, 10444-10450.
- Dietz, G. P. und Bahr, M. (2004) Delivery of bioactive molecules into the cell: the Trojan horse approach, *Mol. Cell Neurosci.* 27, 85-131.

- Dobkin, C. und Bank, A. (1985) Reversibility of IVS 2 missplicing in a mutant human beta-globin gene, *J. Biol. Chem.* 260, 16332-16337.
- Dom, G., Shaw-Jackson, C., Matis, C., Bouffioux, O., Picard, J. J., Prochiantz, A., Mingeot-Leclercq, M. P., Brasseur, R. und Rezsöhazy, R. (2003) Cellular uptake of *Antennapedia* Penetratin peptides is a two-step process in which phase transfer precedes a tryptophan-dependent translocation, *Nucleic Acids Res.* 31, 556-561.
- Dominski, Z. und Kole, R. (1993) Restoration of correct splicing in thalassemic pre-mRNA by antisense oligonucleotides, *Proc. Natl. Acad. Sci. U. S. A* 90, 8673-8677.
- Doyle, D. F., Braasch, D. A., Simmons, C. G., Janowski, B. A. und Corey, D. R. (2001) Inhibition of gene expression inside cells by peptide nucleic acids: Effect of mRNA target sequence, mismatched bases, and PNA length, *Biochemistry* 40, 53-64.
- Drin, G., Cottin, S., Blanc, E., Rees, A. R. und Temsamani, J. (2003) Studies on the internalization mechanism of cationic cell-penetrating peptides, *J. Biol. Chem.* 278, 31192-31201.
- Drin, G., Demene, H., Temsamani, J. und Brasseur, R. (2001a) Translocation of the pAntp peptide and its amphipathic analogue AP-2AL, *Biochemistry* 40, 1824-1834.
- Drin, G., Mazel, M., Clair, P., Mathieu, D., Kaczorek, M. und Temsamani, J. (2001b) Physico-chemical requirements for cellular uptake of pAntp peptide - Role of lipid-binding affinity, *Eur. J. Biochem.* 268, 1304-1314.
- Durlu, Y. K. und Tamai, M. (1997) Transplantation of retinal pigment epithelium using viable cryopreserved cells, *Cell Transplant.* 6, 149-162.
- Egholm, M., Buchardt, O., Christensen, L., Behrens, C., Freier, S. M., Driver, D. A., Berg, R. H., Kim, S. K., Norden, B. und Nielsen, P. E. (1993) PNA hybridizes to complementary oligonucleotides obeying the Watson-Crick hydrogen-bonding rules, *Nature* 365, 566-568.
- Egholm, M., Christensen, L., Dueholm, K. L., Buchardt, O., Coull, J. und Nielsen, P. E. (1995) Efficient pH-independent sequence-specific DNA binding by pseudoisocytosine-containing bis-PNA, *Nucleic Acids Res.* 23, 217-222.
- Eichholtz, T., de Bont, D. B., de Widt, J., Liskamp, R. M. und Ploegh, H. L. (1993) A myristoylated pseudosubstrate peptide, a novel protein kinase C inhibitor, *J. Biol. Chem.* 268, 1982-1986.
- El Andaloussi, S., Holm, T. und Langel, U. (2005) Cell-penetrating peptides: mechanisms and applications, *Curr. Pharm. Des* 11, 3597-3611.
- Elmquist, A., Lindgren, M., Bartfai, T. und Langel, U. (2001) VE-cadherin-derived cell-penetrating peptide, pVEC, with carrier functions, *Exp. Cell Res.* 269, 237-244.
- Erbacher, P., Roche, A. C., Monsigny, M. und Midoux, P. (1996) Putative role of chloroquine in gene transfer into a human hepatoma cell line by DNA/lactosylated polylysine complexes, *Exp. Cell Res.* 225, 186-194.
- Falnes, P. O., Wesche, J. und Olsnes, S. (2001) Ability of the Tat basic domain and VP22 to mediate cell binding, but not membrane translocation of the diphtheria toxin A-fragment, *Biochemistry* 40, 4349-4358.
- Farrera-Sinfreu, J., Giralt, E., Castel, S., Albericio, F. und Royo, M. (2005) Cell-penetrating cis-gamma-amino-l-proline-derived peptides, *J. Am. Chem. Soc.* 127, 9459-9468.

- Faulstich, H. und Wieland, T. (1996) New aspects of amanitin and phalloidin poisoning, *Adv. Exp. Med. Biol.* 391, 309-314.
- Fawell, S., Seery, J., Daikh, Y., Moore, C., Chen, L. L., Pepinsky, B. und Barsoum, J. (1994) Tat-mediated delivery of heterologous proteins into cells, *Proc. Natl. Acad. Sci. U. S. A.* 91, 664-668.
- Feener, E. P., Shen, W. C. und Ryser, H. J. (1990) Cleavage of disulfide bonds in endocytosed macromolecules. A processing not associated with lysosomes or endosomes, *J. Biol. Chem.* 265, 18780-18785.
- Ferber, D. (2001) Gene therapy. Safer and virus-free?, *Science* 294, 1638-1642.
- Fernandez-Carneado, J., Kogan, M. J., Castel, S. und Giralt, E. (2004a) Potential peptide carriers: amphipathic proline-rich peptides derived from the N-terminal domain of gamma-zein, *Angew. Chem. Int. Ed Engl.* 43, 1811-1814.
- Fernandez-Carneado, J., Kogan, M. J., Pujals, S. und Giralt, E. (2004b) Amphipathic peptides and drug delivery, *Biopolymers* 76, 196-203.
- Fernandez-Carneado, J., Van Gool, M., Martos, V., Castel, S., Prados, P., de Mendoza, J. und Giralt, E. (2005) Highly efficient, nonpeptidic oligoguanidinium vectors that selectively internalize into mitochondria, *J. Am. Chem. Soc.* 127, 869-874.
- Fields, G. B. und Noble, R. L. (1990) Solid-Phase Peptide-Synthesis Utilizing 9-Fluorenylmethoxycarbonyl Amino-Acids, *International Journal of Peptide and Protein Research* 35, 161-214.
- Finlay, B. B. und Cossart, P. (1997) Exploitation of mammalian host cell functions by bacterial pathogens, *Science* 276, 718-725.
- Fire, A., Xu, S., Montgomery, M. K., Kostas, S. A., Driver, S. E. und Mello, C. C. (1998) Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*, *Nature* 391, 806-811.
- Fischer, P. M., Zhelev, N. Z., Wang, S., Melville, J. E., Fahraeus, R. und Lane, D. P. (2000) Structure-activity relationship of truncated and substituted analogues of the intracellular delivery vector Penetratin, *J. Pept. Res.* 55, 163-172.
- Fischer, R., Fotin-Mleczek, M., Hufnagel, H. und Brock, R. (2005) Break on through to the other side-biophysics and cell biology shed light on cell-penetrating peptides, *Chembiochem.* 6, 2126-2142.
- Fischer, R., Kohler, K., Fotin-Mleczek, M. und Brock, R. (2004) A stepwise dissection of the intracellular fate of cationic cell-penetrating peptides, *J. Biol. Chem.* 279, 12625-12635.
- Fischer, R., Waizenegger, T., Kohler, K. und Brock, R. (2002) A quantitative validation of fluorophore-labelled cell-permeable peptide conjugates: fluorophore and cargo dependence of import, *Biochim. Biophys. Acta* 1564, 365-374.
- Fittipaldi, A., Ferrari, A., Zoppe, M., Arcangeli, C., Pellegrini, V., Beltram, F. und Giacca, M. (2003) Cell membrane lipid rafts mediate caveolar endocytosis of HIV-1 Tat fusion proteins, *J. Biol. Chem.* 278, 34141-34149.
- Flechsler, I., Surovoy, A., Charisse, K., Bayer, E. und Jung, G. (1998) Comparison of antisense vectors and antisense oligonucleotides delivered by means of the new cationic lipids unifectin and maxifectin, *Adv. Exp. Med. Biol.* 451, 469-472.

- Folini, M., Berg, K., Millo, E., Villa, R., Prasmickaite, L., Daidone, M. G., Benatti, U. und Zaffaroni, N. (2003) Photochemical internalization of a peptide nucleic acid targeting the catalytic subunit of human telomerase, *Cancer Res.* 63, 3490-3494.
- Francis, C. L., Ryan, T. A., Jones, B. D., Smith, S. J. und Falkow, S. (1993) Ruffles induced by *Salmonella* and other stimuli direct macropinocytosis of bacteria, *Nature* 364, 639-642.
- Frankel, A. D. und Pabo, C. O. (1988) Cellular uptake of the tat protein from human immunodeficiency virus, *Cell* 55, 1189-1193.
- Fuchs, S. M. und Raines, R. T. (2004) Pathway for polyarginine entry into mammalian cells, *Biochemistry* 43, 2438-2444.
- Futaki, S., Nakase, I., Suzuki, T., Youjun, Z. und Sugiura, Y. (2002) Translocation of branched-chain arginine peptides through cell membranes: flexibility in the spatial disposition of positive charges in membrane-permeable peptides, *Biochemistry* 41, 7925-7930.
- Futaki, S., Suzuki, T., Ohashi, W., Yagami, T., Tanaka, S., Ueda, K. und Sugiura, Y. (2001) Arginine-rich peptides - An abundant source of membrane-permeable peptides having potential as carriers for intracellular protein delivery, *J. Biol. Chem.* 276, 5836-5840.
- Gait, M. J. (2003) Peptide-mediated cellular delivery of antisense oligonucleotides and their analogues, *Cell Mol. Life Sci.* 60, 844-853.
- Gammon, S. T., Villalobos, V. M., Prior, J. L., Sharma, V. und Piwnica-Worms, D. (2003) Quantitative analysis of permeation peptide complexes labeled with Technetium-99m: chiral and sequence-specific effects on net cell uptake, *Bioconjug. Chem.* 14, 368-376.
- Gausepohl, H., Pides, U. und Frank, W. (1992) Peptide Chemistry and Biology, (Smith, J. A. und Rivier, J. E., Eds.), ESCOM, Leiden, 523.
- Gazit, E., Boman, A., Boman, H. G. und Shai, Y. (1995) Interaction of the mammalian antibacterial peptide cecropin P1 with phospholipid vesicles, *Biochemistry* 34, 11479-11488.
- Gewirtz, A. M., Sokol, D. L. und Ratajczak, M. Z. (1998) Nucleic acid therapeutics: state of the art and future prospects, *Blood* 92, 712-736.
- Giles, R. V., Spiller, D. G., Grzybowski, J., Clark, R. E., Nicklin, P. und Tidd, D. M. (1998) Selecting optimal oligonucleotide composition for maximal antisense effect following streptolysin O-mediated delivery into human leukaemia cells, *Nucleic Acids Res.* 26, 1567-1575.
- Good, L. und Nielsen, P. E. (1998) Antisense inhibition of gene expression in bacteria by PNA targeted to mRNA, *Nat. Biotechnol.* 16, 355-358.
- Gras-Masse, H. (2003) Lipid vector for the delivery of peptides towards intracellular pharmacological targets, *J. Mol. Recognit.* 16, 234-239.
- Gratton, J. P., Yu, J., Griffith, J. W., Babbitt, R. W., Scotland, R. S., Hickey, R., Giordano, F. J. und Sessa, W. C. (2003) Cell-permeable peptides improve cellular uptake and therapeutic gene delivery of replication-deficient viruses in cells and in vivo, *Nat. Med.* 9, 357-362.
- Green, I., Christison, R., Voyce, C. J., Bundell, K. R. und Lindsay, M. A. (2003) Protein transduction domains: are they delivering?, *Trends Pharmacol. Sci.* 24, 213-215.
- Green, M. und Loewenstein, P. M. (1988) Autonomous functional domains of chemically synthesized human immunodeficiency virus tat trans-activator protein, *Cell* 55, 1179-1188.

- Gregoriadis, G. (1976) The carrier potential of liposomes in biology and medicine (second of two parts), *N. Engl. J. Med.* 295, 765-770.
- Gryaznov, S. und Chen, J. K. (1994) Oligodeoxyribonucleotide N3'-]P5' Phosphoramidates - Synthesis and Hybridization Properties, *Journal Of The American Chemical Society* 116, 3143-3144.
- Guillot, F. L., Audus, K. L. und Raub, T. J. (1990) Fluid-phase endocytosis by primary cultures of bovine brain microvessel endothelial cell monolayers, *Microvasc. Res.* 39, 1-14.
- Haberland, A., Knaus, T., Zaitsev, S. V., Stahn, R., Mistry, A. R., Coutelle, C., Haller, H. und Bottger, M. (1999) Calcium ions as efficient cofactor of polycation-mediated gene transfer, *Biochim. Biophys. Acta* 1445, 21-30.
- Hällbrink, M., Floren, A., Elmquist, A., Pooga, M., Bartfai, T. und Langel, U. (2001) Cargo delivery kinetics of cell-penetrating peptides, *Biochim. Biophys. Acta* 1515, 101-109.
- Hamilton, S. E., Simmons, C. G., Kathiriya, I. S. und Corey, D. R. (1999) Cellular delivery of peptide nucleic acids and inhibition of human telomerase, *Chem. Biol.* 6, 343-351.
- Hanvey, J. C., Peffer, N. J., Bisi, J. E., Thomson, S. A., Cadilla, R., Josey, J. A., Ricca, D. J., Hassman, C. F., Bonham, M. A., Au, K. G., Carter, S. G., Bruckenstein, D. A., Boyd, A. L., Noble, S. A. und Babiss, L. E. (1992) Antisense and Antigene Properties of Peptide Nucleic-Acids, *Science* 258, 1481-1485.
- Harrison, R. L., Byrne, B. J. und Tung, L. (1998) Electroporation-mediated gene transfer in cardiac tissue, *FEBS Lett.* 435, 1-5.
- Hawiger, J. (1999) Noninvasive intracellular delivery of functional peptides and proteins, *Curr. Opin. Chem. Biol.* 3, 89-94.
- Heasman, J. (2002) Morpholino oligos: making sense of antisense?, *Dev. Biol.* 243, 209-214.
- Helene, C. und Toulme, J. J. (1990) Specific regulation of gene expression by antisense, sense and antigene nucleic acids, *Biochim. Biophys. Acta* 1049, 99-125.
- Herdewijn, P. (2000) Heterocyclic modifications of oligonucleotides and antisense technology, *Antisense Nucleic Acid Drug Dev.* 10, 297-310.
- Herrera-Ruiz, D. und Knipp, G. T. (2003) Current perspectives on established and putative mammalian oligopeptide transporters, *J. Pharm. Sci.* 92, 691-714.
- Horng, T., Barton, G. M. und Medzhitov, R. (2001) TIRAP: an adapter molecule in the Toll signaling pathway, *Nat. Immunol.* 2, 835-841.
- Janowski, B. A., Kaihatsu, K., Huffman, K. E., Schwartz, J. C., Ram, R., Hardy, D., Mendelson, C. R. und Corey, D. R. (2005) Inhibiting transcription of chromosomal DNA with antigene peptide nucleic acids, *Nat. Chem. Biol.* 1, 210-215.
- Järver, P. und Langel, U. (2004) The use of cell-penetrating peptides as a tool for gene regulation, *Drug Discov. Today* 9, 395-402.
- Jensen, K. K., Orum, H., Nielsen, P. E. und Norden, B. (1997) Kinetics for hybridization of peptide nucleic acids (PNA) with DNA and RNA studied with the BIACore technique, *Biochemistry* 36, 5072-5077.
- Jensen, O. N., Kulkarni, S., Aldrich, J. V. und Barofsky, D. F. (1996) Characterization of peptide-oligonucleotide heteroconjugates by mass spectrometry, *Nucleic Acids Res.* 24, 3866-3872.

- Joliot, A., Pernelle, C., Deagostini-Bazin, H. und Prochiantz, A. (1991) Antennapedia homeobox peptide regulates neural morphogenesis, *Proc. Natl. Acad. Sci. U. S. A* 88, 1864-1868.
- Joliot, A. und Prochiantz, A. (2004) Transduction peptides: from technology to physiology, *Nat. Cell Biol.* 6, 189-196.
- Jönsson, U., Fagerstam, L., Ivarsson, B., Johnsson, B., Karlsson, R., Lundh, K., Lofas, S., Persson, B., Roos, H., Ronnberg, I. und . (1991) Real-time biospecific interaction analysis using surface plasmon resonance and a sensor chip technology, *Biotechniques* 11, 620-627.
- Juliano, R. L., Alahari, S., Yoo, H., Kole, R. und Cho, M. (1999) Antisense pharmacodynamics: Critical issues in the transport and delivery of antisense oligonucleotides, *Pharm. Res.* 16, 494-502.
- Kaihatsu, K., Huffman, K. E. und Corey, D. R. (2004a) Intracellular uptake and inhibition of gene expression by PNAs and PNA-peptide conjugates, *Biochemistry* 43, 14340-14347.
- Kaihatsu, K., Janowski, B. A. und Corey, D. R. (2004b) Recognition of chromosomal DNA by PNAs, *Chem. Biol.* 11, 749-758.
- Kang, S. H., Cho, M. J. und Kole, R. (1998) Up-regulation of luciferase gene expression with antisense oligonucleotides: Implications and applications in functional assay developments, *Biochemistry* 37, 6235-6239.
- Kaplan, I. M., Wadia, J. S. und Dowdy, S. F. (2005) Cationic TAT peptide transduction domain enters cells by macropinocytosis, *J. Control Release* 102, 247-253.
- Kapuscinski, J. und Szer, W. (1979) Interactions of 4', 6-diamidine-2-phenylindole with synthetic polynucleotides, *Nucleic Acids Res.* 6, 3519-3534.
- Karlsson, O., Thor, S., Norberg, T., Ohlsson, H. und Edlund, T. (1990) Insulin gene enhancer binding protein Isl-1 is a member of a novel class of proteins containing both a homeo- and a Cys-His domain, *Nature* 344, 879-882.
- Kaznessis, Y. N., Kim, S. und Larson, R. G. (2002) Specific mode of interaction between components of model pulmonary surfactants using computer simulations, *J. Mol. Biol.* 322, 569-582.
- Keller, S., Sauer, I., Strauss, H., Gast, K., Dathe, M. und Bienert, M. (2005a) Membrane-mimetic nanocarriers formed by a dipalmitoylated cell-penetrating peptide, *Angew. Chem. Int. Ed Engl.* 44, 5252-5255.
- Keller, S., Tsamaloukas, A. und Heerklotz, H. (2005b) A quantitative model describing the selective solubilization of membrane domains, *J. Am. Chem. Soc.* 127, 11469-11476.
- Kim, D. T., Mitchell, D. J., Brockstedt, D. G., Fong, L., Nolan, G. P., Fathman, C. G., Engleman, E. G. und Rothbard, J. B. (1997) Introduction of soluble proteins into the MHC class I pathway by conjugation to an HIV tat peptide, *J. Immunol.* 159, 1666-1668.
- Kim, H. J., Greenleaf, J. F., Kinnick, R. R., Bronk, J. T. und Bolander, M. E. (1996) Ultrasound-mediated transfection of mammalian cells, *Hum. Gene Ther.* 7, 1339-1346.
- Knight, D. E. und Scrutton, M. C. (1986) Gaining access to the cytosol: the technique and some applications of electroporation, *Biochem. J.* 234, 497-506.
- Knudsen, H. und Nielsen, P. E. (1996) Antisense properties of duplex- and triplex-forming PNAs, *Nucleic Acids Res.* 24, 494-500.

- Kobayashi, A., Maeda, T. und Maeda, M. (2004) Membrane localization of transporter associated with antigen processing (TAP)-like (ABCB9) visualized in vivo with a fluorescence protein-fusion technique, *Biol. Pharm. Bull.* 27, 1916-1922.
- Koch, T. (1999) in Peptide Nucleic Acids: Protocols and Applications (Nielsen P. E. und Egholm, M., Eds.), 21.
- Kole, R., Vacek, M. und Williams, T. (2004) Modification of alternative splicing by antisense therapeutics, *Oligonucleotides* 14, 65-74.
- Koppelhus, U., Awasthi, S. K., Zachar, V., Holst, H. U., Ebbesen, P. und Nielsen, P. E. (2002) Cell-dependent differential cellular uptake of PNA, peptides, and PNA-peptide conjugates, *Antisense Nucleic Acid Drug Dev.* 12, 51-63.
- Koshkin, A. A., Singh, S. K., Nielsen, P., Rajwanshi, V. K., Kumar, R., Meldgaard, M., Olsen, C. E. und Wengel, J. (1998) LNA (Locked Nucleic Acids): Synthesis of the adenine, cytosine, guanine, 5-methylcytosine, thymine and uracil bicyclonucleoside monomers, oligomerisation, and unprecedented nucleic acid recognition, *Tetrahedron* 54, 3607-3630.
- Kramer, S. D. und Wunderli-Allenspach, H. (2003) No entry for TAT(44-57) into liposomes and intact MDCK cells: novel approach to study membrane permeation of cell-penetrating peptides, *Biochim. Biophys. Acta* 1609, 161-169.
- Krause, E., Beyermann, M., Dathe, M., Rothmund, S. und Bienert, M. (1995) Location of an amphipathic alpha-helix in peptides using reversed-phase HPLC retention behavior of D-amino acid analogs, *Anal. Chem.* 67, 252-258.
- Krause, E., Beyermann, M., Fabian, H., Dathe, M., Rothmund, S. und Bienert, M. (1996) Conformation of a water-soluble beta-sheet model peptide - A circular dichroism and Fourier-transform infrared spectroscopic study of double D-amino acid replacements, *Int. J. Pept. Protein Res.* 48, 559-568.
- Krishan, A. (1990) Rapid DNA content analysis by the propidium iodide-hypotonic citrate method, *Methods Cell Biol.* 33, 121-125.
- Kurreck, J. (2003) Antisense technologies. Improvement through novel chemical modifications, *Eur. J. Biochem.* 270, 1628-1644.
- Lacerra, G., Sierakowska, H., Carestia, C., Fucharoen, S., Summerton, J., Weller, D. und Kole, R. (2000) Restoration of hemoglobin A synthesis in erythroid cells from peripheral blood of thalassemic patients, *Proc. Natl. Acad. Sci. U. S. A.* 97, 9591-9596.
- Langel, Ü. Ed. (2002) Cell-Penetrating Peptides, *Handbook of Cell-Penetrating Peptides*, CRC Press LLC, Boca Raton.
- Lankat-Buttgereit, B. und Tampe, R. (2002) The transporter associated with antigen processing: function and implications in human diseases, *Physiol Rev.* 82, 187-204.
- Larsen, H. J., Bentin, T. und Nielsen, P. E. (1999) Antisense properties of peptide nucleic acid, *Biochim. Biophys. Acta* 1489, 159-166.
- Le, P. U. und Nabi, I. R. (2003) Distinct caveolae-mediated endocytic pathways target the Golgi apparatus and the endoplasmic reticulum, *J. Cell Sci.* 116, 1059-1071.
- Lemaitre, M., Bayard, B. und Lebleu, B. (1987) Specific antiviral activity of a poly(L-lysine)-conjugated oligodeoxyribonucleotide sequence complementary to vesicular stomatitis virus N protein mRNA initiation site, *Proc. Natl. Acad. Sci. U. S. A* 84, 648-652.

- Leonetti, J. P., Degols, G. und Lebleu, B. (1990) Biological activity of oligonucleotide-poly(L-lysine) conjugates: mechanism of cell uptake, *Bioconjug. Chem.* 1, 149-153.
- Levin, A. A. (1999) A review of the issues in the pharmacokinetics and toxicology of phosphorothioate antisense oligonucleotides, *Biochim. Biophys. Acta* 1489, 69-84.
- Lewin, M., Carlesso, N., Tung, C. H., Tang, X. W., Cory, D., Scadden, D. T. und Weissleder, R. (2000) Tat peptide-derivatized magnetic nanoparticles allow in vivo tracking and recovery of progenitor cells, *Nat. Biotechnol.* 18, 410-414.
- Lewis, M. R., Jia, F., Gallazzi, F., Wang, Y., Zhang, J., Shenoy, N., Lever, S. Z. und Hannink, M. (2002) Radiometal-labeled peptide-PNA conjugates for targeting bcl-2 expression: preparation, characterization, and in vitro mRNA binding, *Bioconjug. Chem.* 13, 1176-1180.
- Liang, L., Liu, D. P. und Liang, C. C. (2002) Optimizing the delivery systems of chimeric RNA.DNA oligonucleotides, *Eur. J. Biochem.* 269, 5753-5758.
- Lin, Y. Z., Yao, S. Y., Veach, R. A., Torgerson, T. R. und Hawiger, J. (1995) Inhibition of nuclear translocation of transcription factor NF- kappa B by a synthetic peptide containing a cell membrane-permeable motif and nuclear localization sequence, *J. Biol. Chem.* 270, 14255-14258.
- Lindberg, M., Jarvet, J., Langel, U. und Graslund, A. (2001) Secondary structure and position of the cell-penetrating peptide transportan in SDS micelles as determined by NMR, *Biochemistry* 40, 3141-3149.
- Lindgren, M., Hällbrink, M., Prochiantz, A. und Langel, Ü. (2000) Cell-penetrating peptides, *Trends Pharmacol. Sci.* 21, 99-103.
- Lipinski, C. A., Lombardo, F., Dominy, B. W. und Feeney, P. J. (2001) Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings, *Adv. Drug Deliv. Rev.* 46, 3-26.
- Ljungstrom, T., Knudsen, H. und Nielsen, P. E. (1999) Cellular uptake of adamantyl conjugated peptide nucleic acids, *Bioconjug. Chem.* 10, 965-972.
- Lochmann, D., Jauk, E. und Zimmer, A. (2004) Drug delivery of oligonucleotides by peptides, *Eur. J. Pharm. Biopharm.* 58, 237-251.
- Lok, C. N., Viazovkina, E., Min, K. L., Nagy, E., Wilds, C. J., Damha, M. J. und Parniak, M. A. (2002) Potent gene-specific inhibitory properties of mixed-backbone antisense oligonucleotides comprised of 2'-deoxy-2'-fluoro-D-arabinose and 2'-deoxyribose nucleotides, *Biochemistry* 41, 3457-3467.
- Lu, J., Higashimoto, Y., Appella, E. und Celis, E. (2004) Multiepitope Trojan antigen peptide vaccines for the induction of antitumor CTL and Th immune responses, *J. Immunol.* 172, 4575-4582.
- Lundberg, M., Wikstrom, S. und Johansson, M. (2003) Cell surface adherence and endocytosis of protein transduction domains, *Molecular Therapy* 8, 143-150.
- Luo, D. und Saltzman, W. M. (2000) Synthetic DNA delivery systems, *Nat. Biotechnol.* 18, 33-37.
- Magzoub, M., Eriksson, L. E. und Graslund, A. (2002) Conformational states of the cell-penetrating peptide penetratin when interacting with phospholipid vesicles: effects of surface charge and peptide concentration, *Biochim. Biophys. Acta* 1563, 53-63.

- Mai, J. C., Shen, H., Watkins, S. C., Cheng, T. und Robbins, P. D. (2002) Efficiency of protein transduction is cell type-dependent and is enhanced by dextran sulfate, *J. Biol. Chem.* 277, 30208-30218.
- Maier, M. A., Esau, C. C., Siwkowski, A. M., Wancewicz, E. V., Albertshofer, K., Kinberger, G. A., Kadaba, N. S., Watanabe, T., Manoharan, M., Bennett, C. F., Griffey, R. H. und Swayze, E. E. (2006) Evaluation of basic amphipathic peptides for cellular delivery of antisense peptide nucleic acids, *J. Med. Chem.* 49, 2534-2542.
- Mann, D. A. und Frankel, A. D. (1991) Endocytosis and targeting of exogenous HIV-1 Tat protein, *EMBO J.* 10, 1733-1739.
- Mao, H. Y., Hart, S. A., Schink, A. und Pollok, B. A. (2004) Sortase-mediated protein ligation: A new method for protein engineering, *J. Am. Chem. Soc.* 126, 2670-2671.
- Marcucci, F. und Lefoulon, F. (2004) Active targeting with particulate drug carriers in tumor therapy: fundamentals and recent progress, *Drug Discov. Today* 9, 219-228.
- Marty, C., Meylan, C., Schott, H., Ballmer-Hofer, K. und Schwendener, R. A. (2004) Enhanced heparan sulfate proteoglycan-mediated uptake of cell-penetrating peptide-modified liposomes, *Cell Mol. Life Sci.* 61, 1785-1794.
- Mazel, M., Clair, P., Rousselle, C., Vidal, P., Scherrmann, J. M., Mathieu, D. und Temsamani, J. (2001) Doxorubicin-peptide conjugates overcome multidrug resistance, *Anticancer Drugs* 12, 107-116.
- Mazmanian, S. K., Liu, G., Ton-That, H. und Schneewind, O. (1999) *Staphylococcus aureus* sortase, an enzyme that anchors surface proteins to the cell wall, *Science* 285, 760-763.
- Mehier-Humbert, S. und Guy, R. H. (2005) Physical methods for gene transfer: improving the kinetics of gene delivery into cells, *Adv. Drug Deliv. Rev.* 57, 733-753.
- Melan, M. A. und Sluder, G. (1992) Redistribution and differential extraction of soluble proteins in permeabilized cultured cells. Implications for immunofluorescence microscopy, *J. Cell Sci.* 101 (Pt 4), 731-743.
- Merrifield, R. B. (1962) Peptide Synthesis on A Solid Polymer, *Federation Proceedings* 21, 412-&.
- Meunier, J. C., Mollereau, C., Toll, L., Suaudeau, C., Moisand, C., Alvinerie, P., Butour, J. L., Guillemot, J. C., Ferrara, P., Monsarrat, B. und et, a. (1995) Isolation and structure of the endogenous agonist of opioid receptor-like ORL1 receptor, *Nature* 377, 532-535.
- Midoux, P., Mendes, C., Legrand, A., Raimond, J., Mayer, R., Monsigny, M. und Roche, A. C. (1993) Specific gene transfer mediated by lactosylated poly-L-lysine into hepatoma cells, *Nucleic Acids Res.* 21, 871-878.
- Mitchell, D. J., Kim, D. T., Steinman, L., Fathman, C. G. und Rothbard, J. B. (2000) Polyarginine enters cells more efficiently than other polycationic homopolymers, *J. Pept. Res.* 56, 318-325.
- Morishita, R., Higaki, J., Tomita, N. und Ogihara, T. (1998) Application of transcription factor "decoy" strategy as means of gene therapy and study of gene expression in cardiovascular disease, *Circ. Res.* 82, 1023-1028.
- Mosmann, T. (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays, *J. Immunol. Methods* 65, 55-63.

- Moulton, H. M., Hase, M. C., Smith, K. M. und Iversen, P. L. (2003) HIV Tat peptide enhances cellular delivery of antisense morpholino oligomers, *Antisense Nucleic Acid Drug Dev.* 13, 31-43.
- Moulton, H. M., Nelson, M. H., Hatlevig, S. A., Reddy, M. T. und Iversen, P. L. (2004) Cellular uptake of antisense morpholino oligomers conjugated to arginine-rich peptides, *Bioconjug. Chem.* 15, 290-299.
- Muratovska, A. und Eccles, M. R. (2004) Conjugate for efficient delivery of short interfering RNA (siRNA) into mammalian cells, *FEBS Lett.* 558, 63-68.
- Myou, S., Leff, A. R., Myo, S., Boetticher, E., Tong, J., Meliton, A. Y., Liu, J., Munoz, N. M. und Zhu, X. (2003) Blockade of inflammation and airway hyperresponsiveness in immune-sensitized mice by dominant-negative phosphoinositide 3-kinase-TAT, *J. Exp. Med.* 198, 1573-1582.
- Nielsen, P. E. (2000) Antisense properties of peptide nucleic acid, *Methods Enzymol.* 313, 156-164.
- Nielsen, P. E., Egholm, M., Berg, R. H. und Buchardt, O. (1991) Sequence-selective recognition of DNA by strand displacement with a thymine-substituted polyamide, *Science* 254, 1497-1500.
- Nielsen, P. E., Egholm, M. und Buchardt, O. (1994) Evidence for (PNA)2/DNA triplex structure upon binding of PNA to dsDNA by strand displacement, *J. Mol. Recognit.* 7, 165-170.
- Nishikawa, K., Sawasdikosol, S., Fruman, D. A., Lai, J., Songyang, Z., Burakoff, S. J., Yaffe, M. B. und Cantley, L. C. (2000) A peptide library approach identifies a specific inhibitor for the ZAP-70 protein tyrosine kinase, *Mol. Cell* 6, 969-974.
- Noble, S. A., Bonham, M. A., Bisi, J. E., Bruckenstein, D. A., Brown, P. H., Brown, S. C., Cadilla, R., Gaul, M. D., Hanvey, J. C., Hassman, C. F., Josey, J. A., Luzzio, M. J., Myers, P. M., Pipe, A. J., Ricca, D. J., Su, C. W., Stevenson, C. L., Thomson, S. A., Wiethe, R. W. und Babiss, L. E. (1995) Impact of Biophysical Parameters on the Biological Assessment of Peptide Nucleic-Acids, Antisense Inhibitors of Gene-Expression, *Drug Development Research* 34, 184-195.
- Norton, J. C., Piatyszek, M. A., Wright, W. E., Shay, J. W. und Corey, D. R. (1996) Inhibition of human telomerase activity by peptide nucleic acids, *Nat. Biotechnol.* 14, 615-619.
- Nur-E-Kamal, M. S., Kamal, J. M., Qureshi, M. M. und Maruta, H. (1999) The CDC42-specific inhibitor derived from ACK-1 blocks v-Ha-Ras-induced transformation, *Oncogene* 18, 7787-7793.
- Oehlke, J., Birth, P., Klauschenz, E., Wiesner, B., Beyermann, M., Oksche, A. und Bienert, M. (2002a) Cellular uptake of antisense oligonucleotides after complexing or conjugation with cell-penetrating model peptides, *Eur. J Biochem* 269, 4025-4032.
- Oehlke, J., Krause, E., Wiesner, B., Beyermann, M. und Bienert, M. (1997) Extensive cellular-uptake into endothelial cells of an amphipathic beta-sheet forming peptide, *FEBS Lett.* 415, 196-199.
- Oehlke, J., Lorenz, D., Wiesner, B. und Bienert, M. (2005) Studies on the cellular uptake of substance P and lysine-rich, KLA-derived model peptides, *J. Mol. Recognit.* 18, 50-59.
- Oehlke, J., Scheller, A., Birth, P., Wiesner, B., Beyermann, M., Faulstich, H. und Bienert, M. (2000) Structural requirements for the cellular uptake of peptides, *Proceedings of the 8th Akabori Conference*, 26. -29. 04. 2000, Nagoya, Japan, ed. by Takayuki Shiori 96-100.
- Oehlke, J., Scheller, A., Wiesner, B., Krause, E., Beyermann, M., Klauschenz, E., Melzig, M. und Bienert, M. (1998) Cellular uptake of an alpha-helical amphipathic model peptide with the potential to deliver polar compounds into the cell interior non-endocytically, *Biochim. Biophys. Acta* 1414, 127-139.

- Oehlke, J., Wallukat, G., Wolf, Y., Ehrlich, A., Wiesner, B., Berger, H. und Bienert, M. (2004) Enhancement of intracellular concentration and biological activity of PNA after conjugation with a cell-penetrating synthetic model peptide, *Eur. J. Biochem.* 271, 3043-3049.
- Oehlke, J., Wiesner, B. und Bienert, M. (2002b) Model amphipathic peptides, *Handbook of Cell-Penetrating Peptides*, (Langel, U. Ed.) CRC Press LLC, Boca Raton, 71-92.
- Olsnes, S., Klingenberg, O. und Wiedlocha, A. (2003) Transport of exogenous growth factors and cytokines to the cytosol and to the nucleus, *Physiol Rev.* 83, 163-182.
- Ostenson, C. G., Sandberg-Nordqvist, A. C., Chen, J., Hallbrink, M., Rotin, D., Langel, U. und Efendic, S. (2002) Overexpression of protein-tyrosine phosphatase PTP sigma is linked to impaired glucose-induced insulin secretion in hereditary diabetic Goto-Kakizaki rats, *Biochem. Biophys. Res. Commun.* 291, 945-950.
- Paterson, G. K. und Mitchell, T. J. (2004) The biology of Gram-positive sortase enzymes, *Trends Microbiol.* 12, 89-95.
- Pelkmans, L. und Helenius, A. (2003) Insider information: what viruses tell us about endocytosis, *Curr. Opin. Cell Biol.* 15, 414-422.
- Perry, A. M., Ton-That, H., Mazmanian, S. K. und Schneewind, O. (2002) Anchoring of surface proteins to the cell wall of *Staphylococcus aureus*. III. Lipid II is an in vivo peptidoglycan substrate for sortase-catalyzed surface protein anchoring, *J. Biol. Chem.* 277, 16241-16248.
- Persson, D., Thoren, P. E., Herner, M., Lincoln, P. und Norden, B. (2003) Application of a novel analysis to measure the binding of the membrane-translocating peptide penetratin to negatively charged liposomes, *Biochemistry* 42, 421-429.
- Persson, D., Thoren, P. E. und Norden, B. (2001) Penetratin-induced aggregation and subsequent dissociation of negatively charged phospholipid vesicles, *FEBS Lett.* 505, 307-312.
- Pichon, C., Monsigny, M. und Roche, A. C. (1999) Intracellular localization of oligonucleotides: influence of fixative protocols, *Antisense Nucleic Acid Drug Dev.* 9, 89-93.
- Pietersz, G. A., Li, W. und Apostolopoulos, V. (2001) A 16-mer peptide (RQIKIWFQNRRMKWKK) from antennapedia preferentially targets the Class I pathway, *Vaccine* 19, 1397-1405.
- Pillay, C. S., Elliott, E. und Dennison, C. (2002) Endolysosomal proteolysis and its regulation, *Biochem. J.* 363, 417-429.
- Pohjanpelto, P., Holtta, E., Janne, O. A., Knuutila, S. und Alitalo, K. (1985) Amplification of ornithine decarboxylase gene in response to polyamine deprivation in Chinese hamster ovary cells, *J. Biol. Chem.* 260, 8532-8537.
- Pollack, A. und Ciancio, G. (1990) Cell cycle phase-specific analysis of cell viability using Hoechst 33342 and propidium iodide after ethanol preservation, *Methods Cell Biol.* 33, 19-24.
- Pooga, M., Hällbrink, M., Zorko, M. und Langel, U. (1998a) Cell penetration by transportan, *FASEB J.* 12, 67-77.
- Pooga, M., Land, T., Bartfai, T. und Langel, U. (2001) PNA oligomers as tools for specific modulation of gene expression, *Biomolecular Engineering* 17, 183-192.
- Pooga, M., Soomets, U., Hällbrink, M., Valkna, A., Saar, K., Rezaei, K., Kahl, U., Hao, J. X., Xu, X. J., Wiesenfeld-Hallin, Z., Hokfelt, T., Bartfai, A. und Langel, U. (1998b) Cell penetrating PNA

- constructs regulate galanin receptor levels and modify pain transmission in vivo, *Nat. Biotechnol.* 16, 857-861.
- Prochiantz, A. (1996) Getting hydrophilic compounds into cells: Lessons from homeopeptides, *Curr. Opin. Neurobiol.* 6, 629-634.
- Provoda, C. J. und Lee, K. D. (2000) Bacterial pore-forming hemolysins and their use in the cytosolic delivery of macromolecules, *Adv. Drug Deliv. Rev.* 41, 209-221.
- Ragin, A. D. und Chmielewski, J. (2004) Probing essential residues for cellular uptake with a cationic nuclear localization signal sequence, *J. Pept. Res.* 63, 155-160.
- Reinscheid, R. K., Nothacker, H. P., Bourson, A., Ardati, A., Henningsen, R. A., Bunzow, J. R., Grandy, D. K., Langen, H., Monsma, F.-J. J. und Civelli, O. (1995) Orphanin FQ: a neuropeptide that activates an opioidlike G protein-coupled receptor, *Science* 270, 792-794.
- Rejman, J., Oberle, V., Zuhorn, I. S. und Hoekstra, D. (2004) Size-dependent internalization of particles via the pathways of clathrin- and caveolae-mediated endocytosis, *Biochem. J.* 377, 159-169.
- Richard, J. P., Melikov, K., Brooks, H., Prevot, P., Lebleu, B. und Chernomordik, L. V. (2005) Cellular uptake of unconjugated TAT peptide involves clathrin-dependent endocytosis and heparan sulfate receptors, *J. Biol. Chem.* 280, 15300-15306.
- Richard, J. P., Melikov, K., Vives, E., Ramos, C., Verbeure, B., Gait, M. J., Chernomordik, L. V. und Lebleu, B. (2003) Cell-penetrating peptides. A reevaluation of the mechanism of cellular uptake, *J. Biol. Chem.* 278, 585-590.
- Rittner, K., Benavente, A., Bompard-Sorlet, A., Heitz, F., Divita, G., Brasseur, R. und Jacobs, E. (2002) New basic membrane-destabilizing peptides for plasmid-based gene delivery in vitro and in vivo, *Mol. Ther.* 5, 104-114.
- Rojas, M., Donahue, J. P., Tan, Z. J. und Lin, Y. Z. (1998) Genetic engineering of proteins with cell membrane permeability, *Nat. Biotechnol.* 16, 370-375.
- Ross, M. F., Filipovska, A., Smith, R. A., Gait, M. J. und Murphy, M. P. (2004) Cell-penetrating peptides do not cross mitochondrial membranes even when conjugated to a lipophilic cation: evidence against direct passage through phospholipid bilayers, *Biochem. J.* 383, 457-468.
- Rothbard, J. B., Garlington, S., Lin, Q., Kirschberg, T., Kreider, E., McGrane, P. L., Wender, P. A. und Khavari, P. A. (2000) Conjugation of arginine oligomers to cyclosporin A facilitates topical delivery and inhibition of inflammation, *Nat. Med.* 6, 1253-1257.
- Rothenmund, S., Krause, E., Beyermann, M., Dathe, M., Engelhardt, H. und Bienert, M. (1995) Recognition of alpha-helical peptide structures using high-performance liquid chromatographic retention data for D-amino acid analogues: influence of peptide amphipathicity and of stationary phase hydrophobicity, *J. Chromatogr.* 689, 219-226.
- Rothenmund, S., Krause, E., Ehrlich, A., Bienert, M., Glusa, E. und Verhallen, P. (1994) Determination of peptide hydrophobicity parameters by reversed-phase high-performance liquid chromatography, *J. Chromatogr.* 661, 77-82.
- Rousselle, C., Clair, P., Lefauconnier, J. M., Kaczorek, M., Schermann, J. M. und Temsamani, J. (2000) New advances in the transport of doxorubicin through the blood-brain barrier by a peptide vector-mediated strategy, *Mol Pharmacol* 57, 679-686.
- Ruoslahti, E., Duza, T. und Zhang, L. (2005) Vascular homing peptides with cell-penetrating properties, *Curr. Pharm. Des* 11, 3655-3660.

- Ryser, H. J. und Hancock, R. (1965) Histones and basic polyamino acids stimulate the uptake of albumin by tumor cells in culture, *Science* 150, 501-503.
- Ryser, H. J. und Shen, W. C. (1978) Conjugation of methotrexate to poly(L-lysine) increases drug transport and overcomes drug resistance in cultured cells, *Proc. Natl. Acad. Sci. U. S. A* 75, 3867-3870.
- Sadler, K., Eom, K. D., Yang, J. L., Dimitrova, Y. und Tam, J. P. (2002) Translocating proline-rich peptides from the antimicrobial peptide bactenecin 7, *Biochemistry* 41, 14150-14157.
- Sakai, N. und Matile, S. (2003) Anion-mediated transfer of polyarginine across liquid and bilayer membranes, *J. Am. Chem. Soc.* 125, 14348-14356.
- Salvatella, X., Martinell, M., Gairi, M., Mateu, M. G., Feliz, M., Hamilton, A. D., de Mendoza, J. und Giralt, E. (2004) A tetraguanidinium ligand binds to the surface of the tetramerization domain of protein P53, *Angew. Chem. Int. Ed Engl.* 43, 196-198.
- Sandvig, K. und van Deurs, B. (2002) Transport of protein toxins into cells: pathways used by ricin, cholera toxin and Shiga toxin, *FEBS Lett.* 529, 49-53.
- Sazani, P., Kang, S. H., Maier, M. A., Wei, C. F., Dillman, J., Summerton, J., Manoharan, M. und Kole, R. (2001) Nuclear antisense effects of neutral, anionic and cationic oligonucleotide analogs, *Nucleic Acids Res.* 29, 3965-3974.
- Scarfi, S., Gasparini, A., Damonte, G. und Benatti, U. (1997) Synthesis, uptake, and intracellular metabolism of a hydrophobic tetrapeptide-peptide nucleic acid (PNA)-biotin molecule, *Biochem. Biophys. Res. Commun.* 236, 323-326.
- Scheller, A., Melzig, M. und Oehlke, J. (2002) Induction of caspase-8 in human cells by the extracellular administration of peptides containing a C-terminal SLV sequence, *Lett. Pept. Sci.* 8, 29-34.
- Scheller, A., Oehlke, J., Wiesner, B., Dathe, M., Krause, E., Beyermann, M., Melzig, M. und Bienert, M. (1999) Structural requirements for cellular uptake of alpha-helical amphipathic peptides, *J. Pept. Sci.* 5, 185-194.
- Scheller, A., Wiesner, B., Melzig, M., Bienert, M. und Oehlke, J. (2000) Evidence for an amphipathicity independent cellular uptake of amphipathic cell-penetrating peptides, *Eur. J. Biochem.* 267, 6043-6049.
- Schmid, S. L. und Carter, L. L. (1990) ATP is required for receptor-mediated endocytosis in intact cells, *J. Cell Biol.* 111, 2307-2318.
- Schutze-Redelmeier, M. P., Gournier, H., GarciaPons, F., Moussa, M., Joliot, A. H., Volovitch, M., Prochiantz, A. und Lemonnier, F. A. (1996) Introduction of exogenous antigens into the MHC class I processing and presentation pathway by *Drosophila antennapedia* homeodomain primes cytotoxic T cells in vivo, *Journal of Immunology* 157, 650-655.
- Schwarze, S. R., Ho, A., Vocero, A. A. und Dowdy, S. F. (1999) In vivo protein transduction: Delivery of a biologically active protein into the mouse, *Science* 285, 1569-1572.
- Seth, P., Willingham, M. C. und Pastan, I. (1985) Binding of adenovirus and its external proteins to Triton X-114. Dependence on pH, *J. Biol. Chem.* 260, 14431-14434.
- Shen, W. C., Wan, J. S. und Ekrami, H. (1992) Enhancement of Polypeptide and Protein-Absorption by Macromolecular Carriers Via Endocytosis and Transcytosis .3, *Advanced Drug Delivery Reviews* 8, 93-113.

- Shiraishi, T. und Nielsen, P. E. (2004) Down-regulation of MDM2 and activation of p53 in human cancer cells by antisense 9-aminoacridine-PNA (peptide nucleic acid) conjugates, *Nucleic Acids Res.* 32, 4893-4902.
- Shiraishi, T. und Nielsen, P. E. (2006) Photochemically enhanced cellular delivery of cell penetrating peptide-PNA conjugates, *FEBS Lett.* 580, 1451-1456.
- Shiraishi, T., Pankratova, S. und Nielsen, P. E. (2005) Calcium ions effectively enhance the effect of antisense Peptide nucleic acids conjugated to cationic tat and oligoarginine peptides, *Chem. Biol.* 12, 923-929.
- Sibley, L. D. (2004) Intracellular parasite invasion strategies, *Science* 304, 248-253.
- Sierakowska, H., Sambade, M. J., Agrawal, S. und Kole, R. (1996) Repair of thalassemic human beta-globin mRNA in mammalian cells by antisense oligonucleotides, *Proc. Natl. Acad. Sci. U. S. A* 93, 12840-12844.
- Simmons, C. G., Pitts, A. E., Mayfield, L. D., Shay, J. W. und Corey, D. R. (1997) Synthesis and membrane permeability of PNA-peptide conjugates, *Bioorg. Med. Chem. Lett.* 7, 3001-3006.
- Simons, K. und Ikonen, E. (1997) Functional rafts in cell membranes, *Nature* 387, 569-572.
- Simons, K. und Toomre, D. (2000) Lipid rafts and signal transduction, *Nat. Rev. Mol. Cell Biol.* 1, 31-39.
- Singer, S. J. und Nicolson, G. L. (1972) The fluid mosaic model of the structure of cell membranes, *Science* 175, 720-731.
- Singh, D., Bisland, S. K., Kawamura, K. und Gariepy, J. (1999) Peptide-based intracellular shuttle able to facilitate gene transfer in mammalian cells, *Bioconjug. Chem.* 10, 745-754.
- Siwkowski, A. M., Malik, L., Esau, C. C., Maier, M. A., Wanczewicz, E. V., Albertshofer, K., Monia, B. P., Bennett, C. F. und Eldrup, A. B. (2004) Identification and functional validation of PNAs that inhibit murine CD40 expression by redirection of splicing, *Nucleic Acids Res.* 32, 2695-2706.
- Sjöback, R., Nygren, J. und Kubista, M. (1995) Absorption and Fluorescence Properties of Fluorescein, *Spectrochimica Acta Part A* 51, L7-L21.
- Smith, A. E. (1995) Viral vectors in gene therapy, *Annu. Rev. Microbiol.* 49, 807-838.
- Smith, A. E. und Helenius, A. (2004) How viruses enter animal cells, *Science* 304, 237-242.
- Smith, S. A. und Pretorius, W. A. (2002) The conservative behaviour of fluorescein, *Water Sa* 28, 403-406.
- Smulevitch, S. V., Simmons, C. G., Norton, J. C., Wise, T. W. und Corey, D. R. (1996) Enhancement of strand invasion by oligonucleotides through manipulation of backbone charge, *Nat. Biotechnol.* 14, 1700-1704.
- Snyder, E. L. und Dowdy, S. F. (2004) Cell penetrating peptides in drug delivery, *Pharm. Res.* 21, 389-393.
- Sonawane, N. D., Szoka, F. C., Jr. und Verkman, A. S. (2003) Chloride accumulation and swelling in endosomes enhances DNA transfer by polyamine-DNA polyplexes, *J. Biol. Chem.* 278, 44826-44831.

- Steiner, V., Schar, M., Bornsen, K. O. und Mutter, M. (1991) Retention behaviour of a template-assembled synthetic protein and its amphiphilic building blocks on reversed-phase columns, *J. Chromatogr.* 586, 43-50.
- Steinman, R. M., Mellman, I. S., Muller, W. A. und Cohn, Z. A. (1983) Endocytosis and the recycling of plasma membrane, *J. Cell Biol.* 96, 1-27.
- Stephens, D. J. und Pepperkok, R. (2001) The many ways to cross the plasma membrane, *Proc. Natl. Acad. Sci. U. S. A.* 98, 4295-4298.
- Stetsenko, D. A., Arzumanov, A. A., Korshun, V. A. und Gait, M. J. (2000) Peptide conjugates of oligonucleotides as enhanced antisense agents, *Mol. Biol.* 34, 852-859.
- Stewart, A. J., Pichon, C., Meunier, L., Midoux, P., Monsigny, M. und Roche, A. C. (1996) Enhanced biological activity of antisense oligonucleotides complexed with glycosylated poly-L-lysine, *Mol. Pharmacol.* 50, 1487-1494.
- Surovoy, A., Flechsler, I. und Jung, G. (1998) A novel series of serum-resistant lipoaminoacid compounds for cellular delivery of plasmid DNA, *Adv. Exp. Med. Biol.* 451, 461-467.
- Suwanmanee, T., Sierakowska, H., Lacerra, G., Svasti, S., Kirby, S., Walsh, C. E., Fucharoen, S. und Kole, R. (2002) Restoration of human beta-globin gene expression in murine and human IVS2-654 thalassemic erythroid cells by free uptake of antisense oligonucleotides, *Molecular Pharmacology* 62, 545-553.
- Suzuki, T., Futaki, S., Niwa, M., Tanaka, S., Ueda, K. und Sugiura, Y. (2002) Possible existence of common internalization mechanisms among arginine-rich peptides, *J. Biol. Chem.* 277, 2437-2443.
- Takeshima, K., Chikushi, A., Lee, K. K., Yonehara, S. und Matsuzaki, K. (2003) Translocation of analogues of the antimicrobial peptides magainin and buforin across human cell membranes, *J. Biol. Chem.* 278, 1310-1315.
- Tamilarasu, N., Huq, I. und Rana, T. M. (1999) High affinity and specific binding of HIV-1 TAR RNA by a Tat-derived oligourea, *J. A. Chem. Soc.* 121, 1597-1598.
- Terrone, D., Sang, S. L. W., Roudaia, L. und Silvius, J. R. (2003) Penetratin and related cell-penetrating cationic peptides can translocate across lipid bilayers in the presence of a transbilayer potential, *Biochemistry* 42, 13787-13799.
- Thein, S. L., Hesketh, C., Taylor, P., Temperley, I. J., Hutchinson, R. M., Old, J. M., Wood, W. G., Clegg, J. B. und Weatherall, D. J. (1990) Molecular basis for dominantly inherited inclusion body beta-thalassemia, *Proc. Natl. Acad. Sci. U. S. A.* 87, 3924-3928.
- Theodore, L., Derossi, D., Chassaing, G., Llirbat, B., Kubes, M., Jordan, P., Chneiweiss, H., Godement, P. und Prochiantz, A. (1995) Intraneuronal delivery of protein kinase C pseudosubstrate leads to growth cone collapse, *J. Neurosci.* 15, 7158-7167.
- Thomson, S. A., Josey, J. A., Cadilla, R., Gaul, M. D., Hassman, C. F., Luzzio, M. J., Pipe, A. J., Reed, K. L., Ricca, D. J., Wiethe, R. W. und Noble, S. A. (1995) Fmoc Mediated Synthesis of Peptide Nucleic-Acids, *Tetrahedron* 51, 6179-6194.
- Thoren, P. E., Persson, D., Isakson, P., Goksor, M., Onfelt, A. und Norden, B. (2003) Uptake of analogs of penetratin, Tat(48-60) and oligoarginine in live cells, *Biochem. Biophys. Res. Commun.* 307, 100-107.
- Thoren, P. E. G., Persson, D., Karlsson, M. und Norden, B. (2000) The *Antennapedia* peptide penetratin translocates across lipid bilayers - the first direct observation, *FEBS Lett.* 482, 265-268.

- Tian, J. H., Xu, W., Zhang, W., Fang, Y., Grisel, J. E., Mogil, J. S., Grandy, D. K. und Han, J. S. (1997) Involvement of endogenous orphanin FQ in electroacupuncture-induced analgesia, *Neuroreport* 8, 497-500.
- Torchilin, V. P., Rammohan, R., Weissig, V. und Levchenko, T. S. (2001) TAT peptide on the surface of liposomes affords their efficient intracellular delivery even at low temperature and in the presence of metabolic inhibitors, *Proc. Natl. Acad. Sci. U. S. A* 98, 8786-8791.
- Tripathi, S., Chaubey, B., Ganguly, S., Harris, D., Casale, R. A. und Pandey, V. N. (2005) Anti-HIV-1 activity of anti-TAR polyamide nucleic acid conjugated with various membrane transducing peptides, *Nucleic Acids Res.* 33, 4345-4356.
- Tung, C. H., Mueller, S. und Weissleder, R. (2002) Novel branching membrane translocational peptide as gene delivery vector, *Bioorg. Med. Chem.* 10, 3609-3614.
- Tung, C. H. und Stein, S. (2000) Preparation and applications of peptide-oligonucleotide conjugates, *Bi conjugate Chem.* 11, 605-618.
- Turner, J. J., Williams, D., Owen, D. und Gait, M. J. (2006) Disulfide Conjugation of Peptides to Oligonucleotides and Their Analogs, in *Current Protocols in Nucleic Acid Chemistry* (John Wiley & Sons, Ed.), Hoboken, New York, 4.28.1-4.28.21.
- Turner, J. J., Arzumanov, A. A. und Gait, M. J. (2005a) Synthesis, cellular uptake and HIV-1 Tat-dependent trans-activation inhibition activity of oligonucleotide analogues disulphide-conjugated to cell-penetrating peptides, *Nucleic Acids Res.* 33, 27-42.
- Turner, J. J., Ivanova, G. D., Verbeure, B., Williams, D., Arzumanov, A. A., Abes, S., Lebleu, B. und Gait, M. J. (2005b) Cell-penetrating peptide conjugates of peptide nucleic acids (PNA) as inhibitors of HIV-1 Tat-dependent trans-activation in cells, *Nucleic Acids Res.* 33, 6837-6849.
- Tyagi, M., Rusnati, M., Presta, M. und Giacca, M. (2001) Internalization of HIV-1 Tat requires cell surface heparan sulfate proteoglycans, *J. Biol. Chem.* 276, 3254-3261.
- Umezawa, N., Gelman, M. A., Haigis, M. C., Raines, R. T. und Gellman, S. H. (2002) Translocation of a beta-peptide across cell membranes, *J. Amer. Chem. Soc.* 124, 368-369.
- Vacek, M. M., Ma, H., Gemignani, F., Lacerra, G., Kafri, T. und Kole, R. (2003) High-level expression of hemoglobin A in human thalassemic erythroid progenitor cells following lentiviral vector delivery of an antisense snRNA, *Blood* 101, 104-111.
- Van der Merwe, A. (2001) Surface plasmon resonance, in *Protein-Ligand Interactions: hydrodynamics and calorimetry* (Harding S.E. und Chowdhry B.Z., Eds.), Oxford University Press, New York, 137-170.
- Veenhoff, L. M., Heuberger, E. H. und Poolman, B. (2002) Quaternary structure and function of transport proteins, *Trends Biochem. Sci.* 27, 242-249.
- Verhoeven, A. J., Leusen, J. H., Kessels, G. C., Hilarius, P. M., de Bont, D. B. und Liskamp, R. M. (1993) Inhibition of neutrophil NADPH oxidase assembly by a myristoylated pseudosubstrate of protein kinase C, *J. Biol. Chem.* 268, 18593-18598.
- Vives, E. (2005) Present and future of cell-penetrating peptide mediated delivery systems: "is the Trojan horse too wild to go only to Troy?", *J. Control Release* 109, 77-85.
- Vives, E., Brodin, P. und Lebleu, B. (1997) A truncated HIV-1 Tat protein basic domain rapidly translocates through the plasma membrane and accumulates in the cell nucleus, *J. Biol. Chem.* 272, 16010-16017.

- Vives, E., Richard, J. P., Rispal, C. und Lebleu, B. (2003) TAT peptide internalization: seeking the mechanism of entry, *Curr. Protein Pept. Sci.* 4, 125-132.
- Wadia, J. S., Stan, R. V. und Dowdy, S. F. (2004) Transducible TAT-HA fusogenic peptide enhances escape of TAT-fusion proteins after lipid raft macropinocytosis, *Nat. Med.* 10, 310-315.
- Walev, I., Bhakdi, S. C., Hofmann, F., Djonder, N., Valeva, A., Aktories, K. und Bhakdi, S. (2001) Delivery of proteins into living cells by reversible membrane permeabilization with streptolysin-O, *Proc. Natl. Acad. Sci. U. S. A* 98, 3185-3190.
- Wallukat, G., Nemecz, G., Farkas, T., Kuehn, H. und Wollenberger, A. (1991) Modulation of the beta-adrenergic response in cultured rat heart cells. I. Beta-adrenergic supersensitivity is induced by lactate via a phospholipase A2 and 15-lipoxygenase involving pathway, *Mol. Cell. Biochem.* 102, 35-47.
- Wang, M. und Drucker, D. J. (1995) The LIM domain homeobox gene *isl-1* is a positive regulator of islet cell-specific proglucagon gene transcription, *J. Biol. Chem.* 270, 12646-12652.
- Wang, R. F. und Wang, H. Y. (2002) Enhancement of antitumor immunity by prolonging antigen presentation on dendritic cells, *Nat. Biotechnol.* 20, 149-154.
- Wender, P. A., Mitchell, D. J., Pattabiraman, K., Pelkey, E. T., Steinman, L. und Rothbard, J. B. (2000) The design, synthesis, and evaluation of molecules that enable or enhance cellular uptake: Peptoid molecular transporters, *Proc. Natl. Acad. Sci. U. S. A.* 97, 13003-13008.
- Wender, P. A., Rothbard, J. B., Jessop, T. C., Kreider, E. L. und Wylie, B. L. (2002) Oligocarbamate molecular transporters: design, synthesis, and biological evaluation of a new class of transporters for drug delivery, *J. Am. Chem. Soc.* 124, 13382-13383.
- Wiesner, B., Lorenz, D., Krause, E., Beyermann, M. und Bienert, M. (2002) Measurement of intracellular fluorescence in the presence of a strong extracellular fluorescence using Confocal Laser Scanning Microscopy, *LAMSO* 3, 1-17.
- Williams, E. J., Dunican, D. J., Green, P. J., Howell, F. V., Derossi, D., Walsh, F. S. und Doherty, P. (1997) Selective inhibition of growth factor-stimulated mitogenesis by a cell-permeable Grb2-binding peptide, *J. Biol. Chem.* 272, 22349-22354.
- Wittung, P., Kajanus, J., Edwards, K., Haaima, G., Nielsen, P., Norden, B. und Malmstrom, B. G. (1995) Phospholipid membrane permeability of peptide nucleic acid (vol 365, pg 27, 1995), *FEBS Lett.* 375, 317-320.
- Wolters, J. C., Abele, R. und Tampe, R. (2005) Selective and ATP-dependent translocation of peptides by the homodimeric ATP binding cassette transporter TAP-like (ABCB9), *J. Biol. Chem.* 280, 23631-23636.
- Wyman, T. B., Nicol, F., Zelphati, O., Scaria, P. V., Plank, C. und Szoka, F. C. (1997) Design, synthesis, and characterization of a cationic peptide that binds to nucleic acids and permeabilizes bilayers, *Biochemistry* 36, 3008-3017.
- Yamaguchi, Y., Kasano, M., Terada, T., Sato, R. und Maeda, M. (1999) An ABC transporter homologous to TAP proteins, *FEBS Lett.* 457, 231-236.
- Zaitsev, S., Buchwalow, I., Haberland, A., Tkachuk, S., Zaitseva, I., Haller, H. und Bottger, M. (2002) Histone H1-mediated transfection: role of calcium in the cellular uptake and intracellular fate of H1-DNA complexes, *Acta Histochem.* 104, 85-92.

- Zamaratski, E., Pradeepkumar, P. I. und Chattopadhyaya, J. (2001) A critical survey of the structure-function of the antisense oligo/RNA heteroduplex as substrate for RNase H, *J. Biochem. Biophys. Methods* 48, 189-208.
- Zamecnik, P. C. und Stephenson, M. L. (1978) Inhibition of Rous sarcoma virus replication and cell transformation by a specific oligodeoxynucleotide, *Proc. Natl. Acad. Sci. U. S. A* 75, 280-284.
- Zarovni, N., Monaco, L. und Corti, A. (2004) Inhibition of tumor growth by intramuscular injection of cDNA encoding tumor necrosis factor alpha coupled to NGR and RGD tumor-homing peptides, *Hum. Gene Ther.* 15, 373-382.
- Zatsepin, T. S., Turner, J. J., Oretskaya, T. S. und Gait, M. J. (2005) Conjugates of oligonucleotides and analogues with cell penetrating peptides as gene silencing agents, *Current Pharmaceutical Design* 11, 3639-3654.
- Zhang, X., Ishihara, T. und Corey, D. R. (2000) Strand invasion by mixed base PNAs and a PNA-peptide chimera, *Nucleic Acids Res.* 28, 3332-3338.
- Zhang, X., Wan, L., Pooyan, S., Su, Y., Gardner, C. R., Leibowitz, M. J., Stein, S. und Sinko, P. J. (2004) Quantitative assessment of the cell penetrating properties of RI-Tat-9: evidence for a cell type-specific barrier at the plasma membrane of epithelial cells, *Mol. Pharm.* 1, 145-155.
- Zhu, C. B., Cao, X. D., Xu, S. F. und Wu, G. C. (1997) Orphanin FQ potentiates formalin-induced pain behavior and antagonizes morphine analgesia in rats, *Neurosci. Lett.* 235, 37-40.
- Ziegler, A., Blatter, X. L., Seelig, A. und Seelig, J. (2003) Protein transduction domains of HIV-1 and SIV TAT interact with charged lipid vesicles. Binding mechanism and thermodynamic analysis, *Biochemistry* 42, 9185-9194.
- Zorko, M. und Langel, U. (2005) Cell-penetrating peptides: mechanism and kinetics of cargo delivery, *Adv. Drug Deliv. Rev.* 57, 529-545.
- Zubin, E. M., Romanova, E. A. und Oretskaya, T. S. (2002) Modern approaches to the synthesis of peptide-oligonucleotide conjugates, *Uspekhi Khimii* 71, 273-302.
- Zuhorn, I. S., Kalicharan, R. und Hoekstra, D. (2002) Lipoplex-mediated transfection of mammalian cells occurs through the cholesterol-dependent clathrin-mediated pathway of endocytosis, *J. Biol. Chem.* 277, 18021-18028.