

## **ACKNOWLEDGEMENTS**

The present work was elaborated in the work group of Dr. Erich Lanka, whom I thank for his trust into my skills and his encouragement for my work. Valuable inspirations frequently emerged during discussions with Erich, who has the virtue to arouse one's own ideas without imposing his own opinions. I also appreciated him giving me tasks I wouldn't have dared to address myself to on my own, like the organization of the scientific congress in September 2001 and the creation of the work group's web site. Although these tasks meant additional work, the experience I gained from fulfilling them is immeasurable.

I own my warmest thanks to my wife Lia Andes, who greatly supported me at all times. With her devoted love she gave me the strength and the trust to carry on with the work and to deal with whatsoever came across.

I also thank my colleagues of the lab for their good team spirit and their contribution to a nice working atmosphere: Franca Bläsing, Christian Rabel, Günter Ziegelin, Jan Deneke, Stefan Ehrentraut, Marianne Schlicht - thank you.

I thank Prof. Wolfram Saenger and Prof. Walter Messer of the Free University of Berlin for their willingness to appraise this thesis. I further thank Prof. Hans Lehrach for his support and concern for my work.

I address my special thanks to Franca Bläsing, who is not only the most enjoyable colleague I ever worked with, but was also willing to critically read my manuscripts intended for publication or for the present thesis.

The work at the institute wouldn't have been as pleasant if there wasn't the Electron Microscopy group of Dr. Rudi Lurz. I greatly appreciated the company of Rudi and of Gerhild, with whom it was a pleasure to work with and to spend the time during the breaks.

Last but not least, I am indebted to my parents Horst Schröder and Lieve Seré, who, through their constant love and care, have prepared me most excellently for life. I also thank my brother Holger Schröder for the most inspiring times we spent together.