

Index of Tables

<u>Table</u>	<u>Page</u>
<u>Table 1:</u> Age of control horses and patients in years	38
<u>Table 2:</u> Distribution of sexes among the groups of controls and patients	38
<u>Table 3:</u> Distribution of the different breeds in the groups of controls and patients	39
<u>Table 4:</u> Distribution of the different joint types in the groups of controls and patients	39
<u>Table 5:</u> Criteria to the grading of lameness	40
<u>Table 6:</u> Radiographic views performed for each joint type	41
<u>Table 7:</u> Classification criteria of samples in diagnostic groups based on anamnestic data, clinical and radiographic examination and analysis of synovia	42
<u>Table 8:</u> pH values in synovia in the groups of controls and patients	47
<u>Table 9:</u> Total protein concentration (g/dl) in the groups of control and patients	48
<u>Table 10:</u> Total white blood cell count (cells/ μ l) in the groups of controls and patients	49
<u>Table 11:</u> MPO-Activity (mU/ μ l) in the groups of controls and patients	51
<u>Table 12:</u> Distribution of the different Degrees of synovial viscosity in the groups of controls and patients	52
<u>Table 13:</u> Distribution of the different degrees of lameness in the groups of controls and patients	53
<u>Table 14:</u> MPO-Activity (mU/ μ l) in samples with different degrees of synovial viscosity	57
<u>Table 15:</u> MPO-Activity (mU/ μ l) detected in samples taken from limbs showing different degrees of lameness	58