

## 7. Literaturverzeichnis

Adachi,M., Watanabe-Fukunaga,R., and Nagata,S. (1993). Aberrant transcription caused by the insertion of an early transposable element in an intron of the Fas antigen gene of lpr mice. *Proc.Natl.Acad.Sci.U.S.A* 90, 1756-1760.

Aktas,O., Smorodchenko,A., Brocke,S., Infante-Duarte,C., Topphoff,U.S., Vogt,J., Prozorovski,T., Meier,S., Osmanova,V., Pohl,E., Bechmann,I., Nitsch,R., Zipp,F. (2005). Neuronal damage in autoimmune neuroinflammation mediated by the death ligand TRAIL. *Neuron* 46, 421-432.

Amor,S., Groome,N., Linington,C., Morris,M.M., Dornmair,K., Gardinier,M.V., Matthieu,J.M., Baker,D. (1994). Identification of epitopes of myelin oligodendrocyte glycoprotein for the induction of experimental allergic encephalomyelitis in SJL and Biozzi AB/H mice. *J.Immunol.* 153, 4349-4356.

Arcscott,P.L., Stokes,T., Myc,A., Giordano,T.J., Thompson,N.W., Baker,J.R., Jr. (1999). Fas (CD95) expression is up-regulated on papillary thyroid carcinoma. *J.Clin.Endocrinol.Metab* 84, 4246-4252.

Ashkenazi,A. Dixit,V.M. (1998). Death receptors: signaling and modulation. *Science* 281, 1305-1308.

Ashkenazi,A., Pai,R.C., Fong,S., Leung,S., Lawrence,D.A., Marsters,S.A., Blackie,C., Chang,L., McMurtrey,A.E., Hebert,A., DeForge,L., Koumenis,I.L., Lewis,D., Harris,L., Bussiere,J., Koeppen,H., Shahrokh,Z., Schwall,R.H. (1999). Safety and antitumor activity of recombinant soluble Apo2 ligand. *J.Clin.Invest.* 104, 155-162.

Barnett,M.H. Prineas,J.W. (2004). Relapsing and remitting multiple sclerosis: pathology of the newly forming lesion. *Ann.Neurol.* 55, 458-468.

Bauer,J., Stadelmann,C., Bancher,C., Jellinger,K., Lassmann,H. (1999). Apoptosis of T lymphocytes in acute disseminated encephalomyelitis. *Acta Neuropathol.(Berl.)* 97, 543-546.

- Bjartmar,C., Kidd,G., Mork,S., Rudick,R., Trapp,B.D. (2000). Neurological disability correlates with spinal cord axonal loss and reduced N-acetyl aspartate in chronic multiple sclerosis patients. *Ann.Neurol.* *48*, 893-901.
- Bjartmar,C., Kinkel,R.P., Kidd,G., Rudick,R.A., Trapp,B.D. (2001). Axonal loss in normal-appearing white matter in a patient with acute MS. *Neurology* *57*, 1248-1252.
- Bjartmar,C. Trapp,B.D. (2001). Axonal and neuronal degeneration in multiple sclerosis: mechanisms and functional consequences. *Curr.Opin.Neurol.* *14*, 271-278.
- Bo,L., Vedeler,C.A., Nyland,H., Trapp,B.D., Mork,S.J. (2003a). Intracortical multiple sclerosis lesions are not associated with increased lymphocyte infiltration. *Mult.Scler.* *9*, 323-331.
- Bo,L., Vedeler,C.A., Nyland,H.I., Trapp,B.D., Mork,S.J. (2003b). Subpial demyelination in the cerebral cortex of multiple sclerosis patients. *J Neuropathol Exp Neurol* *62*, 723-732.
- Bonavida,B., Ng,C.P., Jazirehi,A., Schiller,G., Mizutani,Y. (1999). Selectivity of TRAIL-mediated apoptosis of cancer cells and synergy with drugs: the trail to non-toxic cancer therapeutics (review). *Int.J.Oncol.* *15*, 793-802.
- Bonetti,B. Raine,C.S. (1997). Multiple sclerosis: oligodendrocytes display cell death-related molecules in situ but do not undergo apoptosis. *Ann.Neurol.* *42*, 74-84.
- Brabb,T., Goldrath,A.W., von Dassow,P., Paez,A., Liggitt,H.D., Goverman,J. (1997). Triggers of autoimmune disease in a murine TCR-transgenic model for multiple sclerosis. *J Immunol.* *159*, 497-507.
- Bruck,W., Kuhlmann,T., Stadelmann,C. (2003). Remyelination in multiple sclerosis. *J Neurol Sci.* *206*, 181-185.
- Bustin,S.A. (2000). Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *J.Mol.Endocrinol.* *25*, 169-193.
- Cardullo,R.A., Agrawal,S., Flores,C., Zamecnik,P.C., Wolf,D.E. (1988). Detection of nucleic acid hybridization by nonradiative fluorescence resonance energy transfer. *Proc.Natl.Acad.Sci.U.S.A* *85*, 8790-8794.
- Chien,A., Edgar,D.B., Trela,J.M. (1976). Deoxyribonucleic acid polymerase from the extreme thermophile *Thermus aquaticus*. *J.Bacteriol.* *127*, 1550-1557.

- Chinnaiyan,A.M., Orth,K., O'Rourke,K., Duan,H., Poirier,G.G., Dixit,V.M. (1996). Molecular ordering of the cell death pathway. Bcl-2 and Bcl-xL function upstream of the CED-3-like apoptotic proteases. *J.Biol.Chem.* 271, 4573-4576.
- Claudio,L., Kress,Y., Factor,J., Brosnan,C.F. (1990). Mechanisms of edema formation in experimental autoimmune encephalomyelitis. The contribution of inflammatory cells. *Am.J.Pathol.* 137, 1033-1045.
- Cleveland,J.L. Ihle,J.N. (1995). Contenders in FasL/TNF death signaling. *Cell* 81, 479-482.
- Compston,D.A.S. Sadovnick,A.D. (1992). Epidemiology and Genetics of Multiple Sclerosis. *Curr.Opinion Neurol.Neurosurgery* 5, 175-181.
- Cretney,E., Takeda,K., Yagita,H., Glaccum,M., Peschon,J.J., Smyth,M.J. (2002). Increased susceptibility to tumor initiation and metastasis in TNF-related apoptosis-inducing ligand-deficient mice. *J.Immunol.* 168, 1356-1361.
- Cretney,E., Uldrich,A.P., Berzins,S.P., Strasser,A., Godfrey,D.I., Smyth,M.J. (2003). Normal thymocyte negative selection in TRAIL-deficient mice. *J.Exp.Med.* 198, 491-496.
- Critchfield,J.M., Racke,M.K., Zuniga-Pflucker,J.C., Cannella,B., Raine,C.S., Goverman,J., Lenardo,M.J. (1994). T cell deletion in high antigen dose therapy of autoimmune encephalomyelitis. *Science* 263, 1139-1143.
- De Stefano,N., Matthews,P.M., Fu,L., Narayanan,S., Stanley,J., Francis,G.S., Antel,J.P., Arnold,D.L. (1998). Axonal damage correlates with disability in patients with relapsing-remitting multiple sclerosis. Results of a longitudinal magnetic resonance spectroscopy study. *Brain* 121 ( Pt 8), 1469-1477.
- Degli-Esposti,M.A., Dougall,W.C., Smolak,P.J., Waugh,J.Y., Smith,C.A., Goodwin,R.G. (1997a). The novel receptor TRAIL-R4 induces NF-kappaB and protects against TRAIL-mediated apoptosis, yet retains an incomplete death domain. *Immunity* 7, 813-820.
- Degli-Esposti,M.A., Smolak,P.J., Walczak,H., Waugh,J., Huang,C.P., DuBose,R.F., Goodwin,R.G., Smith,C.A. (1997b). Cloning and characterization of TRAIL-R3, a novel member of the emerging TRAIL receptor family. *J.Exp.Med.* 186, 1165-1170.
- Demjen,D., Klussmann,S., Kleber,S., Zuliani,C., Stieltjes,B., Metzger,C., Hirt,U.A., Walczak,H., Falk,W., Essig,M., Edler,L., Krammer,P.H., Martin-Villalba,A. (2004). Neutralization of CD95

ligand promotes regeneration and functional recovery after spinal cord injury. *Nat.Med.* 10, 389-395.

Desbarats,J., Birge,R.B., Mimouni-Rongy,M., Weinstein,D.E., Palerme,J.S., Newell,M.K. (2003). Fas engagement induces neurite growth through ERK activation and p35 upregulation. *Nat.Cell Biol.* 5, 118-125.

Diem R, Tschirne A, Bähr M (2003). Decreased amplitudes in multiple sclerosis patients with normal visual acuity: a VEP study. *J Clin Neurosci* 7, 69-71.

Diestel,A., Aktas,O., Hackel,D., Hake,I., Meier,S., Raine,C.S., Nitsch,R., Zipp,F., Ullrich,O. (2003). Activation of microglial poly(ADP-ribose)-polymerase-1 by cholesterol breakdown products during neuroinflammation: a link between demyelination and neuronal damage. *J Exp Med* 198, 1729-740.

Dörr,J., Bechmann,I., Waiczies,S., Aktas,O., Walczak,H., Krammer,P.H., Nitsch,R., Zipp,F. (2002a). Lack of tumor necrosis factor-related apoptosis-inducing ligand but presence of its receptors in the human brain. *J.Neurosci.* 22, RC209.

Dörr,J., Waiczies,S., Wendling,U., Seeger,B., Zipp,F. (2002b). Induction of TRAIL-mediated glioma cell death by human T cells. *J.Neuroimmunol.* 122, 117-124.

Dowling,P., Shang,G., Raval,S., Menonna,J., Cook,S., Husar,W. (1996). Involvement of the CD95 (APO-1/Fas) receptor/ligand system in multiple sclerosis brain. *J.Exp.Med.* 184, 1513-1518.

Ehrlich,S., Infante-Duarte,C., Seeger,B., Zipp,F. (2003). Regulation of soluble and surface-bound TRAIL in human T cells, B cells, and monocytes. *Cytokine* 24, 244-253.

Enari,M., Hug,H., Nagata,S. (1995). Involvement of an ICE-like protease in Fas-mediated apoptosis. *Nature* 375, 78-81.

Evangelou,N., Konz,D., Esiri,M.M., Smith,S., Palace,J., Matthews,P.M. (2000). Regional axonal loss in the corpus callosum correlates with cerebral white matter lesion volume and distribution in multiple sclerosis. *Brain* 123 ( Pt 9), 1845-1849.

Ferguson,B., Matyszak,M.K., Esiri,M.M., Perry,V.H. (1997). Axonal damage in acute multiple sclerosis lesions. *Brain* 120 ( Pt 3), 393-399.

- Fillmore,P.D., Brace,M., Troutman,S.A., Blankenhorn,E.P., Diehl,S., Rincon,M., Teuscher,C. (2003). Genetic Analysis of the Influence of Neuroantigen-Complete Freund's Adjuvant Emulsion Structures on the Sexual Dimorphism and Susceptibility to Experimental Allergic Encephalomyelitis. *Am.J.Pathol.* 163, 1623-1632.
- Fowlkes,B.J., Schwartz,R.H., Pardoll,D.M. (1988). Deletion of self-reactive thymocytes occurs at a CD4+8+ precursor stage. *Nature* 334, 620-623.
- Ganter,P., Prince,C., Esiri,M.M. (1999). Spinal cord axonal loss in multiple sclerosis: a post-mortem study. *Neuropathol.Appl.Neurobiol.* 25, 459-467.
- Ginzinger,D.G. (2002). Gene quantification using real-time quantitative PCR: an emerging technology hits the mainstream. *Exp.Hematol.* 30, 503-512.
- Goverman,J., Woods,A., Larson,L., Weiner,L.P., Hood,L., Zaller,D.M. (1993). Transgenic mice that express a myelin basic protein-specific T cell receptor develop spontaneous autoimmunity. *Cell* 72, 551-560.
- Hahne,M., Renno,T., Schroeter,M., Irmeler,M., French,L., Bornard,T., MacDonald,H.R., Tschopp,J. (1996). Activated B cells express functional Fas ligand. *Eur.J.Immunol.* 26, 721-724.
- Haines,J.L., Terwedow,H.A., Burgess,K., Percak-Vance,M.A., Rimmler,J.B., Martin,E.R., Oksenberg,J.R., Lincoln,R., Zhang,D.Y., Banatao,D.R., Gatto,N., Goodkin,D.E., Hauser,S.L. (1998). Linkage of the MHC to familial multiple sclerosis suggests genetic heterogeneity. *Hum.Molec.Genet.* 7, 1229-1234.
- Hein,T. Hopfenmuller,W. (2000). [Projection of the number of multiple sclerosis patients in Germany]. *Nervenarzt* 71, 288-294.
- Higuchi,R., Fockler,C., Dollinger,G., Watson,R. (1993). Kinetic PCR analysis: real-time monitoring of DNA amplification reactions. *Biotechnology (N.Y.)* 11, 1026-1030.
- Hilliard,B., Wilmen,A., Seidel,C., Liu,T.S., Goke,R., Chen,Y. (2001). Roles of TNF-Related Apoptosis-Inducing Ligand in Experimental Autoimmune Encephalomyelitis. *J.Immunol.* 166, 1314-1319.
- Hirata,H., Hibasami,H., Yoshida,T., Morita,A., Ohkaya,S., Matsumoto,M., Sasaki,H., Uchida,A. (1998). Differentiation and apoptosis without DNA fragmentation in cultured Schwann cells derived from wallerian-degenerated nerve. *Apoptosis.* 3, 353-360.

Hobom,M., Storch,M.K., Weissert,R., Maier,K., Radhakrishnan,A., Kramer,B., Bähr,M., Diem,R. (2004). Mechanisms and Time Course of Neuronal Degeneration in Experimental Autoimmune Encephalomyelitis. *Brain Pathol* 14, 148-157.

Holland,P.M., Abramson,R.D., Watson,R., Gelfand,D.H. (1991). Detection of specific polymerase chain reaction product by utilizing the 5'-3' exonuclease activity of *Thermus aquaticus* DNA polymerase. *Proc.Natl.Acad.Sci.U.S.A* 88, 7276-7280.

Hueber,A.O., Zornig,M., Bernard,A.M., Chautan,M., Evan,G. (2000). A dominant negative Fas-associated death domain protein mutant inhibits proliferation and leads to impaired calcium mobilization in both T-cells and fibroblasts. *J.Biol.Chem.* 275, 10453-10462.

Ishiguro,T., Saitoh,J., Yawata,H., Yamagishi,H., Iwasaki,S., Mitoma,Y. (1995). Homogeneous quantitative assay of hepatitis C virus RNA by polymerase chain reaction in the presence of a fluorescent intercalater. *Anal.Biochem.* 229, 207-213.

Itoh,N. Nagata,S. (1993). A novel protein domain required for apoptosis. Mutational analysis of human Fas antigen. *J.Biol.Chem.* 268, 10932-10937.

Janeway,C.A. (1999). T-cell development: a role for self-peptides in positive selection. *Curr.Biol.* 9, R342-R345.

Jo,M., Kim,T.H., Seol,D.W., Esplen,J.E., Dorko,K., Billiar,T.R., Strom,S.C. (2000). Apoptosis induced in normal human hepatocytes by tumor necrosis factor-related apoptosis-inducing ligand. *Nat.Med.* 6, 564-567.

Ju,S.T., Panka,D.J., Cui,H., Ettinger,R., El-Khatib,M., Sherr,D.H., Stanger,B.Z., Marshak-Rothstein,A. (1995). Fas (CD95)/FasL interactions required for programmed cell death after T cell-activation. *Nature* 373, 444-448.

Kabat,E.A., Wolf,A., and Bezer,A.E. (1947). The rapid production of acute disseminated encephalomyelitis in rhesus monkeys by injection of heterologous and homologous brain tissue with adjuvants. *J Exp Med* 117-130.

Kamradt,T., Soloway,P.D., Perkins,D.L., Gefter,M.L. (1991). Pertussis toxin prevents the induction of peripheral T cell anergy and enhances the T cell response to an encephalitogenic peptide of myelin basic protein. *J Immunol.* 147, 3296-3302.

Kawasaki,E. (1989). Amplification of RNA sequences via complementary DNA (cDNA). Amplifications 4-6.

Kerr,J.F., Wyllie,A.H., Currie,A.R. (1972). Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br.J.Cancer* 26, 239-257.

Kerschensteiner,M., Gallmeier,E., Behrens,L., Leal,V.V., Misgeld,T., Klinkert,W.E., Kolbeck,R., Hoppe,E., Oropeza-Wekerle,R.L., Bartke,I., Stadelmann,C., Lassmann,H., Wekerle,H., Hohlfeld,R. (1999). Activated human T cells, B cells, and monocytes produce brain-derived neurotrophic factor in vitro and in inflammatory brain lesions: a neuroprotective role of inflammation? *J Exp Med* 189, 865-870.

Kischkel,F.C., Hellbardt,S., Behrmann,I., Germer,M., Pawlita,M., Krammer,P.H., Peter,M.E. (1995). Cytotoxicity-dependent APO-1 (Fas/CD95)-associated proteins form a death-inducing signaling complex (DISC) with the receptor. *EMBO J.* 14, 5579-5588.

Kornek,B. Lassmann,H. (2003). Neuropathology of multiple sclerosis- new concepts. *Brain Research Bulletin* 61, 321-326.

Kornek,B., Storch,M.K., Weissert,R., Wallstroem,E., Stefferl,A., Olsson,T., Linington,C., Schmidbauer,M., Lassmann,H. (2000). Multiple sclerosis and chronic autoimmune encephalomyelitis: a comparative quantitative study of axonal injury in active, inactive, and remyelinated lesions. *Am.J.Pathol.* 157, 267-276.

Kurtzke,J.F. (1985). Epidemiology of multiple sclerosis. In *Handbook of Clinical Neurology, Demyelinating Diseases*, P. J. Vinken, G. W. Bruyn, H. L. Klawans, J. C. Koetsier, eds. (Amsterdam, New York: Elsevier Sci.), pp. 259-287.

Kurtzke,J.F. (2000). Multiple sclerosis in time and space - geographic clues to cause. *J.Neurovirol. Suppl.* 2, 134-140.

Lamhamedi-Cherradi,S.E., Zheng,S.J., Maguschak,K.A., Peschon,J., Chen,Y.H. (2003). Defective thymocyte apoptosis and accelerated autoimmune diseases in TRAIL-/- mice. *Nat.Immunol.* 4, 255-260.

Lassmann,H., Bruck,W., Lucchinetti,C. (2001). Heterogeneity of multiple sclerosis pathogenesis: implications for diagnosis and therapy. *Trends Mol.Med* 7, 115-121.

- Lenardo,M.J., Chan,K.M., Hornung,F., McFarland,H.F., Siegel,R., Wang,J., Zheng,L. (1999). Mature T lymphocyte apoptosis--immune regulation in a dynamic and unpredictable antigenic environment. *Annu.Rev.Immunol.* 17, 221-253.
- Leutenegger,C.M., Higgins,J., Matthews,T.B., Tarantal,A.F., Luciw,P.A., Pedersen,N.C., North,T.W. (2001). Real-time TaqMan PCR as a specific and more sensitive alternative to the branched-chain DNA assay for quantitation of simian immunodeficiency virus RNA. *AIDS Res.Hum.Retroviruses* 17, 243-251.
- Leverkus,M., Walczak,H., McLellan,A., Fries,H.W., Terbeck,G., Brocker,E.B., and Kampgen,E. (2000). Maturation of dendritic cells leads to up-regulation of cellular FLICE-inhibitory protein and concomitant down-regulation of death ligand-mediated apoptosis. *Blood* 96, 2628-2631.
- Lewen,A., Li,G.L., Nilsson,P., Olsson,Y., Hillered,L. (1995). Traumatic brain injury in rat produces changes of beta-amyloid precursor protein immunoreactivity. *Neuroreport* 6, 357-360.
- Lie,Y.S. Petropoulos,C.J. (1998). Advances in quantitative PCR technology: 5' nuclease assays. *Curr.Opin.Biotechnol.* 9, 43-48.
- Livak,K.J. Schmittgen,T.D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25, 402-408.
- Lovas,G., Szilagyi,N., Majtenyi,K., Palkovits,M., Komoly,S. (2000). Axonal changes in chronic demyelinated cervical spinal cord plaques. *Brain* 123 ( Pt 2), 308-317.
- Lucchinetti,C., Bruck,W., Parisi,J., Scheithauer,B., Rodriguez,M., Lassmann,H. (1999). A quantitative analysis of oligodendrocytes in multiple sclerosis lesions. A study of 113 cases. *Brain* 122 ( Pt 12), 2279-2295.
- Lucchinetti,C.F., Brück,W., Parisi,J., Scheithauer,B., Rodriguez,M., Lassmann,H. (2000). Heterogeneity of multiple sclerosis lesions: implications for the pathogenesis of demyelination. *Ann.Neurol.* 47, 707-717.
- Lucchinetti,C.F., Bruck,W., Rodriguez,M., Lassmann,H. (1996). Distinct patterns of multiple sclerosis pathology indicates heterogeneity on pathogenesis. *Brain Pathol* 6, 259-274.
- Lunemann,J.D., Waiczies,S., Ehrlich,S., Wendling,U., Seeger,B., Kamradt,T., Zipp,F. (2002). Death ligand TRAIL induces no apoptosis but inhibits activation of human (auto)antigen-specific T cells. *J.Immunol.* 168, 4881-4888.



- Malipiero,U., Frei,K., Spanaus,K.S., Agresti,C., Lassmann,H., Hahne,M., Tschopp,J., Eugster,H.P., Fontana,A. (1997). Myelin oligodendrocyte glycoprotein-induced encephalomyelitis is chronic/relapsing in perforin knockout mice, but monophasic in Fas- and Fas ligand-deficient *lpr* and *gld* mice. *Eur.J.Immunol.* 27, 3151-3160.
- Mariani,S.M. Krammer,P.H. (1998). Surface expression of TRAIL/Apo-2 ligand in activated mouse T and B cells. *Eur.J.Immunol.* 28, 1492-1498.
- Martin,R. McFarland,H.F. (1995). Immunological aspects of experimental allergic encephalomyelitis and multiple sclerosis. *Crit.Rev.Clin.Lab.Sci.* 32, 121-182.
- Martin,R., McFarland,H.F., McFarlin,D.E. (1992a). Immunological aspects of demyelinating diseases. *Annu.Rev.Immunol.* 10, 153-187.
- Martin,R., Utz,U., Coligan,J.E., Richert,J.R., Flerlage,M., Robinson,E., Stone,R., Biddison,W.E., McFarlin,D.E., McFarland,H.F. (1992b). Diversity in fine specificity and T cell receptor usage of the human CD4+ cytotoxic T cell response specific for the immunodominant myelin basic protein peptide 87-106. *J.Immunol.* 148, 1359-1366.
- Martinez-Lorenzo,M.J., Anel,A., Garnen,S., Monleon,I., Lasierra,P., Larrad,L., Pineiro,A., Alava,M.A., Naval,J. (1999). Activated human T cells release bioactive Fas ligand and APO2 ligand in microvesicles. *J.Immunol.* 163, 1274-1281.
- Matysiak,M., Jurewicz,A., Jaskolski,D., Selmaj,K. (2002). TRAIL induces death of human oligodendrocytes isolated from adult brain. *Brain* 125, 2469-2480.
- Meyer,R., Weissert,R., Diem R, Storch,M.K., de Graaf,K.L., Kramer,B., Bähr M (2001). Acute Neuronal Apoptosis in a Rat Model of Multiple Sclerosis. *J.Neurosci.* 21, 6214-6220.
- Miura,Y., Misawa,N., Kawano,Y., Okada,H., Inagaki,Y., Yamamoto,N., Ito,M., Yagita,H., Okumura,K., Mizusawa,H., Koyanagi,Y. (2003). Tumor necrosis factor-related apoptosis-inducing ligand induces neuronal death in a murine model of HIV central nervous system infection. *Proc.Natl.Acad.Sci.U.S.A* 100, 2777-2782.
- Morgan,I.M. (1946). Allergic encephalomyelitis in monkeys in response to injection of normal monkey cord. *J.Bakteriol.* 614-615.

- Muzio,M. (1996). FLICE, a novel FADD-homologous ICE/CED-like protease, is recruited to the CD95 (Fas/APO-1) death-inducing signaling complex (DISC). *Cell* 85, 817-827.
- Nagata,S. (1997). Apoptosis by Death Factor. *Cell* 88, 355-365.
- Nagata,S. (2000). Apoptotic DNA fragmentation. *Exp.Cell Res.* 256, 12-18.
- Nagata,S. Golstein,P. (1995). The Fas death factor. *Science* 267, 1449-1456.
- Nitsch,R., Bechmann,I., Deisz,R.A., Haas,D., Lehmann,T.N., Wendling,U., Zipp,F. (2000). Massive cell death induced by tumor-necrosis factor-related apoptosis-inducing ligand (TRAIL) in adult human brain tissue. *Lancet* 356, 827-828.
- Okuda,Y., Sakoda,S., Fujimura,H., Nagata,S., Yanagihara,T., Bernard,C.C. (2000). Intrathecal Administration of Neutralizing Antibody against Fas Ligand Suppresses the Progression of Experimental Autoimmune Encephalomyelitis. *Biochem.biophys.Res.Comm.* 275, 164-168.
- Olsson,T., Wei Zhi,W., HÜjeberg,B., Kostulas,V., Yu-Ping,J., Anderson,G., Ekre,H.P., Link,H. (1990). Autoreactive T Lymphocytes in Multiple Sclerosis Determined by Antigen-Induced Secretion of Interferon-g. *J.Clin.Invest.* 86, 981-985.
- Pan,G., O'Rourke,K., Chinnaiyan,A.M., Gentz,R., Ebner,R., Ni,J., Dixit,V.M. (1997b). The receptor for the cytotoxic ligand TRAIL. *Science* 276, 111-113.
- Paterson,P. (1960). Transfer of allergic encephalomyelitis in rats by means of lymph node cells. *J Exp Med* 111, 119-135.
- Pender,M.P. (1998). Genetically determined failure of activation-induced apoptosis of autoreactive T cells as a cause of multiple sclerosis. *Lancet* 351, 978-981.
- Pitti,R.M., Marsters,S.A., Ruppert,S., Donahue,C.J., Moore,A., Ashkenazi,A. (1996). Induction of apoptosis by Apo-2 ligand, a new member of the tumor necrosis factor cytokine family. *J.Biol.Chem.* 271, 12687-12690.
- Pollak,Y., Ovadia,H., Goshen,I., Gurevich,R., Monsa,K., Avitsur,R., Yirmiyya,R. (2000). Behavioral aspects of experimental autoimmune encephalomyelitis. *J.Neuroimmunol.* 104, 31-36.

Pollak,Y., Ovadia,H., Orion,E., Weidenfeld,J., Yirmiya,R. (2003). The EAE-associated behavioral syndrome: I. Temporal correlation with inflammatory mediators. *J.Neuroimmunol.* *137*, 94-99.

Reddy,H., Narayanan,S., Woolrich,M., Mitsumori,T., Lapierre,Y., Arnold,D.L., Matthews,P.M. (2002). Functional brain reorganization for hand movement in patients with multiple sclerosis: defining distinct effects of injury and disability. *Brain* *125*, 2646-2657.

Rivers,T.M., Berry,G.P., Sprunt,D.H. (1933). Observations on attempts to produce acute encephalomyelitis in monkeys. *J Exp Med* *39*.

Sabelko,K.A., Kelly,K.A., Nahm,M.H., Cross,A.H., Russell,J.H. (1997). Fas and Fas ligand enhance the pathogenesis of experimental allergic encephalomyelitis, but are not essential for immune privilege in the central nervous system. *J.Immunol.* *159*, 3096-3099.

Sabelko-Downes,K.A., Cross,A.H., Russell,J.H. (1999). Dual role for Fas ligand in the initiation of and recovery from experimental allergic encephalomyelitis. *J.Exp.Med.* *189*, 1195-1205.

Sahara,S., Aoto,M., Eguchi,Y., Imamoto,N., Yoneda,Y., Tsujimoto,Y. (1999). Acinus is a caspase-3-activated protein required for apoptotic chromatin condensation. *Nature* *401*, 168-173.

Schneider,P., Bodmer,J.L., Thome,M., Hofmann,K., Holler,N., Tschopp,J. (1997). Characterization of two receptors for TRAIL. *FEBS Lett.* *416*, 329-334.

Schneider,P., Olson,D., Tardivel,A., Browning,B., Lugovskoy,A., Gong,D., Dobles,M., Hertig,S., Hofmann,K., Van Vlijmen,H., Hsu,Y.M., Burkly,L.C., Tschopp,J., Zheng,T.S. (2003). Identification of a new murine tumor necrosis factor receptor locus that contains two novel murine receptors for tumor necrosis factor-related apoptosis-inducing ligand (TRAIL). *J.Biol.Chem.* *278*, 5444-5454.

Selvin,P.R. (1995). Fluorescence resonance energy transfer. *Methods Enzymol.* *246*, 300-334.

Sheridan,J.P., Marsters,S.A., Pitti,R.M., Gurney,A., Skubatch,M., Baldwin,D., Ramakrishnan,L., Gray,C.L., Baker,K., Wood,W.I., Goddard,A.D., Godowski,P., Ashkenazi,A. (1997). Control of TRAIL-induced apoptosis by a family of signaling and decoy receptors. *Science* *277*, 818-821.

Simonet,W.S., Lacey,D.L., Dunstan,C.R., Kelley,M., Chang,M.S., Luthy,R., Nguyen,H.Q., Wooden,S., Bennet,L., Boone,T., Shimamoto,G., DeRose,M., Elliot,R., Colombero,A., Tan,H.L., Trail,G., Sullivan,J., Davy,E., Bucay,N., Renshaw-Gegg,L., Hughes,T.M., Hill,D., Pattison,W.,

Campbell,P., Sanders,S., Van,G., Tarpley,J., Derby,P., Lee,R., Program,A.E., Boyle,W.J. (1997). Osteoprotegerin: a novel secreted protein involved in the regulation of bone density. *Cell* 89, 309-319.

Smyth,M.J., Cretney,E., Takeda,K., Wiltrott,R.H., Sedger,L.M., Kayagaki,N., Yagita,H., Okumura,K. (2001). Tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) contributes to interferon gamma-dependent natural killer cell protection from tumor metastasis. *J.Exp.Med.* 193, 661-670.

Sobel,R.A., Greer,J.M., Isaac,J., Fondren,G., Lees,M.B. (1994). Immunolocalization of proteolipid protein peptide 103-116 in myelin. *J.Neurosci.Res.* 37, 36-43.

Song,K., Chen,Y., Goke,R., Wilmen,A., Seidel,C., Goke,A., Hilliard,B. (2000). Tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) is an inhibitor of autoimmune inflammation and cell cycle progression. *J.Exp.Med.* 191, 1095-1103.

Stadelmann,C., Kerschensteiner,M., Misgeld,T., Bruck,W., Hohlfeld,R., Lassmann,H. (2002). BDNF and gp145trkB in multiple sclerosis brain lesions: neuroprotective interactions between immune and neuronal cells? *Brain* 125, 75-85.

Stanger,B.Z., Leder,P., Lee,T.H., Kim,E., Seed,B. (1995). RIP: a novel protein containing a death domain that interacts with Fas/APO-1 (CD95) in yeast and causes cell death. *Cell* 81, 513-523.

Stuart,G. Krikorian,K. (1928). The neuro-paralytic accidents of anti-rabies treatment. *Ann.Trop.Med.Parasitol* 327-377.

Suda,T., Takahashi,T., Golstein,P., Nagata,S. (1993). Molecular cloning and expression of the Fas ligand, a novel member of the tumor necrosis factor family. *Cell* 75, 1169-1178.

Suter,T., Biollaz,G., Gatto,D., Bernasconi,L., Herren,T., Reith,W., Fontana,A. (2003). The brain as an immune privileged site: dendritic cells of the central nervous system inhibit T cell activation. *Eur.J.Immunol.* 33, 2998-3006.

Tabi,Z., McCombe,P.A., Pender,M.P. (1995). Antigen-specific down-regulation of myelin basic protein-reactive T cells during spontaneous recovery from experimental autoimmune encephalomyelitis: further evidence of apoptotic deletion of autoreactive T cells in the central nervous system. *Int.Immunol.* 7, 967-973.