

## 6. Literaturverzeichnis

[1] Tan, BB, Flaherty, KR, Kazerooni, EA und Iannettoni, MD. The solitary pulmonary nodule. *Chest* (2003); 123:89S-96S.

[2] Lillington, G. Management of solitary pulmonary nodules. *Dis Month* (1991); 37:274-318.

[3] Siegelman, SS, Khouri, NF, Leo, FP, Fishman, EK, Braverman, RM und Zerhouni, EA. Solitary pulmonary nodules: CT assessment. *Radiology* (1986); 160:307-312.

[4] Siegelman, SS, Zerhouni, EA, Leo, FP, Khouri, NF und Stitik, FP. CT of the solitary pulmonary nodule. *AJR Am J Roentgenol* (1980); 135:1-13.

[5] Khouri, NF, Meziane, MA, Zerhouni, EA, Fishman, EK und Siegelman, SS. The solitary pulmonary nodule. Assessment, diagnosis, and management. *Chest* (1987); 91:128-133.

[6] Higgins, GA, Shields, TW und Keehn, RJ. The solitary pulmonary nodule. Ten-year follow-up of veterans administration-armed forces cooperative study. *Arch Surg* (1975); 110:570-575.

[7] Ray, JF, III, Lawton, BR, Magnin, GE et al. The coin lesion story: update 1976. Twenty years' experience with thoracotomy for 179 suspected malignant coin lesions. *Chest* (1976); 70:332-336.

[8] Swensen, SJ, Jett, JR, Payne, WS, Viggiano, RW, Pairolero, PC und Trastek, VF. An integrated approach to evaluation of the solitary pulmonary nodule. *Mayo Clin Proc* (1990); 65:173-186.

[9] Toomes, H, Delphendahl, A, Manke, HG und Vogt-Moykopf, I. The coin lesion of the lung. A review of 955 resected coin lesions. *Cancer* (1983); 51:534-537.

[10] Ost, D und Fein, A. Evaluation and management of the solitary pulmonary nodule. *Am J Respir Crit Care Med* (2000); 162:782-787.

[11] Diaz-Jiménez, JP. Diagnostic approach of solitary pulmonary nodules. In: European Respiratory Society (Ed.). European Respiratory Monograph. Vol. 3. European Respiratory Society Journals Ltd., Sheffield (1998); S. 181-192.

[12] Pretreatment evaluation of non-small-cell lung cancer. The American Thoracic Society and The European Respiratory Society. Am J Respir Crit Care Med (1997); 156:320-332.

[13] National Collaborating Centre for Acute Care and National Institute for Clinical Excellence. Lung Cancer. The Diagnosis and Treatment of Lung Cancer. London (2005); <http://www.nice.org.uk/CG024NICEguideline>.

[14] Schönfeld, N. Differential Diagnosis. In: Sulier, J-P and Fry, WA (Eds.). Malignant Tumors of the Lung. Springer-Verlag, Berlin, Heidelberg, New York (2004); S. 131-138.

[15] Henschke, CI, Yankelevitz, DF, Naidich, DP et al. CT screening for lung cancer: suspiciousness of nodules according to size on baseline scans. Radiology (2004); 231:164-168.

[16] Swensen, SJ, Jett, JR, Hartman, TE et al. CT screening for lung cancer: five-year prospective experience. Radiology (2005); 235:259-265.

[17] Henschke, CI, Yankelevitz, DF, Libby, DM, Pasmantier, MW, Smith, JP und Miettinen, OS. Survival of patients with stage I lung cancer detected on CT screening. N Engl J Med (2006); 355:1763-1771.

[18] Unger, M. A pause, progress, and reassessment in lung cancer screening (Editorial). N Engl J Med (2006); 355:1822-1824.

[19] Edell, ES und Sanderson, DR. History of bronchoscopy. In: Prakash, U (Ed.). Bronchoscopy Raven Press, Ltd., New York (1994); S. 7-11.

[20] Andersen, HA, Fontana, RS und Harrison, EG. Transbronchoscopic lung biopsy in diffuse pulmonary disease. Dis Chest (1965); 48:187-192.

[21] Brandt, HJ. Beiträge zur Technik der Materialgewinnung. In: Grunze, H (Ed.). Klinische Zytologie der Thoraxkrankheiten Ferdinand Enke, Stuttgart (1955).

[22] Tanaka, M, Satoh, M, Kawanami, O und Aihara, K. A new bronchofiberscope for the study of diseases of very peripheral airways. *Chest* (1984); 85:590-594.

[23] Rodrigues, JC und Feinsilver, SH. Indications and contraindications for fiberoptic bronchoscopy. In: Feinsilver, SH und Fein, AM (Eds.). *Textbook of Bronchoscopy*. Williams & Wilkins, Baltimore (1995); S. 3-10.

[24] Cortese, DA und McDougall, JC. Bronchoscopy in peripheral and central lesions. In: Prakash, UBS (Ed.). *Bronchoscopy*. Raven Press Ltd., New York (1994); S. 135-140.

[25] Zavala, DC. Pulmonary hemorrhage in fiberoptic transbronchial biopsy. *Chest* (1976); 70:584-588.

[26] Pereira, W, Jr., Kovnat, DM und Snider, GL. A prospective cooperative study of complications following flexible fiberoptic bronchoscopy. *Chest* (1978); 73:813-816.

[27] Thomas, M, Gatzemeier, U, Goerg, R et al. Empfehlungen zur Diagnostik des Bronchialkarzinoms. Deutsche Gesellschaft für Pneumologie. *Pneumologie* (2000); 54:361-371.

[28] Zwirewich, CV, Vedal, S, Miller, RR und Muller, NL. Solitary pulmonary nodule: high-resolution CT and radiologic-pathologic correlation. *Radiology* (1991); 179:469-476.

[29] Zerhouni, EA, Stitik, FP, Siegelman, SS et al. CT of the pulmonary nodule: a cooperative study. *Radiology* (1986); 160:319-327.

[30] Bittner, RC und Felix, R. Magnetic resonance (MR) imaging of the chest: state-of-the-art. *Eur Respir J* (1998); 11:1392-1404.

[31] Gould, MK, Maclean, CC, Kuschner, WG, Rydzak, CE und Owens, DK. Accuracy of positron emission tomography for diagnosis of pulmonary nodules and mass lesions: a meta-analysis. *JAMA* (2001); 285:914-924.

[32] Hellwig, D, Ukena, D, Paulsen, F, Bamberg, M und Kirsch, CM. Metaanalyse zum Stellenwert der Positronen-Emissions-Tomographie mit F-18-Fluorodesoxyglukose

(FDG-PET) bei Lungentumoren. Diskussionsbasis der deutschen Konsensuskonferenz Onko-PET 2000. *Pneumologie* (2001); 55:367-377.

[33] Gemeinsamer Bundesausschuss. Bekanntmachung eines Beschlusses des Gemeinsamen Bundesausschusses zur Positronenemissionstomographie. (2005); [http://www.g-ba.de/cms/front\\_content.php?idcat=147](http://www.g-ba.de/cms/front_content.php?idcat=147)

[34] Robbins, HM, Sweet, ME, Jefferson, SE, Solomon, DA und Goldman, AL. The determination of resectability of lung cancer by fiberoptic bronchoscopy. *Arch Intern Med* (1981); 141:649-650.

[35] Loddenkemper, R und Schönfeld, N. Role of endoscopy in the preoperative assessment of bronchial carcinoma. *Monaldi Arch Chest Dis* (1994); 49:138-143.

[36] Baaklini, WA, Reinoso, MA, Gorin, AB, Sharafkaneh, A und Manian, P. Diagnostic yield of fiberoptic bronchoscopy in evaluating solitary pulmonary nodules. *Chest* (2000); 117:1049-1054.

[37] Chechani, V. Bronchoscopic diagnosis of solitary pulmonary nodules and lung masses in the absence of endobronchial abnormality. *Chest* (1996); 109:620-625.

[38] Mak, VH, Johnston, ID, Hetzel, MR und Grubb, C. Value of washings and brushings at fiberoptic bronchoscopy in the diagnosis of lung cancer. *Thorax* (1990); 45:373-376.

[39] Wallace, JM und Deutsch, AL. Flexible fiberoptic bronchoscopy and percutaneous needle lung aspiration for evaluating the solitary pulmonary nodule. *Chest* (1982); 81:665-671.

[40] Fletcher, EC und Levin, DC. Flexible fiberoptic bronchoscopy and fluoroscopically guided transbronchial biopsy in the management of solitary pulmonary nodules. *West J Med* (1982); 136:477-483.

[41] Lai, RS, Lee, SS, Ting, YM, Wang, HC, Lin, CC und Lu, JY. Diagnostic value of transbronchial lung biopsy under fluoroscopic guidance in solitary pulmonary nodule in an endemic area of tuberculosis. *Respir Med* (1996); 90:139-143.

[42] Loddenkemper, R, Schaberg, T, Mai, J, Preussler, H und Brandt, H-J. Bronchial evaluation of peripheral localised lung lesions: Accuracy of Different Biopsy Techniques. *Eur J Respir Dis Suppl* (1983); 64:461-464.

[43] McDougall, JC und Cortese, DA. Transbronchoscopic lung biopsy for localized pulmonary disease. *Sem Resp Med* (1981); 3:30-34.

[44] Arroliga, AC und Matthay, RA. The role of bronchoscopy in lung cancer. *Clin Chest Med* (1993); 14:87-98.

[45] Herth, FJ, Ernst, A und Becker, HD. Endobronchial ultrasound-guided transbronchial lung biopsy in solitary pulmonary nodules and peripheral lesions. *Eur Respir J* (2002); 20:972-974.

[46] Kurimoto, N, Miyazawa, T, Okimasa, S et al. Endobronchial ultrasonography using a guide sheath increases the ability to diagnose peripheral pulmonary lesions endoscopically. *Chest* (2004); 126:959-965.

[47] Herth, FJ, Eberhardt, R, Becker, HD und Ernst, A. Endobronchial ultrasound-guided transbronchial lung biopsy in fluoroscopically invisible solitary pulmonary nodules: a prospective trial. *Chest* (2006); 129:147-150.

[48] Asahina, H, Yamazaki, K, Onodera, Y et al. Transbronchial biopsy using endobronchial ultrasonography with a guide sheath and virtual bronchoscopic navigation. *Chest* (2005); 128:1761-1765.

[49] Solomon, SB, White, P, Jr., Wiener, CM, Orens, JB und Wang, KP. Three-dimensional CT-guided bronchoscopy with a real-time electromagnetic position sensor: a comparison of two image registration methods. *Chest* (2000); 118:1783-1787.

[50] Hautmann, H, Schneider, A, Pinkau, T, Peltz, F und Feussner, H. Electromagnetic catheter navigation during bronchoscopy: validation of a novel method by conventional fluoroscopy. *Chest* (2005); 128:382-387.

[51] Schwarz, Y, Greif, J, Becker, HD, Ernst, A und Mehta, A. Real-time electromagnetic navigation bronchoscopy to peripheral lung lesions using overlaid CT images: the first human study. *Chest* (2006); 129:988-994.

[52] Sinner, WN. Pulmonary neoplasms diagnosed with transthoracic needle biopsy. *Cancer* (1979); 43:1533-1540.

[53] Berquist, TH, Bailey, PB, Cortese, DA und Miller, WE. Transthoracic needle biopsy: accuracy and complications in relation to location and type of lesion. *Mayo Clin Proc* (1980); 55:475-481.

[54] Westcott, JL. Direct percutaneous needle aspiration of localized pulmonary lesions: result in 422 patients. *Radiology* (1980); 137:31-35.

[55] Khouri, NF, Stitik, FP, Erozan, YS et al. Transthoracic needle aspiration biopsy of benign and malignant lung lesions. *Am J Roentgenol* (1985); 144:281-288.

[56] Colquhoun, SD, Rosenthal, DL und Morton, DL. Role of percutaneous fine-needle aspiration biopsy in suspected intrathoracic malignancy. *Ann Thorac Surg* (1991); 51:390-393.

[57] Collins, CD, Breatnach, E und Nath, PH. Percutaneous needle biopsy of lung nodules following failed bronchoscopic biopsy. *Eur J Radiol* (1992); 15:49-53.

[58] Levine, MS, Weiss, JM, Harrell, JH, Cameron, TJ und Moser, KM. Transthoracic needle aspiration biopsy following negative fiberoptic bronchoscopy in solitary pulmonary nodules. *Chest* (1988); 93:1152-1155.

[59] Ohno, Y, Hatabu, H, Takenaka, D et al. CT-guided transthoracic needle aspiration biopsy of small (< or = 20 mm) solitary pulmonary nodules. *Am J Roentgenol* (2003); 180:1665-1669.

[60] Di Donna, A, Bazzocchi, M, Dolcet, F und Springolo, E. [CT-guided transthoracic needle aspiration of solitary lung lesions. Personal experience in 118 cases]. *Radiol Med (Torino)* (1995); 89:287-294.

[61] Golfieri, R, Sbrozzi, F, de Santis, F et al. [Clinical role of CT-guided transthoracic needle biopsy in the diagnosis of solitary pulmonary nodules]. *Radiol Med (Torino)* (1998); 95:329-337.

[62] Tanaka, M. Advances and usefulness of ultra-thin bronchofiberscopes. *Keio J Med* (1996); 45:296-300.

[63] Hasegawa, S, Hitomi, S, Murakawa, M und Mori, K. Development of an ultrathin fiberscope with a built-in channel for bronchoscopy in infants. *Chest* (1996); 110:1543-1546.

[64] Tohda, Y, Muraki, M, Iwanaga, T, Kubo, H, Haraguchi, R und Fukuoka, M. Applicability of ureteropelvic fiberscopy in the diagnosis of peripheral lung cancer. *Journal of Bronchology* (1999); 6:251-256.

[65] Rooney, CP, Wolf, K und McLennan, G. Ultrathin bronchoscopy as an adjunct to standard bronchoscopy in the diagnosis of peripheral lung lesions. A preliminary report. *Respiration* (2002); 69:63-68.

[66] Asano, F, Matsushita, T, Kondo, H, Saito, Y, Seko, A und Ishihara, Y. Transbronchial diagnosis of a pulmonary peripheral small lesion using an ultrathin bronchoscope with virtual bronchoscopic navigation. *Journal of Bronchology* (2002); 9:108-111.

[67] Asano, F, Matsuno, Y, Komaki, C et al. [CT-guided transbronchial diagnosis using ultrathin bronchoscope for small peripheral pulmonary lesions]. *Nihon Kokyuki Gakkai Zasshi* (2002); 40:11-16.

[68] Shinagawa, N, Yamazaki, K, Onodera, Y et al. CT-guided transbronchial biopsy using an ultrathin bronchoscope with virtual bronchoscopic navigation. *Chest* (2004); 125:1138-1143.

[69] Asano, F, Matsuno, Y, Shinagawa, N et al. A virtual bronchoscopic navigation system for pulmonary peripheral lesions. *Chest* (2006); 130:559-566.

[70] Schuurmans, MM, Michaud, GC, Diacon, AH und Bolliger, CT. Use of an ultrathin bronchoscope in the assessment of central airway obstruction. *Chest* (2003); 124:735-739.

[71] Oki, M, Saka, H, Sako, C et al. Cavitating invasive pulmonary aspergillosis visualized and diagnosed by ultrathin bronchoscopy. *Chest* (2006); 129:475-479.

[72] Oki, M, Saka, H, Kitagawa, C et al. Visceral pleural perforation in two cases of ultrathin bronchoscopy. *Chest* (2005); 127:2271-2273.