

L I T E R A T U R V E R Z E I C H N I S

- Aruoma,O.I., Halliwell,B., Hoey,B.M., and Butler,J. (1989): The antioxidant action of N-acetylcysteine: its reaction with hydrogen peroxide, hydroxyl radical, superoxide, and hypochlorous acid. *Free Radic.Biol.Med.*, 6:593-597.
- Asahi,M., Fujii,J., Suzuki,K., Seo,H.G., Kuzuya,T., Hori,M., Tada,M., Fujii,S., and Taniguchi,N. (1995): Inactivation of glutathione peroxidase by nitric oxide. Implication for cytotoxicity. *J.Biol.Chem.* , 270:21035-21039.
- Assaf,Y., Beit-Yannai,E., Shohami,E., Berman,E., and Cohen,Y. (1997): Diffusion- and T2-weighted MRI of closed-head injury in rats: a time course study and correlation with histology. *Magn Reson.Imaging*, 15:77-85.
- Baethmann A, Kempfski OS (1997): Pathophysiologie des Hirnödems. *Zentralbl Neurochir* 58:20-29.
- Baethmann A, Maier-Hauff K, Schürer L (1989): Release of glutamate and free fatty acids in vasogenic brain edema. *J Neurosurg* 70:578-91.
- Baethmann A, Maier-Hauff K, Kempfski O, Unterberg A, Wahl M, Schürer L (1988): Mediators of brain edema and secondary brain damage. *Crit Care Med* 16:972-978.
- Baldassarre,S. and Vincent,J.L. (1997): Coagulopathy induced by hydroxyethyl starch. *Anesth.Analg.*, 84:451-453.
- Baskaya,M.K., Dogan,A., Temiz,C., and Dempsey,R.J. (2000): Application of 2,3,5-triphenyltetrazolium chloride staining to evaluate injury volume after controlled cortical impact brain injury: role of brain edema in evolution of injury volume. *J.Neurotrauma*, 17:93-99.
- Baskaya,M.K., Rao,A.M., Dogan,A., Donaldson,D., and Dempsey,R.J. (1997): The biphasic opening of the blood-brain barrier in the cortex and hippocampus after traumatic brain injury in rats. *Neurosci.Lett.* , 226:33-36.
- Bederson JB, Pitts LH, Germano SM, Nishimura MC, Davis RL, Bartkowski HM (1986): Evaluation of 2,3,5-triphenyltetrazolium chloride as a detection and quantification of experimental cerebral infarction on rats. *Stroke* 17, 1304-1308
- Bergamini,S., Rota,C., Canali,R., Staffieri,M., Daneri,F., Bini,A., Giovannini,F., Tomasi,A., and Iannone,A. (2001): N-acetylcysteine inhibits in vivo nitric oxide production by inducible nitric oxide synthase. *Nitric.Oxide.*, 5:349-360.
- Bickler,P.E., Warner,D.S., Stratmann,G., and Schuyler,J.A. (2003): gamma-Aminobutyric acid-A receptors contribute to isoflurane neuroprotection in organotypic hippocampal cultures. *Anesth.Analg.*, 97:564-71, table.

- Bickler,P.E., Buck,L.T., and Hansen,B.M. (1994): Effects of isoflurane and hypothermia on glutamate receptor-mediated calcium influx in brain slices. *Anesthesiology*, 81:1461-1469.
- Biewener A, Holch M, Müller U, Veitinger A, Erfurt C, Zwipp H (2000): Einfluss von logistischem und medizinischem Rettungsaufwand auf die Letalität nach schwerem Trauma. *Unfallchirurg* 103: 137-143
- Bouillon B, Raum M, Fach H, et al. (1999): The incidence and outcome of severe trauma – Design and first results of an epidemiological study in an urban area. *Restorative Neurology and Neuroscience*, 14: 85-92
- Brown et al., Nitric oxide regulates mitochondrial respiration and cell functions by inhibiting cytochrome oxidase, *FEBS Lett.* 369 (1995) 136-139
- Bryan,R.M., Jr., Cherian,L., and Robertson,C. (1995): Regional cerebral blood flow after controlled cortical impact injury in rats. *Anesth.Analg.*, 80:687-695.
- Buga,G.M., Gold,M.E., Fukuto,J.M., and Ignarro,L.J. (1991): Shear stress-induced release of nitric oxide from endothelial cells grown on beads. *Hypertension*, 17:187-193.
- Bullock R (1997): Experimental drug therapies for head injury in: Narayan RK, Wilberger JE, Povlishock JT, (eds.), *Neurotrauma*, New York: McGraw-Hill, 375-391.
- Bullock R, Chesnut RM, Clifton G (1996): Guidelines for the management of severe head injury. *J Neurotrauma* 13:639-734.
- Bullock R, Katake A, Stewart L (1994): The glutamate antagonist CGS 19755 lowers ICP and improves CPP in severely head injured humans. in: Nagai H, Kamiya K, Ishiis I, (eds.), *Intracranial Pressure IX*, Tokyo: Springer Verlag, 238-241.
- Bullock R, Fujisawa H (1992): The role of glutamate antagonist for the treatment of CNS injury. *J Neurotrauma* 9:443-461.
- Bullock,R., Maxwell,W.L., Graham,D.I., Teasdale,G.M., and Adams,J.H. (1991): Glial swelling following human cerebral contusion: an ultrastructural study. *J.Neurol.Neurosurg.Psychiatry*, 54:427-434.
- Bullock R, Statham P, Patterson J, Wyper D, Hadley D, Teasedale E (1990): The time course of vasogenic brain edema after focal human head injury - evidence from SPECT mapping of blood brain barrier defects. *Acta Neurochir Suppl* 51:286-288.
- Bullock R, Mendelow AD, Teasdale GM, Graham DI (1984): Intracranial haemorrhage induced at arterial pressure in the rat. Part 1: Description of technique, ICP changes and neuropathological findings. *Neurol Res* 6:184-8.
- Chen,S., Pickard,J.D., and Harris,N.G. (2003): Time course of cellular pathology after controlled cortical impact injury. *Exp.Neurol.*, 182:87-102.

- Cherian L, Robertson CS, Goodman JC (1996): Secondary insults increase Injury after Controlled Cortical Impact in rats. *J Neurotrauma* 13:371-383.
- Cherian L, Robertson CS, Contant CF, Jr., Bryan RM, Jr. (1994): Lateral cortical impact injury in rats: cerebrovascular effects of varying depth of cortical deformation and impact velocity. *J Neurotrauma* 11:573-85.
- Chesnut RM, Marshall LF, Klauber MR, et al. (1993): The role of secondary brain injury in determining outcome from severe head injury. *J Trauma* 34(2): 216-222
- Chesnut RM, Marshall SB, Piek J, et al. (1993): Early and late systemic hypotension as a frequent and fundamental source of cerebral ischemia following severe brain injury in the Traumatic Coma Data Bank. *Acta Neurochir Suppl* 59:121-125
- Christen,S., Schaper,M., Lykkesfeldt,J., Siegenthaler,C., Bifrare,Y.D., Banic,S., Leib,S.L., and Tauber,M.G. (2001): Oxidative stress in brain during experimental bacterial meningitis: differential effects of alpha-phenyl-tert-butyl nitron and N-acetylcysteine treatment. *Free Radic.Biol.Med.*, 31:754-762.
- Corcoran,G.B. and Wong,B.K. (1986): Role of glutathione in prevention of acetaminophen-induced hepatotoxicity by N-acetyl-L-cysteine in vivo: studies with N-acetyl-D-cysteine in mice. *J.Pharmacol.Exp.Ther.*, 238:54-61.
- Crow,J.P. and Beckman,J.S. (1995): The role of peroxynitrite in nitric oxide-mediated toxicity. *Curr.Top.Microbiol.Immunol.* , 196:57-73.
- Cserr HF, Patlak CS (1991): Regulation of brain volume under isosmotic and anisosmotic conditions. *Adv Comp Environ Physiol* 9:61-80.
- Cserr,H.F., DePasquale,M., Nicholson,C., Patlak,C.S., Pettigrew,K.D., and Rice,M.E. (1991): Extracellular volume decreases while cell volume is maintained by ion uptake in rat brain during acute hypernatremia. *J.Physiol*, 442:277-295.
- Cuzzocrea,S., Mazzon,E., Costantino,G., Serraino,I., De Sarro,A., and Caputi,A.P. (2000): Effects of n-acetylcysteine in a rat model of ischemia and reperfusion injury. *Cardiovasc.Res.*, 47:537-548.
- Cuzzocrea,S., Mazzon,E., Costantino,G., Serraino,I., Dugo,L., Calabro,G., Cucinotta,G., De Sarro,A., and Caputi,A.P. (2000): Beneficial effects of n-acetylcysteine on ischaemic brain injury. *Br.J.Pharmacol.*, 130:1219-1226.
- Dempsey,R.J., Baskaya,M.K., and Dogan,A. (2000): Attenuation of brain edema, blood-brain barrier breakdown, and injury volume by ifenprodil, a polyamine-site N-methyl-D-aspartate receptor antagonist, after experimental traumatic brain injury in rats. *Neurosurgery*, 47:399-404.
- Dewitt DS, Hayes RL, Lyeth BG (1988): Effects of traumatic brain injury on cerebral blood flow an metabolism: Autordiographic studies. *Anesth Rev* 15:31-32.

- Dieterich HJ, Reutershan J, Felbinger TW, Eltzhig HK (2003). Penetration of Intravenous Hydroxyethyl Starch into the Cerebrospinal Fluid in Patients with Impaired Blood-Brain Barrier Function. *Anesth Analg* 96: 1150-1154
- Dieterich,H.J. (2003): Recent developments in European colloid solutions. *J.Trauma*, 54:S26-S30
- Dixon CE, Hayes RL (1997): Fluid percussion and cortical impact models of traumatic brain injury. in: Narayan RK, Wilberger JE, Povlishock JT, (eds.), *Neurotrauma*, New York: McGraw Hill, 1337-1346.
- Dixon CE, Clifton GL, Lighthall JW, Yaghmai AA, Hayes RL (1991): A controlled cortical impact model of traumatic brain injury in the rat. *J Neurosci Methods* 39:253-262.
- Dixon CE, Lyeth BG, Povlishock JT, et al. (1987): A fluid percussion model of experimental brain injury in the rat. *J Neurosurg* 67:110-119.
- Duvdevani R, Roof RL, Fulop Z, Hoffman SW, Stein DG (1995): Blood-brain barrier breakdown and edema formation following frontal cortical contusion: does hormonal status play a role? *J Neurotrauma* 12:65-75.
- Ellis,E.F., Dodson,L.Y., and Police,R.J. (1991): Restoration of cerebrovascular responsiveness to hyperventilation by the oxygen radical scavenger n-acetylcysteine following experimental traumatic brain injury. *J.Neurosurg.*, 75:774-779.
- Farr,S.A., Poon,H.F., Dogrukol-Ak,D., Drake,J., Banks,W.A., Eyerman,E., Butterfield,D.A., and Morley,J.E. (2003): The antioxidants alpha-lipoic acid and N-acetylcysteine reverse memory impairment and brain oxidative stress in aged SAMP8 mice. *J.Neurochem.*, 84:1173-1183.
- Ferber H: Beziehungen von Elimination, Metabolisierung und Wirkung von Hydroxyethylstärke in Abhängigkeit vom Substitutionsgrad un dem Massenmittel (M_w). Habilitationsschrift. Johann-Wolfgang-Goethe-Universität Frankfurt a.M., 1985
- Ferber,H.P., Nitsch,E., and Forster,H. (1985): Studies on hydroxyethyl starch. Part II: Changes of the molecular weight distribution for hydroxyethyl starch types 450/0.7, 450/0.5, 450/0.3, 300/0.4, 200/0.7, 200/0.5, 200/0.3 and 200/0.1 after infusion in serum and urine of volunteers. *Arzneimittelforschung.*, 35:615-622.
- Foda MA, Marmarou A (1994): A new model of diffuse brain injury in rats. Part II: Morphological characterization. *J Neurosurg* 80:301-313.
- Forbes ML, Hendrich KS, Kochanek PM, et al. (1997): Assesment of cerebral blood flow and CO₂ reactivity after controlled cortical impact by perfusion magnetic resonance imaging using arterial spin-labeling in rats. *J Cereb Blood Flow Metab* 17:865-874.

- Forbes, M.L., Hendrich, K.S., Schiding, J.K., Williams, D.S., Ho, C., DeKosky, S.T., Marion, D.W., and Kochanek, P.M. (1997): Perfusion MRI assessment of cerebral blood flow and CO₂ reactivity after controlled cortical impact in rats. *Adv. Exp. Med. Biol.*, 411:7-12.
- Gahm, C., Holmin, S., and Mathiesen, T. (2000): Temporal profiles and cellular sources of three nitric oxide synthase isoforms in the brain after experimental contusion. *Neurosurgery*, 46:169-177.
- Galley, H.F., Howdle, P.D., Walker, B.E., and Webster, N.R. (1997): The effects of intravenous antioxidants in patients with septic shock. *Free Radic. Biol. Med.*, 23:768-774.
- Gazitúa S, Scott JB, Swindall B et al (1971) Resistance responses to local changes in plasma osmolality in three vascular beds *Am J Physiol* 220: 384-391
- Gilgun-Sherky Y, Rosenbaum Z, Melamed E, Offen D (2002) Antioxidant therapy in acute central nervous system injury: current state. *Pharmacol Rev* 54:271-284
- Groeneveld, A.B., den Hollander, W., Straub, J., Nauta, J.J., and Thijs, L.G. (1990): Effects of N-acetylcysteine and terbutaline treatment on hemodynamics and regional albumin extravasation in porcine septic shock. *Circ. Shock*, 30:185-205.
- Gunnar, W., Jonasson, O., Merlotti, G., Stone, J., and Barrett, J. (1988): Head injury and hemorrhagic shock: studies of the blood brain barrier and intracranial pressure after resuscitation with normal saline solution, 3% saline solution, and dextran-40. *Surgery*, 103:398-407.
- Hall ED (1997): Injury mechanisms and therapies: Free radicals and lipid peroxidation. in: Narayan RK, Wilberger JE, Povlishock JT, (eds.), *Neurotrauma*, New York: McGraw Hill, 1405-1419.
- Hall ED, Braugher JM (1991): Free radicals in CNS injury. in: Waxman SG, (eds.), *Molecular and Cellular Approaches to the Treatment of Brain Diseases*, New York: Raven Press, 81-105.
- Hammerle AF, Weinstabl C, Mayer N, Germann P, Steltzer H (1992). Decrease of intracranial pressure following a combination of hypertonic saline and hydroxyethyl starch. In: Baron JF (ed) Plasma volume expansion. Arnette Blackwell, Paris, S 231-233
- Harada, H., Kelly, P.J., Cole, D.J., Drummond, J.C., and Patel, P.M. (1999): Isoflurane reduces N-methyl-D-aspartate toxicity in vivo in the rat cerebral cortex. *Anesth. Analg.*, 89:1442-1447.
- Harrison, P.M., Wendon, J.A., Gimson, A.E., Alexander, G.J., and Williams, R. (1991): Improvement by acetylcysteine of hemodynamics and oxygen transport in fulminant hepatic failure. *N. Engl. J. Med.*, 324:1852-1857.

- Hendrich,K.S., Kochanek,P.M., Melick,J.A., Schiding,J.K., Statler,K.D., Williams,D.S., Marion,D.W., and Ho,C. (2001): Cerebral perfusion during anesthesia with fentanyl, isoflurane, or pentobarbital in normal rats studied by arterial spin-labeled MRI. *Magn Reson.Med.*, 46:202-206.
- Henry,P.J., Horowitz,J.D., and Louis,W.J. (1989): Determinants of in vitro nitroglycerin tolerance induction and reversal: influence of dose regimen, nitrate-free period, and sulfhydryl supplementation. *J.Cardiovasc.Pharmacol.*, 14:31-37.
- Horber,F.F., Krayer,S., Rehder,K., and Haymond,M.W. (1988): Anesthesia with halothane and nitrous oxide alters protein and amino acid metabolism in dogs. *Anesthesiology*, 69:319-326.
- Hrapkiewicz K, Medina L, Holmes DD, Clinical Laboratory Animal Medicine: An Introduction (1998), 2nd edition, Iowa State University, Ames, IA (USA) ISBN 0-8138-2555-5, pp 31-56
- Iadecola,C., Zhang,F., Casey,R., Clark,H.B., and Ross,M.E. (1996): Inducible nitric oxide synthase gene expression in vascular cells after transient focal cerebral ischemia. *Stroke*, 27:1373-1380.
- Jennett,B. (1998): Epidemiology of head injury. *Arch.Dis.Child*, 78:403-406.
- Jennett,B. (1996): Epidemiology of head injury. *J.Neurol.Neurosurg.Psychiatry*, 60:362-369.
- Jennett,B. and MacMillan,R. (1981): Epidemiology of head injury. *Br.Med.J.(Clin.Res.Ed)*, 282:101-104.
- Jevtovic-Todorovic,V., Beals,J., Benshoff,N., and Olney,J.W. (2003): Prolonged exposure to inhalational anesthetic nitrous oxide kills neurons in adult rat brain. *Neuroscience*, 122:609-616.
- Jevtovic-Todorovic,V., Todorovic,S.M., Mennerick,S., Powell,S., Dikranian,K., Benshoff,N., Zorumski,C.F., and Olney,J.W. (1998): Nitrous oxide (laughing gas) is an NMDA antagonist, neuroprotectant and neurotoxin. *Nat.Med.*, 4:460-463.
- Kamada K, Houkin K, Hida K, Iwasaki Y, Abe H (1995): Serial changes in metabolism and histology in the cold-injury trauma rat brain model--proton magnetic resonance imaging and spectroscopy study. *Neurol Med Chir Tokyo* 35:1-7.
- Kawamata T, Katayama Y, Maeda T, et al. (1997): Antioxidant, OPC 14117, attenuates edema formation and behavioral deficits following Cortical Contusion in rats. *Acta Neurochir Suppl* 70:191-193.
- Kellie G (1824): The kind of appearances observed in the dissection of two or three individuals ... with some reference to the pathology of the brain. *Trans Med Chi Edin* 1:84-169.

- Kempinski, O., Obert, C., Mainka, T., Heimann, A., and Strecker, U. (1996): "Small volume resuscitation" as treatment of cerebral blood flow disturbances and increased ICP in trauma and ischemia. *Acta Neurochir. Suppl (Wien.)*, 66:114-117.
- Kempinski OS, Volk C (1994): Neuron-glia interaction during injury and edema of the CNS. *Acta Neurochir. Suppl (Wien.)*, 60:7-11.
- Khan, M., Sekhon, B., Jatana, M., Giri, S., Gilg, A.G., Sekhon, C., Singh, I., and Singh, A.K. (2004): Administration of N-acetylcysteine after focal cerebral ischemia protects brain and reduces inflammation in a rat model of experimental stroke. *J. Neurosci. Res.*, 76:519-527.
- Kien ND, Kramer GC, White DA (1991): Acute hypotension caused by rapid hypertonic saline infusion in anaesthetized dogs, *Anesth Analg* 73: 597-602
- Kimelberg HK (1995): Current concepts of brain edema. Review of laboratory investigations. *J Neurosurg* 83:1051-1059.
- Kimelberg HK, Ransom BR (1986): Physiological and pathological aspects of astrocytic swelling. in: Fedoroff S, Vernadakis A, (eds.), *Astrocytes*, Orlando, Fla: Academic Press, 129-166.
- Kita H, Marmarou A (1994): The cause of acute brain swelling after the closed head injury in rats. *Acta Neurochir Suppl (Wien.)* 60:452-455.
- Klatzo I (1994): Evolution of brain edema concepts. *Acta Neurochir Suppl (Wien.)*, 60:3-6.
- Klatzo I, Wisniewski HM, Steinwall O, Streicher E (1967): Dynamics of cold injury edema. in: Klatzo I, Seitelberger F, (eds.), *Brain Edema*, New York: Springer, 554-563.
- Klatzo I (1967): Presidential address. Neuropathological aspects of brain edema. *J Neuropath Exp Neurol* 26:1-14.
- Klatzo I, Piraux A, Laskowski EJ (1958): The relationship between edema, blood-brain-barrier, and tissue elements in a local brain injury. *J Neuropathol Exp Neurol* 17:548-564.
- Knuckey, N.W., Palm, D., Primiano, M., Epstein, M.H., and Johanson, C.E. (1995): N-acetylcysteine enhances hippocampal neuronal survival after transient forebrain ischemia in rats. *Stroke*, 26:305-310.
- Kochanek, P.M., Marion, D.W., Zhang, W., Schiding, J.K., White, M., Palmer, A.M., Clark, R.S., O'Malley, M.E., Styren, S.D., Ho, C., and . (1995): Severe controlled cortical impact in rats: assessment of cerebral edema, blood flow, and contusion volume. *J. Neurotrauma*, 12:1015-1025.
- Kontos, H.A. and Povlishock, J.T. (1986): Oxygen radicals in brain injury. *Cent. Nerv. Syst. Trauma*, 3:257-263.

- Kreimeier U, Thiel M, Messmer K (1996) Hypertonic-hyperoncotic solutions In: Risberg B (Hrsg) Trauma Care – an update. Pharmacia & Upjohn Sverige Ab, Stockholm, S 142-153
- Kreimeier,U. and Messmer,K. (1996): [Prehospital fluid resuscitation]. *Anaesthetist*, 45:884-899.
- Kreimeier U, Messmer K (1988) Small-volume resuscitation. In: Kox WJ, Gamble J (eds) Fluid resuscitation. Baillière 's Clinical Anaesthesiology, vol 2. Baillière Tindall. London, S 545-577
- Kroppenstedt,S.N., Kern,M., Thomale,U.W., Schneider,G.H., Lanksch,W.R., and Unterberg,A.W. (1999): Effect of cerebral perfusion pressure on contusion volume following impact injury. *J.Neurosurg.*, 90:520-526.
- Kroppenstedt,S.N., Stroop,R., Kern,M., Thomale,U.W., Schneider,G.H., and Unterberg,A.W. (1999): Lubeluzole following traumatic brain injury in the rat. *J.Neurotrauma*, 16:629-637.
- Kroppenstedt,S.N., Schneider,G.H., Thomale,U.W., and Unterberg,A.W. (1998): Neuroprotective properties of aptiganel HCL (Cerestat) following controlled cortical impact injury. *Acta Neurochir.Suppl*, 71:114-116.
- Kroppenstedt,S.N., Schneider,G.H., Thomale,U.W., and Unterberg,A.W. (1998): Protective effects of aptiganel HCl (Cerestat) following controlled cortical impact injury in the rat. *J.Neurotrauma*, 15:191-197.
- Lenz,C., Frietsch,T., Futterer,C., Rebel,A., van Ackern,K., Kuschinsky,W., and Waschke,K.F. (1999): Local coupling of cerebral blood flow to cerebral glucose metabolism during inhalational anesthesia in rats: desflurane versus isoflurane. *Anesthesiology*, 91:1720-1723.
- Lenzlinger,P.M., Morganti-Kossmann,M.C., Laurer,H.L., and McIntosh,T.K. (2001): The duality of the inflammatory response to traumatic brain injury. *Mol.Neurobiol.*, 24:169-181.
- Lighthall,J.W., Dixon,C.E., and Anderson,T.E. (1989): Experimental models of brain injury. *J.Neurotrauma*, 6:83-97.
- Lighthall JW (1988): Controlled cortical impact: a new experimental brain injury model. *J Neurotrauma* 5:1-15.
- Maas,A.I., Dearden,M., Teasdale,G.M., Braakman,R., Cohadon,F., Iannotti,F., Karimi,A., Lapierre,F., Murray,G., Ohman,J., Persson,L., Servadei,F., Stocchetti,N., and Unterberg,A. (1997): EBIC-guidelines for management of severe head injury in adults. European Brain Injury Consortium. *Acta Neurochir.(Wien.)*, 139:286-294.
- Marmarou A, Foda MA, van den Brink W, Campbell J, Kita H, Demetriadou K (1994): A new model of diffuse brain injury in rats. Part I: Pathophysiology and biomechanics. *J Neurosurg* 80:291-300.

- Marmarou,A. (1994): Traumatic brain edema: an overview. *Acta Neurochir.Suppl (Wien.)*, 60:421-424.
- Marshall,L.F. (2000): Head injury: recent past, present, and future. *Neurosurgery*, 47:546-561.
- Masson,F., Thicoipe,M., Mokni,T., Aye,P., Erny,P., and Dabadie,P. (2003): Epidemiology of traumatic comas: a prospective population-based study. *Brain Inj.*, 17:279-293.
- Masson,F., Thicoipe,M., Aye,P., Mokni,T., Senjean,P., Schmitt,V., Dessalles,P.H., Cazaugade,M., and Labadens,P. (2001): Epidemiology of severe brain injuries: a prospective population-based study. *J.Trauma*, 51:481-489.
- Mazzoni,M.C., Borgstrom,P., Arfors,K.E., and Intaglietta,M. (1988): Dynamic fluid redistribution in hyperosmotic resuscitation of hypovolemic hemorrhage. *Am.J.Physiol*, 255:H629-H637
- McIntosh TK, Vink R, Noble L, et al. (1989): Traumatic brain injury in the rat: characterization of a lateral fluid-percussion model. *Neuroscience* 28:233-244.
- Messmer,K. and Kreimeier,U. (1989): Microcirculatory therapy in shock. *Resuscitation*, 18 Suppl:S51-S61
- Mittlmeier,T., Vollmar,B., Menger,M.D., Schewior,L., Raschke,M., and Schaser,K.D. (2003): Small volume hypertonic hydroxyethyl starch reduces acute microvascular dysfunction after closed soft-tissue trauma. *J.Bone Joint Surg.Br.*, 85:126-132.
- Monroe A (1783): Observation on the structure and function of the nervous system. in: *Creech and Johnson*, Edinburgh, 5-6.
- Murray,G.D., Teasdale,G.M., Braakman,R., Cohadon,F., Dearden,M., Iannotti,F., Karimi,A., Lapiere,F., Maas,A., Ohman,J., Persson,L., Servadei,F., Stocchetti,N., Trojanowski,T., and Unterberg,A. (1999): The European Brain Injury Consortium survey of head injuries. *Acta Neurochir.(Wien.)*, 141:223-236.
- Narayan,R.K., Michel,M.E., Ansell,B., Baethmann,A., Biegon,A., Bracken,M.B., Bullock,M.R., Choi,S.C., Clifton,G.L., Contant,C.F., Coplin,W.M., Dietrich,W.D., Ghajar,J., Grady,S.M., Grossman,R.G., Hall,E.D., Heetderks,W., Hovda,D.A., Jallo,J., Katz,R.L., Knoller,N., Kochanek,P.M., Maas,A.I., Majde,J., Marion,D.W., Marmarou,A., Marshall,L.F., McIntosh,T.K., Miller,E., Mohberg,N., Muizelaar,J.P., Pitts,L.H., Quinn,P., Riesenfeld,G., Robertson,C.S., Strauss,K.I., Teasdale,G., Temkin,N., Tuma,R., Wade,C., Walker,M.D., Weinrich,M., Whyte,J., Wilberger,J., Young,A.B., and Yurkewicz,L. (2002): Clinical trials in head injury. *J.Neurotrauma*, 19:503-557.
- Nolte,D., Bayer,M., Lehr,H.A., Becker,M., Krombach,F., Kreimeier,U., and Messmer,K. (1992): Attenuation of postischemic microvascular disturbances in striated muscle by hyperosmolar saline dextran. *Am.J.Physiol*, 263:H1411-H1416

- Oberg,P.A. (1990): Laser-Doppler flowmetry. *Crit Rev.Biomed.Eng*, 18:125-163.
- Pappius HM (1974): Fundamental aspects of brain edema. in: Vinken PJ, Bruyn GW, (eds.), *Handbook Clin Neurol*; American Elsevier, 167-185.
- Perri,B.R., Smith,D.H., Murai,H., Sinson,G., Saatman,K.E., Raghupathi,R., Bartus,R.T., and McIntosh,T.K. (1997): Metabolic quantification of lesion volume following experimental traumatic brain injury in the rat. *J.Neurotrauma*, 14:15-22.
- Petroianu,G.A., Maleck,W.H., Koetter,K.P., Liu,J., and Schmitt,A. (2003): Effect of in vitro hemodilution with hydroxyethyl starch and dextran on the activity of plasma clotting factors. *Crit Care Med.*, 31:250-254.
- Piek,J. and Jantzen,J.P. (2000): [Recommendations for early care of patients with skull and brain trauma in multiple injuries (prepared by the German Interdisciplinary Union for Intensive and Trauma Care (DIVI) on November 5, 1999]. *Zentralbl.Neurochir.*, 61:50-56.
- Qureshi,A.I., Suarez,J.I., Bhardwaj,A., Mirski,M., Schnitzer,M.S., Hanley,D.F., and Ulatowski,J.A. (1998): Use of hypertonic (3%) saline/acetate infusion in the treatment of cerebral edema: Effect on intracranial pressure and lateral displacement of the brain. *Crit Care Med.*, 26:440-446.
- Rapoport SI (1979): Roles of cerebrovascular permeability, brain compliance, and hydraulic conductivity in vasogenic brain edema. in: Popp AJ, Bourke RS, Nelson LR, al. e, (eds.), *Neural Trauma*, New York: Raven Press, 51-62.
- Sarrafzadeh,A.S., Thomale,U.W., Kroppenstedt,S.N., and Unterberg,A.W. (2000): Neuroprotective effect of melatonin on cortical impact injury in the rat. *Acta Neurochir.(Wien.)*, 142:1293-1299.
- Sarrafzadeh A, Thomale U-W, Kroppenstedt S-N, Unterberg A (1997): The role of melatonin following Controlled Cortical Impact Injury. *Zentralbl Neurochir Suppl* P37:56.
- Schiller,H.J., Reilly,P.M., and Bulkley,G.B. (1993): Tissue perfusion in critical illnesses. Antioxidant therapy. *Crit Care Med.*, 21:S92-102.
- Schneider GH, Hennig S, Lanksch WR, Unterberg A (1994): Dynamics of posttraumatic brain swelling following a cryogenic injury in rats. *Acta Neurochir Suppl* 60:437-439.
- Shackford,S.R., Bourguignon,P.R., Wald,S.L., Rogers,F.B., Osler,T.M., and Clark,D.E. (1998): Hypertonic saline resuscitation of patients with head injury: a prospective, randomized clinical trial. *J.Trauma*, 44:50-58.
- Shima K, Marmarou A (1991): Evaluation of brain stem dysfunction following severe fluid percussion head injury to the cat. *J Neurosurg* 72:270-277.

- Skarphedinsson JO, Hårding H, Thorén P: Repeated measurements of cerebral blood flow in rats. Comparisons between the hydrogen clearance method and laser Doppler flowmetry. *Acta Physiol Scand* 134: 133-142, 1988
- Smith,A.L. and Marque,J.J. (1976): Anesthetics and cerebral edema. *Anesthesiology*, 45:64-72.
- Smith ML, Auer RN, Siesjo BK (1984): The density and distribution of ischemic brain injury in the rat following 2-10 min of forebrain ischemia. *Acta Neuropathol* 64:319-332.
- Smith SL (1996): Mild pre- and posttraumatic hypothermia attenuates blood-brain barrier damage following Controlled Cortical Impact Injury in the rat. *J Neurotrauma* 13:1-9.
- Smith SL, Andrus PK, Zhang JR, Hall ED (1994): Direct measurement of hydroxyl radicals, lipid peroxidation, and blood-brain barrier disruption following unilateral cortical impact head injury in the rat. *J Neurotrauma* 11:393-404.
- Soares HD, Thomas M, Cloherty K, McIntosh TK (1992): Development of prolonged focal cerebral edema and regional cation changes following experimental brain injury in the rat. *J Neurochem* 58:1845-1852.
- Stamler,J., Mendelsohn,M.E., Amarante,P., Smick,D., Andon,N., Davies,P.F., Cooke,J.P., and Loscalzo,J. (1989): N-acetylcysteine potentiates platelet inhibition by endothelium-derived relaxing factor. *Circ.Res.*, 65:789-795.
- Statistisches Bundesamt Deutschland – Gesundheitswesen: Gesundheitszustand und Behandlungsanlässe von Krankenhauspatienten 2003: (accessed August 6, 2005, at <http://www.destatis.de/basis/d/gesu/gesutab11.php>)
- Statler,K.D., Kochanek,P.M., Dixon,C.E., Alexander,H.L., Warner,D.S., Clark,R.S., Wisniewski,S.R., Graham,S.H., Jenkins,L.W., Marion,D.W., and Safar,P.J. (2000): Isoflurane improves long-term neurologic outcome versus fentanyl after traumatic brain injury in rats. *J.Neurotrauma*, 17:1179-1189.
- Staub F, Winkler A, Haberstock J, et al. (1996): Swelling, intracellular acidosis, and damage of glial cells. *Acta Neurochir Suppl* 66:56-62.
- Staub F, Winkler A, Peters J, Kempfski OS, Kachel V, Baethmann A (1994): Swelling, acidosis and irreversible damage of glial cells from exposure to arachidonic acid in vitro. *J Cereb Blood Flow Metab* 14:1030-1039.
- Stoffel,M., Rinecker,M., Plesnila,N., Eriskat,J., and Baethmann,A. (2001): Role of nitric oxide in the secondary expansion of a cortical brain lesion from cold injury. *J.Neurotrauma*, 18:425-434.
- Stogermuller,B., Stark,J., Willschke,H., Felfernig,M., Hoerauf,K., and Kozek-Langenecker,S.A. (2000): The effect of hydroxyethyl starch 200 kD on platelet function. *Anesth.Analg.*, 91:823-827.

- Stover, J.F., Sakowitz, O.W., Kroppenstedt, S.N., Thomale, U.W., Kempfski, O.S., Flugge, G., and Unterberg, A.W. (2004): Differential effects of prolonged isoflurane anesthesia on plasma, extracellular, and CSF glutamate, neuronal activity, 125I-Mk801 NMDA receptor binding, and brain edema in traumatic brain-injured rats. *Acta Neurochir.(Wien.)*, 146:819-830.
- Stover, J.F. and Unterberg, A.W. (2000): Increased cerebrospinal fluid glutamate and taurine concentrations are associated with traumatic brain edema formation in rats. *Brain Res.*, 875:51-55.
- Stover, J.F., Kroppenstedt, S.N., Thomale, U.W., Kempfski, O.S., and Unterberg, A.W. (2000): Isoflurane doubles plasma glutamate and increases posttraumatic brain edema. *Acta Neurochir.Suppl*, 76:375-378.
- Stover, J.F., Schoning, B., Beyer, T.F., Woiciechowsky, C., and Unterberg, A.W. (2000): Temporal profile of cerebrospinal fluid glutamate, interleukin-6, and tumor necrosis factor-alpha in relation to brain edema and contusion following controlled cortical impact injury in rats. *Neurosci.Lett.*, 288:25-28.
- Stroop R, Thomale U-W, Pauser S, et al. (1998): Magnetic resonance imaging studies with cluster algorithm for characterization of brain edema after controlled cortical impact injury (CCII). *Acta Neurochir Suppl* 71:303-305.
- Sutton RL, Hovda DA, Adelson PD, Benzel EC, Becker DP (1994): Metabolic changes following cortical contusion: relationships to edema and morphological changes. *Acta Neurochir Suppl* 60:446-448.
- Sutton RL, Lescaudron L, Stein DG (1993): Unilateral cortical contusion injury in the rat: vascular disruption and temporal development of cortical necrosis. *J Neurotrauma* 10:135-149.
- Szabo, C. (1996): Physiological and pathophysiological roles of nitric oxide in the central nervous system. *Brain Res.Bull.*, 41:131-141.
- Szabo, C. (1995): Alterations in nitric oxide production in various forms of circulatory shock. *New Horiz.*, 3:2-32.
- Thomale, U.W., Kroppenstedt, S.N., Beyer, T.F., Schaser, K.D., Unterberg, A.W., and Stover, J.F. (2002): Temporal profile of cortical perfusion and microcirculation after controlled cortical impact injury in rats. *J.Neurotrauma*, 19:403-413.
- Thomale, U.W., Schaser, K., Kroppenstedt, S.N., Unterberg, A.W., and Stover, J.F. (2002): CNS. *Acta Neurochir Suppl* 60:7-11.
- Thomale, U.W., Schaser, K., Kroppenstedt, S.N., Unterberg, A.W., and Stover, J.F. (2002): Cortical hypoperfusion precedes hyperperfusion following controlled cortical impact injury. *Acta Neurochir.Suppl*, 81:229-231.
- Thomale, U.W., Schaser, K.D., Unterberg, A.W., and Stover, J.F. (2001): Visualization of rat pial microcirculation using the novel orthogonal polarized spectral (OPS) imaging after brain injury. *J.Neurosci.Methods*, 108:85-90.

- Thomale U: Evaluation des tierexperimentellen Modells einer traumatischen kortikalen Kontusion (Controlled Cortical Impact Injury) für Therapiestudien. Dissertationsschrift, Humboldt-Universität Berlin, 2001
- Tyurin, V.A., Tyurina, Y.Y., Borisenko, G.G., Sokolova, T.V., Ritov, V.B., Quinn, P.J., Rose, M., Kochanek, P., Graham, S.H., and Kagan, V.E. (2000): Oxidative stress following traumatic brain injury in rats: quantitation of biomarkers and detection of free radical intermediates. *J. Neurochem.*, 75:2178-2189.
- Unterberg AW, Stroop R, Thomale UW, Kiening KL, Päuser S, Vollmann W (1996): Characterization of brain edema following controlled cortical impact injury in rats. *Acta Neurochir Suppl* 70:106-108.
- Unterberg A, Schneider GH, Gottschalk J, Lanksch WR (1994): Development of traumatic brain edema in old versus young rats. *Acta Neurochir Suppl* 60:431-433.
- Unterberg AW, Lanksch WR (1994): Umfrage zur Behandlung von Patienten mit schwerem Schädel-Hirn-Trauma in Neurochirurgischen Kliniken Deutschlands. (pers. Mitteilung).
- Unterberg A, Wahl M, Baethmann A (1984): Effects of bradykinin on permeability and diameter of pial vessels in vivo. *J Cereb Blood Flow Metab* 4:574-585.
- Van den Brink WA, Marmarou A, Avezaat CJJ (1990): Brain edema in experimental closed head injury in the rat. *Acta Neurochir Suppl* 51:261-262.
- Van Landeghem F, Schreiber S, Unterberg AW, von Deimling A, Stover JF (2003): Differential Concentration-Dependent Effects of Prolonged Norepinephrine Infusion on Intraparenchymal Hemorrhage and Cortical Contusion in Brain-Injured Rats. *J Neurotrauma* 20:1327-1337
- Velasco, I.T., Pontieri, V., Rocha e Silva, and Lopes, O.U. (1980): Hyperosmotic NaCl and severe hemorrhagic shock. *Am. J. Physiol.*, 239:H664-H673
- Verweij, B.H., Muizelaar, J.P., Vinas, F.C., Peterson, P.L., Xiong, Y., and Lee, C.P. (2000): Impaired cerebral mitochondrial function after traumatic brain injury in humans. *J. Neurosurg.*, 93:815-820.
- Vorbrodt AW, Lossinsky AS, Wisniewski HM, et al. (1985): Ultrastructural observations on the transvascular route of protein removal in vasogenic brain edema. *Acta Neuropathol* 66:265-273. Cortical hypoperfusion precedes hyperperfusion following controlled cortical impact injury. *Acta Neurochir. Suppl*, 81:229-231.
- Weidhase, R., Faude, K., and Weidhase, R. (1998): [Hydroxyethyl starch--an interim report]. *Anaesthesiol. Reanim.*, 23:4-14.
- Xiong, Y., Peterson, P.L., and Lee, C.P. (1999): Effect of N-acetylcysteine on mitochondrial function following traumatic brain injury in rats. *J. Neurotrauma*, 16:1067-1082.

Xiong,Y., Gu,Q., Peterson,P.L., Muizelaar,J.P., and Lee,C.P. (1997): Mitochondrial dysfunction and calcium perturbation induced by traumatic brain injury. *J.Neurotrauma*, 14:23-34.

Ytrebo,L.M., Korvald,C., Nedredal,G.I., Elvenes,O.P., Nielsen Grymyr,O.J., and Revhaug,A. (2001): N-acetylcysteine increases cerebral perfusion pressure in pigs with fulminant hepatic failure. *Crit Care Med.*, 29:1989-1995.