

# Bibliography

- [1] F. F. Crim. Selective excitation studies of unimolecular reaction dynamics. *Ann. Rev. Phys. Chem.*, **35**, 657 (1984).
- [2] R. J. Gordon and S. A. Rice. Active control of the dynamics of atoms and molecules. *Ann. Rev. Phys. Chem.*, **48**, 601 (1997).
- [3] A. H. Zewail. Laser femtochemistry. *Science*, **242**, 1645 (1988).
- [4] R. N. Zare. Laser Control of Chemical Reactions. *Science*, **279**, 1875 (1998).
- [5] A. Sinha, M. C. Hsiao and F. F. Crim. Bond-selected bimolecular chemistry:  $\text{H} + \text{HOD}(4\nu_{\text{OH}}) \rightarrow \text{OD} + \text{H}_2$ . *J. Chem. Phys.*, **92**, 6333 (1990).
- [6] A. Sinha, M. C. Hsiao and F. F. Crim. Controlling bimolecular reactions: mode and bond selected reaction of water with hydrogen atoms. *J. Chem. Phys.*, **94**, 4928 (1991).
- [7] R. B. Metz, J. D. Thoemke, J. M. Pfeiffer and F. F. Crim. Selectively breaking either bond in the bimolecular reaction of HOD with hydrogen atoms. *J. Chem. Phys.*, **99**, 1744 (1993).
- [8] M. J. Bronikowski, W. R. Simpson, B. Girard and R. N. Zare. Bond-specific chemistry: OD:OH product ratios for the reactions  $\text{H} + \text{HOD}(100)$  and  $\text{H} + \text{HOD}(001)$ . *J. Chem. Phys.*, **95**, 8647 (1991).
- [9] M. J. Bronikowski, W. R. Simpson and R. N. Zare. Effect of reagent vibration on the  $\text{H} + \text{HOD}$  reaction: an example of bond-specific chemistry. *J. Phys. Chem.*, **97**, 2194 (1993).
- [10] P. Brumer and M. Shapiro. Control of unimolecular reactions using coherent light. *Chem. Phys. Lett.*, **126**, 541 (1986).
- [11] A. H. Zewail. Femtochemistry—atomic-scale dynamics of the chemical bond using ultrafast lasers (Nobel paper). *Ang. Chem. Int. Ed.*, **39**, 2586 (2000).

- [12] D. J. Tannor and S. A. Rice. Control of selectivity of chemical reaction via control of wave packet evolution. *J. Chem. Phys.*, **83**, 5013 (1985).
- [13] D. J. Tannor, R. Kosloff and S. A. Rice. Coherent pulse sequence induced control of selectivity of reactions: exact quantum mechanical calculations. *J. Chem. Phys.*, **85**, 5805 (1986).
- [14] E. D. Potter, J. L. Herek, S. Pedersen, Q. Liu and A. H. Zewail. Femtosecond laser control of a chemical reaction. *Nature*, **355**, 66 (1992).
- [15] S. Shi, A. Woody and H. Rabitz. Optimal control of selective vibrational excitations in harmonic linear chain molecules. *J. Chem. Phys.*, **88**, 6870 (1988).
- [16] A. P. Peirce, M. Dahleh and H. Rabitz. Optimal control of quantum mechanical systems: existence, numerical approximations, and applications. *Phys. Rev. A*, **37**, 4950 (1988).
- [17] R. S. Judson and H. Rabitz. Teaching lasers to control molecules. *Phys. Rev. Lett.*, **68**, 1500 (1992).
- [18] W. S. Warren, H. Rabitz and M. Dahleh. Coherent control of quantum dynamics: the dream is alive. *Science*, **259**, 1581 (1993).
- [19] A. Assion, T. Baumert, M. Bergt, T. Brixner, B. Kiefer, V. Seyfried, M. Strehle and G. Gerber. Control of chemical reactions by feedback-optimized phase-shaped femtosecond laser pulses. *Science*, **282**, 919 (1998).
- [20] R. J. Levis, G. M. Menkir and H. Rabitz. Selective bond dissociation and rearrangement with optimally tailored, strong-field laser pulses. *Science*, **292**, 709 (2001).
- [21] J. L. Herek, W. Wohlleben, R. J. Codgell, D. Zeidler and M. Motzkus. Quantum control of energy flow in light harvesting. *Nature*, **417**, 533 (2002).
- [22] C. Daniel, J. Full, L. González, C. Lupulescu, J. Manz, A. Merli, Š. Vajda and L. Wöste. Deciphering the reaction dynamics underlying optimal control laser fields. *Science*, **299**, 536 (2003).
- [23] J. Z. Zhang, D. G. Imre and J. H. Frederick. HOD spectroscopy and photodissociation: selectivity in OH/OD bond breaking. *J. Phys. Chem.*, **93**, 1840 (1989).
- [24] D. G. Imre and J. Zhang. Dynamics and selective bond breaking in photodissociation. *Chem. Phys.*, **139**, 89 (1989).
- [25] F. F. Crim. State- and bond-selected unimolecular reactions. *Science*, **249**, 1387 (1990).

- [26] R. L. Vander Wal, J. L. Scott, F. F. Crim, K. Weide and R. Schinke. An experimental and theoretical study of the bond selected dissociation of HOD. *J. Chem. Phys.*, **94**, 3548 (1991).
- [27] B. Just, J. Manz and G. K. Paramonov. Series of ultrashort infrared laser pulses with analytical shapes for selective vibrational excitation: model simulations for  $\text{OH}(\nu=0) \rightarrow \text{OH}(\nu \leq 10)$ . *Chem. Phys. Lett.*, **193**, 429 (1992).
- [28] W. Jakubetz, E. Kades and J. Manz. State-selective excitation of molecules by means of optimized infrared laser pulses. *J. Phys. Chem.*, **97**, 12609 (1993).
- [29] M. V. Korolkov, J. Manz and G. K. Paramonov. Theory of laser control of vibrational transitions and chemical reactions by ultrashort infrared laser pulses. *Adv. Chem. Phys.*, **101**, 327 (1997).
- [30] S. Chelkowski, A. D. Bandrauk and P. B. Corkum. Efficient molecular dissociation by a chirped ultrashort infrared laser pulse. *Phys. Rev. Lett.*, **65**, 2355 (1990).
- [31] B. Amstrup and N. E. Henriksen. Control of HOD photodissociation dynamics via bond-selective infrared multiphoton excitation and a femtosecond ultraviolet laser pulse. *J. Chem. Phys.*, **97**, 8285 (1992).
- [32] B. Amstrup and N. E. Henriksen. Two-pulse laser control of bond-selective fragmentation. *J. Chem. Phys.*, **105**, 9115 (1996).
- [33] A. Sinha, M. C. Hsiao and F. F. Crim. IR + UV laser pulse control of momenta directed to specific products: quantum simulations for  $\text{HOD}^* \rightarrow \text{H} + \text{OD}$  versus  $\text{HO} + \text{D}$ . *Phys. Chem. Chem. Phys.*, **5**, 4806 (2003).
- [34] E. Charron, A. Giusti-Suzor, and F. H. Mies. Fragment angular distribution in one- and two-color photodissociation by strong laser fields. *Phys. Rev. A*, **49**, R641 (1994).
- [35] E. Charron, A. Giusti-Suzor, and F. H. Mies. Coherent control of isotope separation in  $\text{HD}^+$  photodissociation by strong fields. *Phys. Rev. Lett.*, **75**, 2815 (1995).
- [36] M. Machholm and N. E. Henriksen. Off-resonant vibrational excitation: orientational dependence and spatial control of photofragments. *J. Chem. Phys.*, **113**, 7838 (2000).
- [37] M. Machholm and N. E. Henriksen. Two-pulse laser control for selective photofragment orientation. *J. Chem. Phys.*, **111**, 3051 (1999).
- [38] B. Friedrich and D. Herschbach. Spatial orientation of molecules in strong electric fields and evidence for pendular states. *Nature*, **353**, 412 (1991).

- [39] B. Friedrich and D. Herschbach. Alignment and trapping of molecules in intense laser fields. *Phys. Rev. Lett.*, **74**, 4623 (1995).
- [40] H. J. Loesch and A. Remscheid. Reactive scattering of K from oriented and isotropic CH<sub>3</sub>I molecules. *J. Phys. Chem.*, **95**, 8194 (1990).
- [41] H. J. Loesch and A. Remscheid. Brute force in molecular reaction dynamics: a novel technique for measuring steric effects. *J. Chem. Phys.*, **93**, 4779 (1990).
- [42] B. Friedrich and D. Herschbach. On the possibility of orienting rotationally cooled polar molecules in an electric field. *Z. Phys. D*, **18**, 153 (1991).
- [43] M. J. J. Vrakking and S. Stolte. Coherent control of molecular orientation. *Chem. Phys. Lett.*, **271**, 209 (1997).
- [44] K. H. Kramer and R. B. Bernstein. Focusing and orientation of symmetric-top molecules with the electric six-pole field. *J. Chem. Phys.*, **42**, 767 (1965).
- [45] T. D. Hain, R. M. Moision and T. J. Curtiss. Hexapole state-selection and orientation of asymmetric top molecules: CH<sub>2</sub>F<sub>2</sub>. *J. Chem. Phys.*, **111**, 6797 (1999).
- [46] B. Friedrich and D. Herschbach. Enhanced orientation of polar molecules by combined electrostatic and nonresonant induced dipole forces. *J. Chem. Phys.*, **111**, 6157 (1999).
- [47] H. Sakai, S. Minemoto, H. Nanjo, H. Tanji and T. Suzuki. Controlling the orientation of polar molecules with combined electrostatic and pulsed, nonresonant laser fields. *Phys. Rev. Lett.*, **90**, 083001 (2003).
- [48] N. E. Henriksen. Molecular alignment and orientation in short pulse laser fields. *Chem. Phys. Lett.*, **312**, 196 (1999).
- [49] C. M. Dion, A. D. Bandrauk, O. Atabek, A. Keller, H. Umeda and Y. Fujimura. Two-frequency IR laser orientation of polar molecules. Numerical simulations for HCN. *Chem. Phys. Lett.*, **302**, 215 (1999).
- [50] T. Seideman. Rotational excitation and molecular alignment in intense laser fields. *J. Chem. Phys.*, **103**, 7887 (1995).
- [51] T. Seideman. New means of spatially manipulating molecules with light. *J. Chem. Phys.*, **111**, 4397 (1999).
- [52] T. Seideman. Revival structure of aligned rotational wave packets. *Phys. Rev. Lett.*, **83**, 4971 (1999).

- [53] T. Seideman. On the dynamics of rotationally broad, spatially aligned wave packets. *J. Chem. Phys.*, **115**, 5965 (2001).
- [54] J. J. Larsen, H. Sakai, C. P. Safvan, I. W.-Larsen and H. Stapelfeldt. Aligning molecules with intense nonresonant laser fields. *J. Chem. Phys.*, **111**, 7774 (1999).
- [55] J. J. Larsen, K. Hald, N. Bjerre, H. Stapelfeldt and T. Seideman. Three dimensional alignment of molecules using elliptically polarized laser fields. *Phys. Rev. Lett.*, **85**, 2470 (2000).
- [56] E. Péronne, M. D. Poulsen, H. Stapelfeldt, C. Z. Bisgaard, E. Hamilton and T. Seideman. Nonadiabative laser-induced alignment of iodobenzene molecules. *Phys. Rev. A*, **70**, 063410 (2004).
- [57] A. Ben Haj-Yedder, A. Auger, C. M. Dion, E. Cancés, A. Keller, C. Le Bris and O. Atabek. Numerical optimization of laser fields to control molecular orientation. *Phys. Rev. A*, **66**, 063401 (2002).
- [58] M. Machholm and N. E. Henriksen. Field-free orientation of molecules. *Phys. Rev. Lett.*, **87**, 193001 (2001).
- [59] C. M. Dion, A. Keller and O. Atabek. Orienting molecules using half-cycle pulses. *Eur. Phys. J. D*, **14**, 249 (2001).
- [60] D. You, R. R. Jones, P. H. Bucksbaum and D. R. Dykaar. Generation of high-power sub-single-cycle 500-fs electromagnetic pulses. *Opt. Lett.*, **18**, 290 (1993).
- [61] C. M. Dion, A. Ben Haj-Yedder, E. Cancés, C. Le Bris, A. Keller and O. Atabek. Optimal laser control of orientation: the kicked molecule. *Phys. Rev. A*, **65**, 063408 (2002).
- [62] J. Almlöf. Ab initio calculation of the vibrational structure and equilibrium geometry in  $\text{HF}_2^-$  and  $\text{DF}_2^-$ . *Chem. Phys. Lett.*, **17**, 49 (1972).
- [63] S. A. Barton and W. R. Thorson. Vibrational dynamics of hydrogen bonds. I.  $\text{FHF}^-$  system. *J. Chem. Phys.*, **71**, 4263 (1979).
- [64] C. L. Janssen, W. D. Allen and H. F. Schaefer III. The infrared spectrum of the hydrogen bifluoride anion: unprecedented variation with level of theory. *Chem. Phys. Lett.*, **131**, 352 (1986).
- [65] V. C. Epa, J. H. Choi, M. Klobukowski and W. R. Thorson. Vibrational dynamics of the bifluoride ion. I. Construction of a model potential surface. *J. Chem. Phys.*, **92**, 466 (1990).

- [66] V. C. Epa and W. R. Thorson. Vibrational dynamics of the bifluoride ion. II. Adiabatic separation and proton dynamics. *J. Chem. Phys.*, **92**, 473 (1990).
- [67] V. C. Epa and W. R. Thorson. Vibrational dynamics of the bifluoride ion. III. F–F ( $\nu_1$ ) eigenstates and vibrational intensity calculations. *J. Chem. Phys.*, **93**, 3773 (1990).
- [68] R. B. Metz and D. M. Neumark. Adiabatic three-dimensional simulations of the  $\text{IHI}^-$ ,  $\text{BrHI}^-$ , and  $\text{BrHBr}^-$  photoelectron spectra. *J. Chem. Phys.*, **97**, 962 (1992).
- [69] K. Yamashita, K. Morokuma and C. Leforestier. Theoretical study of the highly vibrationally excited states of  $\text{FHF}^-$ : *ab initio* potential energy surface and hyper-spherical formulation. *J. Chem. Phys.*, **99**, 8848 (1993).
- [70] A. B. McCoy, R. B. Gerber and M. A. Ratner. A quantitative approximation for the quantum dynamics of hydrogen transfer: Transition state dynamics and decay in  $\text{ClHCl}^-$ . *J. Chem. Phys.*, **101**, 1975 (1994).
- [71] J. E. Del Bene and M. J. T. Jordan. A comparative study of vibrational anharmonicity in the bihalide anions  $\text{XHX}^-$ : X = F, Cl, Br. *Spectrochim. Acta*, **55**, 719 (1999).
- [72] A. Kaledin, S. Skokov, J. M. Bowman and K. Morokuma. Theoretical study of the photodetachment spectroscopy of the  $\text{IHBr}$  and  $\text{IDBr}$  anions. *J. Chem. Phys.*, **113**, 9479 (2000).
- [73] J. W. Larson and T. B. McMahon. Strong hydrogen bonding in gas-phase anions. An ion cyclotron resonance determination of fluoride binding energetics to Brønsted acids from gas-phase fluoride exchange equilibrium measurements. *J. Am. Chem. Soc.*, **105**, 2944 (1983).
- [74] J. W. Larson and T. B. McMahon. Gas-phase bihalide and pseudobihalide ions. An ion cyclotron resonance determination of hydrogen bond energies in  $\text{XHY}^-$  species (X, Y = F, Cl, Br, CN). *Inorg. Chem.*, **23**, 2029 (1984).
- [75] G. Caldwell and P. Kebarle. The hydrogen bond energies of the bihalide ions  $\text{XHX}^-$  and  $\text{YHX}^-$ . *Can. J. Chem.*, **63**, 1399 (1985).
- [76] M. Heni and E. Illenberger. The stability of the bifluoride ion ( $\text{HF}_2^-$ ) in the gas phase. *J. Chem. Phys.*, **83**, 6056 (1985).
- [77] K. Kawaguchi and E. Hirota. Infrared diode laser study of the hydrogen bifluoride anion:  $\text{FHF}^-$  and  $\text{FDF}^-$ . *J. Chem. Phys.*, **84**, 2953 (1986).

- [78] K. Kawaguchi and E. Hirota. Diode laser spectroscopy of the  $\nu_3$  and  $\nu_2$  bands of FHF<sup>-</sup> in 1300 cm<sup>-1</sup> region. *J. Chem. Phys.*, **87**, 6838 (1987).
- [79] K. Kawaguchi. Gas-phase infrared spectroscopy of ClHCl<sup>-</sup>. *J. Chem. Phys.*, **88**, 4186 (1988).
- [80] R. B. Metz, T. Kitsopoulos, A. Weaver and D. M. Neumark. Study of the transition state region in the Cl + HCl reaction by photoelectron spectroscopy of ClHCl<sup>-</sup>. *J. Chem. Phys.*, **88**, 1463 (1988).
- [81] S. E. Bradforth, A. Weaver, D. W. Arnold, R. B. Metz and D. M. Neumark. Examination of the Br + HI, Cl + HI, and F + HI hydrogen abstraction reactions by photoelectron spectroscopy of BrHI<sup>-</sup>, ClHI<sup>-</sup>, and FHI<sup>-</sup>. *J. Phys. Chem.*, **92**, 7205 (1990).
- [82] I. M. Waller, T. N. Kitsopoulos and D. M. Neumark. Threshold photodetachment spectroscopy of the I + HI transition-state region. *J. Phys. Chem.*, **94**, 2240 (1990).
- [83] S. E. Bradforth, D. W. Arnold, R. B. Metz, A. Weaver and D. M. Neumark. Spectroscopy of the transition state: hydrogen abstraction reactions of fluorine. *J. Phys. Chem.*, **95**, 8066 (1991).
- [84] D. M. Neumark. Transition-state spectroscopy via negative ion photodetachment. *Acc. Chem. Res.*, **26**, 33 (1993).
- [85] Z. Liu, H. Gómez and D. M. Neumark. Photoelectron spectroscopy of clustered transition state precursors IHI<sup>-</sup>·Ar and BrHI<sup>-</sup>·Ar. *Chem. Phys. Lett.*, **332**, 65 (2000).
- [86] Z. Liu, H. Gómez and D. M. Neumark. Transition state spectroscopy of the I + HI reaction in clusters: photoelectron spectroscopy of IHI<sup>-</sup>·Ar<sub>n</sub> ( $n = 1 - 15$ ). *Faraday Discuss.*, **118**, 221 (2001).
- [87] P. A. Frey, S. A. Whitt and J. B. Tobin. A low-barrier hydrogen bond in the catalytic triad of serine proteases. *Science*, **264**, 1994 (1994).
- [88] T. N. Kitsopoulos, I. M. Waller, J. G. Loeser and D. M. Neumark. High resolution threshold photodetachment spectroscopy of negative ions. *Chem. Phys. Lett.*, **159**, 300 (1989).
- [89] A. Osterwalder, M. J. Nee, J. Zhou and D. M. Neumark. High resolution photodetachment spectroscopy of negative ions via slow photoelectron imaging. *J. Chem. Phys.*, **121**, 6317 (2004).

- [90] J. E. Del Bene, M. J. T. Jordan, S. A. Perera and R. J. Bartlett. Vibrational effects on the F–F spin-spin coupling constant ( $^{2h}J_{F-F}$ ) in  $FHF^-$  and  $FDF^-$ . *J. Phys. Chem. A*, **105**, 8399 (2001).
- [91] B. S. Ault. Infrared matrix isolation studies of the  $M^+HF_2^-$  and  $M^+XHF^-$  ion pairs for  $X = Cl, Br,$  and  $I$ . *J. Phys. Chem.*, **83**, 837 (1979).
- [92] J. J. Sloan, D. G. Watson, J. M. Williamson and J. S. Wright. Direct vs complex reaction dynamics for  $F + OH \rightarrow HF + O$ . *J. Phys. Chem.*, **75**, 1190 (1981).
- [93] R. N. Dixon and H. Tachikawa. The photodetachment spectrum of  $OHF^-$ : the influence of vibration at a transition state. *Mol. Phys.*, **97**, 195 (1999).
- [94] S. Gómez-Carrasco, L. González-Sánchez, A. Aguado, M. Paniagua, O. Roncero, M. L. Hernández and J. M. Alvariño. Dynamics and kinetics of the  $F + OH$  reaction on the ground triplet potential energy surface. *Chem. Phys. Lett.*, **383**, 25 (2004).
- [95] L. González-Sánchez, S. Gómez-Carrasco, A. Aguado, M. Paniagua, M. L. Hernández, O. Roncero and J. M. Alvariño. Photodetachment spectrum of  $OHF^-$ : three-dimensional study of the heavy-light-heavy resonances. *J. Chem. Phys.*, **121**, 309 (2004).
- [96] L. González-Sánchez, S. Gómez-Carrasco, A. Aguado, M. Paniagua, M. L. Hernández, O. Roncero and J. M. Alvariño. Transition state dynamics of  $OHF$  on several electronic states: photodetachment spectrum of  $OHF^-$  and conical intersections. *J. Chem. Phys.*, **121**, 9865 (2004).
- [97] K. P. Huber and G. Herzberg. Molecular spectra and molecular structure. IV. Constants of diatomic molecules. Van Nostrand Reinhold Company, New York (1979).
- [98] M. Born and R. Oppenheimer. Zur Quantentheorie der Molekeln. *Ann. Phys.*, **84**, 457 (1927).
- [99] A. Szabo and N. S. Ostlund. Modern quantum chemistry: introduction to advanced electronic structure theory. Macmillan, New York (1982).
- [100] C. C. J. Roothaan. New developments in molecular orbital theory. *Rev. Mod. Phys.*, **23**, 69 (1951).
- [101] G. G. Hall. The molecular orbital theory of chemical valency. VIII A method of calculating ionization potentials. *Proc. Roy. Soc. (London)*, **A205**, 541 (1951).



- [102] J. A. Pople and R. K. Nesbet. Self-consistent orbitals for radicals. *J. Chem. Phys.*, **22**, 571 (1954).
- [103] F. Jensen. Computational Chemistry. Wiley, Chichester (1999).
- [104] J. K. L. MacDonald. Successive approximations by the Rayleigh-Ritz variation method. *Phys. Rev.*, **43**, 830 (1933).
- [105] J. A. Pople, M. Head-Gordon and K. Raghavachari. Quadratic configuration interaction. A general technique for determining electron correlation energy. *J. Chem. Phys.*, **87**, 5968 (1987).
- [106] J. Čížek and J. Paldus. Coupled cluster approach. *Physica Scripta*, **21**, 251 (1980).
- [107] C. Møller and M. S. Plesset. Note on an approximation treatment for many-electron systems. *Phys. Rev.*, **46**, 618 (1934).
- [108] M. L. Leininger, W. D. Allen H. F. Schaefer III and C. D. Sherrill. Is Møller-Plesset perturbation theory a convergent *ab initio* method? *J. Chem. Phys.*, **112**, 9213 (2000).
- [109] I. A. Levine. Quantum chemistry. Prentice Hall, New Jersey (2000).
- [110] G. Herzberg. Molecular spectra and molecular structure. I. Spectra of diatomic molecules. Van Nostrand Reinhold Company, New York (1950).
- [111] R. Meyer. Trigonometric interpolation method for one-dimensional quantum-mechanical problems. *J. Chem. Phys.*, **52**, 2053 (1970).
- [112] C. C. Marston and G. G. Balint-Kurti. The Fourier grid Hamiltonian method for bound state eigenvalues and eigenfunctions. *J. Chem. Phys.*, **91**, 3571 (1989).
- [113] G. G. Balint-Kurti, R. N. Dixon and C. C. Marston. Grid methods for solving the Schrödinger equation and time-dependent quantum dynamics of molecular photofragmentation and reactive scattering processes. *Int. Rev. in Phys. Chem.*, **11**, 317 (1992).
- [114] D. O. Harris, G. G. Engerholm and W. D. Gwinn. Calculation of matrix elements for one-dimensional quantum-mechanical problems and the application to anharmonic oscillators. *J. Chem. Phys.*, **43**, 1515 (1965).
- [115] J. C. Light, I. P. Hamilton and J. V. Lill. Generalized discrete variable approximation in quantum mechanics. *J. Chem. Phys.*, **82**, 1400 (1985).
- [116] N. Balakrishnan, C. Kalyanaraman and N. Sathyamurthy. Time-dependent quantal approach to reactive scattering and related processes. *Phys. Rep.*, **280**, 79 (1997).

- [117] R. Kosloff and D. Kosloff. A Fourier method solution for the time dependent Schrödinger equation: A study of the reaction  $H^+ + H_2$ ,  $D^+ + HD$ , and  $D^+ + H_2$ . *J. Chem. Phys.*, **79**, 1823 (1983).
- [118] W. H. Press, W. T. Vetterling, S. A. Teukolsky and B. P. Flannery. Numerical recipes in Fortran: the art of scientific computing, 2nd Ed. Cambridge University Press, Cambridge (1992).
- [119] *qmbound*, WAVEPACKET by B. Schmidt, Berlin (1997).
- [120] E. Merzbacher. Quantum mechanics. John Wiley & Sons, New York (1970).
- [121] S. Mukamel. Principles of nonlinear optical spectroscopy. Oxford University Press, Oxford (1995).
- [122] C. Cohen-Tannoudji, B. Liu and F. Laloë. Quantum mechanics. Wiley, New York (1977).
- [123] R. Loudon. The quantum theory of light. Oxford University Press, Oxford (1984).
- [124] E. J. Heller. Quantum corrections to classical photodissociation models. *J. Chem. Phys.*, **68**, 2066 (1978).
- [125] G. K. Paramonov and V. A. Savva. Resonance effects in molecule vibrational excitation with picosecond laser pulses. *Phys. Lett. A*, **97**, 340 (1983).
- [126] A. D. Bandrauk. Molecules in intense laser fields: fundamentals. In J. Manz and L. Wöste, editors, Femtosecond Chemistry. VCH, Weinheim (1995).
- [127] L. V. Keldysh. Ionization in the field of a strong electromagnetic wave. *Sov. Phys. JETP*, **20**, 1307 (1965).
- [128] C. Hirlimann. Pulsed optics. In C. Rulliere, editor, Femtosecond Laser Pulses. Springer, Berlin (1998).
- [129] M. Nisoli, G. Sansone, S. Stagira, S. De Silvestri, C. Vozzi, M. Pascolini, L. Polletto, P. Villoresi and G. Tondello. Effects of carrier-envelope phase differences of few-optical cycle light pulses in single-shot high-order-harmonic spectra. *Phys. Rev. Lett.*, **91**, 213905 (2003).
- [130] D. J. Jones, S. A. Diddams, J. K. Ranka, A. Stenz, R. S. Windeler, J. L. Hall and S. T. Cundiff. Carrier-envelope phase control of femtosecond mode-locked lasers and direct optical frequency synthesis. *Science*, **288**, 635 (2000).

- [131] S. Chelkowski and A. D. Bandrauk. Asymmetries in strong-field photoionization by few-cycle laser pulses: kinetic-energy spectra and semiclassical explanation of the asymmetries of fast and slow electrons. *Phys. Rev. A*, **71**, 053815 (2005).
- [132] A. Bandrauk. Private communication.
- [133] R. Schinke. Photodissociation Dynamics. Cambridge University Press, Cambridge (1993).
- [134] J. J. Sakurai and S. F. Tuan, ed. Modern quantum mechanics, revised edition. Addison-Wesley Publishing Company, Bonn (1994).
- [135] S. A. Rice and M. Zhao. Optical control of molecular dynamics. Wiley Interscience, New York (2000).
- [136] H. Tal-Ezer and R. Kosloff. An accurate and efficient scheme for propagating the time dependent Schrödinger equation. *J. Chem. Phys.*, **81**, 3967 (1984).
- [137] C. J. Lanczos. An iterative method for the solution of the eigenvalue problem of linear differential and integral operators. *J. Res. Natl. Bur. Stand.*, **45**, 255 (1950).
- [138] M. D. Feit, J. A. Fleck Jr. and A. Steiger. Solution of the Schrödinger Equation by a Spectral Method. *J. Comput. Phys.*, **47**, 412 (1982).
- [139] M. D. Feit and J. A. Fleck Jr. Solution of the Schrödinger equation by a spectral method II: Vibrational energy levels of triatomic molecules. *J. Chem. Phys.*, **78**, 301 (1983).
- [140] A. D. Bandrauk and H. Shen. Improved exponential split-operator method for solving the time-dependent Schrödinger equation. *Chem. Phys. Lett.*, **176**, 428 (1991).
- [141] *qmpropa*, WAVEPACKET by B. Schmidt, Berlin (1997).
- [142] R. N. Zare. Angular momentum. Wiley Interscience, New York (1988).
- [143] P. W. Atkins und R. S. Friedman. Molecular quantum mechanics. Oxford University Press, Oxford (1997).
- [144] G. Herzberg. Molecular spectra and molecular structure. II. Infrared and Raman spectra of polyatomic molecules. Van Nostrand Reinhold Company, New York (1945).
- [145] T. Seideman. Private communication.

- [146] J. Ortigoso, M. Rodríguez, M. Gupta and B. Friedrich. Time evolution of pendular states created by the interaction of molecular polarizability with a pulsed nonresonant laser field. *J. Chem. Phys.*, **110**, 3870 (1999).
- [147] C. M. Dion, A. Keller, O. Atabek and A. D. Bandrauk. Laser-induced alignment dynamics of HCN: roles of the permanent dipole moment and the polarizability. *Phys. Rev. A*, **59**, 1382 (1999).
- [148] T. Seideman. Shaping molecular beams with intense light. *J. Chem. Phys.*, **107**, 10420 (1997).
- [149] B. Friedrich and D. Herschbach. Polarization of molecules induced by intense nonresonant laser fields. *J. Phys. Chem.*, **99**, 15686 (1995).
- [150] B. Friedrich, N. H. Nahler and U. Buck. The pseudo-first-order Stark effect and the orientation of HXeI molecules. **50**, 2677 (2003).
- [151] M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, V. G. Zakrzewski, J. A. Jr. Montgomery, R. E. Stratmann, J. C. Burant, S. Dapprich, J. M. Milliam, A. D. Daniels, K. N. Kudin, M. C. Strain, O. Farkas, J. Tomasi, V. Barone, M. Cossi, R. Cammi, B. Mennucci, C. Pomelli, C. Adamo, S. Clifford, J. Ochterski, G. A. Petersson, P. Y. Ayala, Q. Cui, K. Morokuma, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. Cioslowski, J. V. Ortiz, A. G. Baboul, B. B Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. Gomperts, R. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, C. González, M. Challacombe, P. M. W. Gill, B. Johnson, W. Chen, M. W. Wong, J. L. Andres, M. Head-Gordon, E. S. Replogle and J. A. Pople, GAUSSIAN 98, Revision A.7, Gaussian, Inc., Pittsburgh PA, 1998.
- [152] D. E. Woon and T. H. Dunning, Jr. Gaussian basis sets for use in correlated molecular calculations. IV. Calculation of static electrical response properties. *J. Chem. Phys.*, **100**, 2975 (1994).
- [153] J. Full, L. González and J. Manz. Neutral-to-ionic ZEKE transition dipole couplings beyond Koopmans' picture: application to femtosecond pump-probe spectroscopy. *Chem. Phys.*, **314**, 143 (2005).
- [154] K. Kawaguchi and E. Hirota. Annual review of the Institute of Molecular Science, p. 27, item II-A-2. Myodaiji (1988).
- [155] R. D. Hunt and L. Andrews. FTIR spectra of the  $\text{HF}_2^-$  and  $\text{H}_2\text{F}_3^-$  anions isolated in solid argon and neon. *J. Chem. Phys.*, **87**, 6819 (1987).

- [156] G. L. Cote and H. W. Thompson. Infrared spectra and the solid state. III. Potassium bifluoride. *Proc. R. Soc. London Ser. A*, **210**, 206 (1951).
- [157] P. Dawson. Spectroscopic detection of the hydrogen bond isotope effect in  $\text{KHF}_2$ . *J. Chem. Soc. Faraday Trans. 2*, **68**, 1448 (1972).
- [158] Fortran subroutines for mathematical applications. Visual Numerics, Inc., Stuttgart (1997).
- [159] V. Engel and H. Metiu. Vibrational coherence effects in the pump-probe studies of photochemical predissociation. *J. Chem. Phys.*, **95**, 3444 (1991).
- [160] J. Manz, B. Reischl, T. Schröder, F. Seyl and B. Warmuth. On the laser-femtochemistry approach to coherent molecular vibrations: model simulations for  $\text{Ni}[\text{C}_2\text{D}_4] \rightarrow \text{Ni} + \text{C}_2\text{D}_4$ . *Chem. Phys. Lett.*, **198**, 483 (1992).
- [161] *rigid rotor*, by N. Elghobashi, Berlin (1997).
- [162] D. A. McQuarrie. Statistical mechanics. Harper & Row, New York (1973).
- [163] C. Manzoni, G. Cerullo, and S. De Silvestri. Ultrabroadband self-phase-stabilized pulses by difference-frequency generation. *Opt. Lett.*, **29**, 2668 (2004).
- [164] A. Apolonski, A. Poppe, G. Tempea, C. Spielmann, T. Udem, R. Holzwarth, T. W. Hänsch and F. Krausz. Controlling the phase evolution of few-cycle light pulses. *Science*, **288**, 635 (2000).
- [165] G. G. Paulus, F. Grasbon, H. Walther, P. Villoresi, M. Nisoli, S. Stagira, E. Priori and S. De Silvestri. Absolute-phase phenomena in photoionization with few-cycle pulses. *Nature*, **414**, 182 (2001).
- [166] G. Sansone, C. Vozzi, S. Stagira, M. Pascolini, L. Poletto, P. Villoresi, G. Tondello, S. De Silvestri and M. Nisoli. Observation of carrier-envelope phase phenomena in the multi-optical-cycle regime. *Phys. Rev. Lett.*, **92**, 113904 (2004).
- [167] R. R. Fontana. Spin-orbit and spin-spin interactions in diatomic molecules. I. Fine structure of  $\text{H}_2$ . *Phys. Rev.*, **125**, 220 (1962).
- [168] B. Friedrich. Private communication.
- [169] N. Elghobashi and L. González. A theoretical anharmonic study of the infrared absorption spectra of  $\text{FHF}^-$ ,  $\text{FDF}^-$ ,  $\text{OHF}^-$ , and  $\text{ODF}^-$  anions. In preparation.
- [170] N. Elghobashi, L. González and J. Manz. Quantum model simulations of symmetry breaking and control of bond selective dissociation of  $\text{FHF}^-$  using IR + UV laser pulses. *J. Phys. Chem.*, **120**, 8002 (2004).

- [171] N. Elghobashi, L. González and J. Manz. Quantum model simulations for isotope effects of IR + UV laser pulses on symmetry and selective hydrogen bond breaking. *Z. Phys. Chem.*, **217**, 1577 (2003).
- [172] N. Elghobashi and J. Manz. Separating the photofragments of randomly oriented symmetric reactants by IR + UV laser pulses: quantum simulations for  $\text{FHF}^- \rightarrow \text{F} + \text{FH} + \text{e}$ . *Israel J. Chem.*, **43**, 293 (2003).
- [173] N. Elghobashi, L. González and T. Seideman. Molecular orientation and selective dissociation in  $\text{OHF}^-$  using half-cycle laser pulses. In preparation.
- [174] N. Elghobashi and L. González. Breaking the strong and weak bonds of  $\text{OHF}^-$  using few-cycle IR + UV laser pulses. *Phys. Chem. Chem. Phys.*, **6**, 4071 (2004).
- [175] U. Banin, A. Bartana, and R. Kosloff. Impulsive excitation of coherent vibrational motion ground surface dynamics induced by intense short pulses. *J. Chem. Phys.*, **101**, 8461 (1994).
- [176] D. A. McQuarrie. Quantum chemistry. Oxford Science Books, Oxford (1983).