

7. Literatur

- Abildsnes J, Tømmerås, BÅ (2000)** Impacts of experimental habitat fragmentation on ground beetles (Coleoptera, Carabidae) in a boreal spruce forest. *Annales Zoologici Fennici* **37**: 201-212
- Alaruikka D, Kotze DJ, Matveinen K, Niemelä J (2003)** Carabid beetle and spider assemblages along a forested urban-rural gradient in southern Finland. *Journal of Insect Conservation* **6**: 195-206
- Alberti M, Marzluff J M, Shulenberger E, Bradley G, Ryan C, Zumbrunnen C (2003)** Integrating Humans into Ecology: Opportunities and Challenges for Studying Urban Ecosystems. *Bioscience* **53(12)**: 1169-1179.
- Alibert P, Moureau B, Dommergues J-L, David B (2001)** Differentiation at a microgeographical scale within two species of ground beetle, *Carabus auronitens* and *C. nemoralis* (Coleoptera, Carabidae): a geometrical morphometric approach. *Zoologica Scripta* **30**: 299-311
- Alstad DN, Edmunds GF Jr, Weinstein LH (1982)** Effects of Air Pollutants on Insect Populations. *Annual Review of Entomology* **27**: 369-384
- Andersen LW, Fog K, Damgaard C (2004)** Habitat fragmentation causes bottlenecks and inbreeding in the European tree frog (*Hyla arborea*). *Proceedings of the Royal Society of London B* **271**: 1293-1302
- Andreassen HP, Halle S, Ims RA (1996)** Optimal width of movement corridors for root voles: not too narrow and not too wide. *Journal of Applied Ecology* **33**: 63-70
- Andrèn H (1994)** Effects of habitat fragmentation on birds and mammals in landscapes with different proportions of suitable habitat: a review. *Oikos* **71**: 355-366
- Andrewartha HG, Birch LC (1954)** *The Distribution and Abundance of Animals*. The University of Chicago Press - Chicago, Illinois
- Angold PG, Sadler JP, Hill MO, Pullin A, Rushton S, Austin K, Small EC, Wood B, Wadsworth R, Sanderson R, Thompson K (2006)** Biodiversity in urban habitat patches. *Science of the Total Environment* **360**: 196– 204
- Anthrop M (2000)** Changing patterns in the urbanized countryside of Western Europe. *Landscape Ecology* **15**: 257-270
- Arndt E, Trautner J (2004)** Carabini. In: Freude H, Harde KW, Lohse GA, Klausnitzer B, Müller-Motzfeld G (Hrsg.) *Die Käfer Mitteleuropas. Band 2*. Spektrum-Verlag - Heidelberg, Berlin, pp. 30-60
- Arnold CL, Gibbons C (1996)** Impervious surface coverage: The emergence of a key environmental indicator. *Journal of the American Planning Association* **62(2)**: 243-258

- Assing V (1988)** Die Kurzflügelkäferfauna (Coleoptera: Staphylinidae) ausgewählter Grün-, Ruderal- und Kleingartenflächen im Stadtgebiet Hannovers: Ein Beitrag zur Faunistik und Ökologie einer Großstadt. *Berichte der naturhistorischen Gesellschaft Hannover* **130**: 111-131
- Assing V (1992)** Die Kurzflügelkäfer (Coleoptera: Staphylinidae) eines urbanen Inselbiotopkomplexes in Hannover. 3. Beitrag zur Faunistik und Ökologie der Staphyliniden von Hannover. *Berichte der Naturhistorischen Gesellschaft Hannover* **134**: 173-187
- Assmann T (1999)** The ground beetle fauna of ancient and recent woodlands in the lowlands of north-west Germany (Coleoptera, Carabidae). *Biodiversity and Conservation* **8**: 1499-1517
- Assmann T, Nolte O, Reuter H (1994)** Postglacial colonization of middle Europe by *Carabus auronitens* as revealed by population genetics (Coleoptera, Carabidae) In: Desender K, Dufrene M, Loreau M, Luff ML, Maelfait JP (Hrsg.) *Carabid beetles: ecology and evolution*. Kluwer Academic Publishers – Dordrecht, Boston, London, pp. 3-9
- Assmann T, Weber F (1997)** On the allozyme differentiation of *Carabus punctatoauratus* Germar (Coleoptera, Carabidae). *Journal of Zoological Systematics and Evolutionary Research* **35**: 33-43
- Awise JC, Hamrick JL (1996)** *Conservation genetics*. Chapman and Hall-New York
- Baars M A (1979)** Patterns of Movement of Radioactive Carabid Beetles. *Oecologia* **44**: 125-140
- Backhaus K, Erichson B, Plinke W, Weiber R (1990)** *Multivariate Analysemethoden*. Springer Verlag - Berlin
- Balbi J Freiherr von (1748-1749)** Geographische Special Chartre von der Mittelmarck, eines Theils des Unter Barnim-Havelaend-Zauch- und Teltauischen Creises. Handgezeichnete Karte in der Kartensammlung der Staatsbibliothek Berlin
- Bale JS (1991)** Insects at low temperature: a predictable relationship? *Functional Ecology* **5**: 291-298
- Balloux F, Brünnner H, Lugon-Moulin N, Hausser J, Goudet J (2000)** Microsatellites can be misleading: an empirical and simulation study. *Evolution* **54(4)**: 1414-1422
- Balloux F, Lugon-Moulin N (2002)** The estimation of population differentiation with microsatellite markers. *Molecular Ecology* **11**: 155-165
- Barndt D, Brase S, Glauche M, Gruttke H, Kegel B, Platen R, Winkelmann H (1991)** Die Laufkäferfauna von Berlin (West) – mit Kennzeichnung und Auswertung der verschollenen und gefährdeten Arten. In: (Auhagen A, Platen R, Sukopp H (Hrsg.) *Rote Listen der gefährdeten Pflanzen und Tiere in Berlin.*, Sonderband **S6** von: Landschaftentwicklung und Umweltforschung. Technische Universität Berlin, pp. 243-275

- Basedow T, Rzehak H, Dickler E (1990)** Untersuchungen zur Flugaktivität epigäischer Raubarthropoden mittels Licht- und Fensterfallen. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* **7**: 386-394
- Baum KA, Haynes KJ, Dilleuth FP, Cronin J T (2004)** The matrix enhances the effectiveness of corridors and stepping stones. *Ecology* **85(10)**: 2671-2676
- Baur B, Erhardt A (1995)** Habitat fragmentation and habitat alterations: principal threats to most animal and plant species. *Gaia* **4**: 221-226
- Berggren Å, Birath B, Kindvall O (2002)** Effect of Corridors and Habitat Edges on Dispersal Behaviour, Movements Rates, and Movement Angles in Roesel's Bush-Cricket (*Metrioptera roeseli*). *Conservation Biology* **16(6)**: 1562-1569
- BNatschG (2002)** Gesetz über Naturschutz und Landschaftspflege. *Bundesgesetzblatt Teil I* **22**: 1193-1218
- Bolger DT, Suarez AV, Crooks KR, Morrison SA, Case TJ (2000)** Arthropods in urban habitat fragments in southern California: Area, age, and edge effects. *Ecological Applications* **10(4)**: 1230-1248
- Bowne DR, Peles JD, Barrett GW (1999)** Effects of landscape spatial structure on movement patterns of the hispid cotton rat (*Sigmodon hispidus*). *Landscape Ecology* **14**: 53-65.
- Brandmayr P, Zetto Brandmayr T (1994)** The evolutionary history of the genus *Abax* (Coleoptera, Carabidae). In: Desender K, Dufrene M, Loreau M, Luff ML, Maelfait JP (Hrsg.) *Carabid beetles: ecology and evolution*. Kluwer Academic Publishers – Dordrecht, Boston, London, pp. 19-24
- Brouat C, Mondor-Genson G, Audiot P, Sennedot F, Lesobre L, Rasplus J-Y (2002)** Isolation and characterization of microsatellite loci in the ground beetle *Carabus nemoralis* (Coleoptera, Carabidae) *Molecular Ecology Notes* **2**: 119-120
- Brouat C, Sennedot F, Audiot P, Leblois R, Rasplus J-Y (2003)** Fine-scale genetic structure of two carabid species with contrasted levels of habitat specialization. *Molecular Ecology* **12**: 1731-1745
- Burger M-L (1971)** Proprioceptive Course Control in Millipedes (*Schizophyllum sabulosum*). *Zeitschrift für vergleichende Physiologie* **71**: 219-254
- Butterweck MD (1998)** *Metapopulationsstudien an Waldlaufkäfern (Coleoptera: Carabidae) Einfluß von Korridoren und Trittsteinbiotopen*. Wissenschaft und Technik Verlag – Berlin
- Carreiro MM, Tripler CE (2005)** Forest Remnants Along Urban-Rural Gradients: Examining Their Potential for Global Change Research. *Ecosystems* **8**: 568-582
- CBIF (2006)** Canadian Biodiversity Information facility. <http://www.cbif.gc.ca>
- Charrier S, Petit S, Burel F (1997)** Movements of *Abax parallelepipedus* (Coleoptera, Carabidae) in woody habitats of a hedgerow network landscape: a radio-tracing study. *Agriculture, Ecosystems & Environment* **61**: 133-144

- Coates DJ (1991)** Gene flow along corridors. In: Saunders DA, Hobbs RJ (Hrsg.) *Nature Conservation*. Surrey Beatty and Sons – Chipping Norton Australia, pp. 408-409
- Cole LJ, McCracken DI, Foster GN, Aitken MN (2001)** Using *Collembola* to assess the risks of applying metal-rich sewage sludge to agricultural land in western Scotland. *Agriculture, Ecosystems and Environment* **83**: 177-189
- Coleman DC, Crossley DA Jr., Hendrix PF (2004)** *Fundamentals of Soil Ecology*. Elsevier Academic Press – Amsterdam
- Collinge SK (2000)** Effects of grassland fragmentation on insect species loss, colonization, and movement patterns. *Ecology* **81**: 2211-2226
- Colwell RK (2004)** EstimateS: Statistical estimation of species richness and shared species from samples. Version 7. <http://purl.oclc.org/estimates>.
- Cornelius R (1995)** *Geschichte der Waldentwicklung*. Monitoringprogramm Naturhaushalt Heft 3 der Senatsverwaltung für Stadtentwicklung und Umweltschutz Berlin, Kulturbuch-Verlag – Berlin
- Czechowski W (1982)** Occurrence of carabids (Coleoptera, Carabidae) in the urban greenery of Warsaw according to the land utilization and cultivation. *Memorabilia Zoologica* **39**: 3-108
- Davis BNK (1978)** Urbanisation and the diversity of insects. In: Mound LA, Waloff N (Hrsg.) *Diversity of Insect Faunas*. Symposia of the Royal Entomological Society of London 9 Blackwell – Oxford, pp. 126-138
- Davis BNK (1979)** The ground arthropods of London gardens. *London Naturalist* **58**: 15-24
- Davis BNK (1982)** Habitat diversity and invertebrates in urban areas. In: Bornkamm R, Lee JA, Seaward MRD (Hrsg.) *Urban Ecology*. Blackwell – Oxford, pp. 49-63
- Decker (1816- ca. 1824)** *Landaufnahme durch den preussischen Generalstab unter Leitung des Maj. Decker*. Handgezeichnete Karten in der Kartensammlung der Staatsbibliothek Berlin
- Deichsel R (2004)** Winter activity of epigeic arthropods in a central European oak forest. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* **14**: 501-504
- Deichsel R (2006)** Species change in an urban setting - ground and rove beetles (Coleoptera: Carabidae and Staphylinidae) in Berlin. *Urban Ecosystems* **9**: 161-178
- De Lattin G (1967)** *Grundriss der Zoogeographie*. Gustav Fischer Verlag – Jena
- Den Boer PJ (1970)** On the Significance of Dispersal Power for Populations of Carabid-Beetles (Coleoptera, Carabidae). *Oecologia* **4**: 1-28
- Den Boer PJ (1981)** On the survival of populations in a heterogeneous and variable environment. *Oecologia* **50**: 39-53
- Den Boer PJ (1990)** The survival value of dispersal in terrestrial arthropods. *Biological Conservation* **54**: 175-192

- Desender K (1989)** Heritability of wing development and body size in a carabid beetle, *Pogonus chalceus* Marsham, and its evolutionary significance. *Oecologia* **78**: 513-520
- Desender K, Dufrene M, Loreau M, Luff ML, Maelfait JP (1994)** *Carabid Beetles – Ecology and Evolution*. Kluwer Academic Publishers – Dordrecht
- Desender K, Small E, Gaublomme E, Verdyck P (2005)** Rural-urban gradients and the population genetic structure of woodland ground beetles. *Conservation Genetics* **6**: 51-62
- Desender K, Verdyck P, Gaublomme E, Dhuyvetter H, Rasplus J-Y (2002)** Extreme genetic differentiation and isolation by non-distance in *Carabus auronitens* in relation to forest historical ecology in Flanders (Belgium). In: Szyszko J, Den Boer PJ, Bauer T (Hrsg.) *How to protect or what we know about Carabid Beetles*. Warsaw Agricultural University Press – Warschau, pp. 227-235
- Dhuyvetter H, Gaublomme E, Desender K (2004)** Genetic differentiation and local adaption in the salt-marsh beetle *Pogonus chalceus*: a comparison between allozyme and microsatellite loci. *Molecular Ecology* **13**: 1065-1074
- Diamond JM, May RM (1981)** Island biogeography and the design of natural reserves. In: May RM (Hrsg.) *Theoretical Ecology*. Blackwell Science – Oxford, pp. 282-252
- Diamond J, Case TJ (1986)** *Community Ecology*. Harper & Row – New York
- Díaz JA, Pérez-Tris J, Tellería JL, Carbonell R, Santos T (2005)** Reproductive Investment of a Lacertid Lizard in Fragmented Habitat. *Conservation Biology* **19(5)**: 1578-1585
- Dieckmann U, O’Hara B, Weisser W (1999)** The evolutionary ecology of dispersal. *Trends in Ecology and Evolution* **14**: 88-90
- Dingle H (1961)** Correcting behaviour in boxelder bugs. *Ecology* **42**: 207-211
- Drees C, Weber F (2001)** Hecken als Korridorhabitate in einer fragmentierten Kulturlandschaft: Untersuchungen an *Carabus auronitens* F. (Col. Carabidae). *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* **13**: 571-578
- El Mousadik A, Petit R (1996)** High level of genetic differentiation for allelic richness among populations of the argan tree (*Argania spinosa* (L.) Skeels) endemic to Morocco. *Theoretical and Applied Genetics* **92**: 832-839
- Endlicher W, Simon U (2005)** Editorial: Perspectives of Urban Ecology – The Metropolis of Berlin as a Natural and Socioeconomic System. *Die Erde* **136(2)**: 97-102
- Faeth SH, Kane TC (1978)** City parks as Islands for Diptera and Coleoptera. *Oecologia* **32**: 127-133
- FFH-Richtlinie (1992)** Richtlinie 92/43/EWG. *Amtsblatt der Europäischen Union* L 206: 7
- Frankham R (1995)** Conservation genetics. *Annual Review of Genetics* **29**: 305-327
- Frankham R (1996)** Relationship of genetic variation to population size in wildlife. *Conservation Biology* **10**: 1500-1508

- Frankham R, Ballou JD, Briscoe DA (2002)** *Introduction to Conservation Genetics*. Cambridge University Press – Cambridge
- Frankie GW, Ehler LE (1978)** Ecology of insects in urban environments. *Annual Review of Entomology* **23**: 367-387
- Fountain MT, Hopkin SP (2004)** A Comparative Study of the Effects of Metal Contamination on Collembola in the Field and in the Laboratory. *Ecotoxicology* **13**: 573-587
- Garnier S, Alibert P, Audiot P, Prieur B, Rasplus J-Y (2004)** Isolation by distance and sharp discontinuities in gene frequencies: implications for the phylogeography of an alpine insect species, *Carabus solieri*. *Molecular Ecology* **13**: 1883-1897
- Gaston KJ (1996)** *Biodiversity: a biology of numbers and difference*. Blackwell Science – Oxford
- Gautestad AO, Mysterud I (2005)** Intrinsic Scaling Complexity in Animal Dispersion and Abundance. *The American Naturalist* **165(1)**: 44-55
- Gibb H, Hochuli DF (2002)** Habitat fragmentation in an urban environment: large and small fragments support different arthropod assemblages. *Biological Conservation* **106(1)**: 91-100
- Gibbs JP, Stanton EJ (2001)** Habitat fragmentation and arthropod community change: Carrion beetles, phoretic mites, and flies. *Ecological Applications* **11(1)**: 79-85
- Gill DE (1978)** The metapopulation ecology of the red-spotted newt, *Notophthalmus viridescens* (Rafinesque). *Ecological Monographs* **48**: 145-166
- Goudet J, Raymond M, De Meeüs T, Rousset F (1996)** Testing differentiation in diploid populations. *Genetics* **144**: 1933-1940
- Goudet J (2001)** FSTAT, a Programm to Estimate and Test Gene Diversities and Fixation Indices, Version 2.9.3.2. Department of Ecology and Evolution University of Lausanne, Switzerland, <http://www2.unil.ch/popgen/softwares/fstat.htm>
- Grandchamp A-C, Niemelä J, Kotze DJ (2000)** The effects of trampling on assemblages of ground beetles (Coleoptera, Carabidae) in urban forests in Helsinki, Finland. *Urban Ecosystems* **4**: 321-332
- Gotelli NJ, Colwell RK (2001)** Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecology Letters* **4**: 379-391
- Guo S, Thompson E (1992)** Performing the exact test of Hardy-Weinberg proportion for multiple alleles. *Biometrics* **48**: 361-372
- Haddad NM (1999a)** Corridor and distance effects in interpatch movements: a landscape experiment with butterflies. *Ecological Applications* **9(2)**: 612-622
- Haddad NM (1999b)** Corridor Use Predicted from Behaviors at Habitat Boundaries. *The American Naturalist* **153(2)**: 215-227
- Haddad NM, Bowne DR, Cunningham A, Danielson BJ, Levey DJ, Sargent S, Spira T (2003)** Corridor use by diverse taxa. *Ecology* **84(3)**: 609-615

- Halme E, Niemelä J (1993)** Carabid beetles in fragments of coniferous forest. *Annales Zoologici Fennici* **30**: 17-30
- Hanski I, Gilpin ME (1997)** *Metapopulation Biology: Ecology, Genetics, and Evolution*. Academic Press – San Diego
- Hardy OJ, Vekemans X (1999)** Isolation by distance in a continuous population: reconciliation between spatial autocorrelation analysis and population genetics models. *Heredity* **83**: 145-154
- Hartl DL, Clarke AC (1989)** *Principles of Population Genetics*. Sinauer – Sunderland, Massachusetts
- Hastings A, Harrison S (1994)** Metapopulation dynamics and genetics. *Annual Review of Ecology and Systematics* **25**: 167-188
- Hebert C, Danzman RG, Jones MW, Bernatchez L (2000)** Hydrography and population genetic structure in brook charr (*Salvelinus fontinalis*, Mitchill) from eastern Canada. *Molecular Ecology* **9(7)**: 971-982
- Heisswolf A, Reichmann S, Obermaier E, Schröder B, Poethke HJ (2006)** Habitat size, isolation, and quality determine the distribution of a monophagous leaf beetle and its egg parasitoid in a fragmented landscape. *Verhandlungen der Gesellschaft für Ökologie* **36**: 104
- Hewitt GM (1999)** Post-glacial re-colonization of European biota. *Biological Journal of the Linnean Society* **68**: 87-112
- Heydemann B (1957)** Die Biotopstruktur als Raumwiderstand und Raumfülle für die Tierwelt. *Verhandlungen der deutschen zoologischen Gesellschaft Hamburg* **50**: 332-347
- Hockmann P, Menke K, Schlomberg P, Weber F (1992)** Untersuchungen zum individuellen Verhalten (Orientierung und Aktivität) des Laufkäfers *Carabus nemoralis* im natürlichen Habitat. *Abhandlungen aus dem Westfälischen Museum für Naturkunde* **54(4)**: 65-98
- Honěk, A (1997)** The effect of temperature on the activity of Carabidae (Coleoptera) in a fallow field. *European Journal of Entomology* **94**: 97-104
- Horbert M (2000)** Klimatologische Aspekte der Stadt- und Landschaftsplanung. *Schriftenr. Fachber. Umwelt. Gesellsch.* **113**: 1-330
- Huckenbeck B, Wipking W (2000)** (Über-)Leben in Raum und Zeit: Laufkäfer auf Biotopinseln in der Großstadt Köln. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* **12**: 307-310
- Hupfer P, Chmielewski F-M (1990)** *Das Klima von Berlin*. Akademie Verlag – Berlin
- Irmeler U (1998)** Die vertikale Verteilung flugaktiver Käfer (Coleoptera) in drei Wäldern Norddeutschlands. *Faunistisch-Ökologische Mitteilungen* **7**: 387-404
- Irmeler U (2001)** Bedeutung urbaner Brachen für Laufkäfer (Carabidae). *Faunistisch-Ökologische Mitteilungen* **8**: 135-145

- Ishitani M, Kotze DJ, Niemelä J (2003)** Changes in carabid beetle assemblages across an urban-rural gradient in Japan. *Ecography* **26**: 481-489
- Janzen DH (1983)** No park is an island - increase in interference from outside as park size decreases. *Oikos* **41**: 402-410
- Jopp F (2003)** *Empirische Analysen und Modellierungen des Ausbreitungsverhaltens von Wirbellosen aus einem Niedermoor*. Dissertation, Freie Universität Berlin
- Jopp F, Reuter H (2005)** Dispersal of carabid beetles – emergence of distribution patterns. *Ecological Modelling* **186**: 389-405
- Kareiva PM, Shigesada N (1983)** Analyzing Insect Movement as a Correlated Random Walk. *Oecologia* **56**: 234-238
- Kennedy PJ (1994)** The distribution and movement of ground beetles in relation to set-aside arable land. In: Desender K, Dufrene M, Loreau M, Luff ML, Maelfait JP (Hrsg.) *Carabid Beetles – Ecology and Evolution*. Kluwer Academic Publishers – Dordrecht, pp. 439-444
- Keller I, Largiadèr CR (2002)** Identification of one X-linked and five autosomal microsatellite loci in *Carabus violaceus* (Coleoptera, Carabidae) and their applicability to related taxa. *Molecular Ecology Notes* **2**: 290-292
- Keller I, Nentwig W, Largiadèr CR (2004)** Recent habitat fragmentation due to roads can lead to significant genetic differentiation in an abundant flightless ground beetle. *Molecular Ecology* **13**: 2983-2994
- Keller LF, Waller DM (2002)** Inbreeding effects in wild populations. *Trends in Ecology and Evolution* **17(5)**: 230-241
- Kennedy PJ (1994)** The distribution and movement of ground beetles in relation to set-aside arable land. In: Desender K, Dufrene M, Loreau M, Luff ML, Maelfait JP (Hrsg.) *Carabid beetles: ecology and evolution*. Kluwer Academic Publishers – Dordrecht, Boston, London, pp. 439-444
- Kersting T, Hahn W, Nordmeier V, Schlichting HJ (2003)** Automatische VideoANalyse 3.64 (VIANA). <http://didaktik.physik.uni-essen.de/viana>
- Kielhorn K-H (2004)** Entwicklung von Laufkäfergemeinschaften auf forstlich rekultivierten Kippenstandorten des Lausitzer Braunkohlenreviers. *Cottbuser Schriften zu Bodenschutz und Rekultivierung* **22**: 1-189
- Kimura M, Crow JF (1963)** On the maximum avoidance of inbreeding. *Genetic Research* **4**: 399-415
- Klausnitzer B, Richter K (1983)** Presence of an urban gradient demonstrated for carabid associations. *Oecologia* **59**: 79-82
- Klazenga N, de Vries HH (1994)** Walking distances of five differently sized ground beetle species. *Proceedings of the Section Experimental and Applied Entomology of the Netherlands Entomological Society* **5**: 99-100

- Klomann U (1977)** Die Carabidenfauna unterschiedlich belasteter Standorte im Raum Saarbrücken. *Faunistisch-floristische Notizen aus dem Saarland* **9(1-2)**: 12-18 (zitiert in: Nagel (1978) Speziesdiversität und Raumbewertung. *Verhandlungen des Deutschen Geographentages* 41: 486-498)
- Knutsen H, Rukke BA, Jorde PE, Ims RA (2000)** Genetic differentiation among populations of the beetle *Bolitophagus reticulatus* (Coleoptera: Tenebrionidae) in a fragmented and a continuous landscape. *Heredity* **84**: 667-676
- Koivula MJ, Vermeulen HJW (2005)** Highways and forest fragmentation – effects on carabid beetles (Coleoptera, Carabidae). *Landscape Ecology* **20**: 911-926
- Korge H (1991)** Liste der Kurzflügelkäfer (Coleoptera, Staphylinidae) von Berlin (West) mit Kennzeichnung der verschollenen und gefährdeten Arten (Rote Liste). In: Auhagen A, Platen R, Sukopp H (Hrsg.) Rote Listen der gefährdeten Pflanzen und Tiere in Berlin. Extra volume **S6** of: *Landschaftsentwicklung und Umweltforschung*. Technische Universität Berlin, pp. 277-317
- Kowarik I (2003)** Biologische Invasionen. Neophyten und Neozoen in Mitteleuropa. Ulmer Verlag – Stuttgart
- Kowarik I (2005)** Wild Urban Woodlands: Towards a Conceptual Framework. In: Kowarik I, Körner S (Hrsg.) Wild Urban Woodlands. Springer Verlag – Berlin
- Kühn I, Brandl R, Klotz S (2004)** The flora of German cities is naturally species rich. *Evolutionary Ecology Research* **6**: 749-764
- Kuttler W (2004)** Stadtklima. Teil 2: Phänomene und Wirkungen. *Zeitschrift für Umweltchemie und Ökotoxikologie* **16**: 263-274
- Lande R (1995)** Mutation and conservation. *Conservation Biology* **9**: 782-791
- Landgrebe D, Biehl L (2005)** MultiSpec – A Freeware Multispectral Image Data Analysis System. <http://dynamo.ecn.purdue.edu/~biehl/MultiSpec/Index.html>
- Landsberg H (1981)** *The Urban Climate*. International Geophysics Series **28** – New York
- Lapro (2006)** Landscape programme including Nature Conservation. Berlin Senate Department of Urban Development <http://www.stadtentwicklung.berlin.de/umwelt/landschaftsplanung/lapro/de/kontakt/index.shtml>
- Laurance WF, Yensen E (1991)** Predicting the impacts of edge effects in fragmented habitats. *Biological Conservation* **55**: 72-92
- Lauterbach A-W (1964)** Verbreitungs- und aktivitätsbestimmende Faktoren bei Carabiden in sauerländischen Wäldern. *Abhandlungen des Landesmuseums für Naturkunde Münster* **26 (4)**: 1-103
- Leberg PL (1996)** Applications of allozyme electrophoresis in conservation biology. In: Smith TB, Wayne RK (Hrsg.) *Molecular genetic approaches in conservation*. Oxford University Press – New York, Oxford, pp 87-103

- Leimu R, Mutikainen P, Koricheva J, Fischer M (2006)** How general are positive relationships between plant population size, fitness and genetic variation? *Journal of Ecology* **94**: 942-952
- Lemaire C, Allegrucci G, Naciri M, Bahri-Sfar L, Kara H, Bonhomme F (2000)** Do discrepancies between microsatellite and allozyme variation reveal differential selection between sea and lagoon in the sea bass (*Dicentrarchus labrax*). *Molecular Ecology* **9**: 457-467
- Levins R (1970)** Extinction. In: Gesternhaber M (Hrsg.) *Some Mathematical Problems in Biology*. American Mathematical Society - Providence, pp 77–107
- Lindroth CH (1945)** Die fennoskandischen Carabidae, eine tiergeographische Studie. I. Spezieller Teil. *Göteborgs kungliga Vetenskaps- och Vitterhets-Samhälles Handlingar Serie B* **4(1)**: 1-709
- Lindroth CH (1949)** Die Fennoskandischen Carabidae. III. Allgemeiner Teil. *Göteborgs kungliga Vetenskaps- och Vitterhets- Samhälles Handlingar Serie B* **4(3)**: 1-911
- Little CE (1990)** *Greenways for America*. John Hopkins University Press – Baltimore
- Loeschke V, Tomiuk J, Jain SK (1994)** *Conservation genetics*. Birkhäuser – Basel
- Lomolino MV (1994)** An evaluation of alternative strategies for building networks of nature reserves. *Biological Conservation* **69**: 243-249
- Lund U, Agostinelli C (2006)** Circular statistics. A package for the statistical software R. <http://cran.r-project.org/src/contrib/Descriptions/circular.html>
- Lynch M, Conery J, Burger R (1995)** Mutation accumulation and the extinction of small populations. *The American Naturalist* **146**: 489-518
- MacArthur RH, Wilson EO (1967)** *The Theory of Island Biogeography*. Princeton University Press – Princeton, pp. 1-203
- Mader H-J (1979)** Die Isolationswirkung von Verkehrsstraßen auf Tierpopulationen am Beispiel von Arthropoden und Kleinsäugetern der Waldbiozönose. *Schriftenreihe für Landschaftspflege und Naturschutz* **19**: 1-130
- Mader H-J (1980)** Die Verinselung der Landschaft aus tierökologischer Sicht. *Natur und Landschaft* **55**: 91-96
- Mader H-J (1981)** Untersuchungen zum Einfluß der Flächengröße von Inselbiotopen auf deren Funktion als Trittstein oder Refugium. *Natur und Landschaft* **56(7/8)**: 235-242
- Mader H-J (1983)** Warum haben kleine Inselbiotope hohe Artenzahlen? *Natur und Landschaft* **58(10)**: 367-370
- Magura T, Ködöböcz V, Tóthmérés B (2001)** Effects of habitat fragmentation on carabids in forest patches. *Journal of Biogeography* **28**: 129-138
- Magura T, Tóthmérés B, Molnár T (2004)** Changes in carabid beetle assemblages along an urbanisation gradient in the city of Debrecen, Hungary. *Landscape Ecology* **19**: 747-759

- Magurran AE (1988)** *Ecological Diversity and its Measurement*. Princeton University Press – Princeton
- Markgraf A, Basedow T (2002)** Flight activity of predatory Staphylinidae in agriculture in central Germany. *Journal of Applied Entomology* **126**: 79-81
- Mascanzoni D, Wallin H (1986)** The harmonic radar: a new method of tracing insects in the field. *Ecological Entomology* **11**: 387-390
- McDonnell MJ, Pickett STA, Groffman P, Bohlen P, Poyat RV, Zipperer WC, Parmelee RW, Carreiro MM, Medley K (1997)** Ecosystem processes along an urban-to-rural gradient. *Urban Ecosystems* **1**: 21-36
- Miller MP (1997)** Tools for population genetic analyses (TFPGA) 1.3: a Windows program for the analysis of allozyme and molecular population genetic data. Computer software distributed by the author
- Miyashita T, Shinkai A, Chida T (1998)** The effects of forest fragmentation on web spider communities in urban areas. *Biological Conservation* **86**: 357-364
- Mols PJM (1983)** Simulation of the motivation and egg production of the carabid beetle *Pterostichus coerulescens* L.. In: Brandmayr P, Den Boer PJ, Weber F (Hrsg.) *The synthesis of field and laboratory experiments, Report of the 4th Meeting of European Carabidologists, 24-26 September 1981*. Druk Co Pudoc - Wageningen pp. 36-43
- Morales JM, Ellner SP (2002)** Scaling up animal movements in heterogeneous landscapes: the importance of behavior. *Ecology* **83(8)**: 2240-2247
- Moran PAP (1950)** Notes on continuous stochastic phenomena. *Biometrika* **37**: 17-23
- Mossakowski D, Schnauder C, Carius W, Klein B (1995)** Differenzen in der Enzym-Variabilität von Carabus-Arten nahe des früheren Eisrandes. *Mitteilungen der deutschen Gesellschaft für allgemeine und angewandte Entomologie* **9**: 787-792
- Mossakowski D, Stier J (1983)** Vergleichende Untersuchungen zur Laufgeschwindigkeit von Carabiden. In: Brandmayr P, Den Boer PJ, Weber F (Hrsg.) *The synthesis of field and laboratory experiments, Report of the 4th Meeting of European Carabidologists, 24-26 September 1981*. Druk Co Pudoc – Wageningen, pp. 19-33
- Mueller-Dombois D, Ellenberg H (1974)** *Aims and Methods of Vegetation Ecology*. John Wiley & Sons – New York, London, Sydney, Toronto
- Nei M, Muruyama T, Chakroborty R (1975)** The bottleneck effect and genetic variability in populations. *Evolution* **29**: 1-10
- Neumann U (1971)** Die Sukzession der Bodenfauna (Carabidae (Coleoptera), Diplopoda und Isopoda) in den forstlichen Rekultivierungen des rheinischen Braunkohlenreviers. *Pedobiologia* **11**: 193-226
- Niemelä J (1999)** Ecology and urban planning. *Biodiversity and Conservation* **8**: 119-131

- Niemelä J, Kotze DJ, Ashworth A, Brandmayr P, Desender K, New T, Penev L, Samways M, Spence J (2000)** The search for common anthropogenic impacts on biodiversity: a global network. *Journal of Insect Conservation* **4**: 3-9
- Niehues F-J, Hockmann P, Weber F (1996)** Genetics and dynamics of a *Carabus auronitens* metapopulation in the Westphalian Lowlands (Coleoptera, Carabidae). *Ann. Zool. Fennici* **33**: 85-96
- Niemelä J (2001)** Carabid beetles (Coleoptera: Carabidae) and habitat fragmentation: a review. *European Journal of Entomology* **98**: 127-132
- Niemelä J, Kotze DJ, Venn S, Penev L, Stoyanov I, Spence J, Hartley D, Montes de Oca E (2002)** Carabid beetle assemblages (Coleoptera, Carabidae) across urban-rural gradients: an international comparison. *Landscape Ecology* **17**: 387-401
- Oksanen J, Kindt R, O'Hara B (2005)** Vegan – Community Ecology Package. <http://cc.oulu.fi/~jarioksa/>
- Paine RT (1966)** Food web complexity and species diversity. *American Naturalist* **100**: 65-75
- Parker PC, Snow AA, Schug MD, Booton GC, Fuerst PA (1998)** What molecules can tell us about populations: Choosing and using a molecular marker. *Ecology* **79(2)**: 361-382
- Peakall R, Ruibal M, Lindenmayer DB (2003)** Spatial autocorrelation analysis offers new insights into gene flow in the Australian bush rat, *Rattus fuscipes*. *Evolution* **57**: 1182-1195
- Peakall R, Smouse PE (2006)** GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* **6**: 288-295
- Peterson MA (1995)** Phenological isolation, gene flow and developmental differences among low- and high-elevation populations of *Euphilotes enoptes* (Lepidoptera: Lycaenidae). *Evolution* **49**: 446-455
- Peterson MA (1996)** Long-distance gene flow in the sedentary butterfly, *Euphilotes enoptes* (Lepidoptera: Lycaenidae). *Evolution* **50**: 1990-1999
- Petit S (1994)** Diffusion of forest carabid beetles in hedgerow network landscapes. In: Desender K, Dufrene M, Loreau M, Luff ML, Maelfait JP (Hrsg.) *Carabid Beetles – Ecology and Evolution*. Kluwer Academic Publishers – Dordrecht, pp. 337-341
- Pharus (1921)** Pharus Plan Berlin (Grosse Ausgabe mit Vororten). Pharus Verlag - Berlin
- Pither J, Taylor PD (1998)** An experimental assessment of landscape connectivity. *Oikos* **83**: 166–174
- Platt RH, Rowntree RA, Muick PC (1994)** *The Ecological City: Preserving & Restoring Urban Biodiversity*. University of Massachusetts Press – Amherst, pp. 1-291
- Powell JR (1971)** Genetic polymorphisms in varied environments. *Science* **174**: 1035-1036
- Preussische Urmesstischblätter (1831-1871)** Handgezeichnete Karten in der Kartensammlung der Staatsbibliothek Berlin

- Qiagen (2006)** DNeasy Tissue System. http://www.ebiotrade.com/buyf/productsf/qiagen/DNeasy_tissue_system.htm
- R Development Core Team (2005)** R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>
- Read HJ, Martin MH, Rayner JMV (1998)** Invertebrates in Woodlands Polluted by Heavy Metals – An Evaluation Using Canonical Correspondence Analysis. *Water, Air, and Soil Pollution* **106**: 17–42
- Rebele F (1994)** Urban Ecology and Special Features of Urban Ecosystems. *Global Ecology and Biogeography Letters* **4(6)**: 173-187
- Rötzer T, Wittenzeller M, Haeckel H, Nekovar J (2000)** Phenology in central Europe - differences and trends of spring phenophases in urban and rural areas. *International Journal of Biometeorology* **44**: 60-67
- Reed DH, Frankham R (2001)** How closely related are molecular and quantitative measures of genetic diversity: a meta-analysis. *Evolution* **55**: 1095-1103
- Reed DH, Frankham R (2003)** Correlation between Fitness and Genetic Diversity. *Conservation Biology* **17(1)**: 230-237
- Ricketts TH (2001)** The matrix matters: effective isolation in fragmented landscapes. *The American Naturalist* **158**: 87-99
- Reike H-P (2004)** Untersuchungen zum Raum-Zeit-Muster epigäischer Carabidae an der Wald-Offenland-Grenze. *Contributions to Forest Science* **21**: 1-372
- Riecken U, Ries U (1992)** Untersuchung zur Raumnutzung von Laufkäfern (Col.: Carabidae) mittels Radio-Telemetrie. Methodenentwicklung und erste Freilandversuche. *Zeitschrift für Ökologie und Naturschutz* **1**: 147-149
- Riecken U, Raths U (1996)** Use of radio telemetry for studying dispersal and habitat use of *Carabus coriaceus* L.. *Annales Zoologici Fennici* **33**: 109-116
- Riecken U, Raths U (2000)** Radio-telemetrische Untersuchungen zum Raum-Zeit-Verhalten von Laufkäfern am Beispiel von Laufkäfern am Beispiel von *Carabus coriaceus* Linné, 1758 und *C.monilis* Fabricius, 1792. *Angewandte Carabidologie* **2/3**: 49-58
- Rietze J (2001)** Zur Phänologie ausgewählter Carabiden in baden-württembergischen Wäldern. *Angewandte Carabidologie Suppl.* **2**: 105-115
- Rijnsdorp AD (1980)** Pattern of Movement in and Dispersal from a Dutch Forest of *Carabus nemoralis* Hbst. (Coleoptera, Carabidae). *Oecologia* **45**: 274-281
- Rosenberg DK, Noon BR, Meslow EC (1997)** Biological corridors: form, function, and efficacy. *Bioscience* **47**: 677-687
- Rosenberg DK, Noon BR, Megahan JW, Meslow EC (1998)** Compensatory behaviour of *Ensatina eschscholtzii* in biological corridors: a field experiment. *Canadian Journal of Zoology* **76**: 117-133

- Roslin T (2001)** Spatial population structure in a patchily distributed beetle. *Molecular Ecology* **10**: 823-837
- Rousset F (1997)** Genetic differentiation and estimation of gene flow from *F*-statistics under isolation by distance. *Genetics* **145**: 1219-1228
- Rudd WG, Gandour RW (1985)** Diffusion model for insect dispersal. *Journal of Economic Entomology* **78**: 295-301
- Sachs L (1997)** *Angewandte Statistik – Anwendung statistischer Methoden*. Springer – Berlin, Heidelberg, New York
- Sadler JP, Small EC, Fiszpan H, Telfer G, Niemelä J (2006)** Investigating environmental variation and landscape characteristics of an urban–rural gradient using woodland carabid assemblages. *Journal of Biogeography* **33**: 1126-1138
- Samietz J, Berger U (1997)** Evaluation of movement parameters in insects – bias and robustness with regard to resight numbers. *Oecologia* **110**: 40-49
- Saunders DA, Hobbs RJ (1991)** The role of corridors in nature conservation: what do we know and where do we go? In: Saunders DA, Hobbs RJ (Hrsg.) *Nature Conservation*. Surrey Beatty and Sons – Chipping Norton Australia, pp. 408-409
- Schaefer M, Kock K (1979)** Zur Ökologie der Arthropodenfauna einer Stadtlandschaft und ihrer Umgebung. I. Laufkäfer (Carabidae) und Spinnen (Araneida). *Anzeiger für Schädlingskunde, Pflanzenschutz, Umweltschutz* **52**: 85-90
- Schlötterer C, Ritter R, Harr B, Brem G (1998)** High mutation rate of a long microsatellite allele in *Drosophila melanogaster* provides evidence for allele-specific mutation rates. *Molecular Biology and Evolution* **15**: 1269-1274
- Schmettau FWC (1767-1787)** Landaufnahme Section 1-121 Pommern, Mark, Netzedistrikt, Magdeburg, Altmark, Braunschweig. Handgezeichnete Karten in der Kartensammlung der Staatsbibliothek Berlin
- Schneider S, Roessli D, Exoffier C (2000)** ARLEQUIN, Version 2.000: a Software for Population Genetics Data Analysis, Genetics and Biometry Laboratory, University of Geneva, Switzerland, <http://lgb.unige.ch/arlequin>
- Schultz CB (1998)** Dispersal behavior and its implications for reserve design in a rare oregon butterfly. *Conservation Biology* **12**: 284-292
- Segebade R, Schaefer M (1979)** Zur Ökologie der Arthropoden Fauna einer Stadtlandschaft und ihrer Umgebung. II. Pflanzengallen und Pflanzenminen. *Anzeiger für Schädlingskunde, Pflanzenschutz, Umweltschutz* **52**: 117-214
- Shochat E, Stefanov WL, Whitehouse MEA, Faeth SH (2004)** Urbanization and Spider Diversity: Influences of Human Modification of Habitat Structure and Productivity. *Ecological Applications* **14**: 268-280

- Simberloff D, Farr JA, Cox J, Mehlman DW (1992)** Movement corridors: conservation bargains or poor investments? *Conservation Biology* **6**: 493-504
- Slatkin M (1985)** Gene flow in natural populations. *Annual Review of Ecology and Systematics* **16**: 393-430
- Slatkin M, Excoffier L (1996)** Testing for linkage disequilibrium in genotypic data using the EM algorithm. *Heredity* **76**: 377-383
- Smith TB, Wayne RK (1996)** *Molecular genetic approaches in conservation*. Oxford University Press – New York, Oxford
- Smouse PE, Peakall R (1999)** Spatial autocorrelation analysis of multi-allele and multi-locus genetic micro-structure. *Heredity* **82**: 561-573
- Soulé M E, Alberts AC, Bolger DT (1992)** The effects of habitat fragmentation on chaparral plants and vertebrates. *Oikos* **63**: 39–47
- StatSoft (2001)** STATISTICA for Windows 6.0. Tulsa, Oklahoma
- Steiniger H (1979)** Populationsgenetische Adaptionsstrategien von Carabiden in einem urbanen Ökosystem (Coleoptera). *Verhandlungen der Gesellschaft für Ökologie* **7**: 475-481
- Stow AJ, Sunnucks P, Briscoe DA, Gardner MG (2001)** The impact of habitat fragmentation on dispersal of Cunningham's skink (*Egernia cunninghami*): evidence from allelic and genotypic analysis of microsatellites. *Molecular Ecology* **10(4)**: 867-878
- Szyszko J, Den Boer PJ, Bauer T (2002)** *How to protect or what we know about Carabid Beetles*. Warsaw Agricultural University Press – Warsaw
- Taberlet P, Fumigalli L, Wustsaucy AG, Cosson JF (1998)** Comparative phylogeography and postglacial colonization routes in Europe. *Molecular Ecology* **7**: 453-464
- Taylor PD, Fahrig L, Henein K, Merriam G (1993)** Connectivity is a vital element of landscape structure. *Oikos* **68(3)**: 571–573
- Terlutter H (1990)** An Allele Gradient of an Esterase Gene Locus as a Result of Recent Gene Flow: Electrophoretic Investigations of *Carabus auronitens* F. (Col. Carabidae). In: Stork NE (Hrsg.) *The Role of Ground Beetles in Ecology and Environmental Studies*. Intercept – Andover, Hampshire, pp. 359-364
- Thiele H-U (1977)** *Carabid beetles in their environments*. Springer – Berlin, Heidelberg, New York
- Tischendorf L, Wissel C (1997)** Corridors as conduits for small animals: attainable distances depending on movement pattern, boundary reaction and corridor width. *Oikos* **79**: 603-611
- Tonteri T, Haila Y (1990)** Plants in a boreal city: ecological characteristics of vegetation in Helsinki and its surroundings, southern Finland. *Annales Botanici Fennici* **27**: 337-352
- TOP50 (1999)** Amtliche topographische Karten auf CD-ROM. Eine Serie der Deutschen Landesvermessung, Version 2
- Topp W (1972)** Die Besiedelung eines Stadtparks durch Käfer. *Pedobiologia* **12**: 336-346

- Trautner J (1991)** Die Laufkäferfauna des Rosensteinparks und weiterer Grünflächen im Stadtgebiet von Stuttgart (Coleoptera, Carabidae). *Jahreshefte der Gesellschaft für Naturkunde in Württemberg* **146**: 233-258
- Tscharntke T (1992)** Fragmentation of *Phragmites* habitats, minimum viable population size, habitat suitability, and local extinctions of moths, midges, flies, aphids, and birds. *Conservation Biology* **6**: 530-536
- Tscharntke T, Brandl R (2004)** Plant-Insect Interactions in Fragmented Landscapes. *Annual Review of Entomology* **49**: 405-430
- Turin H (2000)** *De Nederlandse Loopkevers, verspreiding en oecologie (Coleoptera: Carabidae)*. – *Nederlandse Fauna 3, Nationaal Natuurhistorisch Museum Naturalis*. KNNV Uitgeverij & EIS-Nederland – Leiden
- UEIS (2005)** Berlin Urban and Environmental Information System. http://www.stadtentwicklung.berlin.de/umwelt/umweltatlas/edua_index.shtml
- Usher MB, Field J, Bedford S (1993)** Biogeography and diversity of ground-dwelling arthropods in farm woodlands. *Biodiversity Letters* **1**: 54-62
- Van der Wurff, AWG, Gols R, Ernsting G, Van Straalen NM (2005)** Population genetic structure of *Orchesella cincta* (Collembola; Hexapoda) in NW Europe, as revealed by microsatellite markers. *Pedobiologia* **49**: 167-174
- Van Dongen S, Backeljau T, Matthysen E, Dhondt AA (1998)** Genetic population structure of the winter moth (*Operophtera brumata* L.) (Lepidoptera, Geometridae) in a fragmented landscape. *Heredity* **80**: 92-100
- Vermeulen HJW (1994)** Corridor function of a road verge for dispersal of stenotopic heathland ground beetles Carabidae. *Biological Conservation* **69**: 339-349
- Vernon P, Vannier G (2002)** Evolution of freezing susceptibility and freezing tolerance in terrestrial arthropods. *Comptes Rendus Biologies* **325 (12)**: 1185-1190
- Wallin H (1986)** Habitat choice of some field inhabiting carabid beetles (Coleoptera: Carabidae) studied by recapture of marked individuals. *Ecological Entomology* **11**: 457-466
- Wallin H, Ekbom B (1988)** Movements of carabid beetles inhabiting cereal fields: a field tracing study. *Oecologia* **77**: 39-43
- Wallin H, Ekbom B (1994)** Influence of Hunger Level and Prey Densities on Movement Patterns in Three Species of Pterostichus Beetles (Coleoptera: Carabidae). *Environmental Ecology* **23(5)**: 1171-1181
- Walter R (1992)** *Geologie von Mitteleuropa*. 5. Aufl. Schweizerbart'sche Verlagsbuchhandlung - Stuttgart
- Wang M, Schreiber A (1999)** Population genetics of the woodlouse *Porcellio scaber* Latr. (Isopoda: Oniscoidea) in central Europe: passive dispersal and postglacial range expansion. *Canadian Journal Zoology* **77**: 1337-1370

- Warnaffe GB, Dufrêne M (2004)** To what extent can management variables explain species assemblages. A study of carabid beetles in forests. *Ecography* **27**: 701-714
- Watts PC, Rouquette R, Saccheri J, Kemp SJ, Thompson DJ (2004)** Molecular and ecological evidence for small-scale isolation by distance in an endangered damselfly, *Coenagrion mercuriale*. *Molecular Ecology* **13**: 2931-2945
- Weigmann G (1982)** The colonization of ruderal biotopes in the city of Berlin by arthropods. In: Bornkamm R, Lee JA, Seaward MRD (Hrsg.) *Urban Ecology*. Blackwell – Oxford, pp. 75-82
- Weir BS, Cockerham CC (1984)** Estimating F-statistics for the analysis of population structure. *Evolution* **38**: 1358-1370
- Weißpflug H (1999)** Das Landeswaldgesetz wird erlassen. *Berlinische Monatsschrift* **8(1)**: 47-49
- Weller B, Ganzhorn JU (2004)** Carabid beetle community composition, body size, and fluctuating asymmetry along an urban-rural gradient. *Basic and Applied Ecology* **5**: 193-201
- Welling M (1990)** Dispersal of ground beetles (Col., Carabidae) in arable land. *Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent* **55(2b)**: 483-491
- Wendler G (1999)** Fortbewegung und sensomotorische Integration. In: Dettner K, Peters W (Hrsg.) *Lehrbuch der Entomologie*. Gustav Fischer Verlag - Stuttgart, Jena, Lübeck, Ulm
- White MA, Nemani RR, Thornton PE, Running SW (2002)** Satellite evidence of phenological differences between urbanized and rural areas of the eastern United State deciduous broadleaf forest. *Ecosystems* **5**: 260-277
- Wiens JA, Crist TO, Milne BT (1993)** On Quantifying Insect Movements. *Environmental Entomology* **22(4)**: 709-715
- Wiens JA, Milne BT (1989)** Scaling of 'landscapes' in landscape ecology, or, landscape ecology from a beetle's perspective. *Landscape Ecology* **3(2)**: 87-96
- Wilcove DS, McLellan CH, Dobson AP (1986)** Habitat fragmentation in the temperate zone. In: Soule M (Hrsg.) *Conservation Biology*. Sinauer – Sunderland, Massachusetts, pp. 237-256
- Williamson P, Evans PR (1973)** A preliminary study of the effects of high levels of inorganic lead on soil fauna. *Pedobiologia* **13**: 16-21
- Wilson EO, Willis EO (1975)** Applied biogeography. In: Cody ML, Diamond JM (Hrsg.) *Ecology and evolution of communities*. Belknap – Cambridge, Massachusetts, pp. 522-534
- Wood BC, Pullin AS (2002)** Persistence of species in a fragmented urban landscape: The importance of dispersal ability and habitat availability for grassland butterflies. *Biodiversity and Conservation* **11**: 1451-1468
- Wright S (1951)** The genetical structure of populations. *Annals of Eugenics* **15**: 323-354

-
- Young A, Boyle T, Brown T (1996)** The population genetic consequences of habitat fragmentation for plants. *Trends in Ecology and Evolution* **11**: 413-418
- Zar JH (1999)** *Biostatistical Analysis*. Prentice Hall – Upper Saddle River
- Zerbe S (1989)** Testing of movement pattern, specific to species, along corridors taking ground beetles (Carabidae) and locusts (Acrididae) as an example. *Landschaft und Stadt* **21(3)**: 100-103