

7. Literaturverzeichnis

1. Adams DH, Shaw S. Leucocyte-endothelial interactions and regulation of leucocyte migration. *Lancet* 1994; 343: 831-836
2. Allen JS, Eisma R, La Freniere D. Characterization of the eosinophil chemokine RANTES in nasal polyps. *Ann Otol Rhinol Laryngol* 1998; 107(5): 416-420
3. Asadullah K, Sabat R, Friedrich M, Volk HD, Sterry W. Interleukin-10: an important immunoregulatory cytokine with major impact on psoriasis. *Curr Drug Targets Inflamm Allergy* 2004; 3(2): 185-192
4. Asero R, Bottazzi G. Nasal polyposis: a study of its association with airborne allergen hypersensitivity. *Ann Allergy Asthma Immunol* 2001; 86(3): 283-285
5. Atsuta S, Majima Y. Nasal mucociliary clearance of chronic sinusitis in relation to rheologic properties of nasal mucos. *Ann Otol Rhinol Laryngol* 2000; 107: 47-51
6. Bachert C. Die Schleimhaut der oberen Atemwege- zur Pathophysiologie der Entzündung. *Eur Arch Oto-Rhino-Laryngology* 1995; Suppl. 1: 156-206
7. Bachert C, Hauser U, Prem B, Rudack C. IL-5 synthesis is upregulated in human nasal polyp tissue. *J Allergy Clin Immunol* 1997; 99: 837-842
8. Bachert C, Wagenmann M, Rudack C, van Cauwenberge P. The role of cytokines in infectious sinusitis and nasal polyposis. *Allergy* 1998; 53: 2-13
9. Bachert C, Gevaert P, Holtappels G, van Cauwenberge P. Total and specific IgE in nasal polyps is related to local eosinophilic inflammation. *J Allergy Clin Immunol* 2001; 107(4): 607-614
10. Bachert C; Hörmann K, Mösges R, Rasp G, Riechelmann H, Müller R, Luckhaupt H, Stuck BA, Rudack C. An update on the diagnosis and treatment of sinusitis and nasal polyposis. *Allergy* 2003; 58(3):176-191
11. Baggolini M. Chemokines and leukocyte traffic. *Nature* 1998; 392: 565-568
12. Bartels J, Schluter C, Richter E, Christophers E, Schröder JM. Human dermal fibroblasts express eotaxin: molecular cloning, mRNA expression, and identification of eotaxin sequence variants. *Biochem Biophys Res Commun* 1996; 225(3): 1045-1051
13. Bartels J, Maune S, Meyer JE, et al. Increased eotaxin-mRNA expression in non-atopic and atopic nasal polyps: comparison to RANTES and MCP-3 expression. *Rhinology* 1997; 35(4): 171-174
14. Bergler W, Adam S, Gross HJ. Age-dependent altered proportions in subpopulations of tonsillar lymphocytes. *Clin Exp Immunol* 1999; 116(1): 9-18.
15. Berkout TA, Sarau HM, Moores K et al. Cloning, in vitro expression, and functional characterization of a novel human CC chemokine of the monocyte chemotactic protein (MCP) family (MCP-4) that binds and signals through the CC chemokine receptor 2B. *J Biol Chem* 1997; 272(26): 16404-16413
16. Berlin C, Berg EL, Briskin MJ, Butcher EC. Alpha 4 beta 7 integrin mediates lymphocyte binding to the mucosal vascular addressin MAdCAM-1. *Cell* 1993; 74(1): 185-195
17. Bertrand CP, Ponath PD. CCR3 blockade as a new therapie for asthma. *Exp Opin Invest Drugs* 2000; 9: 43-52
18. Bocchino V, Bertorelli G, Bertrand CP, Ponath PD, Olivieri D. Eotaxin and CCR3 are up-regulated in exacerbations of chronic bronchitis. *Allergy* 2002; 57(1): 17-22
19. Bolger WE, Butzin CA., Parsons DS. Paranasal sinus bony anatomic variations and mucosal abnormalities: CT analysis for endoscopic sinus surgery. *Laryngoscope* 1991; 101: 56-64

20. Brandtzaeg P. Immune functions and immunopathology of palatine and nasopharyngeal tonsils. In: Bernstein JM, Ogra P (editors). Immunology of the ear. New York: Raven Press, 1987: 63-106.
21. Brandtzaeg P, Halstensen TS. Immunology and immunopathology of tonsils. *Adv Otorhinolaryngol* 1992; 47: 64-75
22. Brandtzaeg P. The B-cell development in tonsillar lymphoid follicles. *Acta Otolaryngol* 1996; 523: 55-59
23. Brandtzaeg P, Farstad IN, Haraldsen. Regional specialization in the mucosal immune system: primed cells do not always home along the same track. *Immunol Today* 1999; 20(6): 267-277
24. Brandtzaeg P. Immunology of tonsils and adenoids: everything the ENT surgeon needs to know. *Int J Pediatr Otorhinolaryngol* 2003; 67 Suppl 1: 69-76
25. Brown DC, Gatter KC. Ki67 protein: the immaculate deception? *Histopathology* 2002; 40(1): 2-11.
26. Brustein M, Kraal G, Mebius RE, Watson SR. Identification of a soluble form of a ligand for the lymphocyte homing receptor. *J Exp Med* 1992; 176(5): 1415-1419
27. Caplin, I, Haynes TJ, Spann J. Are nasal polyps an allergic phenomenon? *Ann Allergy* 1971; 29: 631 -634
28. Cassatella MA, Meda L, Bonora S, Constantin G. Interleukin 10 (IL-10) inhibits the release of proinflammatory cytokines from human polymorphonuclear leukocytes. Evidence for an autocrine role of tumor necrosis factor and IL-1 beta in mediating the production of IL-8 triggered by lipopolysaccharide. *J Exp Med* 1993; 178(6): 2207-2211
29. Caversaccio M, Hartnell A, Calnan D et al. The role of chemokines in nasal polyps. *Schweiz Med Wochenschr* 2000; Suppl 125 :92-95
30. Caversaccio M, Uggolini M, Hartnell A, Conroy D. Chronic immunological rhinosinusitis: general aspects, cytokines, chemokines and possible therapeutic consequences. *Otorhinolaryngol Nova* 2003;12(2): 52-62
31. Chensue SW, Warmington K, Ruth JH et al. Mycobacterial and schistosomal antigen-elicited granuloma formation in IFN-gamma and IL-4 knockout mice: analysis of local and regional cytokine and chemokine networks. *J Immunol* 1997; 159(7): 3565-3573
32. Chomarat P, Rissoan MC, Banchereau J, Miossec P. Interferon gamma inhibits interleukin 10 production by monocytes. *J Exp Med* 1993; 177(2): 523-527
33. Chomczynski P, Sacchi N. Single step method of RNA isolation by guanidine thiocyanate-phenol-chloroform extraction. *Anal Biochem* 1987; 162: 156-159
34. Costa JJ, Matossian K, Resnick MB, Beil WJ, Wong DT, Gordon JR, Dvorak AM. Human eosinophils can express the cytokines tumor necrosis factor-alpha and macrophage inflammatory protein-1 alpha. *J Clin Invest* 1993; 91 (6): 2673-2684
35. Coste A., Rateau JG, Escudier E. Increased epithelial cell proliferation in nasal polyps. *Arch Otolaryngol Head Neck Surg* 1996; 122: 432-436
36. Coyle AJ, Ackerman SJ, Burch R, Irvin CG. Human eosinophil-granule major basic protein and synthetic polycations induce airway hyperresponsiveness in vivo dependent on bradykinin generation. *J Clin Invest* 1995; 95(4): 1735-1740
37. Cunningham JM, Chiu EJ, Landgraf JM, Gliklich RE. The health impact of chronic recurrent rhinosinusitis in children. *Arch Otolaryngol. Head Neck Surg* 2000; 126: 1363-1368
38. del Pozo MA, Sanchez- Mateos P, Sanchez F. Cellular polarization induced by chemokines: a mechanism for leucocyte recruitment. *Immunol Today* 1996; 17: 127-131

39. DeMaeyer E, DeMaeyer-Guipnard J. Interferons. In: Thomson AW (editor). *The cytokine handbook*. London: Academic Press, 1994: 265-288
40. Drake-Lee AB. Histamine and its release from nasal polyps: preliminary communication. *J Roy Soc Med* 1984; 77: 120-124
41. Dulkys Y, Schramm G, Kimmig D et al. Detection of mRNA for eotaxin-2 and eotaxin-3 in human dermal fibroblasts and their distinct activation profile on human eosinophils. *J Invest Dermatol* 2001; 116(4): 498-505
42. Ebisawa M, Bochner BS, Georas SN, Schleimer RP. Eosinophil transendothelial migration induced by cytokines. I. Role of endothelial and eosinophil adhesion molecules in IL-1 beta-induced transendothelial migration. *J Immunol* 1992; 149 (12): 4021-4028
43. Ebisawa M, Liu MC, Yamada T, Kato M, Lichtenstein LM, Bochner BS, Schleimer RP. Eosinophil transendothelial migration induced by cytokines. II. Potentiation of eosinophil transendothelial migration by eosinophil-active cytokines. *J Immunol* 1994; 152 (9): 4590-4596
44. Ebisawa M, Yamada T, Bickel C, Klunk D, Schleimer RP. Eosinophil transendothelial migration induced by cytokines. III. Effect of the chemokine RANTES. *J Immunol* 1994; 153 (5): 2153-2160
45. Elsner J, Kapp A. The chemokine network in eosinophil activation. *Allergy Asthma Proc* 2001; 22(3): 139-148
46. Elsner J, Kapp A. Aktivierung und Modulation eosinophiler Granulozyten durch Chemokine:Ihre Bedeutung für die allergische Entzündung. *Allergologie* 2001; 24; 9: 368-389
47. Elsner J, Kapp A. Eosinophile Granulozyten im Netzwerk der Chemokine. *Allergologie* 2000; 23: 59-72
48. Elsner J, Petering H, Kimmig D, Wells TN, Kapp A. The CC chemokine receptor antagonist Met-RANTES inhibits eosinophil effector functions. *Int Arch Allergy Immunol* 1999; 1118: 462-465
49. Elsner J, Oppermann M, Czech W. C3a activates the respiratory burst in human polymorphonuclear neutrophilic leukocytes via pertussis toxin-sensitive G-proteins. *Blood* 1994; 83(11): 3324-3331
50. Fokkens W, Holm A, Rijntjes E, Mulder P, Vroom T. Characterization and quantification of cellular infiltrates in nasal mucosa of patients with grass pollen allergy, non-allergic patients with nasal polyps and controls. *Int Arch Allergy Immunol* 1990; 93: 66-72
51. Forssmann U, Uguccioni P, Loetscher CA, Dahinden H, Baggioini M. Eotaxin-2, a novel CC- chemokine that is selective for the chemokine receptor CCR3, and acts like eotaxin on human eosinophil and basophil leukocytes. *J Exp Med* 1997; 185: 2171-2176
52. Fuchs E, Gronemeyer W. Pollinosis (Heufieber-Pollenallergie). In: Fuchs E., Schulz KH (Hrsg). *Manuale allergologicum*. Dustri-Verlag, München-Deisenhofen 1988: 1 -22
53. Fujisawa T, Kato Y, Atsuta J et al. Chemokine production by the BEAS-2B human bronchial epithelial cells: differential regulation of eotaxin, IL-8, and RANTES by TH2- and TH1-derived cytokines. *J Allergy Clin Immunol* 2000; 105(1): 126-133
54. Gebert A. The role of M cells in the protection of mucosal membranes. *Histochem Cell Biol* 1997; 108: 455-470
55. Gerber BO, Zanni MP, Uguccioni M, Yawalkar N, Maggiolini M, Moser B. Functional expression of the eotaxin receptor CCR3 in T lymphocytes co-localizing with eosinophils. *Curr Biol* 1997; 7(11): 836-843
56. Giampiero P, Paolo B, Eleonora N, Domenico S, Giuseppe T, Giuseppa F, Rita PL, Giovanna F. Intranasal treatment with lysine acetylsalicylate in patients with nasal polyposis. *Ann Allergy* 1991; 67: 588 -592

57. Hamaguchi Y, Suzumura H, Arima S, Sakakura Y. Quantitation and immunocytological identification of interleukin-1 in nasal polyps from patients with chronic sinusitis. *Int Arch Allergy Immunol* 1994; 104 (2): 155-159
58. Hamilos DL, Leung DY, Wood R, Meyers A, Stephens JK, Barkans J, Meng Q, Cunningham L, Bean DK, Kay AB. Chronic hyperplastic sinusitis: association of tissue eosinophilia with mRNA expression of granulocyte-macrophage colony-stimulating factor and interleukin-3. *J Allergy Clin Immunol* 1993; 92: 39-48
59. Hamilos DL, Leung DY, Wood R, Cunningham L, Bean DK, Yasrue Z, Schotman E, Hamid Q. Evidence for distinct cytokine expression in allergic versus nonallergic chronic sinusitis. *J Allergy Clin Immunol* 1995; 96: 537-544
60. Hamilos DL, Leung DY, Wood R, Bean DK, Song YL, Schotman E, Hamid Q. Eosinophil infiltration in nonallergic chronic hyperplastic sinusitis with nasal polyposis is associated with endothelial VCAM-1 upregulation and expression of TNF-alpha. *Amer J Respir Cell Mol Biol* 1996; 15: 443-450
61. Hamilos DL, Leung DY, Huston DP et al. GM-CSF, IL-5 and RANTES immunoreactivity and mRNA expression in chronic hyperplastic sinusitis with nasal polyposis (NP). *Clin Exp Allergy* 1998; 28(9): 1145-1152
62. Hellquist HB. Nasal polyps update. *Histopathology*. *Allergy Asthma Proc* 1996; 17(5): 237-242
63. Hilka MB, Koch T, Laszig R. Late results of endonasal ethmoid bone operation with special reference of polypous sinusitis. *HNO* 1992; 40: 165-169
64. Hirschberg VG. Anaphylactoid reaction to aspirin (1902) (classical article). *Allergy Proc* 1990; 11: 249-252
65. Horuk R, Ng HP. Chemokine receptor antagonists. *Med Res Rev* 2000; 20: 155-168
66. Humbert M, Ying S, Corrigan C et al. Bronchial mucosal expression of the genes encoding chemokines RANTES and MCP-3 in symptomatic atopic and nonatopic asthmatics: relationship to the eosinophil-active cytokines interleukin IL-5, granulocyte macrophage-colony-stimulating factor and IL-3. *Am J Respir Cell Mol Biol* 1997; 16(1): 1-8
67. Inoue H, Fukuizumi T, Tsujisawa T, Uchiyama C. Simultaneous induction of specific IgA-producing cells in major and minor salivary glands after tonsillar application of antigen. *Oral Microbiol Immunol* 1999; 14(1): 21-26.
68. Ishida K, Zhu BL, Maeda H. Novel approach to quantitative reverse transcription PCR assay of mRNA component in autopsy material using the TaqMan fluorogenic detection system: dynamics of pulmonary surfactant apoprotein. *A. Forensic Sci Int* 2000; 113(1-3): 127-131
69. Jahnson FL, Haye R, Gran E, Brandtzaeg P, Johansen FE. Glucocorticosteroids inhibit mRNA expression for eotaxin, eotaxin-2, and monocyte-chemotactic protein-4 in human airway inflammation with eosinophilia. *J Immunol* 1999; 163(3): 1545-1551
70. Jose PJ, Griffits-Johnson DA, Collins PD, Walsh DT, Moqbel K, Totty NF, Truong O, Hsuan JJ, Williams TJ. Eotaxin: a potent eosinophil chemoattractant cytokine detected in a guinea pig model of allergic airways inflammation. *J Exp Med* 1993; 179: 881-887
71. Kaldenbach T, Schäfer D, Gosepath J, Bittinger F, Klimek L, Mann WJ. Die Bedeutung eosinophiler Granulozyten in Beziehung zu Allergie und Aspirin-Intoleranz bei Patienten mit Sinusitis polyposa. *Laryngo-Rhino-Otol* 1999; 78: 429-433
72. Kaliner M. Treatment of sinusitis in the next millennium. *Allergy Asthma Proc* 1998; 19(4): 181-184
73. Kaliner M. Medical management of sinusitis. *Am J Med Sci* 1998; 316: 21-28
74. Kaplan AP. Chemokines, chemokine receptors and allergy. *Int Arch Allergy Immunol* 2001; 124(4): 423-431

75. Kapp A, Zeck-Kapp G, Czech W, Schöpf E. The chemokine RANTES is more than a chemoattractant: characterization of its effect on human eosinophil oxidative metabolism and morphology in comparison with IL-5 and GM-CSF. *J Invest Dermatol* 1994; 102: 906-914
76. Kasama T, Strieter RM, Lukacs NW, Kunkel SL. Regulation of neutrophil-derived chemokine expression by IL-10. *J Immunol* 1994; 152(7): 3559-3569
77. Kim JT, Schimming AW, Kita H. Ligation of FcRII (CD32) pivotally regulates survival of human eosinophils. *J Immunol* 1999; 162: 4253-4259
78. Kim YK, Uno M, Hamilos DL, Beck L, Bochner B, Schleimer R. Immunolocalization of CD34 in nasal polyposis. Effect of topical corticosteroids. *Am J Respir Cell Mol Biol* 1999; 20(3): 388-397
79. Kitaura M, Nakajama T, Imai T, Harada S, Combadiere C, Tiffany HL, Murphy PM, Yoshie O. Molecular cloning of human eotaxin, an eosinophil-specific CC chemokine, and identification of a specific receptor, CC chemokine receptor 3. *J Biol Chem* 1996; 271: 7725-7730
80. Kitaura M, Suzuki N, Imai T, et al. Molecular cloning of a novel human CC chemokine (Eotaxin-3) that is a functional ligand of CC chemokine receptor 3. *J Biol Chem* 1999; 274(39): 27975-27980
81. Klapan I, Culo F, Culig J, Miljenko V. Arachidonic acid metabolites and sinonasal polyposis. I. Possible prognostic value. *Am J Otolaryngol* 1995; 16(6): 396-402
82. Korsrud FR, Brandtzaeg P. Immune systems of human nasopharyngeal and palatine tonsils: histomorphometry of lymphoid components and quantification of immunoglobulin-producing cells in health and disease. *Clin Exp Immunol* 1980; 39(2): 361-370
83. Korsrud FR, Brandtzaeg P. Influence of tonsillar disease on the expression of J chain by immunoglobulin-producing cells in human palatine and nasopharyngeal tonsils. *Scand J Immunol* 1981; 13(3): 281-287
84. Lamkhioued B, Abdelilah SG, Hamid Q, Renzi MP. The CCR3 receptor is involved in eosinophil differentiation and is up-regulated by Th2 cytokines in CD34+ progenitor cells. *J Immunol* 2003; 170(1): 537-47
85. Larsen P, Tos M. Origin of nasal polyps. *Laryngoscope* 1991; 101: 305-312
86. Larsen PL, Tingsgaard PK, Harcourt J, Tos M. Nasal polyps and their relation to polyps/hypertrophic polypoid mucosa in the paranasal sinuses: a macro-, endo-, and microscopic study of autopsy materials. *Am J Rhinol* 1998; 12(1): 45-51
87. Lee CH, Rhee CS, Min YG. Cytokine gene expression in nasal polyps. *Ann Otol Rhinol Laryngol* 1998; 107(8): 665-670
88. Leutenegger CM, Mislin CN, Sigrist B, et al. Quantitative real-time PCR for the measurement of feline cytokine mRNA. *Vet Immunol Immunopathol* 1999; 71(3-4): 291-305
89. Liu Y, Hamaguchi Y, Taya M, Sakakura Y. Quantification of interleukin-1 in nasal polyps from patients with chronic sinusitis. *Eur Arch Otorhinolaryngol* 1993; 250: 123-125
90. Liu C., Shun C, Hsu M. Lymphocyte subsets and antigen-specific IgE antibody in nasal polyps. *Ann Allergy* 1994; 72: 19-24
91. Luger EO, Achatz G, Meco C, Radbruch A, Olze H. Differential expression of IgA-positive plasmacells in palatine tonsils of Betv1-allergic patients and healthy individuals. *Clin Exp Allergy Int* 2004, submitted
92. Luger EO, Achatz G, Zemlin M, Olze H, Meco C, Krenn V, Burkhardt H, Radbruch A, Berek C. Allergen-specific IgG and IgA memory B cells in healthy and allergic individuals. *Man. in prep.*
93. Lund VJ, Mackay IS. Staging in rhinosinusitis. *Rhinology* 1993; 31: 183-184
94. Lund VJ, Kennedy DW. Quantification for staging sinusitis. *Ann. Otol. Rhinol. Laryngol.* 1995; 167(Suppl.): 17-21

95. Luster AD. Chemokines-chemotactic cytokines that mediate inflammation. *N Engl J Med* 1998; 338(7): 436-445
96. Luttmann W, Opfer A, Dauer E, Foerster M, Matthys H, Virchow JC. Differential regulation of CD95 (Fas/APO-1) expression in human blood eosinophils. *Eur J Immunol* 1998; 28(7): 2057-2065
97. MacLennan IC. Germinal centers. *Annu Rev Immunol* 1994; 12: 117-139
98. MacLennan IC, Gulbranson-Judge A, Toellner KM. The changing preference of T and B cells for partners as T-dependent antibody responses develop. *Immunol Rev* 1997; 156: 53-66
99. Matthews BL, Burke AJ. Recirculation of mucus via accessory ostia causing chronic maxillary sinus disease. *Otolaryngol Head Neck Surg* 1997; 117: 422-423
100. Maune S, Rudert H, Heissenberg C, Kuechler T. Concepts for measuring quality of life in patients with chronic sinusitis. *Laryngorhinootologie* 1999; 78: 475-480
101. Maune S, Meyer JE, Pods R, Bartels J, Schröder JM. MCP-4 Genexpression in nasaler Schleimhaut. *Allergologie* 1999; 10: 606-610
102. Maune S; Meyer JE. The role of CC chemokines in chronic sinusitis. *Otorhinolaryngol Nova* 2003; 12(2): 63-70
103. McKenzie AN, Sanderson CJ. Interleukin 5. *Chem Immunol* 1992; 51: 135-152
104. Middleton E, Reed CE, Ellis EF, Jr. Adkinson FR, Yunginger JW, Busse WW. Allergy principles and practice. 4 St. Louis Mosby 1993 (nach Kaldenbach 1999)
105. Min YG, Lee CH, Rhee CS. Inflammatory cytokine expression on nasal polyps developed in allergic and infectious rhinitis. *Acta Otolaryngol* 1997; 117(2): 302-306
106. Minshall EM, Cameron L, Lavigne F et al. Eotaxin mRNA and protein expression in chronic sinusitis and allergen-induced nasal responses in seasonal allergic rhinitis. *Am J Respir Cell Mol Biol* 1997; 17(6): 683-690
107. Miyamasu M, Misaki Y, Yamaguchi M, Yamamoto K, Morita Y, Matsushima K, Nakajima T, Hirai K. Regulation of human eotaxin generation by Th1/Th2-derived cytokines. *Int Arch Allergy Immunol* 2000; 122 Suppl 1: 54-58
108. Mochizuki M, Bartels J, Mallet AI, Christophers E, Schroeder JM. IL-4 induces eotaxin: a possible mechanism of selective eosinophil recruitment in helminth infection and atopy. *J Immunol* 1998; 160(1): 60-68
109. Moneret-Vautrin DA, Hsieh V, Wayoff W, Guyot JL, Mouton C, Maria Y. Nonallergic rhinitis with eosinophilia syndrome a precursor of the triad: nasal polyposis, intrinsic asthma, and intolerance to aspirin. *Ann Allergy* 1990; 64: 513-518
110. Mosmann TR, Coffman RL. TH1 and TH2 cells. Different pattern of lymphokine secretion lead to different functional properties. *Ann Rev Immunol* 1989; 7: 145-173
111. Mosmann TR, Moore KW. The role of IL-10 in crossregulation of T(H)1 and T(H)2 responses. *Parasitol Today* 1991; 7(3): 49-53
112. Mosmann TR. Interleukin 10. In: Thomson AW (editor) *The cytokine handbook*. Academic Press, London 1994: 223-238
113. Motegi Y, Kita H. Interaction with secretory component stimulates effector functions of human eosinophils but not of neutrophils. *J Immunol* 1998; 161: 4340-4346
114. Mularkey MF. Eosinophilic non-allergic rhinitis and vasomotoric rhinitis. In: Settipane G A. (editor). *Rhinitis*. Ocean Side Publ. Rhode Island, Rhode Island 1991: 169-172

115. Mygind N. Nasal polyposis. *J Allergy Clin Immunol* 1990; 86: 827-829
116. Naclerio, R. M.: Pathophysiology of perennial allergic rhinitis. *Allergy* 1997; 52 (Suppl. 36): 7-13
117. Nakajima T, Yamada H, Likura M et al. Intracellular localization and release of eotaxin from normal eosinophils. *FEBS Lett* 1998; 434(3): 226-30
118. Nave H, Gebert A, Pabst R. Morphology and immunology of the human palatine tonsil. *Anat Embryol* 2001; 204(5): 367-373
119. Nonaka M, Nonaka R, Woolley K, Wooley K; Adelroth E, Miura K, Okhawara Y, Glibetic M, Nakano K, Dolovich J. Distinct immunohistochemical localization of IL-4 in human inflamed airway tissues. IL-4 is localized to eosinophils in vivo and is released by peripheral blood eosinophils. *J Immunol* 1995; 155(6): 3234-3244
120. Nonaka M, Pawankar R, Saji F, Yagi T. Eotaxin synthesis by nasal polyp fibroblasts. *Acta Otolaryngol* 1999; 119(7): 816-820
121. Nouri-Aria KT, Wachholz PA, Francis JN, Durham SR. Grass pollen immunotherapy induces mucosal and peripheral IL-10 responses and blocking IgG activity. *J Immunol* 2004; 172(5): 3252-3259
122. Ogra PL. Mucosal immune response in the ear, nose and throat. *Pediatr Infect Dis J* 2000; 19(5 Suppl): 4-8
123. Ohno I, Lea R, Finotto S, Marshall J, Denburg J, Dolovich J, Gauldie J, Jordana M. Granulocyte/macrophage colony-stimulating factor (GM-CSF) gene expression by eosinophils in nasal polyposis. *Amer J Respir Cell Mol Biol* 1991; 5: 505-510
124. Ohno I, Lea RG, Flanders KC, Clark DA, Banwatt D, Dolovich J, Denburg, Harley CB, Gauldie J, Jordana M. Eosinophils in chronically inflamed human upper airway tissues express transforming growth factor beta 1 gene (TGF beta 1). *J Clin Invest* 1992; 89 (5): 1662-1668
125. Pabst R. The anatomical basis for the immune function of the gut. *Anat Embryol* 1987; 176(2): 135-144
126. Pabst R. Is BALT a major component of the human lung immune system? *Immunol Today* 1992; 13(4): 119-122
127. Pabst R, Tschemig T. Lymphocytes in the lung: an often neglected cell. Numbers, characterization and compartmentalization. *Anat Embryol* 1995; 192(4): 293-299
128. Perry ME. The specialised structure of crypt epithelium in the human palatine tonsil and its functional significance. *J Anat* 1994; 185: 111-127
129. Perry M, Whyte A. Immunology of the tonsils. *Immunol Today* 1998; 19: 414-421
130. Pods R, Ross D, van Hulst S et al. RANTES, eotaxin and eotaxin-2 expression and production in patients with aspirin triad. *Allergy* 2003; 58(11): 1165-1170
131. Ponath PD, Qin S, Ringler DJ, Clark-Lewis I, Wang J, Kassam N, Smith H, Shi X, Gonzalo JA, Newman W, Gutierrez-Ramos JC, Mackay CR. Cloning of the human eosinophil chemoattractant eotaxin. Expression, receptor binding, and functional properties suggest a mechanism for the selective recruitment of eosinophils. *J Clin Invest* 1996; 97: 604-612
132. Ponikau JU, Sherris DA, Kern EB, Roberts GD. The diagnosis and incidence of allergic fungal sinusitis. *Mayo Clin Proc* 1999; 74(9): 877-84
133. Prem B., Hauser U, Behrendt H, Bachert C. Charakterisierung von antigen-präsentierenden Zellen in der menschlichen Nasenschleimhaut. *Allergologie* 1992; 15: 175
134. Proud D, Naclerio RM, Gwaltney JM, Hendley JO. Kinins are generated in nasal secretions during natural rhinovirus colds. *J Infect Dis* 1990; 161: 120-123

- 135.Rasp G. Die eosinophile Entzündung der Nasenschleimhaut. Laryngo-Rhino-Otol 2002; 81: 491-498
- 136.Richardson MA. Sore throat, tonsillitis and adenoiditis. Med Clin North Am 1999; 83(1): 75-83
- 137.Rosenberg HF, Tiffany HL. Characterization of the eosinophil granule proteins recognized by the activation-specific antibody EG2. J Leukoc Biol 1994; 56(4): 502-506
- 138.Rudack C, Stoll W, Bachert C. Cytokines in nasal polyposis, acute and chronic sinusitis. Amer J Rhinol 1998; 12 (6): 383-388
- 139.Rudack C, Wagmann M, Hauser U, Ganzer U. Zytokinmuster bei verschiedenen Formen der Sinusitis. Laryngo-Rhino-Otol 1998; 77(1): 34-37
- 140.Rudack C, Bachert C. Zytokine und Chemokine bei Nasennebenhöhlenerkrankungen. Laryngo-Rhino-Otol 1999; 78: 481-490
- 141.Rudack C. Die Bedeutung von Zytokinen und Chemokinen für Entzündungen in den oberen Atemwegen. Laryngo-Rhino-Otol 2001; 80: 343-352
- 142.Sackett DL., Rosenberg WM, Gray JA, Haynes RB, Richardson WS. Evidence-based medicine: what is and what isn't. Br Med J 1996; 312: 71-72
- 143.Saji F, Nonaka M, Pawankar R. Expression of RANTES by IL-1 beta and TNF-alpha stimulated nasal polyp fibroblasts. Auris Nasus Larynx 2000; 7(3): 247-252
- 144.Samuelsson B. An elucidation of the arachidonic acid cascade. Discovery of prostaglandins, thromboxane and leucotrienes. Drugs 1987; 33 (Suppl. 1): 2-9
- 145.Schäfer D, Lindenthal U, Wagner M, Bölskei PL, Baenkler HW. Effect of prostaglandin E2 on eicosanoid release by human bronchial biopsy specimens from normal and inflamed mucosa. Thorax 1996; 51: 919-923
- 146.Schapowal A., Schmitz M. Provocationstests bei aspirinsensitivem Asthma und aspirinsensitiver Rhinosinusitis. Orale, inhalative und bronchiale Provokation. Allergologie 1992; 15: 158 -164
- 147.Schleimer RP, Sterbinsky SA, Kaiser J, Bickel CA, Klunk DA, Tomioka K, Newman W, Luscinskas FW, Gimbrone MJ, McIntyre BW. IL-4 induces adherence of human eosinophils and basophils but not neutrophils to endothelium. Association with expression of VCAM-1. J Immunol 1992; 148 (4): 1086-1092
- 148.Schmitz-Schumann M, Juhl E, Costabel U. Analgesic asthma-provocation challenge with acetylsalicylic acid. Atmung Lungenerkrkh 1985; 10: 479
- 149.Schroder JM, Mochizuki M. The role of chemokines in cutaneous allergic inflammation. Biol Chem 1999; 380(7-8): 889-896
- 150.Schwachmann H, Kulczycki L, Mueller H. Nasal polyposis in patients with cystic fibrosis. Amer J Dis Child 1996; 129: 761; 102: 768
- 151.Sehmi R, Cromwell O, Wardlaw J, Moqbel R, Kay B. Interleukin-8 is a chemoattractant for eosinophils purified from subjects with a blood eosinophilia but not from healthy subjects. Clin Exp Allergy 1994; 23: 1027-1031
- 152.Sehmi R, Wood LJ, Watson R, Foley R, Hamid Q, O'Byrne PM et al. Allergen-induced increases in IL-5 receptor-subunit expression on bone marrow-derived CD34+ cells from asthmatic subjects: a novel marker of progenitor cell commitment towards eosinophilic differentiation. J Clin Invest 1997; 100: 2466-2475
- 153.Settipane RA., Constantine HP, Settipane GA. Aspirin intolerance and recurrent urticaria in normal adults and children. Allergy 1980; 35: 149
- 154.Settipane GA., Chafee FH. Nasal polyps. Amer J Rhinol 1987; 1: 119-126

155. Settipane GA. Epidemiology of nasal polyps. *Allergy Asthma Proc* 1996; 17(5): 231-236
156. Shin SH, Park JY, Jeon CH et al. Quantitative analysis of eotaxin and RANTES messenger RNA in nasal polyps: association of tissue and nasal eosinophils. *Laryngoscope* 2000; 110(8): 1353-1357
157. Shinkai A, Yoshisue H, Koike M, Shoji E, Nakagawa S, Nishi T. A novel human CC-chemokine, eotaxin-3, which is expressed in IL-4-stimulated vascular endothelial cells, exhibits potent activity toward eosinophils. *J Immunol* 1999; 163: 1602-1610
158. Simon HU, Blaser K. Inhibition of programmed eosinophil death: a key pathogenic event for eosinophilia? *Immunol Today* 1995; 16 (2): 53-55
159. Simon HU, Yousefi S, Schranz C, Schapowal A, Bachert C, Blaser K.. Direct demonstration of delayed eosinophil apoptosis as a mechanism causing tissue eosinophilia. *J Immunol* 1997; 99: 3902-3908
160. Simson L, Foster PS. Chemokine and cytokine cooperativity: eosinophil migration in the asthmatic response. *Immunol Cell Biol* 2000; 78: 415-422
161. Sladek K., Dworski R, Fitzgerald GA, Buitkus KL, Block FJ, Jr. Marney SR, Sheller R. Allergen-stimulated release of thromboxane A2 and leucotriene E4 in humans. Effect of indomethacin. *Amer Rev Resp Dis* 1990; 141: 1441 –1445
162. Spector SL, Bernstein IL, Li JT, Beiger WK, Kaliner A, Schuller DE, Nicklas RA. Parameters for diagnosis and management of sinusitis. *J Allergy Clin Immunol* 1998; 110(2): 104-144
163. Spits H, de Waal Malefy R. Functional characterization of human IL-10. *Int Arch Allergy Immunol* 1992; 99(1): 8-15
164. Stammberger H., Posawetz W. Functional endoscopic sinus surgery. Concept indications and results of the Messerklinger technique. *Arch Otorhinolaryngol* 1990; 247: 63-76
165. Stoop AE, v.d. Heijden HA, Biewenga J et al. Eosinophils in nasal polyps and nasal mucosa: an immuno-histochemical study. *J Allergy Clin Immunol* 1993; 91(2): 616-22
166. Stoop AE., v.d. Heijden HA, Biewenga J, v.d. Baan S. Lymphocytes and non-lymphoid cells in human nasal polyps. *J. Allergy Clin Immunol.* 1998; 87: 470-475
167. Stoop AE, v.d. Heijden HA, Biewenga J, v.d. Baan S. Clinical aspects and distribution of immunologically active cells in the nasal mucosa of patients with nasal polyps after endoscopic sinus surgery and treatment with topical corticosteroids. *Eur Arch Otorhinolaryngol* 1998; 249 (6): 313-317
168. Subauste MC, Jacoby DB, Richards SM, Proud D. Infection of a human respiratory epithelial cell line with rhinovirus. Induction of cytokine release and modulation of susceptibility to infection by cytokine exposure. *J Clin Invest* 1995; 96: 549-557
169. Surjan L. Tonsils and lymphoepithelial structures in the pharynx as immunobarriers. *Acta Otolaryngol* 1987; 103(5-6): 369-372
170. Szczeklik A, Gryglewski RJ, Czemiawska-Mysik G. Clinical patterns of hypersensitivity to non-steroidal antiinflammatory drugs and their pathogenesis. *J Allergy Clin Immunol* 1977; 60: 296
171. Szczeklik A. The cyclooxygenase theory of aspirin-induced asthma. *Europ Respir J* 1990; 3: 588-593
172. Takatsu K. Interleukin 5. *Curr Opin Immunol* 1992; 4: 299-306
173. Teixeira MM, Wells TN, Lukacs NW, Kunkel SL, Williams TJ, Hellewell PG. Chemokine induced eosinophil recruitment. Evidence of a role for endogenous eotaxin in an vivo allergy model in mouse skin. *J Clin Invest* 1997; 100: 1657-1666
174. Terada N, Maesako K, Hamano N et al. Eosinophil adhesion regulates RANTES production in nasal epithelial cells. *J Immunol* 1997; 158(11): 5464-5470

- 175.Terada N, Hamano N, Nomura T, Numata T, Hirai K. Interleukin-13 and tumour necrosis factor-alpha synergistically induce eotaxin production in human nasal fibroblasts. *Clin Exp Allergy* 2000; 30(3): 348-55
- 176.Terada N, Hamano N, Kim WJ. The kinetics of allergen-induced eotaxin level in nasal lavage fluid: its key role in eosinophil recruitment in nasal mucosa. *Am J Respir Crit Care Med* 2001; 164(4): 575-579
- 177.Teran LM, Mochizuki M, Bartels J, Valencia EL, Hirai K, Schroeder JM. Th1- and Th2-type cytokines regulate the expression and production of eotaxin and RANTES by human lung fibroblasts. *Am J Respir Cell Mol Biol* 1999; 20(4): 777-786
- 178.Tiffany HL, Alkhatib G, Combadiere C, Murphy PM. CC chemokine receptors 1 and 3 are differentially regulated by IL-5 during maturation of eosinophilic HL-60 cells. *J Immunol* 1998; 160(3): 1385-1392
- 179.Till S, Jopling L, Wachholz P, Durham S, Sabroe I. T cell phenotypes of the normal nasal mucosa: induction of Th2 cytokines and CCR3 expression by IL-4. *J Immunol*. 2001; 166(4): 2303-2310
- 180.Tos M, Sasaki Y, Ohnishi M, Larsen P, Drake LA. Pathogenesis of nasal polyps. *Rhinol Suppl* 1992; 14: 181-185
- 181.van Damme J. Interleukin-8 and related molecules. In: Thompson AW (editor). *The cytokine handbook*. London: Academic Press, 1991: 201-214
- 182.van Kempen MJ, Rijkers GT, van Cauwenberge PB. The immune response in adenoids and tonsils. *Int Arch Allergy Immunol* 2000; 122(1): 8-19
- 183.van Cauwenberge PB, Ingels KJ, Bachert C, Wang DY. Microbiology of chronic sinusitis. *Acta Otorhinolaryngol Belg* 1997; 51: 239-246
- 184.Vanchieri C, Ohtoshi T, Cox G, Xaubet A, Abrams JS, Gaudie J, Dolovich J, Denburg JA, Jordana M. Neutrophilic differentiation induced by human upper respiratory airway fibroblast-derived granulocyte/macrophage colony-stimulating factor (GM-CSF). *Amer J Respir Cell Mol Biol* 1991; 4: 11-17
- 185.Wardlaw AJ. Molecular basis for selective eosinophil trafficking in asthma : a multistep paradigm. *J Allerg Clin Immunol* 1999; 104: 917-926
- 186.Weber M, Uguccioni M, Baggioline M, Clark Lewis I, Dahinden CA. Deletion of the NH₂- terminal residue converts monocyte chemotactic protein1 from an activator of basophil mediator release to an eosinophil chemoattractant. *J Exp Med* 1996; 183: 681-685
- 187.Wegner CD, Gundel RH, Reilly P, Haynes N, Letts LG, Rothlein R. Intercellular adhesion molecule-1 (ICAM-1) in the pathogenesis of asthma. *Science* 1990; 149: 452-459
- 188.Weller PF. The immunobiology of eosinophils. *New Engl J Med* 1991; 324(16): 1110-1118
- 189.Wells TNC, Power CA, Lusti-Narasimhan M, Chung C, Peitsch MC, Proudfoot AEI. Selectivity and antagonism of chemokine receptors. *J Leukoc Biol* 1996; 59: 53-60
- 190.Wells TN, Proudfoot AE. Chemokine receptors and their antagonists in allergic lung disease. *Inflamm Res* 1999; 48(7): 353-362
- 191.Widal F, Abrami P, Lermoyez J. First complete description of aspirin idiosyncrasy-asthma-nasal polyposis syndrome (plus urticaria) - 1922 (with a note on aspirin desensitization). *J. Asthma* 1987; 24: 297-300
- 192.Woerly G, Roger N, Loiseau S, Capron M. Expression of Th1 and Th2 immunoregulatory cytokines by human eosinophils. *Int Arch Allergy Immunol* 1999; 118: 95-97
- 193.Wright ED, Frenkel S, Ghaffar O et al. Monocyte chemotactic protein expression in allergy and non-allergy-associated chronic sinusitis. *J Otolaryngol* 1998; 27(5): 281-287
- 194.Wüthrich B. Zur Häufigkeit der Pollenallergie in der Schweiz. In: Ring, J. (Hrsg). *Epidemiologie allergischer Erkrankungen*. MMV Medizin Verlag, Braunschweig 1991: 119 -123

- 195.Yamaguchi Y, Hayashi Y, Sugama Y, Miura Y, Kasahara T, Kitamura S, Torisu M, Mita S, Tominga A, Takatsu K, Suda T. Highly purified murine interleukin-5 (IL-5) stimulates eosinophil function and prolongs in vitro survival. *J Exp Med* 1988; 167: 1737 –1742
- 196.Yawalkar N, Uggioni M, Scharer J, Baggiole M. Enhanced expression of eotaxin and CCR3 in atopic dermatitis. *J Invest Dermatol* 1999; 113(1): 43-48
- 197.Ying S, Meng Q, Taborda-Barata L et al. Human eosinophils express messenger RNA encoding RANTES and store and release biologically active RANTES protein. *Eur J Immunol* 1996; 26(1): 70-76
- 198.Ying S, Meng Q, Zeibecoglou K et al. Eosinophil chemotactic chemokines (eotaxin, eotaxin-2, RANTES, monocyte chemoattractant protein-3 (MCP-3), and MCP-4), and C-C chemokine receptor 3 expression in bronchial biopsies from atopic and nonatopic (intrinsic) asthmatics. *J Immunol* 1999; 163(11): 6321-6329
- 199.Ying S, Robinson DS, Meng Q, et al. C-C chemokines in allergen-induced late-phase cutaneous responses in atopic subjects: association of eotaxin with early 6-hour eosinophils, and of eotaxin-2 and monocyte chemoattractant protein-4 with the later 24-hour tissue eosinophilia, and relationship to basophils and other C-C chemokines (monocyte chemoattractant protein-3 and RANTES). *J Immunol* 1999; 163(7): 3976-3984
- 200.Zeck-Kapp AG, Kapp A. Subcellular mechanisms of eosinophil degranulation. *Int Arch Allergy Immunol* 1995; 107: 345
- 201.Zeck-Kapp G, Kroegel C, Riede UN, Kapp A.. Mechanisms of human eosinophil activation by complement protein C5a and platelet-activating factor: similar functional responses are accompanied by different morphologic alterations. *Allergy* 1995; 50(1): 34-47