

6. Literaturverzeichnis

- Abbott BD, Morgan KS, Birnbaum LS, Pratt RM. TCDD alters the extracellular matrix and basal lamina of the fetal mouse kidney. *Teratology*. 1987;35:335-44.
- Abbott BD, Schmid JE, Pitt JA, Buckalew AR, Wood CR, Held GA, Diliberto JJ. Adverse reproductive outcomes in the transgenic Ah receptor-deficient mouse. *Toxicol Appl Pharmacol*. 1999;155:62-70.
- Arthur MF and Frea JJ. Microbiol activity in soils contaminated with 2,3,7,8-TCDD. *Environ Toxicol Chem*. 1989;7:5-13.
- Atkinson R. Atmospheric lifetimes of dibenzo-p-dioxins and dibenzofurans. *Sci Tot Environ*. 1991;104:17-33.
- Bacci E, Cerejeira MJ, Gaggi G, Chemello D, Calamari, Vighi M. Bioconcentration of organic chemical vapours in plant leaves: the azalea model. *Chemosphere*. 1990;21:525-535.
- Bager Y, Kato Y, Kenne K, Warngard L. The ability to alter the gap junction protein expression outside GST-P positive foci in liver of rats was associated to the tumour promotion potency of different polychlorinated biphenyls. *Chem Biol Interact*. 1997;103:199-212.
- Bannister R, Safe S. Synergistic interactions of 2,3,7,8-TCDD and 2,2',4,4',5,5'-hexachlorobiphenyl in C57BL/6J and DBA/2J mice: role of the Ah receptor. *Toxicology*. 1987;44:159-69.
- Bannister R, Davis D, Zacharewski T, Tizard I, Safe S. Aroclor 1254 as a 2,3,7,8-tetrachlorodibenzo-p-dioxin antagonist: effects on enzyme induction and immunotoxicity. *Toxicology*. 1987;46:29-42.
- Beck H, Eckart K, Mathar W, Wittkowski R. PCDD and PCDF body burden from food intake in the Federal Republic of Germany. *Chemosphere*. 1989;18:417.

- Beck H, Dross A, Mathar W, Wittkowsky R. Influence of different regional emissions and cardboard containers on the levels of PCDD, PCDF and related compounds in milk. *Chemosphere*. 1990;21:789.
- Birnbaum LS, McDonald MM, Blair PC, Clark AM, Harris MW. Differential toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in C57BL/6J mice congenic at the Ah Locus. *Fundam Appl Toxicol*. 1990;15:186-200.
- Bookstaff RC, Kamel F, Moore RW, Bjerke DL, Peterson RE. Altered regulation of pituitary gonadotropin-releasing hormone (GnRH) receptor number and pituitary responsiveness to GnRH in 2,3,7,8-tetrachlorodibenzo-p-dioxin-treated male rats. *Toxicol Appl Pharmacol*. 1990;105:78-92.
- Boyd RL, Tucek CL, Godfrey DI, Izon DJ, Wilson TJ, Davidson NJ, Bean AG, Ladyman HM, Ritter MA, Hugo P. The thymic microenvironment. *Immunol Today*. 1993;14:445-59.
- Brouwer A, Ahlborg UG, Van den Berg M, Birnbaum LS, Boersma ER, Bosveld B, Denison MS, Gray LE, Hagmar L, Holene E, et al. Functional aspects of developmental toxicity of polyhalogenated aromatic hydrocarbons in experimental animals and human infants. *Eur J Pharmacol*. 1995;293:1-40.
- Buchmann A, Stinchcombe S, Korner W, Hagenmaier H, Bock KW. Effects of 2,3,7,8-tetrachloro- and 1,2,3,4,6,7,8-heptachlorodibenzo-p-dioxin on the proliferation of preneoplastic liver cells in the rat. *Carcinogenesis*. 1994;15:1143-50.
- Bumb RR, Crummett WB, Cutie SS, Gledhill JR, Hummel RH, Kagel RO, Lamparski LL, Luoma EV, Miller DL, Nestruck TJ, Shadoff LA, Stehl RH, Woods JS. Trace chemistries of fire: a source of chlorinated dioxins. *Science*. 1980;210:385-90.
- Burbach KM, Poland A, Bradfield CA. Cloning of the Ah-receptor cDNA reveals a distinctive ligand-activated transcription factor. *Proc Natl Acad Sci U S A*. 1992;89:8185-9.

- Burleson GR, Lebec H, Yang YG, Ibanes JD, Pennington KN, Birnbaum LS. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on influenza virus host resistance in mice. *Fundam Appl Toxicol.* 199;29:40-7.
- Buu-Hoi NP, Chanh PH, Sesque G, Azum-Gelade MC, Saint-Ruf G. Organs as targets of "dioxin" (2,3,7,8-tetrachlorodibenzo-p-dioxin) intoxication. *Naturwissenschaften.* 1972;59:174-5.
- Camacho IA, Nagarkatti M, Nagarkatti PS. Evidence for Induction of Apoptosis in T Cells from Murine Fetal Thymus following Perinatal Exposure to 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD). *Toxicol Sci.* 2004;78:96-106.
- Chang AC, Salomon DR, Wadsworth S, Hong MJ, Mojcik CF, Otto S, Shevach EM, Coligan JE. Alpha 3 beta 1 and alpha 6 beta 1 integrins mediate laminin/merosin binding and function as costimulatory molecules for human thymocyte proliferation. *J Immunol.* 1995;154:500-10.
- Chang TJ, Juan CC, Yin PH, Chi CW, Tsay HJ. Up-regulation of beta-actin, cyclophilin and GAPDH in N1S1 rat hepatoma. *Oncol Rep.* 1998;5:469-71.
- CLUA (Chemisches Landesuntersuchungsamt NRW Münster). Jahresberichte 1992/1993.
- Contreiras EC, Lenzi HL, Meirelles MN, Caputo LF, Calado TJ, Villa-Verde DM, Savino W. The equine thymus microenvironment: a morphological and immunohistochemical analysis. *Dev Comp Immunol.* 2004;28:251-64.
- Cook JC, Dold KM, Greenlee WF. An in vitro model for studying the toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin to human thymus. *Toxicol Appl Pharmacol.* 1987;89:256-68.
- Courtney KD, Moore JA. Teratology studies with 2,4,5-trichlorophenoxyacetic acid and 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Toxicol Appl Pharmacol.* 1971;20:396-403.
- CVUA (Chemisches Landes- und Staatliches Veterinäruntersuchungsamt). 1994-1997.

- Davis D, Safe S. Immunosuppressive activities of polychlorinated dibenzofuran congeners: quantitative structure-activity relationships and interactive effects. *Toxicol Appl Pharmacol.* 1988;94:141-9.
- Davis D, Safe S. Dose-response immunotoxicities of commercial polychlorinated biphenyls (PCBs) and their interaction with 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Toxicol Lett.* 1989;48:35-43.
- Davis D, Safe S. Immunosuppressive activities of polychlorinated biphenyls in C57BL/6N mice: structure-activity relationships as Ah receptor agonists and partial antagonists. *Toxicology.* 1990;63:97-111.
- De Heer C, Schuurman HJ, Liem AK, Penninks AH, Vos JG, van Loveren H. Toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) to the human thymus after implantation in SCID mice. *Toxicol Appl Pharmacol.* 1995;134:296-304.
- Dencker L, Hassoun E, d'Argy R, Alm G. Fetal thymus organ culture as an in vitro model for the toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin and its congeners. *Mol Pharmacol.* 1985;27:133-40.
- DeVito MJ, Thomas T, Martin E, Umbreit TH, Gallo MA. Antiestrogenic action of 2,3,7,8-tetrachlorodibenzo-p-dioxin: tissue-specific regulation of estrogen receptor in CD1 mice. *Toxicol Appl Pharmacol.* 1992;113:284-92.
- De Waal EJ, Schuurman HJ, Loeber JG, Van Loveren H, Vos JG. Alterations in the cortical thymic epithelium of rats after in vivo exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD): an (immuno)histological study. *Toxicol Appl Pharmacol.* 1992;115:80-8.
- De Waal EJ, Schuurman HJ, Van Loveren H, Vos JG. Differential effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin, bis(tri-n-butyltin) oxide and cyclosporine on thymus histophysiology. *Crit Rev Toxicol.* 1997;27:381-430.

- De Waal Malefijt R, Leene W, Roholl PJ, Wormmeester J, Hoeben KA. T cell differentiation within thymic nurse cells. *Lab Invest.* 1986;55:25-34.
- Diliberto JJ, Burgin DE, Birnbaum LS. Effects of CYP1A2 on disposition of 2,3,7, 8-tetrachlorodibenzo-p-dioxin, 2,3,4,7,8-pentachlorodibenzofuran, and 2,2',4,4',5,5'-hexachlorobiphenyl in CYP1A2 knockout and parental (C57BL/6N and 129/Sv) strains of mice. *Toxicol Appl Pharmacol.* 1999;159:52-64.
- Eadon G, Kaminsky L, Silkworth J, Aldous K, Hilker D, O'Keefe P, Smith R, Gierthy J, Hawley J, Kim N, et al. Calculation of 2,3,7,8-TCDD equivalent concentrations of complex environmental contaminant mixtures. *Environ Health Perspect.* 1986;70:221-7.
- Elder PK, Schmidt LJ, Ono T, Getz MJ. Specific stimulation of actin gene transcription by epidermal growth factor and cycloheximide. *Proc Natl Acad Sci U S A.* 1984;81:7476-80.
- Elder PK, French CL, Subramaniam M, Schmidt LJ, Getz MJ. Evidence that the functional beta-actin gene is single copy in most mice and is associated with 5' sequences capable of conferring serum- and cycloheximide-dependent regulation. *Mol Cell Biol.* 1988;8:480-5.
- European Commission DG Environment. UK Department of the Environment, Transport and the Regions (DETR). *Compilation of EU Dioxin Exposure and Health Data.* 1999.
- Fava RA, McClure DB. Fibronectin-associated transforming growth factor. *J Cell Physiol.* 1987;131:184-9.
- Fiedler H, Swerev M, Nordsiek H, Dörr G, Hutzinger O. Long-term ambient air measurements of PCDD/PCDF in Southern Germany (1993-1996). *Organohalogen Compd.* 1997;33:93-98.

- Filser JG, Csanády GyA, Gans G, Kreuzer PE. Pharmakokinetik der Dioxin-Belastung von Säuglingen durch Muttermilch - Entwicklung und Validierung eines pharmakokinetischen Langzeitmodells für 2,3,7,8-Tetrachlorodibenz-p-dioxin-Abschlussbericht. Ministerium für Arbeit, Gesundheit und Sozialordnung Baden-Württemberg, Stuttgart.1994.
- Fine JS, Gasiewicz TA, Silverstone AE. Lymphocyte stem cell alterations following perinatal exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Mol Pharmacol.* 1989;35:18-25.
- Fine JS, Silverstone AE, Gasiewicz TA. Impairment of prothymocyte activity by 2,3,7,8-tetrachlorodibenzo-p-dioxin. *J Immunol.* 1990;144:1169-76.
- Fürst P, Fürst C, Wilmers K. Body burden with PCDD and PCDF from food. Banbury Report. 1990;35:133-142.
- Gagelin C, Pierre M, Gavaret JM, Toru-Delbauffe D. Rapid TGF beta 1 effects on actin cytoskeleton of astrocytes: comparison with other factors and implications for cell motility. *Glia.* 1995;13:283-93.
- Gaido KW, Maness SC, Leonard LS, Greenlee WF. 2,3,7,8-Tetrachlorodibenzo-p-dioxin dependent regulation of transforming growth factors-alpha and -beta 2 expression in a human keratinocyte cell line involves both transcriptional and post-transcriptional control. *J Biol Chem.* 1992;267:24591-5.
- Gaido KW, Maness SC. Regulation of gene expression and acceleration of differentiation in human keratinocytes by 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Toxicol Appl Pharmacol.* 1994;127:199-208.
- Gasiewicz TA, Geiger LE, Rucci G, Neal RA. Distribution, excretion, and metabolism of 2,3,7,8-tetrachlorodibenzo-p-dioxin in C57BL/6J, DBA/2J, and B6D2F1/J mice. *Drug Metab Dispos.* 1983;11:397-403.

- Gasiewicz TA and Henty EC. Different forms of the Ah-receptor and possible role in species- and tissue-specific responses to TCDD. *Banbury Report*. 1991;35:321-350.
- Geberhiwot T, Assefa D, Kortessmaa J, Ingerpuu S, Pedraza C, Wondimu Z, Charo J, Kiessling R, Virtanen I, Tryggvason K, Patarroyo M. Laminin-8 (alpha4beta1gamma1) is synthesized by lymphoid cells, promotes lymphocyte migration and costimulates T cell proliferation. *J Cell Sci*. 2001;114:423-33.
- Geyer HJ, Scheunert I, Filser JG, Korte F. Bioconcentration potential (BCP) of 2,3,7,8-tetrachlorodibenzo-p-dioxin in terrestrial organisms including humans. *Chemosphere*. 1986;15:1495-1502.
- Greenberg ME, Hermanowski AL, Ziff EB. Effect of protein synthesis inhibitors on growth factor activation of c-fos, c-myc, and actin gene transcription. *Mol Cell Biol*. 1986;6:1050-7.
- Greenlee WF, Dold KM, Irons RD, Osborne R. Evidence for direct action of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on thymic epithelium. *Toxicol Appl Pharmacol*. 1985;79:112-20.
- Helling CS, Isensee AR, Woolson EA, Ensor PJD, Jones GE, Plimmer JR, Kearney PC. Chlordioxins in pesticides, soils and plants. *J Environ Quality*. 1973;2:1717-178.
- Henry EC, Kende AS, Rucci G, Totleben MJ, Willey JJ, Dertinger SD, Pollenz RS, Jones JP, Gasiewicz TA. Flavone antagonists bind competitively with 2,3,7, 8-tetrachlorodibenzo-p-dioxin (TCDD) to the aryl hydrocarbon receptor but inhibit nuclear uptake and transformation. *Mol Pharmacol*. 1999;55:716-25.
- Hites RA, Foran JA, Carpenter DO, Hamilton MC, Knuth BA, Schwager SJ. Global assessment of organic contaminants in farmed salmon. *Science*. 2004;303:226-9.
- Hoffman RE, Stehr-Green PA, Webb KB, Evans RG, Knutsen AP, Schramm WF, Staake JL, Gibson BB, Steinberg KK. Health effects of long-term exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *JAMA*. 1986;255:2031-8.

- Holladay SD, Lindstrom P, Blaylock BL, Comment CE, Germolec DR, Heindell JJ, Luster MI. Perinatal thymocyte antigen expression and postnatal immune development altered by gestational exposure to tetrachlorodibenzo-p-dioxin (TCDD). *Teratology*. 1991;44:385-93.
- Holsapple MP, Snyder NK, Wood SC, Morris DL. A review of 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced changes in immunocompetence: 1991 update. *Toxicology*. 1991;69:219-55.
- Houk VN. Discussion of Vietnam veteran and the Agent Orange issues. *Banbury Report*. 1991;35:259-275.
- IARC (WHO, Lyon). Polychlorinated dibenzo-para-dioxins and polychlorinated dibenzofurans. IARC monographs on the evaluation of carcinogenic risks to humans. 1997;69.
- Ikuta T, Eguchi H, Tachibana T, Yoneda Y, Kawajiri K. Nuclear localization and export signals of the human aryl hydrocarbon receptor. *J Biol Chem*. 1998;273:2895-904.
- Kamath AB, Nagarkatti PS, Nagarkatti M. Characterization of phenotypic alterations induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin on thymocytes in vivo and its effect on apoptosis. *Toxicol Appl Pharmacol*. 1998;150:117-24.
- Kerkvliet NI, Brauner JA, Matlock JP. Humoral immunotoxicity of polychlorinated diphenyl ethers, phenoxyphenols, dioxins and furans present as contaminants of technical grade pentachlorophenol. *Toxicology*. 1985;36:307-24.
- Kerkvliet NI, Steppan LB, Brauner JA, Deyo JA, Henderson MC, Tomar RS, Buhler DR. Influence of the Ah locus on the humoral immunotoxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin: evidence for Ah-receptor-dependent and Ah-receptor-independent mechanisms of immunosuppression. *Toxicol Appl Pharmacol*. 1990;105:26-36.
- Kerkvliet NI, Oughton JA. Acute inflammatory response to sheep red blood cell challenge in mice treated with 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD): phenotypic and functional analysis of peritoneal exudate cells. *Toxicol Appl Pharmacol*. 1993;119:248-57.



- Kikuchi H, Shibasaki M, Ahmed S, Baba T. Method for evaluation of immunotoxicity of dioxin compounds using human T-lymphoblastic cell line, L-MAT. *Chemosphere*. 2001;43:815-8.
- Kim MG, Lee G, Lee SK, Lolkema M, Yim J, Hong SH, Schwartz RH. Epithelial cell-specific laminin 5 is required for survival of early thymocytes. *J Immunol*. 2000;165:192-201.
- Kjeller LO, Jones KC, Rappe C, Johnston AE. Evidence for increases in the environmental burden of polychlorinated dibenzo-p-dioxins and furans (PCDD/PCDF) over the last century. *Organochlorine Compounds* (ed. Hutzinger O and Fiedler H). 1990;Vol.1:433.
- Kociba RJ, Keyes DG, Beyer JE, Carreon RM, Wade CE, Dittenber DA, Kalnins RP, Frauson LE, Park CN, Barnard SD, Hummel RA, Humiston CG. Results of a two-year chronic toxicity and oncogenicity study of 2,3,7,8-tetrachlorodibenzo-p-dioxin in rats. *Toxicol Appl Pharmacol*. 1978;46:279-303.
- Kremer J, Lai ZW, Esser C. Evidence for the promotion of positive selection of thymocytes by Ah receptor agonist 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Eur J Pharmacol*. 1995;293:413-27.
- Kreuzer PE, Csanady GA, Baur C, Kessler W, Papke O, Greim H, Filser JG. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) and congeners in infants. A toxicokinetic model of human lifetime body burden by TCDD with special emphasis on its uptake by nutrition. *Arch Toxicol*. 1997;71:383-400.
- Kupfer D, Bulger WH. Interactions of chlorinated hydrocarbons with steroid hormones. *Fed Proc*. 1976;35:2603-8.
- Kurl RN, Abraham M, Olnes MJ. Early effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on rat thymocytes in vitro. *Toxicology*. 1993;77:103-14.
- Kwok ESC, Arey J, Atkinson R. Gas-phase atmospheric chemistry of dibenzo-p-dioxin and dibenzofuran. *Environ Sci and Technol*. 1994;28:528-533.

- Kwok ESC, Atkinson R, Arey J. Atmospheric chemical loss processes for PCDDs and PCDFs. *Organhalogen Compd.* 1995;24:163-166.
- Lagrotta-Candido JM, Villa-Verde DM, Vanderlei FH Jr, Savino W. Extracellular matrix components of the mouse thymus microenvironment. V. Interferon-gamma modulates thymic epithelial cell/thymocyte interactions via extracellular matrix ligands and receptors. *Cell Immunol.* 1996;170:235-44.
- Lai ZW, Hundeiker C, Gleichmann E, Esser C. Cytokine gene expression during ontogeny in murine thymus on activation of the aryl hydrocarbon receptor by 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Mol Pharmacol.* 1997;52:30-7.
- Lai ZW, Fiore NC, Gasiewicz TA, Silverstone AE. 2,3,7,8-Tetrachlorodibenzo-p-dioxin and diethylstilbestrol affect thymocytes at different stages of development in fetal thymus organ culture. *Toxicol Appl Pharmacol.* 1998;149:167-77.
- Laios MD, Wyman A, Murante FG, Fiore NC, Staples JE, Gasiewicz TA, Silverstone AE. Cell proliferation arrest within intrathymic lymphocyte progenitor cells causes thymic atrophy mediated by the aryl hydrocarbon receptor. *J Immunol.* 2003;171:4582-91.
- Lang DS, Becker S, Clark GC, Devlin RB, Koren HS. Lack of direct immunosuppressive effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on human peripheral blood lymphocyte subsets in vitro. *Arch Toxicol.* 1994;68:296-302.
- Lang DS, Becker S, Devlin RB, Koren HS. 2,3,7,8-TCDD induces cytochrome P450 enzyme activity but not proliferation or phenotypical changes in human peripheral blood lymphocytes. *Toxicol Lett.* 1996;88:317-25.
- Lawrence BP, Leid M, Kerkvliet NI. Distribution and behavior of the Ah receptor in murine T lymphocytes. *Toxicol Appl Pharmacol.* 1996;138:275-84.

- Lee DC, Barlow KD, Gaido KW. The actions of 2,3,7,8-tetrachlorodibenzo-p-dioxin on transforming growth factor-beta2 promoter activity are localized to the TATA box binding region and controlled through a tyrosine kinase-dependent pathway. *Toxicol Appl Pharmacol.* 1996;137:90-9.
- Leof EB, Proper JA, Getz MJ, Moses HL. Transforming growth factor type beta regulation of actin mRNA. *J Cell Physiol.* 1986;127:83-8.
- Lippert H. *Lehrbuch der Anatomie.* Urban und Schwarzenberg. 4. Auflage, 1996.
- Löffler G und Petrides PE. *Biochemie und Pathobiochemie.* Springer. 6. Auflage, 1997.
- Lundberg K. Dexamethasone and 2,3,7,8-tetrachlorodibenzo-p-dioxin can induce thymic atrophy by different mechanisms in mice. *Biochem Biophys Res Commun.* 1991;178:16-23.
- Lu YF, Santostefano M, Cunningham BD, Threadgill MD, Safe S. Substituted flavones as aryl hydrocarbon (Ah) receptor agonists and antagonists. *Biochem Pharmacol.* 1996;51:1077-87.
- Mably TA, Bjerke DL, Moore RW, Gendron-Fitzpatrick A, Peterson RE. In utero and lactational exposure of male rats to 2,3,7,8-tetrachlorodibenzo-p-dioxin. 3. Effects on spermatogenesis and reproductive capability. *Toxicol Appl Pharmacol.* 1992a;114:118-26.
- Mably TA, Moore RW, Goy RW, Peterson RE. In utero and lactational exposure of male rats to 2,3,7,8-tetrachlorodibenzo-p-dioxin. 2. Effects on sexual behavior and the regulation of luteinizing hormone secretion in adulthood. *Toxicol Appl Pharmacol.* 1992b;114:108-17.
- Matsumura F and Benezet HJ. Studies on the bioaccumulation and microbial degradation of 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Environ Health Perspect.* 1973;5:253-258.
- McConkey DJ, Hartzell P, Duddy SK, Hakansson H, Orrenius S. 2,3,7,8-Tetrachlorodibenzo-p-dioxin kills immature thymocytes by Ca<sup>2+</sup>-mediated endonuclease activation. *Science.* 1988;242:256-9.

- McLachlan MS. Bioaccumulation of hydrophobic chemicals in agricultural food chains. *Environ Sci Technol.* 1996;30:252-259.
- McNulty WP, Nielsen-Smith KA, Lay JO Jr, Lippstreu DL, Kangas NL, Lyon PA, Gross ML. Persistence of TCDD in monkey adipose tissue. *Food Chem Toxicol.* 1982;20:985-6.
- Mocarelli P, Marocchi A, Brambilla P, Gerthoux PM, Colombo L, Mondonico A, Meazza L. Effects of dioxin exposure in humans at Seveso, Italy. *Banbury Report* 1991;35:95-107.
- Mocarelli P, Brambilla P, Gerthoux PM, Patterson DG Jr, Needham LL. Change in sex ratio with exposure to dioxin. *Lancet.* 1996;348:409.
- Morris DL, Snyder NK, Gokani V, Blair RE, Holsapple MP. Enhanced suppression of humoral immunity in DBA/2 mice following subchronic exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *Toxicol Appl Pharmacol.* 1992;112:128-32.
- NATO/CCMS. Scientific basis for the development of the International Toxicity Equivalency Factor (I-TEF) method of risk assessment for complex mixtures of dioxins and related compounds. Report No. 178; 1989.
- Nebert DW, Jensen NM. The Ah locus: genetic regulation of the metabolism of carcinogens, drugs, and other environmental chemicals by cytochrome P-450-mediated monooxygenases. *CRC Crit Rev Biochem.* 1979;6:401-37.
- Needham LL, Patterson DG, Houk VN. Levels of TCDD in selected human populations and their relevance to human risk assessment. *Banbury Report.* 1991;35:229-247.
- Needham LL, Gerthoux PM, Patterson DG Jr, Brambilla P, Turner WE, Beretta C, Pirkle JL, Colombo L, Sampson EJ, Tramacere PL, Signorini S, Meazza L, Carreri V, Jackson RJ, Mocarelli P. Serum dioxin levels in Seveso, Italy, population in 1976. *Teratog Carcinog Mutagen.* 1997-98;17:225-40.

- Neubert R, Jacob-Müller U, Stahlmann R, Helge H, Neubert D. Polyhalogenated dibenzo-p-dioxins and dibenzofurans and the immune system. 1. Effects on peripheral lymphocyte subpopulations of a non-human primate (*Callithrix jacchus*) after treatment with 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *Arch Toxicol.* 1990;64:345-59.
- Neubert R, Jacob-Müller U, Helge H, Stahlmann R, Neubert D. Polyhalogenated dibenzo-p-dioxins and dibenzofurans and the immune system. 2. In vitro effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on lymphocytes of venous blood from man and a non-human primate (*Callithrix jacchus*). *Arch Toxicol.* 1991;65:213-9.
- Neubert R, Golor G, Stahlmann R, Helge H, Neubert D. Polyhalogenated dibenzo-p-dioxins and dibenzofurans and the immune system. 4. Effects of multiple-dose treatment with 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on peripheral lymphocyte subpopulations of a non-human primate (*Callithrix jacchus*). *Arch Toxicol.* 1992;66:250-9.
- Neubert R, Stahlmann R, Korte M, van Loveren H, Vos JG, Golor G, Webb JR, Helge H, Neubert D. Effects of small doses of dioxins on the immune system of marmosets and rats. *Ann N Y Acad Sci.* 1993;685:662-86.
- Neubert R, Golor G, Maskow L, Helge H, Neubert D. Evaluation of possible effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin and other congeners on lymphocyte receptors in *Callithrix jacchus* and man. *Exp Clin Immunogenet.* 1994;11:119-27.
- Neubert R, Golor G, Helge H, Neubert D. Risk assessment for possible effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) and related substances on components and functions of the immune system. *Exp Clin Immunogenet.* 1994;11:163-71.
- Neubert D. Reflections on the assessment of the toxicity of "dioxins" for humans, using data from experimental and epidemiological studies. *Teratog Carcinog Mutagen.* 1997-98;17:157-215.
- Ng SY, Gunning P, Liu SH, Leavitt J, Kedes L. Regulation of the human beta-actin promoter by upstream and intron domains. *Nucleic Acids Res.* 1989;17:601-15.

- Nohara K, Fujimaki H, Tsukumo S, Ushio H, Miyabara Y, Kijima M, Tohyama C, Yonemoto J. The effects of perinatal exposure to low doses of 2,3,7,8-tetrachlorodibenzo-p-dioxin on immune organs in rats. *Toxicology*. 2000;154:123-33.
- Okey AB, Riddick DS, Harper PA. The Ah receptor: mediator of the toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) and related compounds. *Toxicol Lett*. 1994;70:1-22.
- Olson JR, Gasiewicz TA, Neal RA. Tissue distribution, excretion, and metabolism of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in the Golden Syrian hamster. *Toxicol Appl Pharmacol*. 1980;56:78-85.
- Olson JR, Holscher MA, Neal RA. Toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in the golden Syrian hamster. *Toxicol Appl Pharmacol*. 1980b;55:67-78.
- Olson JR. Metabolism and disposition of 2,3,7,8-tetrachlorodibenzo-p-dioxin in guinea pigs. *Toxicol Appl Pharmacol*. 1986;85:263-73.
- Olson JR, McGarrigle BP, Tonucci DA, Schecter A, Henry Eichelberger H. Developmental toxicity of 2,3,7,8-TCDD in the rat and hamster. *Chemosphere* 1990;20:1117-23.
- Oughton JA, Pereira CB, DeKrey GK, Collier JM, Frank AA, Kerkvliet NI. Phenotypic analysis of spleen, thymus, and peripheral blood cells in aged C57B1/6 mice following long-term exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Fundam Appl Toxicol*. 1995;25:60-9.
- Park JY, Shigenaga MK, Ames BN. Induction of cytochrome P4501A1 by 2,3,7,8-tetrachlorodibenzo-p-dioxin or indolo(3,2-b)carbazole is associated with oxidative DNA damage. *Proc Natl Acad Sci U S A*. 1996;93:2322-7.
- Pennise DM and Kamens RM. Atmospheric behavior of polychlorinated dibenzo-p-dioxins and dibenzofurans and the effect of the combustion temperatur. *Environ Sci and Technol*. 1996;30:2832-2842.
- Peterson RE, Seefeld MD, Christian BJ. The wasting syndrome in 2,3,7,8-tetrachlorodibenzo-p-dioxin toxicity: basic features and their interpretation. *Banbury Report*. 1984;18:291-308.

- Peterson RE, Theobald HM, Kimmel GL. Developmental and reproductive toxicity of dioxins and related compounds: cross-species comparisons. *Crit Rev Toxicol.* 1993;23:283-335.
- Piek E, Heldin CH, Ten Dijke P. Specificity, diversity, and regulation in TGF-beta superfamily signaling. *FASEB J.* 1999;13:2105-24.
- Pfister K. Wirkungen von Fluorchinolonen auf Sehnen- und Knorpelgewebe bei Ratten und Reduktion der Gelenkschäden durch Magnesium oder Vitamin E. Institut für klinische Pharmakologie und Toxikologie, Universitätsklinikum Benjamin Franklin, Freie Universität Berlin. 2000.
- Podoll R, Jaber H, Mill T. TCDD: rates of volatilisation and photolysis in the environment. *Environ Sci and Technol.* 1986;20:490-492.
- Pohl H, Holler J. Halogenated aromatic hydrocarbons and toxicity equivalency factors (TEFs) from the public health assessment perspective. *Chemosphere.* 1995;31:2547-59.
- Poland A and Knutson JC. 2,3,7,8-tetrachlorodibenzo-p-dioxin and related halogenated hydrocarbons: examination of the mechanism of toxicity. *Annu Rev Pharmacol Toxicol.* 1982;22:517-554.
- Poiger H, Schlatter C. Pharmacokinetics of 2,3,7,8-TCDD in man. *Chemosphere.* 1986;15:1489-1494.
- Pongratz I, Mason GG, Poellinger L. Dual roles of the 90-kDa heat shock protein hsp90 in modulating functional activities of the dioxin receptor. Evidence that the dioxin receptor functionally belongs to a subclass of nuclear receptors which require hsp90 both for ligand binding activity and repression of intrinsic DNA binding activity. *J Biol Chem.* 1992;267:13728-34.
- Prell RA, Oughton JA, Kerkvliet NI. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on anti-CD3-induced changes in T-cell subsets and cytokine production. *Int J Immunopharmacol.* 1995;17:951-61.

- Prinz B, Krause GHM, Radermacher L. Criteria for evaluation of dioxins in garden soils and garden plants. *Organohalogen Compd.* 1990;1:595.
- Rappe C, Andersson R, Bergqvist PA, Brohede C, Hansson M, Kjeller LO, Lindström G, Marklund S, Nygren M, Swanson SE, Tysklind M, Wiberg K. Overview over the environmental fate of chlorinated dioxins and dibenzofurans. Sources, levels and isomeric pattern in various matrices. *Chemosphere.* 1987;16:1603.
- Rappe C, Kjeller LO, Bruckmann P, Hackhe KH. Identification and quantification of PCDDs in urban air. *Chemosphere.* 1988;17:3.
- Rappe C, Kjeller LO, Kulp SE. Sampling and analysis of PCDDs and PCDFs in surface water and drinking water at 0.001 ppq levels. *Organohalogen Compd.* 1990;2:207.
- Rhile MJ, Nagarkatti M, Nagarkatti PS. Role of Fas apoptosis and MHC genes in 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD)-induced immunotoxicity of T cells. *Toxicology.* 1996;110:153-67.
- Riecke K, Grimm D, Shakibaei M, Kossmehl P, Schulze-Tanzil G, Paul M, Stahlmann R. Low doses of 2,3,7,8-tetrachlorodibenzo-p-dioxin increase transforming growth factor beta and cause myocardial fibrosis in marmosets (*Callithrix jacchus*). *Arch Toxicol.* 2002;76:360-6.
- Riecke K, Schmidt A, Stahlmann R. Effects of 2,3,7,8-TCDD and PCB 126 on human thymic epithelial cells in vitro. *Arch Toxicol.* 2003;77:358-64.
- Rier SE, Martin DC, Bowman RE, Dmowski WP, Becker JL. Endometriosis in rhesus monkeys (*Macaca mulatta*) following chronic exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Fundam Appl Toxicol.* 1993;21:433-41.
- Rowlands JC, Gustafsson JA. Aryl hydrocarbon receptor-mediated signal transduction. *Crit Rev Toxicol.* 1997;27:109-34.



- Ryan JJ, Panopio LG, Lewis DA, Weber DF. Polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans in cows milk packaged in plastic-coated bleached paperboard containers. *J Agric Food Chem.* 1991;39:218.
- Safe S. Polychlorinated biphenyls (PCBs), dibenzo-p-dioxins (PCDDs), dibenzofurans (PCDFs), and related compounds: environmental and mechanistic considerations which support the development of toxic equivalency factors (TEFs). *Crit Rev Toxicol.* 1990;21:51-88.
- Safe SH. Comparative toxicology and mechanism of action of polychlorinated dibenzo-p-dioxins and dibenzofurans. *Annu Rev Pharmacol Toxicol.* 1986;26:371-99.
- Salomon DR, Crisa L, Mojcik CF, Ishii JK, Klier G, Shevach EM. Vascular cell adhesion molecule-1 is expressed by cortical thymic epithelial cells and mediates thymocyte adhesion. Implications for the function of alpha4beta1 (VLA4) integrin in T-cell development. *Blood.* 1997;89:2461-71.
- Savino W, Villa-Verde DM, Lannes-Vieira J. Extracellular matrix proteins in intrathymic T-cell migration and differentiation? *Immunol Today.* 1993;14:158-61.
- Savino W, Vermeulen M, Dardenne M, Lepault F. The thymic extracellular matrix in genetically engineered mice. *Immunol Today.* 1998;19:144-5.
- Savino W, Dalmau SR, Dealmeida VC. Role of extracellular matrix-mediated interactions in thymocyte migration. *Dev Immunol.* 2000;7:279-91.
- Schantz SL, Bowman RE. Learning in monkeys exposed perinatally to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *Neurotoxicol Teratol.* 1989;11:13-9.
- Schechter A, Ryan JJ. Polychlorinated dibenzo-para-dioxin and dibenzofuran levels in human adipose tissues from workers 32 years after occupational exposure to 2,3,7,8-TCDD. *Chemosphere* 1988;17:915-20.

- Schechter A, Pöpke O, Ball M, Dai LC. Chlorinated dioxin and dibenzofuran levels in human blood samples from various countries, including Vietnam, the Soviet Union, the United States and Germany. *Organohalogen Compd.* 1990;1:271.
- Schechter A, Pöpke O, Ball M. Evidence for transplacental transfer of dioxins from mother to fetus: Chlorinated dioxins and dibenzofuran levels in livers of stillborn infants. *Chemosphere.* 1990b;21:1017.
- Schechter A. Dioxins and related chemicals in humans and in the environment. *Banbury Report.* 1991; 35:169-212.
- Schmeissner PJ, Xie H, Smilenov LB, Shu F, Marcantonio EE. Integrin functions play a key role in the differentiation of thymocytes in vivo. *J Immunol.* 2001;167:3715-24.
- Schmidt JV, Bradfield CA. Ah receptor signaling pathways. *Annu Rev Cell Dev Biol.* 1996;12:55-89.
- Schmidt A. Wirkungen von 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) und 3,3',4,4',5-Pentachlorbiphenyl (PCB 126) auf das Thymusepithel des Menschen in vitro. Institut für klinische Pharmakologie und Toxikologie, Universitätsklinikum Benjamin Franklin, Freie Universität Berlin. 2002.
- Shull MM, Ormsby I, Kier AB, Pawlowski S, Diebold RJ, Yin M, Allen R, Sidman C, Proetzel G, Calvin D, et al. Targeted disruption of the mouse transforming growth factor-beta 1 gene results in multifocal inflammatory disease. *Nature.* 1992;359:693-9.
- Silbergeld EK, Gasiewicz TA. Dioxins and the Ah receptor. *Am J Ind Med.* 1989;16:455-74.
- Smialowicz RJ, Riddle MM, Williams WC, Diliberto JJ. Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on humoral immunity and lymphocyte subpopulations: differences between mice and rats. *Toxicol Appl Pharmacol.* 1994;124:248-56.

- Standley LJ and Hites RA. Chlorinated organic contaminants in the atmosphere. In: Organic Contaminants in the Environment: Environmental Pathways and Effects. Ed. Jones KC. Elsevier Applied Science, London.1991.
- Stinchcombe S, Buchmann A, Bock KW, Schwarz M. Inhibition of apoptosis during 2,3,7,8-tetrachlorodibenzo-p-dioxin-mediated tumour promotion in rat liver. *Carcinogenesis*. 1995;16:1271-5.
- Svensson BG, Nilsson A, Hansson M, Rappe C, Akesson B, Skerfving S. Exposur to dioxins and dibenzofurans through the consumption of fish. *N Engl J Med*. 1991;324:8.
- Svensson C, Lundberg K. Immune-specific up-regulation of adseverin gene expression by 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Mol Pharmacol*. 2001;60:135-42.
- Swanson SE, Rappe C, Malmström J, Kringstad KP. Emissions of PCDDs and PCDFs from the pulp industry. *Chemosphere*. 1990;17:681.
- Tomita S, Jiang HB, Ueno T, Takagi S, Tohi K, Maekawa S, Miyatake A, Furukawa A, Gonzalez FJ, Takeda J, Ichikawa Y, Takahama Y. T cell-specific disruption of arylhydrocarbon receptor nuclear translocator (Arnt) gene causes resistance to 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced thymic involution. *J Immunol*. 2003;171:4113-20.
- Utsumi K, Sawada M, Narumiya S, Nagamine J, Sakata T, Iwagami S, Kita Y, Teraoka H, Hirano H, Ogata M, et al. Adhesion of immature thymocytes to thymic stromal cells through fibronectin molecules and its significance for the induction of thymocyte differentiation. *Proc Natl Acad Sci U S A*. 1991;88:5685-9.
- van Baarlen J, Schuurman HJ, Reitsma R, Huber J. Acute thymus involution during infancy and childhood: immunohistology of the thymus and peripheral lymphoid tissues after acute illness. *Pediatr Pathol*. 1989;9:261-75.

- Van Birgelen AP, Van der Kolk J, Fase KM, Bol I, Poiger H, Brouwer A, Van den Berg M. Subchronic dose-response study of 2,3,7,8-tetrachlorodibenzo-p-dioxin in female Sprague-Dawley rats. *Toxicol Appl Pharmacol.* 1995;132:1-13.
- van Logten MJ, Gupta BN, McConnell EE, Moore JA. Role of the endocrine system in the action of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on the thymus. *Toxicology.* 1980;15:135-44.
- van Logten MJ, Gupta BN, McConnell EE, Moore JA. The influence of malnutrition on the toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in rats. *Toxicology.* 1981;21:77-88.
- Vecchi A, Sironi M, Canegrati MA, Recchia M, Garattini S. Immunosuppressive effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin in strains of mice with different susceptibility to induction of aryl hydrocarbon hydroxylase. *Toxicol Appl Pharmacol.* 1983;68:434-41.
- Viluksela M, Stahl BU, Birnbaum LS, Rozman KK. Subchronic/chronic toxicity of 1,2,3,4,6,7,8-heptachlorodibenzo-p-dioxin (HpCDD) in rats. Part II. Biochemical effects. *Toxicol Appl Pharmacol.* 1997;146:217-26.
- Villa-Verde DM, Lagrota-Candido JM, Vannier-Santos MA, Chammas R, Brentani RR, Savino W. Extracellular matrix components of the mouse thymus microenvironment. IV. Modulation of thymic nurse cells by extracellular matrix ligands and receptors. *Eur J Immunol.* 1994;24:659-64.
- Villa-Verde DM, Calado TC, Ocampo JS, Silva-Monteiro E, Savino W. The conveyor belt hypothesis for thymocyte migration: participation of adhesion and de-adhesion molecules. *Braz J Med Biol Res.* 1999;32:569-72.
- Vivinus-Nebot M, Ticchioni M, Mary F, Hofman P, Quaranta V, Rousselle P, Bernard A. Laminin 5 in the human thymus: control of T cell proliferation via  $\alpha 6 \beta 4$  integrins. *J Cell Biol.* 1999;144:563-74.

- Vogel C, Abel J. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on growth factor expression in the human breast cancer cell line MCF-7. *Arch Toxicol.* 1995;69:259-65.
- Vogel C, Donat S, Dohr O, Kremer J, Esser C, Roller M, Abel J. Effect of subchronic 2,3,7,8-tetrachlorodibenzo-p-dioxin exposure on immune system and target gene responses in mice: calculation of benchmark doses for CYP1A1 and CYP1A2 related enzyme activities. *Arch Toxicol.* 1997;71:372-82.
- Vos JG and Luster MI. Immune alterations. Halogenated biphenyls, terphenyls, naphthalens, dibenzodioxins and related products. 1989;265.
- Ward CT and Matsumura F. Fate of 2,3,7,8-tetrachlorodibenzo-p-dioxin in a model aquatic ecosystem. *Arch Environ Contam Toxicol.* 1979;7:349-357.
- Watt SM, Thomas JA, Edwards AJ, Murdoch SJ, Horton MA. Adhesion receptors are differentially expressed on developing thymocytes and epithelium in human thymus. *Exp Hematol.* 1992;20:1101-11.
- White TEK and Gasiewicz TA. The human oestrogen receptor structural gene contains a DNA sequence that binds activated mouse and human Ah receptors: a possible mechanism of oestrogen receptor regulation by 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Biochem Biophys Res Comm.* 1993;193:956-962.
- Whitlock JP JR. Mechanism of dioxin action. Relevance to risk assessment. *Banbury Report.* 1991;35:351-361.
- Wilson CL, Safe S. Mechanisms of ligand-induced aryl hydrocarbon receptor-mediated biochemical and toxic responses. *Toxicol Pathol.* 1998;26:657-71.
- World Health Organisation. International Programme on Chemical Safty (WHO/IPCS). Polychlorinated dibenzo-p-dioxins and dibenzofurans. *Environmental Health Criteria.* 1989;88.

---

Yanagawa Y, Iwabuchi K, Onoe K. Enhancement of stromal cell-derived factor-1alpha-induced chemotaxis for CD4/8 double-positive thymocytes by fibronectin and laminin in mice. *Immunology*. 2001;104:43-9.

Zamir E, Geiger B. Molecular complexity and dynamics of cell-matrix adhesions. *J Cell Sci*. 2001;114:3583-90.