

7. Literaturverzeichnis

- (1) Gehan EA, Freireich EJ. Cancer clinical trials. A rational basis for use of historical controls. *Semin Oncol* 1981 Dec;8(4):430-6.
- (2) Gehan EA. The evaluation of therapies: historical control studies. *Stat Med* 1984 Oct;3(4):315-24.
- (3) WHO Handbook for reporting results of cancer treatment. Geneva, 1979.
- (4) Miller AB, Hoogstraten B, Staquet M, Winkler A. Reporting results of cancer treatment. *Cancer* 1981 Jan 1;47(1):207-14.
- (5) Therasse P, Arbuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, et al. New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst* 2000 Feb 2;92(3):205-16.
- (6) Green S, Weiss GR. Southwest Oncology Group standard response criteria, endpoint definitions and toxicity criteria. *Invest New Drugs* 1992 Nov;10(4):239-53.
- (7) Buyse M, Thirion P, Carlson RW, Burzykowski T, Molenberghs G, Piedbois P. Relation between tumour response to first-line chemotherapy and survival in advanced colorectal cancer: a meta-analysis. *Meta-Analysis Group in Cancer. Lancet* 2000 Jul 29;356(9227):373-8.
- (8) Li D, Xie K, Wolff R, Abbruzzese JL. Pancreatic cancer. *Lancet* 2004 Mar 27;363(9414): 109-57.
- (9) Gehan EA, Tefft MC. Will there be resistance to the RECIST (Response Evaluation Criteria in Solid Tumors)? *J Natl Cancer Inst* 2000 Feb 2;92(3):179-81.
- (10) Therasse P, Eisenhauer EA, Verweij J. RECIST revisited: A review of validation studies on tumour assessment. *Eur J Cancer* 2006 May;42(8):1031-9.
- (11) Husband JE, Hawkes DJ, Peckham MJ. CT estimations of mean attenuation values and volume in testicular tumors: a comparison with surgical and histologic findings. *Radiology* 1982 Aug;144(3):553-8.
- (12) Mazumdar M, Smith A, Debroy PP, Schwartz LH. A theoretical approach to choosing the minimum number of multiple tumors required for assessing treatment response. *J Clin Epidemiol* 2005 Feb;58(2):150-3.

- (13) Schwartz LH, Mazumdar M, Brown W, Smith A, Panicek DM. Variability in response assessment in solid tumors: effect of number of lesions chosen for measurement. *Clin Cancer Res* 2003 Oct 1;9(12):4318-23.
- (14) Negrier S, Ollivier L, Di Stefano-Louineau D, Escudier B, Savary J, Lasset C, et al. [Reliability of the response rate in oncology: analysis of the causes for variation]. *Bull Cancer* 2000 Dec;87(12):927-34.
- (15) Park JH, Nazarian LN, Halpern EJ, Feld RI, Lev-Toaff AS, Parker L, et al. Comparison of unenhanced and contrast-enhanced spiral CT for assessing interval change in patients with colorectal liver metastases. *Acad Radiol* 2001 Aug;8(8):698-704.
- (16) Park JO, Lee SI, Song SY, Kim K, Kim WS, Jung CW, et al. Measuring response in solid tumors: comparison of RECIST and WHO response criteria. *Jpn J Clin Oncol* 2003 Oct;33(10):533-7.
- (17) Prasad SR, Saini S, Sumner JE, Hahn PF, Sahani D, Boland GW. Radiological measurement of breast cancer metastases to lung and liver: comparison between WHO (bidimensional) and RECIST (unidimensional) guidelines. *J Comput Assist Tomogr* 2003 May;27(3):380-4.
- (18) Sohaib SA, Turner B, Hanson JA, Farquharson M, Oliver RT, Reznick RH. CT assessment of tumour response to treatment: comparison of linear, cross-sectional and volumetric measures of tumour size. *Br J Radiol* 2000 Nov;73(875):1178-84.
- (19) Trillet-Lenoir V, Freyer G, Kaemmerlen P, Fond A, Pellet O, Lombard-Bohas C, et al. Assessment of tumour response to chemotherapy for metastatic colorectal cancer: accuracy of the RECIST criteria. *Br J Radiol* 2002 Nov;75(899):903-8.
- (20) Mazumdar M, Smith A, Schwartz LH. A statistical simulation study finds discordance between WHO criteria and RECIST guideline. *J Clin Epidemiol* 2004 Apr;57(4):358-65.
- (21) Fritschy P, Schneekloth G. [Practical problems of determining liver volume from computed tomographic transverse sections]. *Rofo* 1983 Apr;138(4):453-7.
- (22) Breiman RS, Beck JW, Korobkin M, Glennly R, Akwari OE, Heaston DK, et al. Volume determinations using computed tomography. *AJR Am J Roentgenol* 1982 Feb;138(2):329-33.
- (23) Disler DG, Marr DS, Rosenthal DI. Accuracy of volume measurements of computed tomography and magnetic resonance imaging phantoms by three-dimensional reconstruction and preliminary clinical application. *Invest Radiol* 1994 Aug;29(8):739-45.

- (24) Helmberger H, Bautz W, Sendler A, Fink U, Gerhardt P. [Volumetry of abdominal tumors. Problems--feasibility]. *Radiologe* 1995 Sep;35(9):587-91.
- (25) Mahr A, Levegrun S, Bahner ML, Kress J, Zuna I, Schlegel W. Usability of semiautomatic segmentation algorithms for tumor volume determination. *Invest Radiol* 1999 Feb;34(2):143-50.
- (26) Van HL, Haven F, Bellon E, Baert AL, Bosmans H, Feron M, et al. Factors influencing the accuracy of volume measurements in spiral CT: a phantom study. *J Comput Assist Tomogr* 1997 Mar;21(2):332-8.
- (27) Yim PJ, Vora AV, Raghavan D, Prasad R, McAullife M, Ohman-Strickland P, et al. Volumetric analysis of liver metastases in computed tomography with the fuzzy C-means algorithm. *J Comput Assist Tomogr* 2006 Mar;30(2):212-20.
- (28) Dachman AH, MacEneaney PM, Adedipe A, Carlin M, Schumm LP. Tumor size on computed tomography scans: is one measurement enough? *Cancer* 2001 Feb 1;91(3):555-60.
- (29) Hopper KD, Kasales CJ, Egli KD, TenHave TR, Belman NM, Potok PS, et al. The impact of 2D versus 3D quantitation of tumor bulk determination on current methods of assessing response to treatment. *J Comput Assist Tomogr* 1996 Nov;20(6):930-7.
- (30) Pfeifer T, Haussler MD, Tomczak R, Rilinger N, Zeitler H, Laasch HU, et al. [The value of tumor volumetry as opposed to bidimensional determination of tumor size during follow-up of hepatic metastases from colorectal carcinoma]. *Rofa* 1992 Dec;157(6):548-51.
- (31) Prasad SR, Jhaveri KS, Saini S, Hahn PF, Halpern EF, Sumner JE. CT tumor measurement for therapeutic response assessment: comparison of unidimensional, bidimensional, and volumetric techniques initial observations. *Radiology* 2002 Nov;225(2):416-9.
- (32) Warren KE, Patronas N, Aikin AA, Albert PS, Balis FM. Comparison of one-, two-, and three-dimensional measurements of childhood brain tumors. *J Natl Cancer Inst* 2001 Sep 19;93(18):1401-5.
- (33) Kasper HU, Drebber U, Dries V, Dienes HP. Liver metastases: incidence and histogenesis. *Z. Gastroenterol.* 2005 Okt;43(10): 1149-57.
- (34) Arbeitsgemeinschaft Bevölkerungsbezogener Krebsregister in Deutschland: Krebs in Deutschland, Häufigkeiten und Trends. Saarbrücken 2002. 3. aktualisierte Auflage
- (35) Ballantyne GH, Quin J. Surgical treatment of liver metastases in patients with colorectal cancer. *Cancer* 1993 Jun 15;71(12 Suppl):4252-66.

- (36) Cunningham D, Findlay M. The chemotherapy of colon cancer can no longer be ignored. *Eur J Cancer* 1993;29A(15):2077-9.
- (37) Stangl R, tendorf-Hofmann A, Charnley RM, Scheele J. Factors influencing the natural history of colorectal liver metastases. *Lancet* 1994 Jun 4;343(8910):1405-10.
- (38) Ercolani G, Grazi GL, Ravaioli M, Ceson M, Gardini A, Varotti G, Del Gaudio M, Nardo B, Cavallari A, Liver resection for multiple colorectal metastases: influence of parenchymal involvement and total tumor volume, vs number or location, on long-term survival. *Arch Surg*. 2002 Oct;137(10):1187-92.
- (39) Leonard GD, Brenner B, Kemeny NE. Neoadjuvant chemotherapy before liver resection for patients with unresectable liver metastases from colorectal carcinoma. *J Clin Oncol* 2005 Mar 20;23(9):2038-48.
- (40) Cady B, Jenkins RL, Steele GD, Jr., Lewis WD, Stone MD, McDermott WV, et al. Surgical margin in hepatic resection for colorectal metastasis: a critical and improvable determinant of outcome. *Ann Surg* 1998 Apr;227(4):566-71.
- (41) Saltz LB, Cox JV, Blanke C, Rosen LS, Fehrenbacher L, Moore MJ, et al. Irinotecan plus fluorouracil and leucovorin for metastatic colorectal cancer. Irinotecan Study Group. *N Engl J Med* 2000 Sep 28;343(13):905-14.
- (42) Tournigand C, Andre T, Achille E, Lledo G, Flesh M, Mery-Mignard D, et al. FOLFIRI followed by FOLFOX6 or the reverse sequence in advanced colorectal cancer: a randomized GERCOR study. *J Clin Oncol* 2004 Jan 15;22(2):229-37.
- (43) Bismuth H, Adam R, Levi F, Farabos C, Waechter F, Castaing D, et al. Resection of nonresectable liver metastases from colorectal cancer after neoadjuvant chemotherapy. *Ann Surg* 1996 Oct;224(4):509-20.
- (44) Giacchetti S, Itzhaki M, Gruia G, Adam R, Zidani R, Kunstlinger F, et al. Long-term survival of patients with unresectable colorectal cancer liver metastases following infusional chemotherapy with 5-fluorouracil, leucovorin, oxaliplatin and surgery. *Ann Oncol* 1999 Jun;10(6):663-9.
- (45) Hurwitz H, Fehrenbacher L, Novotny W, Cartwright T, Hainsworth J, Heim W, et al. Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N Engl J Med* 2004 Jun 3;350(23):2335-42.
- (46) Ruers T, Bleichrodt RP. Treatment of liver metastases, an update on the possibilities and results. *Eur J Cancer* 2002 May;38(7):1023-33.

- (47) Curley SA, Izzo F, Delrio P, Ellis LM, Granchi J, Vallone P, et al. Radiofrequency ablation of unresectable primary and metastatic hepatic malignancies: results in 123 patients. *Ann Surg* 1999 Jul;230(1):1-8.
- (48) Wood TF, Rose DM, Chung M, Allegra DP, Foshag LJ, Bilchik AJ. Radiofrequency ablation of 231 unresectable hepatic tumors: indications, limitations, and complications. *Ann Surg Oncol* 2000 Sep;7(8):593-600.
- (49) Helmberger T, Rau H, Linke R, Reiser M. [Diagnosis and staging of liver metastases with imaging methods]. *Chirurg* 1999 Feb;70(2):114-22.
- (50) Barton JB, Langdale LA, Cummins JS et al, McAndrew MR, Saba AK. The utility of routine preoperative computed tomography scanning in the management of veterans with colon cancer. *Am J Surg* 2002 May;183(5):499-503.
- (51) Neumaier CE, Cittadini G, Grasso A, Dahmane M. Role of ultrasonography in the staging of gastrointestinal neoplasms. *Semin Surg Oncol* 2001 Mar;20(2):86-90.
- (52) Soyer P, Pocard M, Boudiaf M, Abitbol M, Hamzi L, Panis Y, et al. Detection of hypovascular hepatic metastases at triple-phase helical CT: sensitivity of phases and comparison with surgical and histopathologic findings. *Radiology* 2004 May;231(2):413-20.
- (53) Kuszyk BS, Bluemke DA, Urban BA, Choti MA, Hruban RH, Sitzmann JV, et al. Portal-phase contrast-enhanced helical CT for the detection of malignant hepatic tumors: sensitivity based on comparison with intraoperative and pathologic findings. *AJR Am J Roentgenol* 1996 Jan;166(1):91-5.
- (54) Valls C, Andia E, Sanchez A, Guma A, Figueras J, Torras J, et al. Hepatic metastases from colorectal cancer: preoperative detection and assessment of resectability with helical CT. *Radiology* 2001 Jan;218(1):55-60.
- (55) Weg N, Scheer MR, Gabor MP. Liver lesions: improved detection with dual-detector-array CT and routine 2.5-mm thin collimation. *Radiology* 1998 Nov;209(2):417-26.
- (56) Baron RL, Oliver JH, III, Dodd GD, III, Nalesnik M, Holbert BL, Carr B. Hepatocellular carcinoma: evaluation with biphasic, contrast-enhanced, helical CT. *Radiology* 1996 May;199(2):505-11.
- (57) Paulson EK, McDermott VG, Keogan MT, DeLong DM, Frederick MG, Nelson RC. Carcinoid metastases to the liver: role of triple-phase helical CT. *Radiology* 1998 Jan;206(1):143-50.
- (58) Kopka L, Grabbe E. [Biphasic liver diagnosis with multiplanar-detector spiral CT]. *Radiologe* 1999 Nov;39(11):971-8.

- (59) Soyer P, Bluemke DA, Hruban RH, Sitzmann JV, Fishman EK. Hepatic metastases from colorectal cancer: detection and false-positive findings with helical CT during arterial portography. *Radiology* 1994 Oct;193(1):71-4.
- (60) Ward J, Naik KS, Guthrie JA, Wilson D, Robinson PJ. Hepatic lesion detection: comparison of MR imaging after the administration of superparamagnetic iron oxide with dual-phase CT by using alternative-free response receiver operating characteristic analysis. *Radiology* 1999 Feb;210(2):459-66.
- (61) Reske SN, Kotzerke J. FDG-PET for clinical use. Results of the 3rd German Interdisciplinary Consensus Conference, "Onko-PET III", 21 July and 19 September 2000. *Eur J Nucl Med* 2001 Nov;28(11):1707-23.
- (62) Zwirerich CV, Vedal S, Miller RR, Muller NL. Solitary pulmonary nodule: high-resolution CT and radiologic-pathologic correlation. Analysis of interobserver and intraobserver variability in CT tumor measurements. *Radiology* 1991 May; 179 (2): 469-76. *AJR*
- (63) Henschke CI, McCauley DI, Yankelevitz DF, Naidich DP, McGuinness G, Miettinen OS, Libby DM, Pasmantier MW, Koizumi J, Altorki NK, Smith JP. Early Lung Cancer Action Project: overall design and findings from baseline screening. *Lancet* 1999 Jul;354(9173):99-105.
- (64) Revel MP, Lefort C, Bissery A, Bienvenu M, Aycard L, Chatellier G, et al. Pulmonary nodules: preliminary experience with three-dimensional evaluation. *Radiology* 2004 May;231(2):459-66.
- (65) Yankelevitz DF, Gupta R, Zhao B, Henschke CI. Small pulmonary nodules: evaluation with repeat CT--preliminary experience. *Radiology* 1999 Aug;212(2):561-6.
- (66) Revel MP, Bissery A, Bienvenu M, Aycard L, Lefort C, Frija G. Are two-dimensional CT measurements of small noncalcified pulmonary nodules reliable? *Radiology* 2004 May;231(2):453-8.
- (67) Henschke CI, Naidich DP, Yankelevitz DF, McGuinness G, McCauley DI, Smith JP, Libby D, Pasmantier M, Vazquez M, Koizumi J, Flieder D, Altorki N, Miettinen OS. Early lung cancer action project: initial findings on repeat screenings. *Cancer* 2001 Jul; 92(1): 153-9.
- (68) Yankelevitz DF, Reeves AP, Kostis WJ, Zhao B, Henschke CI. Small pulmonary nodules: volumetrically determined growth rates based on CT evaluation. *Radiology* 2000 Oct;217(1):251-6.

- (69) Marten K, Auer F, Schmidt S, Kohl G, Rummeny EJ, Engelke C. Inadequacy of manual measurements compared to automated CT volumetry in assessment of treatment response of pulmonary metastases using RECIST criteria. *Eur Radiol* 2006 Apr;16(4):781-90.
- (70) Jennings SG, Winer-Muram HT, Tarver RD, Farber MO. Lung tumor growth: assessment with CT--comparison of diameter and cross-sectional area with volume measurements. *Radiology*. 2004 Jun;231(3):866-71.
- (71) Marten K, Funke M, Engelke C. Flat panel detector-based volumetric CT: prototype evaluation with volumetry of small artificial nodules in a pulmonary phantom. *J Thorac Imaging*. 2004 Jul;19(3):156-63.
- (72) Goo JM, Tongdee T, Tongdee R, Yeo K, Hildebolt CF, Bae KT. Volumetric measurement of synthetic lung nodules with multi-detector row CT: effect of various image reconstruction parameters and segmentation thresholds on measurement accuracy. *Radiology* 2005 Jun;235(3):850-6.
- (73) Ko JP, Rusinek H, Jacobs EL, Babb JS, Betke M, McGuinness G, et al. Small pulmonary nodules: volume measurement at chest CT--phantom study. *Radiology* 2003 Sep;228(3):864-70.
- (74) Wormanns D, Kohl G, Klotz E, Marheine A, Beyer F, Heindel W, et al. Volumetric measurements of pulmonary nodules at multi-row detector CT: in vivo reproducibility. *Eur Radiol* 2004 Jan;14(1):86-92.
- (75) Marten K, Auer F, Schmidt S, Kohl G, Rummeny EJ, Engelke C. Inadequacy of manual measurements compared to automated CT volumetry in assessment of treatment response of pulmonary metastases using RECIST criteria. *Eur Radiol*. 2006 Apr;16(4):781-90.
- (76) Bolte H, Riedel C, Jahnke T, Inan N, Freitag S, Kohl G, Heller M, Biederer J. Reproducibility of computer-aided volumetry of artificial small pulmonary nodules in ex vivo porcine lungs. *Invest Radiol*. 2006 Jan;41(1):28-35.
- (77) Honda O, Sumikawa H, Johkoh T, Tomiyama N, Mihara N, Inoue A, Tsubamoto M, Natsag J, Hamada S, Nakamura H. Computer-assisted lung nodule volumetry from multi-detector row CT: influence of image reconstruction parameters. *Eur J Radiol*. 2007 Apr;62(1):106-13.
- (78) Winer-Muram HT, Jennings SG, Meyer CA, Liang Y, Aisen AM, Tarver RD, McGarry RC. Effect of varying CT section width on volumetric measurement of lung tumors and application of compensatory equations. *Radiology*. 2003 Oct;229(1):184-94.

- (79) Zhao B, Schwartz LH, Moskowitz CS, Wang L, Ginsberg MS, Cooper CA, Jiang L, Kalaigian JP. Pulmonary metastases: effect of CT section thickness on measurement--initial experience. *Radiology*. 2005 Mar;234(3):934-9.
- (80) Wedegartner U, Lorenzen M, Nagel HD, Koops A, Weber C, Nolte-Ernsting C, et al. [Image quality of thin- and thick-slice MSCT reconstructions in low-contrast objects (liver lesions) with equal doses]. *Rofo* 2004 Nov;176(11):1676-82.
- (81) Kopka L, Rogalla P, Hamm B. [Multislice CT of the abdomen--current indications and future trends]. *Rofo* 2002 Mar;174(3):273-82.
- (82) Catalano C, Laghi A, Fraioli F, Pediconi F, Napoli A, Danti M, et al. Pancreatic carcinoma: the role of high-resolution multislice spiral CT in the diagnosis and assessment of resectability. *Eur Radiol* 2003 Jan;13(1):149-56.
- (83) Pannu HK, Horton KM, Fishman EK. Thin section dual-phase multidetector-row computed tomography detection of peritoneal metastases in gynecologic cancers. *J Comput Assist Tomogr* 2003 May;27(3):333-40.
- (84) Cohnen M, Vogt C, Aurich V, Beck A, Haussinger D, Modder U. Multi-slice CT-colonography in low-dose technique--Preliminary results. *Rofo* 2002 Jul;174(7):835-8.
- (85) Wessling J, Fischbach R, Domagk D, Lugerling N, Neumann E, Heindel W. Colorectal polyps: Detection with multi-slice CT colonography. *Rofo* 2001 Dec;173(12):1069-71.
- (86) Van HL, Van CE, Vergote I, Baert AL, Bellon E, Dupont P, et al. Size quantification of liver metastases in patients undergoing cancer treatment: reproducibility of one-, two-, and three-dimensional measurements determined with spiral CT. *Radiology* 1997 Mar;202(3):671-5.
- (87) Bland JM, Altman DG. Comparing two methods of clinical measurement: a personal history. *Int J Epidemiol* 1995;24 Suppl 1:S7-14.
- (88) Bland JM, Altman DG. Comparing methods of measurement: why plotting difference against standard method is misleading. *Lancet* 1995 Oct 21;346(8982):1085-7.
- (89) Jaffe CC. Measures of response: RECIST, WHO, and new alternatives. *J Clin Oncol*. 2006 Jul 10;24(20):3245-51.