

8 Literaturverzeichnis

- Agoulnik, I.Y., Cho, Y., Niederberger, C., Kieback, D.G. und Cooney, A.J. (1998) Cloning, expression analysis and chromosomal localization of the human nuclear receptor gene GCNF. *FEBS Lett.* **424**, 73-78
- Alting-Mees et al. (1994) *Strategies* **7**, 70-72
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D.J. (1990) Basic local alignment search tool *J. Mol. Biol.* **215**, 403-410
- Artelt et al. (1989) Vectors for efficient expression in mammalian fibroblastoid, myeloid and lymphoid cells via transfection or infection. *Gene* **68**, 213-219
- Auwerx, J. et al. (1999) A unified nomenclature system for the nuclear receptor superfamily. *Cell* **97**, 161-163
- Bailey, P.J., Dowhan, D.H., Franke, K., Burke, L.J., Downes, M. und Muscat, G.E. (1997) Transcriptional repression by COUP-TF II is dependent on the C-terminal domain and involves the N-CoR variant, RIP13delta1. *J. Steroid Biochem. Mol. Biol.* **63**, 165-174
- Baniahmad, A., Kohne, A.C. und Renkawitz, R. (1992) A transferable silencing domain is present in the thyroid hormone receptor, in the v-erbA oncogene product and in the retinoic acid receptor. *EMBO J.*, **11**, 1015-1023
- Barretino, D., Vivanco-Ruiz, M.d.M. und Stunnenberg, H.G. (1994) Characterization of the ligand-dependent transactivation domain of thyroid hormone receptor. *EMBO J.* **13**, 3039-3049
- Bartel, P.L., Chien, C.-T., Sternglanz R. & Fields, S. (1993) Using the two-hybrid system to detect protein-protein interactions. In *Cellular Interactions in Development: A Practical Approach*, D.A. Hartley, ed., Oxford University Press, Oxford, 153-179
- Bauer, J., Deely, D., Braman, J., Viola, J. & Weiner, M.P. (1992) *Strategies* **5**, 62-64

- Bauer. U.M., Schneider-Hirsch, S., Reinhardt, S., Benavente, R. und Maelicke A. (1998) The murine nuclear orphan receptor GCNF is expressed in the XY body of primary spermatocytes. *FEBS Lett.* **439**, 208-214
- Bauer. U.M., Schneider-Hirsch, S., Reinhardt, S., Pauly. T., Maus, A., Wang, F., Heiermann, R., Rentrop, M. und Maelicke A. (1997) Neuronal cell nuclear factor- a nuclear receptor possibly involved in the control of neurogenesis and neuronal differentiation. *Eur. J. Biochem.* **249**, 826-837
- Becker-Andre, M., Wiesenberg, I., Schaeren, W.N., Andre, E., Missbach, M., Saurat, J.H. und Carlberg, C. (1994) Pineal gland hormone melatonin binds and activates an orphan of the nuclear receptor superfamily. *J. Biol. Chem.* **269**, 28531-28534
- Behr JP, Demeneix B, Loeffler JP, et al. (1989) Efficient gene transfer into mammalian primary endocrine cells with lipopolyamine-coated DNA. *Proc.Natl. Acad. Sci. USA* **86**, 6982-6986
- Blanco, J.C.G., Minucci, S., Lu, J.M., Yang, X.J., Walker, K.K., Chen, H.W., Evans, R.M., Nakatani, Y und Ozato, K. (1998) The histone acetylase PCAF is a nuclear receptor coactivator. *Gene Dev.* **12**, 1638-1651
- Boi, Y., Adam, M., Rushmore, T. und Kennedy, B.P. (1993) Enantiomeric activation of the peroxisome proliferator-activated receptor. *J. Biol. Chem.* **268**, 5530-5534
- Borgmeyer, U. (1997) Dimeric binding of the mouse germ cell nuclear factor. *Eur. J. Biochem.* **244**, 120-127
- Bourgout, W., Ruff, M., Chambon, P., Gronemeyer, H. und Moras, D. (1995) Crystal structure of the ligand-binding domain of the human nuclear receptor RXR α . *Nature* **375**, 377-382
- Braat, A.K., Zandbergen, M.A., De Vries, E., Van Der Burg, B., Bogerd, J. und Goos, H. J. (1999) Cloning an expression of the zebrafish germ cell nuclear factor. *Mol. Reprod. Dev.* **53**, 369-375
- Bradford, M.M., (1976) A rapid and sensitive method for quantification of microgram quantities of protein utilizing the principle of protein dye-binding. *Anal. Biochem.* **72**, 246-264

- Bullock, W.O., Fernandez, J.M. & Short, J.M. (1987), *Biotechniques* **5**, 376-379
- Burbach, J.P.H., Da Silva, S.L., Cox, J.J. et al. (1994) Repression of estrogen-dependent stimulation of the oxytocin gene by COUP-TF I. *J. Biol. Chem.* **269**, 15046-15053
- Cavailles, V., Deuvois, S., L'Horset, F., Lopez, G., Hoarse, S., Kushner, P.J. und Parker, M.G. (1995) Nuclear factor RIP140 modulates transcriptional activation by the estrogen receptor. *EMBO J.* **14**; 3741-3751
- Chalfie, M., Tu, Y., Euskirchen, G., Ward, W. & Prasher, D.C. (1994) Green Fluorescent Protein as marker for gene expression. *Science* **263**, 802-805
- Chen, F., Cooney, A.J., Wang, Y., Law, B.W. und O'Malley, B.W. (1994) Cloning of a novel orphan receptor (GCNF) expressed during germ cell development. *Mol. Endocrinol.* **8**, 1434-1444
- Chen, J.D., Evans, R.M. (1995) A transcriptional co-repressor that interacts with nuclear hormone receptors. *Nature* **377**, 454-457
- Chittenden, T., Lupton, S. & Levine, A.J. (1989) Functional limits of oriP, the Epstein Barr Virus Plasmid Origin of Replication. *J. of Virology* **63**, 3016-3025
- Cooney, A.J., Katz, D., Hummelke, G.C. und Jackson, K.J. (1999) *Am. Zool.* **39**, 796-806
- Cooney, Al J., Hummelke, G.C., Herman, T., Chem, F. und Jackson, K. J. (1998) Germ cell nuclear factor is a response element specific repressor of transcription. *Biochem. Biophys. Res. Commun.* **245**, 94-100
- Cormack, B.P., Valdivia, R. & Falkow, S. (1996) FACS-optimized mutants of the green fluorescent protein (GFP). *Gene* **173**, 33-38
- Danielian, P.S., White, R., Lees, J.A. und Parker, M.G. (1992) Identification of a conserved region required for hormone dependent transcriptional activation by steroid hormone receptors *Embo J.* **11**, 1025-1033

- Danielian, P.S., White, R., Lees, J.A. und Parker, M.G. (1992) Identification of a conserved region required for hormone dependent transcriptional activation by steroid hormone receptors. *EMBO J.* **11**, 1025-1033
- Darimont, B.D., Wagner, R.L., Apriletti, J.W., Stallcup, M.R., Kushner, P.J., Baxter, J.D., Fletterick, R.J. und Yamamoto K.R. (1998) Structure and specificity of nuclear receptor-coactivator interactions. *Genes Dev.* **12**, 3343-3356
- Davis, I.J., Lau, L.F. (1994) Endocrine and neurogenic regulation of the orphan nuclear receptors Nur77 and Nurr-1 in the adrenal glands. *Mol. Cell. Biol.* **14**, 3469-3483
- Dreyer, C. and Ellinger-Ziegelbauer, H. (1996) Retinoic acid receptors and nuclear orphan receptors in the development of *Xenopus laevis*. *Int. J. Dev. Biol.* **40**, 255-262
- Enmark, E. und Gustafsson, J.-A. (1996) Orphan nuclear receptors- the first eight years. *Mol. Endocrinol.* **10**, 1293-1307
- Evans, R.M. (1988) The steroid and thyroid hormone receptor superfamily. *Science* **240**, 889-895
- Felgner, P.L. & Holm, M. (1989) Cationic liposome mediated transfection. *Focus* **11**, 21-25
- Felgner, P.L., Gadek, T.R., Holm, M., Roman, R., Chan, H.W., Wenzel, M., Northrop, J.P., Ringold, G.M. & Danielsen, M. (1987) Lipofection: a high efficient, lipid-mediated DNA-transfection procedure. *Proc. Natl. Acad. Sci. USA* **84**, 7413 ff
- Feng, W., Ribeiro, R.C., Wagner, R.L., Nguyen, H., Apriletti, J.W., Fletterick, R.J., Baxter, J.D., Kushner, P.J. und West, B.L. (1998) Hormone-dependent coactivator binding to a hydrophobic cleft on nuclear receptors. *Science* **280**, 1747-1749
- Forman, B.M., Goode, E., Chen, J., Oro, A.E., Bradley, D.J., Perlmann, T., Noonan, D.J., Burka, L.T., McMorris, T., Lamph, W.W., Evans, R.M. und Weinberger, C. (1995) Identification of a nuclear receptor that is activated by farnesol metabolites. *Cell* **81**, 687-693

- Freedman, L.P. (1999) Increasing the complexity of coactivation in nuclear receptor signalling. *Cell* **97**, 5-8
- Funder, J.W. (1993) Mineralocorticoids, glucocorticoids, receptors and response elements. *Science* **259**, 1132-1133
- Gietz, D., St. Jean, A., Woods, R.A. & Schiestl, R.H. (1992) Improved method for high efficiency transformation of intact yeast cells. *Nucleic Acids Res.* **20**, 1425
- Glass, C.K. (1994) Differential recognition of target genes by nuclear receptor monomers, dimers, and heterodimers. *Endocr. Rev.* **15**, 391-407
- Glass, C.K. und Rosenfeld, M.G. (2000) The coregulator exchange in transcriptional functions of nuclear receptors. *Genes and Dev.* **14**, 121-141
- Göttlicher, M., Demoz, A., Svensson, D., Tollet, P., Berge, R.K. und Gustafsson, J.-A. (1992) Fatty acids activate a chimera of the clofibrilic acid-activated receptor and the glucocorticoid receptor. *Proc. Nat. Acad. Sci. USA* **89**, 4653-4657
- Green, S. und Chambon, P. (1988) Nuclear receptors enhance our understanding of transcription regulation. *Trends Genet.* **4**, 309-314
- Greschik, H. und Schüle, R. (1998) Germ cell nuclear factor: an orphan receptor with unexpected properties. *J. Mol. Med.* **76**, 800-810
- Gronemeyer, H. (1991) Transcription activation by estrogen and progesterone receptors. *A. Rev. Genet.* **25**, 89-123
- Gronemeyer, H. und Laudet, V. (1995) Nuclear receptors. *Protein profile* **2**, 1173-1308
- Gronemeyer, H. und Moras, D. (1995) How to finger DNA. *Nature* **377**, 397-403
- Haendler, B., Hofer-Warbinek, R. und Hofer, E. (1987) Complementary DNA for human T-cell cyclophilin. *EMBO J.* **6**, 947-950
- Hanahan, D. (1983) Studies on transformation of *Escherichia coli* with Plasmids. *J. Mol. Biol.* **166**, 557 ff

- Harding, H. und Lazar, M. (1993) The orphan receptor Rev-ErbA α activates transcription via a novel response element. *Mol. Cell. Biol.* **13**, 3113-3121
- Harper, J.W., Adami, G.R., Wei, N., Keyomarsi, K. & Elledge S.J. (1993) The p21 cdk-interacting protein Cip-1 is a potent inhibitor of G1 cyclin-dependent kinases. *Cell* **75**, 805-816
- Heidaran, M.A. und Kistler, W.S. (1987) Transcriptional and translational regulation of the mRNA for transition protein 1 in the rat. *J. Biol. Chem.* **269**, 13309-13315
- Heinzel, T., Lavinsky, R.M., Mullen, T.M., Soderstrom, M., Laherty, C.D., Torchia, J., Yang, W.M., Brard, G., Ngo, S.D., Davie, J.R. et al. (1997) A complex containing N-CoR, mSin3 and histone deacetylase mediates transcriptional repression. *Nature* **387**, 43-48
- Heinzer, C., Süsens, U., Schmitz, T.P. und Borgmeyer, U. (1998) Retinoids induce differential expression and DNA binding of the mouse germ cell nuclear factor in P19 embryonal carcinoma cells. *Biol. Chem.* **379**, 349-359
- Hirose, T., O'Brian, D.A., Jetten, A.M. (1995) RTR: a new member of the nuclear receptor superfamily that is highly expressed in murine testis. *Gene* **152**, 247-251
- Hollenberg, S.M. und Evans, R.M. (1988) Multiple and cooperative trans-activation domains of the human glucocorticoid receptor. *Cell* **55**, 899-906
- Hörlein, A.J., Naar, A.M., Heinzel, T., Torchia, J., Gloss, B., Kurokawa, R., Ryan, A., Kamel, Y., Soderstrom, M., Glass, C.K., Rosenfeld, M.G. (1995) Ligand-independent repression by the thyroid hormone receptor mediated by a nuclear receptor co-repressor. *Nature* **377**, 451-454
- Hu, I. und Lazar, M.A. (2000) Transcriptional Repression by Nuclear Hormone Receptors. *Trends Endocrinol. Metab.* **11**, 6-10
- Hu, X. und Lazar, M.A. (1999) The CoRNR motif controls the recruitment of corepressors by nuclear hormone receptors. *Nature* **402**, 93-96

- Hummelke, G.C., Meistrich, M.L. und Cooney, A.J.(1998) Mouse protamine genes are candidate targets for the novel orphan nuclear receptor, germ cell nuclear factor. *Mol. Reprod. Dev.* **50**, 396-405
- Izumao, S. und Mahdavi, V. (1988) Thyroid hormone receptor α isoforms generate by alternative splicing differentially activate myosin HC gene transcription. *Nature* **334**,539-542
- Jensen, F.C. (1964) . *Proc. Nat. Acad. Sci.* **53**, 53
- Jenster, G. (1998) Coactivators and corepressors as mediators of nuclear receptor function: An update. *Mol. Cell. Endocrinol.* **143**, 1-7
- Jenster, G., Spencer, T.E., Burcin M.M., Tsai, S.Y., Tsai, M.J. und O'Malley, B.W. (1997) Steroid receptor induction of gene transcription: A two-step model. *Proc. Natl. Acad. Sci. USA* **94**, 7879-7884
- Johnson, P.A., Peschon, J.J., Yelick, P.C., Palmiter, R.D. und Hecht, N.B. (1988) Sequence homologies in the mouse protamine 1 and 2 genes. *Biochem. Biophys. Acta* **950**, 45-53
- Joos, T.O., David, R., Dreyer, C. (1996) xGCNF, a nuclear orphan receptor is expressed during neurulation in *Xenopus laevis*. *Mech. Dev.* **60**, 45-57
- Kadonaga, J.T. (1998) Eukaryotic transcription: an interlaced network of transcription factors and chromatin-modifying machines. *Cell* **92**, 307-313
- Kapelle, M., Krätzschar, J., Husemann, M. und Schleuning, W.D. (1997) cDNA cloning of two closely related forms of human germ cell nuclear factor (GCNF). *Biochim. Biophys. Acta* **1352**, 13-17
- Kastner, P., Mark, M. und Chambon, P. (1995) Nonsteroid nuclear receptors: what are genetic studies telling us about their role in real life? *Cell* **83**, 859-869
- Katz, D., Niederberger, C., Slaughter, G.R. und Cooney, A.J. (1997) Characterization of germ cell-specific expression of the orphan nuclear receptor, germ cell nuclear factor. *Endocrinology* **138**, 4364-4372

- Kleene, K.C. (1996) Patterns of translational regulation in the mammalian testis. *Mol. Rep. Dev.* **43**, 268-281
- Kleene, K.C., Distel, R.J. und Hecht, N.B. (1984) Translational regulation and deadenylation of a protamine mRNA during spermatogenesis in the mouse. *Dev. Biol.* **105**, 71-79
- Kliwer, S.A., Umesono, K., Noonan, D.J., Heyman, R.A. und Evans, R.M. (1992) Convergence of 9-cis retinoic acid and peroxisome proliferator signalling pathways through heterodimer formation of their receptors. *Nature* **358**, 771-774
- Kliwer, S.A., Umesonso, K., Mangelsdorf, D.J. und Evans, R.M. (1992) Retinoid X receptor interacts with nuclear receptors in retinoic acid, thyroid hormone and vitamin D3 signalling. *Nature* **355**, 446-449
- Korzus, E., Torchia, J., Rose, D.W., Xu, L., Kurokawa, R., McInerney, E.M., Mullen, T.M., Glass, C.K. und Rosenfeld, M.G. (1998) Transcription factor-specific requirements for coactivators and their acetyltransferase functions. *Science* **279**, 703-707
- Kostrouch, Z., Kostrouchova, M. und Rall, J.E. (1995) Steroid/thyroid hormone receptor genes in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. USA* **92**, 156-159.
- Kozak, M. (1989) The scanning model for translation: an update. *J. Biol. Chem.* **108**, 229-241
- Krätzschmar, J., Haendler, B., Eberspächer, U., Roosterman, D., Donner, P., Schleuning, W.-D. (1996) The human cysteine-rich secretory protein (CRISP) family. Primary structure and tissue distribution of CRISP-1, CRISP-2 and CRISP-3. *Eur. J. Biochem.* **236**, 827-836
- Laemmly, U.K. (1970) Cleavage of structural proteins during assembly of the head of bacteriophage T4. *Nature* **227**, 680-685
- Lala, D.S., Rice, D.A. und Parker, K.L. (1992) Steroidogenic factor 1, a key regulator of steroidogenic enzyme expression is the mouse homolog of fushi-tarazu factor 1. *Mol. Endocrinol.* **6**, 1249-1258

- Laudet, V. (1997) Evolution of the nuclear receptor superfamily: early diversification from an ancestral orphan receptor. *J. Mol. Endocrinol.* **19**, 207-226
- Laudet, V., Hanni, C., Coll, J., Catzeflis, F. und Stehelin, D. (1992) Evolution of the nuclear receptor gene superfamily. *EMBO J.* **11**, 1003-1013
- Lazar, M.A., Hodin, R.A. und Chin, W.W. (1989) Human carboxyl-terminal variant of α -type *c-erbA* inhibits trans-activation by thyroid hormone receptors without binding thyroid hormone. *Proc. Natl. Acad. Sci. USA* **86**, 7771-7774
- Lei, W., Hirose, T., Zhang, L.X., Adachi, H., Spinella, M.J., Dmitrovsky, E. und Jetten, A.M. (1997) Cloning of the human orphan receptor germ cell nuclear factor/retinoid receptor-related testis-associated receptor and its differential regulation during embryonal carcinoma cell differentiation. *J. Mol. Endocrinol.* **18(2)**, 167-176
- Leid, M., Kastner, P. und Chambon, P. (1992) Multiplicity generates diversity in the retinoic acid signalling pathways. *Trends Biochem. Sci.* **17**, 427-433
- Levin, A.A., Stuzenbecker J.L., Kamer, S., Bosakowski, T., Huselton, C., Allenby, G., Speck, J., Kratzeisen, C.L., Rosenberger M., Lovey, A., Grippo, J.F. (1992) 9-cis Retinoic Acid stereoisomer binds and activates the nuclear receptor RXR α . *Nature* **355**, 359-361
- Macpherson, I.A. und Stoker M.G.P. (1962) *Virology*, **16**, 147
- Mangelsdorf, D.J. und Evans, R.M. (1995) The RXR heterodimers and orphan receptors. *Cell* **83**, 841-850
- Mangelsdorf, D.J. und Willy, P.J. (1998) Nuclear orphan receptors: The search for novel ligands and signalling pathways. In hormones and signalling, O'Malley, B., ed.: Academic Press, pp. 307-358
- Mangelsdorf, D.J., Thummel, C., Beato, M., Herrlich, P., Schütz, G., Umesono, K., Blumberg, B., Kastner, P., Mark, M., Chambon, P. und Evans, R.M. (1995) The nuclear receptor superfamily: the second decade. *Cell* **83**, 835-839

- McBurney, M.W. und Rogers, B.J. (1982) Isolation of male embryonic carcinoma cells and their chromosome replication patterns. *Dev. Biol.*, **89**, 503-508
- McEwan, I., Dahlman-Wright, K., Almlöf, T., Ford, J., Wright, A. und Gustafsson, J.-A. (1995) Mechanisms of transcriptional activation by nuclear receptors: studies on the human glucocorticoid receptor τ_1 transactivation domain. *Mutat. Res.* **333**, 15-22
- McKenna, N.J., Lanz, R.B. und O'Malley, B.W. (1999) Nuclear receptor coregulators: cellular and molecular biology. *Endocrine reviews*, **20(3)**, 321-344
- McKenna, N.J., Xu, J., Nawaz, Z., Tsai, S.Y., Tsai, M.J. und O'Malley, B.W. (1999) Nuclear receptor coactivators: multiple enzymes, multiple complexes, multiple functions. *J. Steroid Biochem. Mol. Biol.* **69**, 3-12
- Meier, C.A. (1997) Regulation of gene expression by nuclear hormone receptors. *J. recep. sign. transd. res.* **17**, 319-335
- Meistrich, M.L. (1988) Histone and basic nuclear protein transitions in mammalian spermatogenesis. In: Hnilica, L., Stein, G., Stein, J., eds. (1988) Histones and other basic nuclear proteins. Boca Raton: CRC Press; 165-182
- Meistrich, M.L. Histone and basic nuclear protein transitions in mammalian spermatogenesis. In: Hnilica, L., Stein, G., Stein, J., eds. (1988) Histones and other basic nuclear proteins. *Boca Raton: CRC Press*, 165-182
- Nagy, L., Kao, H.Y., Chakravarti, D., Lin, R.J., Hassig, C.A., Ayer, D.E., Schreiber, S.L. und Evans, R.M. (1997) Nuclear receptor repression mediated by a complex containing SMRT, mSin3A and histone deacetylase. *Cell* **89**, 373-380
- Nagy, L., Kao, H.Y., Love, J.D., Li, C., Banayo, E., Gooch, J.T., Krishna, V., Chatterjee, K., Evans, R.M. und Schwabe, J.W. (1999) Mechanism of corepressor binding and release from nuclear hormone receptors. *Genes and Dev.* **13**, 3209-3216
- Napoli, J.L., und Race, K.R. (1987) The biosynthesis of retinoic acid from retinol by rat tissues *in vitro*. *Arch. Biochem. Biophys.* **255**, 95-101

- Nolte, R.T., Wisely, G.B., Westin, S., Cobb, J.E., Lambert, M.H., Kurokawa, R., Rosenfeld, M.G., Willson, T.M., Glass, C.K. und Milburn, M.V. (1998) Ligand binding and co-activator assembly of the peroxisome proliferator-activated receptor- γ . *Nature* **395**, 137-143
- Ogryzko, V.V., Schiltz, R.L., Russanova, V., Howard, B.H. und Nakatani, Y. (1996) The transcriptional coactivators p300 and CBP are histone acetyltransferases. *Cell* **87**, 953-959
- Parker, M.G. (1993) Steroid and related receptors. *Curr. Opin. Cell Biol.* **5**, 499-504
- Pratt, W.B. (1993) The role of heat shock proteins in regulating the function, folding and trafficking of the glucocorticoid receptor. *J. Biol. Chem.* **268**, 21455-21458
- Renaud, J.P et al. (1995) Crystal structure of the RAR γ ligand-binding domain bound to *all-trans* retinoic acid. *Nature* **378**, 681-689
- Rizutto, R., Brini, M., Pizzo, P., Murgia, M. und Pozzan, T. (1995) Chimeric green fluorescent protein as a tool for visualizing subcellular organelles in living cells. *Curr. Biol.* **5**, 635-642
- Sande, S. und Privalsky, M.L. (1996) Identification of TRACs (T3 receptor-associating cofactors), a family of cofactors that associate with, and modulate the activity of, nuclear hormone receptors. *Mol. Endocrinol.* **10**, 813-825
- Sanger, F., Nickler, S. und Coulson, A.R. (1977) DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**, 5463-5467
- Schäfer, M., Nayernia, K., Engel, W. und Schäfer, U. (1995) Translational control in spermatogenesis. *Dev. Bio.* **172**, 344-352
- Schmitz, T.P., Süsens, U. und Borgmeyer, U. (1999) DNA binding, protein interaction and differential expression of the human germ cell nuclear factor. *Biochim. Biophys. Acta* **1446**, 173-180
- Seol, W., Mahon, M.J., Lee, Y.K. und Moore, D.D (1996) Two receptor interacting domains in the nuclear hormone receptor corepressor RIP13/N-CoR. *Mol. Endocrinol.* **10**, 1646-1655

- Shigeta, H., Zuo, W., Yang, N., DiAugustine, R. und Teng, C.T. (1997) The mouse estrogen receptor-related orphan receptor alpha 1: molecular cloning and estrogen responsiveness. *J. Mol. Endocrinol.* **19**, 299-309
- Short, J.M., Fernandez J.M., Sorge, J.A. und Huse W.D. (1988) λ -ZAP: a bacteriophage λ expression vector with *in vivo* excision properties. *Nucleic Acids Res.* **16**, 7583-7600
- Sladek, F.M., Zhong, W.M., Lai, E. und Darnell, J.J. (1990) Liver-enriched transcription factor HNF-4 is a novel member of the steroid hormone receptor superfamily. *Genes Dev.* **4**, 2353-2365
- Smith, D.B. und Johnson, K.S. (1988) Single-step purification of polypeptides expressed in *Escherichia coli* as fusions with glutathione S-transferase. *Gene* **67**, 31-40
- Spencer, T.E., Jenster, G., Burcin, M.M., Allis, C.D., Zhou, J.X., Mizzen, C.A., McKenna, M.J. Onate. S.A., Tsai, S.Y., Tsai, M.J. und O'Malley, B.W. (1997) Steroid receptor coactivator-1 is a histone acetyltransferase. *Nature* **389**, 194-198
- Struhl, K. (1998) Histone acetylation and transcriptional regulatory mechanisms. *Genes Dev.* **12**, 599-606
- Süsens, U. und Borgmeyer, U. (1996) Characterization of the human germ cell nuclear factor gene. *Biochim. Biophys. Acta* **1309(3)**, 179-182
- Süsens, U., Aguiluz, J.B., Evans, R.M. und Borgmeyer, U. (1997) The germ cell nuclear factor mGCNF is expressed in the developing nervous system. *Cev. Neurosci.* **19**, 410-420
- Takehita, A., Yen, P.M., Misiti, S., Cordona, G.R., Liu, Y. und Chin, W.W. (1996) Molecular cloning and properties of a full-length putative thyroid hormone receptor co-activator. *Endocrinology* **137**, 3594-3597
- Umesono, K Murakami, K.K., Thompson, C.C. und Evans, R.M. (1991) Direct repeats as selective response elements for the thyroid hormone, retinoic acid and vitamin D3 receptors. *Cell* **65**, 1255-1266

- Umesono, K. und Evans, R.M. (1989) Determinants of target gene specificity for steroid/thyroid hormone receptors. *Cell* **57**, 1139-1146
- Venables, J.P. und Eperon, I.C. (1999) The roles of RNA-binding proteins in spermatogenesis and male fertility. *Curr. Opin. Gen. Dev.* **9**, 346-354
- Voegel, J.J., Heine, M.J.D., Zechel, C., Chambon, P., Gronemeyer, H. (1996) TIF2, a 160 kDa transcriptional mediator for the ligand-dependent activation function AF-2 of nuclear receptors. *EMBO J.* **15**; 3667-3675
- Weinberger, C. und Bradley, D.J. (1990) Gene regulation by receptors binding lipid-soluble substances. *Ann. Rev. Physiol.* **52**, 823-844
- Wilson, T.E., Fahmer, T.J. und Milbrandt, J. (1993) The orphan receptors NGFI-B and steroidogenic factor 1 establish monomer binding as a third paradigm of nuclear receptor-DNA interaction. *Mol. Cell. Biol.* **13**, 5794-5804
- Wong, J., Shi, Y.-B. und Wolffe, A.P. (1995) A role for nucleosome assembly in both silencing and activation of the *Xenopus* TR β A gene by the thyroid hormone receptor. *Genes Dev.* **9**, 2696-2711
- Wong, J., Shi, Y.-B. und Wolffe, A.P. (1997) Determinants of chromatin disruption and transcriptional regulation instigated by the thyroid hormone receptor: hormone-regulated chromatin disruption is not sufficient for transcriptional activation
- Wurtz, J.-M., Bourguet, J.-P., Renaud, J.-P., Vivat, V., Chambon, P., Moras, D. und Gronemeyer, H. (1996) A canonical structure for the ligand-binding domain of nuclear receptors. *Nat. Struct. Biol.* **3**, 87-94
- Xu, L., Glass, C.K. und Rosenfeld, M.G. (1999) Coactivator and corepressor complexes in nuclear receptor function. *Curr. Opin. Genet. Dev.* **9**, 140-147
- Yan, Z.H., Medvedev, A., Hirose, T., Gotoh, H. und Jetten, A.M. (1997) Characterization of the response element and DNA binding properties of the nuclear orphan receptor Germ Cell Nuclear Factor/Retinoid Receptor-related testis-associated receptor. *J. Biol. Chem.* **272**, 10565-10572

- Yanisch-Perron, C., Vieira, J. und Messing, J. (1985) Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* **33**, 103-119
- Yu, V.C., Delsert, C., Andersen, B., Holloway, J.M., Devary, O.V., Naar, A.M., Kim, S.Y., Boutin, J.M., Glass, C.K. und Rosenfeld, M.G. (1991) RXR β : a coregulator that enhances binding of retinoic acid, thyroid hormone and vitamin D receptors to their cognate response elements. *Cell* **67**, 1251-1266
- Zamir, I., Harding, H.P., Atkins, G.B., Horlein, A., Glass, C.K., Rosenfeld, M.G. und Lazar M.A. 1996 Anuclear hormone receptor corepressor mediates transcriptional silencing by receptors with different repression domains. *Mol. Cell. Biol.* **16**, 5458-5465
- Zanaria, E., Muscatelli, F., Bardoni, B., Strom, T.M., Guioli, S., Guo, W., Lalli, E., Moser, C., Walker, A.P., McCabe, E.R.B., Meitinger, T., Monaco, A.P., Sassone-Corsi, P. und Camerino, G. (1994) An unusual member of the nuclear hormone receptor superfamily responsible for X-linked adrenal hypoplasia congenita. *Nature* **372**, 635-641
- Zhang, J., Hu, X. und Lazar, M.A. (1999) A novel role for helix 12 of retinoid X receptor in regulating repression. *Mol. Cell Biol.* **19**, 6448-6457
- Zhang, Y.L., Akmal, K.M., Tsuruta, J.K., Shang, Q., Hirose, T., Jetten, A.M., Kim, K. H. und O' Brian, D. A. (1998) Expression of germ cell nuclear factor GCNF (RTR) during spermatogenesis. *Mol. Reprod. Dev.* **50**, 93-102