

## 6. Literatur

Ala, Y.; Morin, D.; Mouillac, B.; Sabatier, N.; Vargas, R.; Cotte, N.; Dechaux, M.; Antignac, C.; Arthus, M.F.; Lonergan, M.; Turner, M.S.; Balestre, M.N.; Alonso, G.; Hibert, M.; Barberis, C.; Hendy, G.N.; Bichet, D.G.; Jard, S.:

Functional Studies of Twelve Mutant V2 Vasopressin Receptors Related to Nephrogenic Diabetes insipidus: Molecular Basis of a Mild Clinical Phenotype.

J.Am.Soc.Nephrol. 9 (1998) 1861-1872

Albertazzi, E.; Zancheta, D.; Barbier, P.; Faranda, S.; Frattini, A.; Vezzoni, P.; Procaccio, M.; Bettinelli, A.; Guzzi, F.; Parenti, M.; Chini, B.:

Nephrogenic Diabetes Insipidus: Functional Analysis of New AVPR2 Mutations Identified in Italian Families.

J. Am.Soc.Nephrol. 11 (2000) 1033-1043

Alfaidy, N. ; Blot-Chabaud, M. ; Bonvalet, J.P. ; Farman, N.:

Vasopressin potentiates mineralocorticoid selectivity by stimulating 11 beta hydroxysteroid dehydrogenase in rat collecting duct

J.Clin.Invest. 100 (1997) 2437-2442

Andersen-Beckh, B.; Dehe, M.; Schülein, R.; Wiesner, B.; Rutz, C.; Liebenhoff, U.; Rosenthal, W.; Oksche, A.:

Polarized expression of the vasopressin V2 receptor in Madin Darby canine kidney cells.

Kidney Int. 56 (1999) 517-527

Anfinsen, C.B.:

Principles that govern the folding of protein chains.

Science 181 (1973) 223-230

Aslund, F.; Berndt, K.D.; Holmgren, A.:

Redox Potentials of Glutaredoxins and Other Thiol-Disulfide Oxidoreductases of the Thioredoxin Superfamily Determined by Direct Protein-Protein Redox Equilibria. J.Biol.Chem. 272 (1997) 30780-30786

Asman, Y.W.; Dong, M.; Ganguli, S.; Hadac, E.M.; Miller, L.J.:

Structural Insights into the Amino-Terminus of the Secretin Receptor: I. Status of Cysteine and Cystine Residues.

Mol.Pharmacol. 58 (2000) 911-919

Attramadal, H.; Arriza, J. L.; Aoki, C.; Dawson, T. M.; Codina, J.; Kwatra, M. M.; Snyder, S. H.; Caron, M. G.; Lewkowitz, R. J.:

Beta-arrestin2, a novel member of the arrestin/beta-arrestin gene family.

J. Biol. Chem. 267 (1992) 17882-17890

Bai, M.; Trivedi, S.; Brown, E.M.:

Dimerization of the extracellular calcium-sensing receptor (CaR) on the cell surface of CaR-transfected HEK293 cells.

J.Biol.Chem. 273 (1998) 23605-23610

Bargmann, C.I.:

Neurobiology of the *Caenorhabditis elegans* genome.

Science 282 (1998) 2028-2033

Bichet, D.G.; Birnbaumer, M.; Lonergan, M.; Arthus, M.F.; Rosenthal, W.; Goodyer, P.; Nivet, H.; Benoit, S.; Giampietro, P.; Simonetti, S.; Fish, A.; Whitley, C.B.; Jaeger, P.; Gertner, J.; New, M.; DiBona, F.J.; Kaplan, B.S.; Robertson, G.L.; Hendy, G.N.; Fujiwara, T.M.; Morgan, K.:

Nature and Recurrence of AVPR2 Mutations in X-linked nephrogenic Diabetes insipidus.  
Am.J.Hum.Genet. 55 (1994) 278-286

Birnbaumer, M.; Seibold, A.; Gilbert, S.; Ishido, M.; Barberis, C.; Antaramian, A.; Brabet, P.; Rosenthal, W.:

Molecular cloning of the receptor for human antidiuretic hormone.

Nature 357 (1992) 333-335

Bockaert, J.; Pin, J.P.:

Molecular tinkering of G protein-coupled receptors: an evolutionary success.

EMBO J. 18 (1999) 1723-1729

Bose, S.; Seetharam, B.:

Effect of Disulfide Bonds of Transcobalamin II Receptor on its Activity and Basolateral Targeting in Human Intestinal Epithelial Caco-2 Cells.

J.Biol.Chem. 272 (1997) 20920-20928

Bourne, H.:

How receptors talk to trimeric G proteins.

Current Opinion in Cell Biology 9 (1997) 134-142

Brandt, W.; Golbraikh, A.; Täger, M.; Lendeckel, U.:

A molecular mechanism for the cleavage of a disulfide bond as the primary function of agonist binding to G-protein-coupled receptors based on theoretical calculations supported by experiments.

Eur.J.Biochem. 261 (1999) 89-97

Charlton, J.A., Baylis, P.H.:

Stimulation of rat renal medullary Na<sup>+</sup>/K<sup>(+)</sup>-ATPase by arginine vasopressin is mediated by the V2 receptor.

J.Endocrinol. 127 (1990) 213-216

Cook, J.V.F.; Eidne, K.A.:

An Intramolecular Disulfide Bond between Conserved Extracellular Cysteines in the Gonadotropin-Releasing Hormone Receptor Is Essential for Binding and Activation.  
Endocrinology 138 (1997) 2800-2806

Cotte, N.; Balestre, M.N.; Phalipou, S.; Hibert, M.; Manning, M.; Barberis, C.; Mouillac, B.: Identification of Residues Responsible for the Selective Binding of Peptide Antagonists and Agonists in the V2 Vasopressin Receptor.  
J.Biol.Chem. 273 (1998) 29462-29468

Creighton, T.E.: (1993) Proteins, Structures and Molecular Properties, W.H. Freeman and Company, New York, p.98

D'Angelo, D.D.; Eubank, J.J.; Davis, M.G.; Dorn, G.W.:  
Mutagenic Analysis of Platelet Thromboxane Receptor Cysteines. Roles in ligand binding and receptor-effector coupling.  
J.Biol.Chem. 271 (1996) 6233-6240

Davidson, F.F.; Loewen, P.C.; Khorona, H.G.:  
Structure and function in rhodopsin : Replacement by alanine of cysteine residues 110 and 187, components of a conserved disulfide bond in rhodopsin, affects the light-activated metarhodopsin II state.  
Proc.Natl.Acad.Sci.USA 91 (1994) 4029-4033

Devreotes, P.N.:  
G protein-linked signaling pathways control the developmental program of *Dictyostelium*. Neuron 12 (1994) 235-241

Dixon, R.A.; Sigal, I.S.; Candelore, M.R.; Register, R.B.; Scattergood, W.; Rands, E.; Strader, C.D.:  
Structural features required for ligand binding to the beta-adrenergic receptor.  
EMBO J. 6 (1987) 3269-3275

Dohlman, H.G.; Thorner, J.; Caron, M.G.; Lefkowitz, R.J.:  
Model systems for the study of seven-transmembrane-segment receptors.  
Ann.Rev.Biochem. 60 (1991) 653-688

Elling, C.E.; Raffetseder, U.; Nielsen, S.M.; Schwartz, T.W.:  
Disulfide Bridge Engineering in the Tachykinin NK<sub>1</sub> Receptor.  
Biochemistry 39 (2000) 667-675

Ellis, R.J.:  
The general concept of molecular chaperones.  
Philos.Trans.R.Soc.Lond.B Biol. Sci. 339 (1993): 257-261

Farrens, D.L.; Altenbach, C.; Yang, K.; Hubbell, W.L.; Khorona, H.G.:  
Requirement of rigid-body motion of transmembrane helices for light activation of rhodopsin. Science 274 (1996) 768-770

Freedman, R.B.; Hirst, T.R.; Tuite, M.F.:  
Protein Disulphide Isomerase: Building Bridges in Protein Folding.  
Trends Biochem. Sci. 19 (1994) 331-336

Gilbert, H.F.: (1994) The formation of native disulfide bonds, in Pain, R.H. (Ed.): Mechanism of Protein folding,  
IRL Press, p.81-103

Glocker, M.O.; Arbogast, B.; Milley, R.; Cowgill, C.; Deinzer, M.L.:  
Disulfide linkages in the in vitro refolded intermediates of recombinant human macrophage-colony-stimulating factor: Analysis of the sulfhydryl alkylation of free cysteine residues by fast-atom bombardment mass spectrometry.  
Proc.Natl.Acad.Sci. USA 91 (1994) 5868-5872

Hwa, J.; Reeves, P.J.; Klein-Seetharaman, J.; Davidson, F.; Khorona, H.G.:  
Structure and function in rhodopsin: Further elucidation of the role of the intradiscal cysteines, Cys-110, -185, and -187, in rhodopsin folding and function.  
Proc.Natl.Acad.Sci. 96 (1999) 1932-1935

Ji, T.H.; Grossmann, M.; Ji, I.:  
G Protein-coupled Receptors. I. Diversity of Receptor-Ligand Interactions.  
J.Biol.Chem. 273 (1998) 17299-17302

Jordan, B.A.; Devi, L.A.:  
G-protein-coupled receptor heterodimerization modulates receptor function.  
Nature 399 (1999) 697-700

Kaiser, C.; Ferro-Novick,S.:  
Transport from the endoplasmic reticulum to the Golgi  
Curr.Opin.Cell.Biol 10 (1998) 477-482

Karnik, S.S.; Sakmar, T.P.; Chen, H.B.; Khorona, H.G.:  
Cysteine residues 110 and 187 are essential for the formation of correct structure in bovine rhodopsin.  
Proc.Natl.Acad.Sci.USA 85 (1988) 8459-8463

Kim, G.H.; Ecelbarger, C.A.; Mitchell, C.; Packer, R.K.; Wade, J.B.; Knepper, M.A.:  
Vasopressin increases Na-K-2Cl cotransporter expression in thick ascending limb of Henle's loop.  
Am. J. Physiol. 276 (Renal Physiol. 45) (1999) F96-F103

Klußmann, E.; Maric, K.; Rosenthal, W.:  
The mechanisms of aquaporin control in the renal collecting duct.  
Rev. Physiol. Biochem. Pharmacol. 141 (2000) 33-95

Kojro,E.; Eich, P.; Gimpl, G.; Fahrenholz, F.:  
Direct Identification of an Extracellular Agonist Binding Site in the Renal V2 Vasopressin Receptor.

Biochemistry 32 (1993) 13537-13544

Kono, M.; Yu, H.; Oprian, D.D.:  
Disulfide Bond Exchange in Rhodopsin.  
Biochemistry 37 (1998) 1302-1305

Krause, G.; Hermosilla, R.; Oksche, A.; Rutz, C.; Rosenthal, W.; Schülein, R.:  
Molecular and conformational features of a transport-relevant domain in the C-terminal tail of the  
vasopressin V(2) receptor.  
Mol. Pharmacol. 57 (2000) 232-242

Kuner, R.; Kohr, G.; Grunewald, S.; Eisenhardt, G.; Bach, A.; Kornau, H.C.:  
Role of heteromer formation in GABA<sub>B</sub> receptor function.  
Science 283 (1999) 74-77

Kurtenbach, E.; Curtis, C.A.M.; Pedder, E.K.; Aitken, A.; Harrison, A.C.M.; Hulme, E.C.:  
Muscarinic Acetylcholine Receptors. Peptide sequencing identifies residues involved in antagonist  
binding and disulfide bond formation.  
J.Biol.Chem. 265 (1990) 13702-13708

Lomize, A.L.; Pogozheva, I.D.; Mosberg, H.I.:  
Structural organization of G-protein-coupled receptors.  
J.Comput.Aided Mol.Des. 13 (1999) 325-353

Maiti, N.R.; Sturewicz, W.K.  
The Role of Disulfide Bridge in the Folding and Stability of the Recombinant Human Prion Protein.  
J.Biol.Chem. 276 (2001) 2427-2431

Moro, S.; Hoffmann, C.; Jakobson, K.A.:  
Role of the Extracellular Loops of G Protein-Coupled Receptors in Ligand Recognition: A  
Molecular Modeling Study of the Human P2Y<sub>1</sub> Receptor.  
Biochemistry 38 (1999) 3498-3507

New, D.C.; Wong, J.T.:  
The evidence for G-protein-coupled receptors and heterotrimeric G proteins in protozoa and  
ancestral metazoa.  
Biol.Signals Recept. 7 (1998) 98-108

Noda, K.; Saad, Y.; Graham, R.M.; Karnik, S.S.:  
The High Affinity State of the  $\beta_2$ -Adrenergic Receptor Requires Unique Interactions between  
Conserved and Non-conserved Extracellular Loop Cysteines.  
J.Biol.Chem. 269 (1994) 6743-6752

Oakley, R.H.; Laprote, S.A.; Holt, J.A.; Caron, M.G.; Barak, L.S.:  
Differential affinities of visual arrestin,  $\beta$ Arrestin1, and  $\beta$ Arrestin2 for G protein-coupled receptors  
delineate two major classes of receptors.  
J. Biol. Chem. 275 (2000) 17201-17210

Okamoto,T.; Sekiyama, N.; Otsu, M.; Shimada, Y.; Sato, A.; Nakanishi, S.; Jingami, H.: Expression and Purification of the Extracellular Ligand Binding Region of Metabotropic Glutamate Receptor Subtype 1.

J.Biol.Chem. 273 (1998) 13089-13096

Oksche, A.; Boese, G.; Horstmeyer, A.; Ferkert, J.; Beyermann, M.; Bienert, M.; Rosenthal, W.: Late endosomal/lysosomal targeting and lack of recycling of the ligand-occupied endothelin B receptor.

Mol.Pharmacol. 57 (2000) 1104-1113

Oksche, A.; Rosenthal, W. (1998)

The molecular basis of nephrogenic diabetes insipidus.

J.Mol.Med. 76 (1998) 326-337

Pace, C.N.; Grimsley, G.R.; Thomson, J.A.; Barnett, B.J.:

Conformational stability and activity of ribonuclease T1 with zero, one, and two intact disulfide bonds.

J.Biol.Chem. 263 (1988) 11820-11825

Palczewski, K.; Kumazaka, T.; Hori, T.; Behnke, C.A.; Motoshima, H.; Fox, B.A.;

Le-Trong,I.; Teller, D.C.; Okada, T.; Stenkamp, R.E.; Yamamoto, M.; Miyano, M.:

Crystal Structure of Rhodopsin: A G Protein-Coupled Receptor.

Science 289 (2000) 739-745

Pan, Y.; Metzenberg, A.; Das, S.; Jing, B.; Gitschier, J.:

Mutation in the V2 vasopressin receptor gene are associated with X-linked nephrogenic diabetes insipidus.

Nat Genet. 2 (1992) 103-106

Pan, Y.; Wilson, P.; Gitschier, J.:

The Effect of Eight V2 Vasopressin Receptor Mutations on Stimulation of Adenylyl Cyclase and Binding to Vasopressin.

J.Biol.Chem. 269 (1994) 31933-31937

Pavo, I.; Fahrenholz, F.:

Differential inactivation of vasopressin receptor subtypes in isolated membranes and intact cells by N-ethylmaleimide.

FEBS Lett. 272 (1990) 205-208

Plakidou-Dymock, S.; Dymock, D.; Hooley, R.:

A higher plant seven-transmembrane receptor that influences sensitivity to cytokinins.

Curr.Biol. 8 (1998) 315-324

Qu, S.J.; Fan, H.Z.; Kilinc, C.; Pownall, H.J.:

Role of Cysteine Residues in Human Plasma Phospholipid Transfer Protein.

J.Protein Chem. 18(1999) 193-198

Rapoport, T.A.:

Protein transport across the ER membrane.

TIBS 15 (1990) 355-358

Robbins, M.J.; Ciruela, F.; Rhodes, A.; McIlhinney, R.A.J.:

Characterization of the dimerization of metabotropic glutamate receptors using an N-terminal truncation of mGluR1alpha.

J Neurochem. 72 (1999) 2539-2547

Rosenthal, W.; Seibold, A.; Antaramian, A.; Lonergan, M.; Arthus, M.F.; Hendy, G.N.; Birnbaumer, M.; Bichet, D.G.:

Molecular identification of the gene responsible for congenital nephrogenic diabetes insipidus.

Nature 359 (1992) 233-235

Sadeghi, H.M.; Innamorati, G.; Dagarag, M.; Birnbaumer, M.:

Palmitoylation of the V2 vasopressin receptor.

Mol.Pharmacol. 52 (1997a) 21-29

Sadeghi, H.M.; Innamorati, G.; Birnbaumer, M.:

An X-linked NDI mutation reveals a requirement for cell surface V2R expression.

Mol.Endocrinol. 11 (1997b) 706-713

Sadeghi, H.; Birnbaumer, M.:

O-Glycosylation of the V2 vasopressin receptor.

Glycobiology 9 (1999) 731-737

Savarese, T.M.; Wang, C.D.; Fraser, C.M.: Site-directed Mutagenesis of the Rat m Muscarinic Acetylcholine Receptor. Role of conserved cysteines in receptor function.

J.Biol.Chem.267 (1992) 11439-11448

Schöneberg, T.; Schulz,A.; Biebermann, H.; Grüters, A.; Grimm, T.; Hübschmann, K.; Filler, G.; Gudermann, T.; Schultz, G.:

V2 Vasopressin Receptor Dysfunction in Nephrogenic Diabetes Insipidus Caused by Different Molecular Mechanisms.

Hum. Mutat. 12 (1998) 196-205

Schülein, R.; Rutz, C.; Rosenthal, W.:

Membrane targeting and determination of transmembrane topology of the human vasopressin V2 receptor.

J.Biol.Chem. 271 (1996b) 28844-28852

Schülein, R.; Hermosilla, R.; Oksche, A.; Dehe, M.; Wiesner, B.; Krause, G.; Rosenthal, W.:

A dileucine sequence and an upstream glutamate residue in the intracellular carbocyl terminus of the vasopressin V2 receptor are essential for the cell surface transport in COS.M6 cells.

Mol.Pharmacol. 54 (1998) 525-535

Schülein, R.; Lorenz, D.; Oksche, A.; Wiesner, B.; Hermosilla, R.; Ebert, J.; Rosenthal, W.: Polarized cell surface expression of the green fluorescent protein-tagged vasopressin V2 receptor in Madin Darby canine kidney cells.  
FEBS-Lett. 441 (1998) 170-176

Schulz, A.; Grosse, R.; Schultz, G.; Gundermann, T.; Schöneberg, T.: Structural implication for receptor oligomerization from functional reconstitution studies of mutant V2 vasopressin receptors.  
J.Biol.Chem. 275 (2000) 2381-2389

Seibold, A.; Brabet, P.; Rosenthal, W.; Birnbaumer, M.: Structure and chromosomal localization of the human antidiuretic hormone receptor gene. Am. J. Hum. Genet 51 (1992) 1078-83

Stillman, B.A.; Breyer, M.D.; Breyer, R.M.: Importance of the Extracellular Domain for Prostaglandin EP<sub>2</sub> Receptor Function. Mol.Pharmacol. 56 (1999) 545-551

Struthers, M.; Yu, H.; Oprian, D.D.:G Protein-Coupled Receptor Activation: Analysis of a Highly Constrained, "Straitjacketed" Rhodopsin.  
Biochemistry 39 (2000) 7938-7942

Sung, C.H.; Schneider, B.G.; Agarwal, N.; Papermaster, D.S.; Nathans, J.: Functional heterogeneity of mutant rhodopsins responsible for autosomal dominant retinitis pigmentosa.  
Proc.Natl.Acad.Sci.USA 88 (1991) 8840-8844

Trombetta, E.S.; Helenius, A.: Lectins as chaperones in glycoprotein folding.  
Curr.Opin.Struct.Biol. 8 (1998) 587-592

Tsukaguchi, H.; Matsubara, H.; Teketani, S.; Mori, Y.; Seido, T.; Inada, M.: Binding-, Intracellular Transport-, and Biosynthesis-defective Mutants of Vasopressin Type 2 Receptor in Patients with X-linked Nephrogenic Diabetes Insipidus.  
J.Clin.Invest. 96 (1995) 2043-2050

Unger, V.M.; Hargrave, P.A.; Baldwin, J.M.; Schertler, G.F.X.: Arrangement of rhodopsin transmembrane alpha-helices.  
Nature 389 (1997) 203-206

van den Ouwehand, A.M.W.; Dreesen, J.C.F.M.; Verdijk, M.; Knoers, N.V.M.; Monnens, L.A.H.; Rocchi, M.; van Oost, B.A.: Mutations in the vasopressin type 2 receptor gene (AVPR2) associated with nephrogenic diabetes insipidus.  
Nat. Genet. 2 (1992) 99-102

Wedemeyer, W.J.; Welker, E.; Narayan, M.; Scheraga, H.A.:  
Disulfide Bonds and Protein Folding.  
Biochemistry 39 (2000) 4207-4216

Wildin, R.S.; Antush, M.J.; Bennett, R.L.; Schoof, J.M.; Scott, C.R.:  
Heterogenous AVPR2 Gene Mutations in Congenital Nephrogenic Diabetes Insipidus.  
Am.J.Hum.Genet. 55 (1994) 266-277

Wright, M.; Drummond, G.I.:  
Inactivation of the beta-adrenergic receptor in skeletal muscle by dithiols.  
Biochem. Pharmacol. 32 (1983) 509-515

Yamamoto, T.; Bishop, R.W.; Brown, M.S.; Goldstein, J.L.; Russel, D.W.:  
Deletion in cysteine rich region of LDL receptor impedes transport to cell surface in WHHL rabbit.  
Science 232 (1986) 1230-1237

Yokoyama, K.; Yamauchi, A.; Izumi, M.; Itoh, T.; Ando, A.; Imai, E.; Kamada, T.; Ueda, N.:  
A low-affinity vasopressin V2-receptor gene in a kindred with X-linked nephrogenic diabetes insipidus.  
J.Am.Soc.Nephrol. 7 (1996) 410-414

Zeng, F.Y.; Soldner, A.; Schöneberg, T.; Wess, J.:  
Conserved Extracellular Cysteine Pair in the M<sub>3</sub> Muscarinic Acetylcholine Receptor Is Essential for Proper Receptor Cell Surface Localization but Not for G Protein Coupling. J.Neurochem. 72 (1999) 2404-2414

Zeng, F.; Wess, J.  
Molecular Aspects of Muscarinic Receptor Dimerization.  
Neuropsychopharmacology 23 (2000) 19-31

Zhang, P.; Johnson, P.S.; Zöllner, C.; Wang, W.; Wang, Z.; Montes, A.E.; Seidleck, B.K.; Blaschak, C.J.; Surratt, C.K.:  
Mutation of human μ opioid receptor extracellular “disulfide cysteine” residues alters ligand binding but does not prevent receptor targeting to the cell plasma membrane.  
Brain Res.Mol.Brain Res. 72 (1999) 195-204