

9. Literaturliste

- ¹ Braun J, Bollow M, Remlinger G et al: Prevalence of spondyloarthropathy in HLA-B27-positive and negative blood donors. *Arthritis Rheum* 1998; 41: 58-67
- ² Francois RJ: *Les rachis dans la Spondylarthrite Ankylosante*. Editions Arscia Brussels; 1976.
- ³ Shichikawa K, Tsujimoto M, Nishioka J et al: Histopathology of early sacroiliitis and enthesitis in ankylosing spondylitis. *Advances in Inflammation Research*. 1985, Vol. 9. *The spondyloarthropathies*, Ziff M, Cohen SB, eds. New York, Raven Press
- ⁴ Friedburg H, Meske S, Hennig J et al: Die Kernspintomographie des Sakroiliacalgelenkes: eine neue Untersuchungstechnik zur Differentialdiagnose rheumatischer Erkrankungen im lumbosacralen Übergang. *Radiologie* 1987; 27: 130-134
- ⁵ König H, Deimling M, Sauter R, Vogt M: Cartilage disorders: comparison of spin-echo, chess and flash sequence MR images. *Radiology* 1987; 64: 753-758
- ⁶ Ahlström H, Feltelius N, Nyman R, Hällgren R: Magnetic resonance imaging of sacroiliac joint inflammation. *Arthritis Rheum* 1990; 33: 1763-1769
- ⁷ Wittram C, Whitehouse GH, Williams JW, Bucknall RC: A comparison of MR and CT in suspected sacroiliitis. *J Comput Assist Tomogr* 1996; 20: 68-72
- ⁸ Oostveen J, Prevo R, den Boer J, van de Laar M: Early detection of sacroiliitis on magnet resonance imaging and subsequent development of sacroiliitis on plain radiography. A prospective, longitudinal study. *J Rheumatol* 1999; 26: 1953-8
- ⁹ Iovane A, Midiri M, Finazzo M et al: Magnetic resonance imaging of seronegative sacroiliitis. [Article in Italian, abstract]. *Radiol Med (Torino)* 1998; 96: 185-9
- ¹⁰ Remy M, Bouillet P, Bertin P et al: Evaluation of magnetic resonance imaging for the detection of sacroiliitis in patients with early seronegative spondylarthropathy. *Rev Rhum Engl Ed* 1996; 63: 577-83
- ¹¹ Docherty P, Mitchell MJ, MacMillan L et al: Magnetic resonance imaging in the detection of sacroiliitis. *J Rheumatol* 1992; 19: 393-401
- ¹² Puhakka KB, Jurik AG, Egund N et al: Imaging of abnormalities by MR in comparison with radiography and CT. *Acta Radiol* 2003; 44: 218-229
- ¹³ Bloch F: Nuclear induction. *Phys Rev* 1946; 70: 460-474
- ¹⁴ Purcell EM, Torrey HC, Pound RV: Resonance absorption by nuclear magnetic moments in a solid. *Phys Rev* 1946; 69: 37-38
- ¹⁵ Blum U, Buitrago-Tellez C, Mundinger A et al: Magnetic resonance imaging (MRI) for detection of active sacroiliitis – a prospective study comparing conventional radiography, scintigraphy and contrast enhanced MRI. *J Rheumatol* 1996; 23: 2107-2115
- ¹⁶ Schild HH: *MRI made easy (leicht gemacht)*. Schering Aktiengesellschaft, Berlin 1990
- ¹⁷ Köchli VD, Marinček B: *Wie funktioniert MRI?* Springer, Berlin Heidelberg 1994
- ¹⁸ Gadian DG: *Nuclear Magnetic Resonance and its Applications for Living Systems*. Oxford University Press, 1982
- ¹⁹ Frahm J, Haase A, Matthaei D: Rapid three-dimensional MR imaging using the FLASH technique. *J Comput Assist Tomogr*. 1986; 10: 363-8.

- ²⁰ Dwyer AJ, Frank JA, Sank VJ et al: Short-Ti inversion-recovery pulse sequence: analysis and initial experience in cancer imaging. *Radiology*. 1988; 168: 827-36.
- ²¹ Vahlensieck M, Seelos K, Träber F et al: Magnetresonanztomographie mit schneller STIR-Technik: Optimierung und Vergleich mit anderen Sequenzen an einem 0.5 Tesla-System. *Fortschr. Röntgenstr.* 1993; 159: 288-294
- ²² Becker ED, Farrar TC: Fourier transform spectroscopy. *Science*. 1972; 178: 361-8.
- ²³ Kumar A, Welti D, Ernst RR: NMR-Fourier-Zeugmatography. *J Magn Reson* 1975, 18: 69
- ²⁴ Vogl TJ: Einfluß der Kernspintomographie auf den menschlichen Organismus. Enke-Verlag, Stuttgart 1988
- ²⁵ Vogl TJ, Paulus W, Fuchs A et al: Influence of magnetic resonance imaging on evoked potentials and nerve conduction velocities in humans. *Invest Radiol* 1991; 26: 432-437
- ²⁶ Vogl TJ, Krimmel K, Fuchs A, Lissner J: Influence of magnetic resonance imaging on human body core and intravascular temperature. *Med Phys* 1988; 15: 562-566
- ²⁷ Carr DH, Brown J, Bydder GM et al.: Intravenous chelated gadolinium as a contrast agent in NMR imaging of cerebral tumours. *Lancet*. 1984; 1: 484-6.
- ²⁸ „Magnevist“ Gebrauchsinformation, Schering AG, Berlin, 1994
- ²⁹ Bellamy N, Park W, Rooney PJ: What do we know about the sacroiliac joint? *Semin Arth Rheum* 1983; 12: 218-313
- ³⁰ Ball J: The enthesopathy of ankylosing Spondylitis. *Brit J Rheum* 1983; 22 (suppl.): 25-28
- ³¹ McGonagle D, Khan MA, Marzo-Ortega H et al: Enthesitis in spondyloarthropathy. *Curr Opin Rheumatol* 1999; 11: 244-250
- ³² Francois RJ, Gardner DL, Degraeve EJ, Bywaters EGL: Histopathologic evidence that sacroiliitis in ankylosing spondylitis is not merely enthesitis. *Arthritis Rheum* 2000; 43: 2011-2024
- ³³ Braun J, Bollow M, Remlinger G et al: Prevalence of spondylarthropathies in HLA B27-positive and -negative blood donors. *Arthritis Rheum* 1998; 41: 58-67
- ³⁴ Underwood MR, Dawes P: Inflammatory back pain in primary care. *Br J Rheumatol* 1995; 34: 1074-1077
- ³⁵ Zink A, Braun J, Listing A et al: Disability and handicap in rheumatoid arthritis and ankylosing spondylitis – results of the German rheumatological database. *J Rheumatol* 2000; 27: 613-622
- ³⁶ Gran JT, Skomsvoll JF: The outcome of ankylosing spondylitis: a study of 100 patients. *Br J Rheumatol* 1997; 36: 766-71
- ³⁷ Braun J, Tuszewski M, Ehlers S et al: Nested polymerase chain reaction strategy simultaneously targeting DNA sequences of multiple bacterial species in inflammatory joint diseases. II. Examination of sacroiliac and knee joint biopsies of patients with spondyloarthropathies and other arthritides. *J Rheumatol*. 1997; 24: 1101-5
- ³⁸ Poole AR: The histopathology of ankylosing spondylitis: are there unifying hypotheses? *Am J Med Sci* 1998; 316: 228-233
- ³⁹ Braun J, Bollow M, Neure L et al: Use of immunohistologic and in situ hybridization techniques in the examination of sacroiliac joint biopsy specimens from patients with ankylosing spondylitis. *Arthritis Rheum* 1995; 38: 499-505
- ⁴⁰ Beutler BA: The role of TNF in health and disease. *J Rheumatol* 1999; 26: 16-21
- ⁴¹ Ribbens C, Andre B, Kaye O et al: Synovial fluid matrix metalloproteinase-3 levels are increased in inflammatory arthritides whether erosive or not. *Rheumatology (Oxford)*. 2000; 39: 1357-65

- ⁴² Cunnane G, Fitzgerald O, Beeton C, Cawston TE, Bresnihan B: Early joint erosions and serum levels of matrix metalloproteinase 1, matrix metalloproteinase 3, and tissue inhibitor of metalloproteinases 1 in rheumatoid arthritis. *Arthritis Rheum.* 2001; 44: 2263-74
- ⁴³ Fraser A, Fearon U, Reece R, Emery P, Veale DJ. Matrix metalloproteinase 9, apoptosis, and vascular morphology in early arthritis. *Arthritis Rheum.* 2001 Sep;44(9):2024-8.
- ⁴⁴ Calin A, Fries JF: Striking prevalence of ankylosing spondylitis in "healthy" w27 positive males and females. A controlled study. *N Engl J Med* 1975; 293: 835-839
- ⁴⁵ Mielants H, Veys EM, de Vos M et al: The evolution of spondyloarthropathies in relation to gut history. *J Rheumatol* 1995; 22: 2266-2284
- ⁴⁶ Calin A, Porta J, Fries JF et al: Clinical history as a screening test for ankylosing spondylitis. *JAMA* 1977; 237: 2613-4
- ⁴⁷ Maksymowych WP, Chou CT, Russell AS: Matching prevalence of peripheral arthritis and acute anterior uveitis in individuals with ankylosing spondylitis. *Ann Rheum Dis* 1995; 54: 128-130
- ⁴⁸ Leirisalo-Repo M, Turunen U, Stenman S et al: High frequency of silent inflammatory bowel disease in spondylarthropathy. *Arthritis Rheum* 1994; 37: 23-31
- ⁴⁹ Dougados M, Van der Linden S, Juhlin R, et al: The European Spondylarthropathy Study Group preliminary criteria for the classification of spondylarthropathy. *Arthritis Rheum* 1991; 34: 1218-1227
- ⁵⁰ Gran JT, Husby G: The epidemiology of ankylosing spondylitis. *Arthritis Rheum* 1993; 22: 319-344
- ⁵¹ Van der Linden S, Valkenburg HA, Cats A: Evaluation of diagnostic criteria for ankylosing spondylitis: a proposal for modification of New York criteria. *Arthritis Rheum* 1984; 27: 361-8
- ⁵² Mau W, Zeidler H, Mau R et al: Clinical features and prognosis of patients with possible ankylosing spondylitis. Results of a 10-year followup. *J Rheumatol* 1988; 15: 1109-1114
- ⁵³ Braun J, Sieper J: The sacroiliac Joint in the spondylarthropathies. *Curr Opin Rheum* 1996; 8: 275-287
- ⁵⁴ Bollow M, Braun J, Hamm B et al: Early sacroiliitis in patients with spondyloarthropathy: evaluation with dynamic gadolinium-enhanced MR imaging. *Radiology* 1995, 194: 529-236
- ⁵⁵ Braun J, Bollow M, Seyrekbasan F et al: Computed tomography guided corticosteroid injection of the sacroiliac joint in patients with spondyloarthropathy with sacroiliitis: clinical outcome and followup by dynamic magnetic resonance imaging. *J Rheumatol* 1996; 23: 659-664
- ⁵⁶ Fam AG, Rubenstein JD, Chin Sang H et al: Computed tomography in the diagnosis of early ankylosing spondylitis. *Arthritis Rheum* 1985; 28: 930-937
- ⁵⁷ Yu W, Feng F, Dion E et al: Comparison of radiography, computed tomography and magnetic resonance imaging in the detection of sacroiliitis accompanying ankylosing spondylitis. *Skeletal Radiol* 1998; 27: 311-320
- ⁵⁸ Brandt J, Haibel H, Cornely D et al: Successful treatment of active ankylosing spondylitis with the anti-tumor necrosis factor alpha monoclonal antibody infliximab. *Arthritis Rheum* 2000; 43: 1346-1352
- ⁵⁹ Maksymowych WP, Jaghri GS, Fitzgerald AA et al: A six month randomized, controlled, double-blind, dose-response comparison of intravenous pamidronate (60 mg versus 10 mg) in the treatment of nonsteroidal antiinflammatory drug-refractory ankylosing spondylitis. *Arthritis Rheum* 2002; 46: 766-733.
- ⁶⁰ Haibel H, Brandt J, Rudwaleit M et al: Treatment of active ankylosing spondylitis with pamidronate. *Rheumatology* 2003; 42: 1018-1020
- ⁶¹ Kommission Pharmakotherapie: Stellungnahme der Deutschen Gesellschaft für Rheumatologie zur Therapie der ankylosierenden Spondylitis (AS) mit Radiumchlorid (²²⁴SpondylAT®). *Z Rheumatol* 2001; 60: 84-87

- ⁶² Huang F, Gu J, Zhao W et al: One-year open-label trial of thalidomide in ankylosing spondylitis. *Arthritis Rheum.* 2002; 47: 249-54
- ⁶³ Braun J, Kingsley G, van der Heijde D, Sieper J: On the difficulties of establishing a consensus on the definition of and diagnostic investigations for reactive arthritis. *J Rheumatol* 2000; 27: 2185-92
- ⁶⁴ Bollow M, Braun J, Biedermann T et al: Use of contrast-enhanced MR imaging to detect sacroiliitis in children. *Skeletal Radiol* 1998; 27: 606-616
- ⁶⁵ Bollow M, Loreck D, Banzer D et al: Bildgebende Diagnostik bei Verdacht auf entzündlich rheumatische Achsenskeletterkrankungen (Sacroiliitis). *Z Rheumatol* 1999; 58: 61-70
- ⁶⁶ Sturzenbecher A, Braun J, Paris S et al: MR imaging of septic sacroiliitis. *Skeletal Radiol.* 2000; 29: 439-46
- ⁶⁷ Braun J, Khan MA, Sieper J: Enthesitis and ankylosis in spondyloarthritis: What is the target of the immune response? *Ann Rheum Dis* 2000; 59: 985-994
- ⁶⁸ Ball J: Enthesopathy of rheumatoid and ankylosing spondylitis. The Heberden Oration 1970. *Ann rheum Dis* 1971; 30: 213-223
- ⁶⁹ Benjamin M, Ralphs JR: Fibrocartilage in tendons and ligaments – an adaption to compressive loads. *J Anat* 1998; 193: 481-494
- ⁷⁰ McGonagle D, Khan MA, Marzo-Ortega H et al: Enthesitis in spondyloarthritis. *Curr Opin Rheumatol* 1999; 11: 244-250
- ⁷¹ Francois RJ, Gardner DL, Degraeve EJ, Bywaters EG: Histopathologic evidence that sacroiliitis in ankylosing spondylitis is not merely enthesitis. *Arthritis Rheum* 2000; 43: 2011-24
- ⁷² Garrett S, Jenkinson T, Kennedy LG et al: A new approach to defining disease status in ankylosing spondylitis: The Bath Ankylosing Spondylitis Disease Activity Index. *J Rheumatol* 1994; 21: 2286-2291
- ⁷³ Calin A, Garrett S, Whitelock H et al: A new approach to defining functional ability in ankylosing spondylitis: The Bath Ankylosing Spondylitis Functional Index. *J Rheumatol* 1994; 21: 2281-2285
- ⁷⁴ Wittram C, Whitehouse GH, Williams J et al: A comparison of MR and CT in suspected sacroiliitis. *J Comput Assist Tomogr* 1996; 20: 68-72
- ⁷⁵ Blum U, Buitrago-Tellez C, Munding A et al: Magnetic resonance imaging (MRI) for detection of active sacroiliitis – a prospective study comparing conventional radiography, scintigraphy and contrast enhanced MRI. *J Rheum* 1996; 23: 2107-2115
- ⁷⁶ Yu W, Feng F, Dion E et al: Comparison of radiography, computed tomography and magnetic resonance imaging in the detection of sacroiliitis accompanying ankylosing spondylitis. *Skeletal Radiol* 1998; 27: 311-20
- ⁷⁷ Wittram C, Whitehouse GH, Bucknall RC: Fat suppressed contrast enhanced MR imaging in the assessment of sacroiliitis. *Clin Radiol* 1996; 51: 554-558