

6 LITERATURVERZEICHNIS

Albert, M.L., Pearce, S.F., Francisco, L.M., Sauter, B., Roy, P., Silverstein, R.L., Bhardwaj, N. (1998). Immature dendritic cells phagocytose apoptotic cells via alphavbeta5 and CD36, and cross-present antigens to cytotoxic T lymphocytes. *J. Exp. Med.* 188: 1359-1368.

Alexander, W.S., Hilton, D.J. (2004). The role of suppressors of cytokine signaling (SOCS) proteins in regulation of the immune response. *Annu. Rev. Immunol.* 22: 503-529.

Angenieux, C., Salamero, J., Fricker, D., Cazenave, J.P., Goud, B., Hanau, D., de La Salle, H. (2000). Characterization of CD1e, a third type of CD1 molecule expressed in dendritic cells. *J. Biol. Chem.* 275: 37757-37764.

Anjuere, F., del Hoyo, G.M., Martin, P., Ardavin, C. (2000). Langerhans cells develop from a lymphoid-committed precursor. *Blood* 96: 1633-1637.

Ardavin, C., Martinez del Hoyo, G., Martin, P., Anjuere, F., Arias, C.F., Marin, A.R., Ruiz, S., Parrillas, V., Hernandez, H. (2001). Origin and differentiation of dendritic cells. *Trends Immunol.* 22: 691-700.

Ashworth, J., Kahan, M.C., Breathnach, S.M. (1989). Flow cytometric analysis and sorting of HLA-DR⁺CD1⁺ Langerhans cells. *Br. J. Dermatol.* 121: 11-18.

Austyn, J.M. (1999). Dendritic cells in spleen and lymph node. In: *Dendritic cells*. Eds. Lotze, M.T. und Thomson, A.W., Academic Press, San Diego, pp. 179-204.

Banchereau, J., Paczesny, S., Blanco, P., Bennett, L., Pascual, V., Fay, J., Palucka, A.K. (2003). Dendritic cells: controllers of the immune system and a new promise for immunotherapy. *Ann. N. Y. Acad. Sci.* 987: 180-187.

Banchereau, J. (2002). The long arm of the immune system. *Sci. Am.* 287: 36-41.

Banchereau, J., Briere, F., Caux, C., Davoust, J., Lebecque, S., Liu, Y.L., Pulendran, B., Palucka, K. (2000). Immunobiology of dendritic cells. *Annu. Rev. Immunol.* 18: 767-811.

Banchereau, J., Steinman, R.M. (1998). Dendritic cells and the control of immunity. *Nature* 392: 245-252.

Bartz, H., Rothoeft, T., Anhenn, O., Bunse, D., Schauer, U. (2003). Large-scale isolation of immature dendritic cells with features of Langerhans cells by sorting CD34+ cord blood stem cells cultured in the presence of TGF-beta1 for cutaneous leukocyte antigen (CLA). *J. Immunol. Meth.* 275: 137-148.

Bauer, J., Bahmer, F.A., Wörl, J., Neuhuber, W., Schuler, G., Fartasch, M. (2001). A strikingly constant ratio exists between Langerhans cells and other epidermal cells in human skin. A stereologic study using the optical disector method and the confocal laser scanning microscope. *J. Invest. Dermatol.* 116: 313-318.

Bjercke, S., Elgo, J., Braathen, L., Thorsby, E. (1984). Enriched epidermal Langerhans cells are potent antigen-presenting cells for T cells. *J. Invest. Dermatol.* 83: 286-289.

Boivin, G.P., O`Toole, B.A., Ormsby, I.E., Diebold, R.J., Eis, M.J., Doetschman, T., Kier, A.B. (1995). Onset and progression of pathological lesions in transforming growth factor- β 1-deficient mice. *Am. J. Pathol.* 146: 276-288.

Bonifaz, L., Bonnyay, D., Mahnke, K., Rivera, M., Nussenzweig, M.C., Steinman, R.M. (2002). Efficient targeting of protein antigen to dendritic cell receptor DEC-205 in the steady state leads to antigen presentation on major histocompatibility complex I products and peripheral CD8 $^{+}$ T cell tolerance. *J. Exp. Med.* 196: 1627-1638.

Borkowski, T.A., Letterio, J.J., Mackall, C.L., Saito, A., Wang, X.J., Roop, D.R., Gress, R.E., Udey, M.C. (1997). A role for TGF β 1 in Langerhans cell biology. *J. Clin. Invest.* 100: 575-581.

Borkowski, T.A., Letterio, J.J., Farr, A.G., Udey, M.C. (1996). A role for endogenous transforming growth factor β 1 in Langerhans cell biology: the skin of transforming growth factor beta 1 null mice is devoid of epidermal Langerhans cells. *J. Exp. Med.* 184: 2417-2422.

Borkowski, T.A., Van Dyke, B.J., Schwarzenberger, K., McFarland, V.W., Farr, A.G., Udey, M.C. (1994). Expression of E-cadherin by murine dendritic cells. E-cadherin as a dendritic cell antigen characteristic of epidermal Langerhans cells and related cells. *Eur. J. Immunol.* 24: 2767-2774.

Bøyum, A. (1964). Separation of white blood cells. *Nature* 204: 793-794.

Broker, T., Riedinger, M., Karjalainen, K. (1997). Targeted expression of major histocompatibility complex (MHC) class II molecules demonstrates that dendritic cells can induce negative but not positive selection by thymocytes in vivo. *J. Exp. Med.* 185: 541-550.

Caux, C., Massacrier, C., Dubois, B., Valladeau, J., Dezutter-Dambuyant, C., Durand, I., Schmitt, D., Saeland, S. (1999). Respective involvement of TGF- β and IL-4 in the development of Langerhans cells and non-Langerhans dendritic cells from CD34 $^{+}$ progenitors. *J. Leukoc. Biol.* 66: 781-791.

Caux, C., Vanbervliet, B., Massacrier, C., Dezutter-Dambuyant, C., de-Saint-Vis, B., Jacquet, C., Yoneda, K., Imamura, S., Schmitt, D., Banchereau, J. (1996). CD34 $^{+}$ hematopoietic progenitors from human cord blood differentiate along two independent dendritic cell pathways in response to GM-CSF+TNF- α . *J. Exp. Med.* 184: 695-706.

Caux, C., Massacrier, C., Vanbervliet, B., Dubois, B., Van Kooten, C., Durand, I., Banchereau, J. (1994). Activation of human dendritic cells through CD40 cross-linking. *J. Exp. Med.* 180: 1263-1272.

Caux, C., Dezutter-Dambuyant, C., Schmitt, D. (1992). GM-CSF and TNF- α cooperate in the generation of dendritic Langerhans cells. *Nature* 360: 258-261.

Cella, M., Engering, A., Pinet, V., Pieters, J., Lanzavecchia, A. (1997). Inflammatory stimuli induce accumulation of MHC class II complexes on dendritic cells. *Nature* 388: 782-787.

Cella, M., Scheidegger, D., Palmer-Lehmann, K., Lane, P., Lanzavecchia, A., Alber, G. (1996). Ligation of CD40 on dendritic cells triggers production of high levels of interleukin-12 and enhances T cell stimulatory capacity: T-T help via APC activation. *J. Exp. Med.* 184: 747-752.

Chen, L.F., Greene, W.C. (2004). Shaping the nuclear action of NF-κB. *Nat. Rev. Mol. Cell Biol.* 5: 392-401.

Chomczynski, P., Sacci, N. (1987). Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* 162: 156-159.

Chow, J.C., Young, D.W., Golenbock, D.T., Christ, W.J., Gusovsky, F. (1999). Toll-like receptor-4 mediates lipopolysaccharide-induced signal transduction. *J. Biol. Chem.* 274: 10689-10692.

Companjen, A.R., van der Wel, L.I., Wie, L., Laman, J.D., Prens, E.R. (2001). A modified ex vivo skin organ culture system for functional studies. *Arch. Dermatol. Res.* 293: 184-190.

Corinti, S., Albanesi, C., la Sala, A., Pastore, S., Girolomoni, G. (2001). Regulatory activity of autocrine IL-10 on dendritic cell functions. *J. Immunol.* 166: 4312-4318.

Cumberbatch, M., Dearman, R.J., Kimber, I. (1997). Langerhans cells require signals from both tumour necrosis factor-alpha and interleukin-1 beta for migration. *Immunology* 92: 388-395.

Cumberbatch, M., Dearman, R.J., Kimber, I. (1996). Constitutive and inducible expression of interleukin-6 by Langerhans cells and lymph node dendritic cells. *Immunology* 87: 513-518.

Cumberbatch, M., Peters, S.W., Gould, S.J., Kimber, I. (1992). Intercellular adhesion molecule-1 (ICAM-1) expression by lymph node dendritic cells: comparison with epidermal Langerhans cells. *Immunol. Lett.* 32: 105-110.

De Graaf, J.H., Tamminga, R.Y.J., Kamps, W.A., Timens, W. (1995). Expression of cellular adhesion molecules in Langerhans cell histiocytosis and normal Langerhans cells. *Am. J. Pathol.* 147: 1161-1171.

Demangel, C., Bertolino, P., Britton, W.J. (2002). Autocrine IL-10 impairs dendritic cell (DC)-derived immune responses to mycobacterial infection by suppressing DC trafficking to draining lymph nodes and local IL-12 production. *Eur. J. Immunol.* 32: 994-1002.

Dezutter-Dambuyant, C., Cordier, G., Schmitt, D., Faure, M., Laquoi, C., Thivolet, J. (1984). Quantitative evaluation of two distinct cell populations expressing HLA-DR antigens in normal human epidermis. *Br. J. Dermatol.* 111: 1-11.

Dzionek, A., Sohma, Y., Nagafune, J., Cella, M., Colonna, M., Facchetti, F., Gunther, G., Johnston, I., Lanzavecchia, A., Nagasaka, T., Okada, T., Vermi, W., Winkels, G., Yamamoto, T., Zysk, M., Yamaguchi, Y., Schmitz, J. (2001). BDCA-2, a novel plasmacytoid dendritic cell-specific type II C-type lectin, mediates antigen capture and is a potent inhibitor of interferon alpha/beta induction. *J. Exp. Med.* 194: 1823-1834.

Elder, J.T., Reynolds, N.J., Cooper, K.D., Griffiths, C.E., Hardas, B.D., Bleicher, P.A. (1993). CD1 gene expression in human skin. *J. Dermatol. Sci.* 6: 206-213.

Enk, A.H., Katz, S.I. (1992). Early molecular events in the induction phase of contact sensitivity. *Proc. Natl. Acad. Sci. USA* 89: 1398-1402.

Esser, C. (1992). Powerful Preselection. In: *Flow Cytometry and Cell Sorting*. Ed. Radbruch, A., Springer-Verlag, Berlin, pp. 133-140.

Figdor, C.G., van Kooyk, Y., Adema, G.J. (2002). C-type lectin receptors on dendritic cells and Langerhans cells. *Nat. Rev. Immunol.* 2: 77-84.

Fithian, E., Kung, P., Goldstein, G., Rubenfeld, M., Fenoglio, C., Edelson, R. (1981). Reactivity of Langerhans cells with hybridoma antibody. *Proc. Natl. Acad. Sci. USA.* 78: 2541-2544.

Frisch, S.M., Francis, H. (1994). Disruption of epithelial cell-matrix interactions induces apoptosis. *J. Cell Biol.* 124: 619-626.

Fritsch, P. (2004). Epidermis. In: *Dermatologie - Venerologie. Grundlagen. Klinik. Atlas.* Ed. Fritsch, P., Springer-Verlag, Berlin, pp. 5-23.

Gatti, E., Velleca, M.A., Biedermann, B.C., Ma, W., Unternaehrer, J., Ebersold, M.W., Medzhitov, R., Pober, J.S., Mellman, I. (2000). Large-scale culture and selective maturation of human Langerhans cells from granulocyte colony-stimulating factor-mobilized CD34⁺ progenitors. *J. Immunol.* 164: 3600-3607.

Geissmann, F., Dieu-Nosjean, M.C., Dezutter, C., Valladeau, J., Kayal, S., Leborgne, M., Brousse, N., Saeland, S., Davoust, J. (2002). Accumulation of immature Langerhans cells in human lymph nodes draining chronically inflamed skin. *J. Exp. Med.* 196: 417-430.

Geissmann, F., Revy, P., Regnault, A., Lepelletier, Y., Dy, M., Brousse, N., Amigorena, S., Hermine, O., Durandy, A. (1999). TGF-β1 prevents the noncognate maturation of human dendritic Langerhans cells. *J. Immunol.* 162: 4567-4575.

Geissmann, F., Prost, C., Monnet, J.P., Dy, M., Brousse, N., Hermine, O. (1998). TGF-β1, in the presence of GM-CSF and IL-4 induces differentiation of human peripheral blood monocytes into dendritic LCs. *J. Exp. Med.* 187: 961-966.

Gerosa, F., Baldani-Guerra, B., Nisii, C., Marchesini, V., Carra, G., Trinchieri, G. (2002). Reciprocal activating interaction between natural killer cells and dendritic cells. *J. Exp. Med.* 195: 327-333.

Gilliet, M., Boonstra, A., Paturel, C., Antonenko, S., Xu, X.L., Trinchieri, G., O'Garra, A., Liu, Y.J. (2002). The development of murine plasmacytoid dendritic cell precursors is differentially regulated by FLT3-ligand and granulocyte/macrophage colony-stimulating factor. *J. Exp. Med.* 195: 953-958.

Girolomoni, G., Caux, C., Dezutter-Dambuyant, C., Lebecque, S., Ricciardi-Castagnoli, P. (2002). Langerhans cells: still a fundamental paradigm for studying the immunobiology of dendritic cells. *Trends Immunol.* 23: 6-8.

Granucci, F., Vizzardelli, C., Pavelka, N., Feau, S., Persico, M., Virzi, E., Rescigno, M., Moro, G., Ricciardi-Castagnoli, P. (2001). Inducible IL-2 production by dendritic cells revealed by global gene expression analysis. *Nat. Immunol.* 2: 882-888.

Hacker, C., Kirsch, R.D., Ju, X.S., Hieronymus, T., Gust, T.C., Kuhl, C., Jorgas, T., Kurz, S.M., Rose-John, S., Yokota, Y., Zenke, M. (2003). Transcriptional profiling identifies Id2 function in dendritic cell development. *Nat. Immunol.* 4: 380-386.

Hanau, D., Schmitt, D. A., Fabre, M., Cazenave, J. P. (1988). A method for the rapid isolation of human epidermal Langerhans cells using immunomagnetic microspheres. *J. Invest. Dermatol.* 91: 274-279.

Handgretinger, R., Lang, P., Schumm, M., Taylor, G., Neu, S., Koscielnak, E., Niethammer, D., Klingebiel, T. (1998). Isolation and transplantation of autologous peripheral CD34⁺ progenitor cells highly purified by magnetic-activated cell sorting. *Bone Marrow Transplant.* 21: 987-993.

Hashimoto, K. (1970). Lanthanum staining of Langerhans'cell. *Arch. Dermatol.* 102: 280-290.

Hayashi, F., Smith, K.D., Ozinsky, A., Hawn, T.R., Yi, E.C., Goodlett, D.R., Eng, J.K., Akira, S., Underhill, D.M., Aderem, A. (2001). The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5. *Nature* 410: 1099-1103.

Heath, W.R., Carbone, F.R. (2001). Cross-presentation, dendritic cells, tolerance and immunity. *Annu. Rev. Immunol.* 19: 47-64.

Heil, F., Hemmi, H., Hochrein, H., Ampenberger, F., Kirschning, C., Akira, S., Lipford, G., Wagner, H., Bauer, S. (2004). Species-specific recognition of single-stranded RNA via toll-like receptor 7 and 8. *Science* 303: 1526-1529.

Hemmi, H., Takeuchi, O., Kawai, T., Kaisho, T., Sato, S., Sanjo, H., Matsumoto, M., Hoshino, K., Wagner, H., Takeda, K., Akira, S. (2000). A Toll-like receptor recognizes bacterial DNA. *Nature* 408: 740-745. Erratum in: *Nature* 409: 646.

Heufler, C., Topar, G., Koch, F., Trockenbacher, B., Kämpgen, E., Romani, N., Schuler, G. (1992). Cytokine gene expression in murine epidermal cell suspensions: interleukin 1 β and macrophage inflammatory protein 1 α are selectively expressed in Langerhans cells but are differentially regulated in culture. *J. Exp. Med.* 176: 1221-1226.

Heufler, C., Koch, F., Schuler, G. (1988). Granulocyte/macrophage colony-stimulating factor and interleukin 1 mediate the maturation of murine epidermal Langerhans cells into potent immunostimulatory dendritic cells. *J. Exp. Med.* 167: 700-705.

Hochrein, H., O'Keeffe, M., Luft, T., Vandenabeele, S., Grumont, R.J., Maraskovsky, E., Shortman, K. (2000). Interleukin (IL)-4 is a major regulatory cytokine governing bioactive IL-12 production by mouse and human dendritic cells. *J. Exp. Med.* 192: 823-833.

Holt, P.G., Stumbles, P.A., McWilliam, A.S. (1999). Functional studies on dendritic cells in the respiratory tract and related mucosal tissues. *J. Leukoc. Biol.* 66: 272-275.

Huang, F.P., Platt, N., Wykes, M., Major, J.R., Powell, T.J., Jenkins, C.D., MacPherson, G.G. (2000). A discrete subpopulation of dendritic cells transports apoptotic intestinal epithelial cells to T cell areas of mesenteric lymph nodes. *J. Exp. Med.* 191: 435-444.

Hunger, R.E., Sieling, P.A., Ochoa, M.T., Sugaya, M., Burdick, A.E., Rea, T.H., Brennan, P.J., Belisle, J.T., Blauvelt, A., Porcelli, S.A., Modlin, R.L. (2004). Langerhans cells utilize CD1a and langerin to efficiently present nonpeptide antigens to T cells. *J. Clin. Invest.* 113: 701-718.

Inaba, K., Turley, S., Yamaide, F., Iyoda, T., Mahnke, K., Inaba, M., Pack, M., Subklewe, M., Sauter, B., Sheff, D., Albert, M., Bhardwaj, N., Mellman, I., Steinman, R.M. (1998). Efficient presentation of phagocytosed cellular fragments on the major histocompatibility complex class II products of dendritic cells. *J. Exp. Med.* 188: 2163-2173.

Inaba, K., Pack, M., Inaba, M., Sakuta, H., Isdell, F., Steinman, R.M. (1997). High levels of a major histocompatibility complex II-self peptide complex on dendritic cells from lymph node. *J. Exp. Med.* 186: 665-672.

Ishii, M., Terao, Y., Kitajima, J.I., Hamada, T. (1984). Sequential production of Birbeck granules through adsorptive pinocytosis. *J. Invest. Dermatol.* 82: 28-29.

Ito, T., Amakawa, R., Kaisho, T., Hemmi, H., Tajima, K., Uehira, K., Ozaki, Y., Tomizawa, H., Akira, S., Fukuhara, S. (2002). Interferon-alpha and interleukin-12 are induced differentially by Toll-like receptor 7 ligands in human blood dendritic cell subsets. *J. Exp. Med.* 195: 1507-1512.

Iwama, A., Osawa, M., Hirasawa, R., Uchiyama, N., Kaneko, S., Onodera, M., Shibuya, K., Shibuya, A., Vinson, C., Tenen, D.G., Nakauchi, H. (2002). Reciprocal roles for CCAAT/enhancer binding protein (C/EBP) and PU.1 transcription factors in Langerhans cell commitment. *J. Exp. Med.* 195: 547-558.

Janeway, C.A., Travers, P., Walport, M., Shlochik, M. (2001). *Immunobiology. The immune system in health and disease.* 5th ed., Garland Science, New York.

Janssen, G.H., Tangelder, G.J., Oude Egbrink, M.G., Reneman, R.S. (1994). Spontaneous leukocyte rolling in venules in untraumatized skin of conscious and anesthetized animals. *Am. J. Physiol.* 267: H1199-H1204.

Jefford M., Schnurr M., Toy T., Masterman K.A., Shin A., Beecroft T., Tai T.Y., Shortman, K., Shackleton M., Davis I.D., Parente P., Luft T., Chen W., Cebon J., Maraskovsky E. (2003). Functional comparison of DCs generated in vivo with Flt3 ligand or in vitro from blood monocytes: differential regulation of function by specific classes of physiologic stimuli. *Blood* 102: 1753-1763.

Jolles S., Christensen J., Holman M., Klaus G.B., Ager A. (2002). Systemic treatment with anti-CD40 antibody stimulates Langerhans cell migration from the skin. *Clin. Exp. Immunol.* 129: 519-526.

Jonuleit, H., Kuhn, U., Muller, G., Steinbrink, K., Paragnik, L., Schmitt, E., Knop, J., Enk, A.H. (1997). Pro-inflammatory cytokines and prostaglandins induce maturation of potent immunostimulatory dendritic cells under fetal calf serum-free conditions. *Eur. J. Immunol.* 27: 3135-3142.

Kadowaki, N., Ho, S., Antonenko, S., Malefyt, R.W., Kastelein, R.A., Bazan, F., Liu, Y.J. (2001). Subsets of human dendritic cell precursors express different toll-like receptors and respond to different microbial antigens. *J. Exp. Med.* 194: 863-869.

Kamath, A.T., Pooley, J., O`Keefe, M.A., Vremec, D., Zhan, Y., Lew, A.M., D`Amico, A., Wu, L., Tough, D.F., Shortman, K. (2000). The development, maturation, and turnover rate of mouse spleen dendritic cell populations. *J. Immunol.* 165: 6762-6770.

Kang, K., Kubin, M., Cooper, K.D., Lessin, S.R., Trinchieri, G., Rook, A.H. (1996). IL-12 synthesis by human Langerhans cells. *J. Immunol.* 156: 1402-1407.

Kantor, A.B., Gibbons, I., Miltenyi, S., Schmitz, J. (1997). Magnetic cell sorting with colloidal superparamagnetic particles. In: *Cell separation methods and applications*. Eds. Recktenwald, D., Radbruch, A., Marcel Dekker Verlag, New York, pp. 153-173.

Kapsenberg, M.L. (2003). Dendritic cell control of pathogen-driven T-cell polarization. *Nat. Rev. Immunol.* 3: 984-993.

Karin, M., Lin, A. (2002). NFκB at the crossroads of life and death. *Nat. Immunol.* 3: 221-227.

Kawamura, T., Azuma, M., Kayagaki, N., Shimada, S., Yagita, H., Okumura, K. (2000). Fas/Fas ligand-mediated apoptosis of murine Langerhans cells. *J. Dermatol. Sci.* 22: 96-101.

Kawamura, T., Azuma, M., Kayagaki, N., Shimada, S., Yagita, H., Okumura, K. (1999). Fas/Fas ligand-mediated elimination of antigen-bearing Langerhans cells in draining lymph nodes. *Br. J. Dermatol.* 141: 201-205.

Koch, F., Stanzl, U., Jennewein, P., Janke, K., Heufler, C., Kämpgen, E., Romani, N., Schuler, G. (1996). High level IL-12 production by murine dendritic cells: upregulation via MHC class II and CD40 molecules and downregulation by IL-4 and IL-10. *J. Exp. Med.* 184: 741-746.

Kolde, G., Knop, J. (1987). Different cellular reaction patterns of epidermal Langerhans cells after application of contact sensitizing, toxic, and tolerogenic compounds. A comparative ultrastructural and morphometric time-course analysis. *J. Invest. Dermatol.* 89: 19-23.

Krug, A., Towarowski, A., Britsch, S., Rothenfusser, S., Hornung, V., Bals, R., Giese, T., Engelmann, H., Endres, S., Krieg, A.M., Hartmann, G. (2001). Toll-like receptor expression reveals CpG DNA as a unique microbial stimulus for plasmacytoid dendritic cells which synergizes with CD40 ligand to induce high amounts of IL-12. *Eur. J. Immunol.* 31: 3026-3037.

Kupper, T.S., Fuhlbrigge, R.C. (2004). Immune surveillance in the skin: mechanisms and clinical consequences. *Nat. Rev. Immunol.* 4: 211-222.

Langenkamp, A., Messi, M., Lanzavecchia, A., Sallusto, F. (2000). Kinetics of dendritic cell activation: impact on priming of TH1, TH2 and nonpolarized T cells. *Nat. Immunol.* 1: 311-316.

Langerhans, P. (1868). Über die Nerven der menschlichen Haut. *Virchows Arch. (Pathol. Arch.)* 44: 325-327.

Lanzavecchia, A., Sallusto, F. (2002). Progressive differentiation and selection of the fittest in the immune response. *Nat. Rev. Immunol.* 2: 982-987.

Lanzavecchia, A., Sallusto, F. (2001). Regulation of T cell immunity by dendritic cells. *Cell* 106: 263-266.

Lerrick, J.W., Morhenn, V., Chiang, Y.L., Shi, T. (1989). Activated Langerhans cells release tumor necrosis factor. *J. Leukoc. Biol.* 45: 429-433.

Larsen, C.P., Steinman, R.M., Witmer-Pack, M., Hankins, D.F., Morris, P.J., Austyn, J.M. (1990). Migration and maturation of Langerhans cells in skin transplants and explants. *J. Exp. Med.* 172: 1483-1493.

Lee, W.C., Zhong, C., Qian, S., Wan, Y., Gauldie, J., Mi, Z., Robbins, P.D., Thomson, A.W., Lu, L. (1998). Phenotype, function, and in vivo migration and survival of allogeneic dendritic cell progenitors genetically engineered to express TGF- β . *Transplantation* 66: 1810-1817.

Lenzner, S., Peters, J.H. (1992). Harvesting monocytes by adherence. In: *Monoclonal Antibodies*. Eds. Peters, J.H., Baumgarten, H., Springer-Verlag, Berlin, p.172.

Leslie, D.S., Vincent, M.S., Spada, F.M., Das, H., Sugita, M., Morita, C.T., Brenner, M.B. (2002). CD1-mediated gamma/delta T cell maturation of dendritic cells. *J. Exp. Med.* 196: 1575-1584.

Le Varlet, B., Dezutter-Dambuyant, C., Staquet, M.J., Delorme, P., Schmitt, D. (1991). Human epidermal Langerhans cells express integrins of the $\beta 1$ subfamily. *J. Invest. Dermatol.* 96: 518-522.

Leverkus, M., McLellan, A.D., Heldmann, M., Eggert, A.O., Brocker, E.B., Koch, N., Kämpgen, E. (2003). MHC class II-mediated apoptosis in dendritic cells: a role for membrane-associated and mitochondrial signaling pathways. *Int. Immunol.* 8: 993-1006.

Litinskiy, M.B., Nardelli, B., Hilbert, D.M., He, B., Schaffer, A., Casali, P., Cerutti, A. (2002). DCs induce CD40-independent immunoglobulin class switching through BLyS and APRIL. *Nat. Immunol.* 3: 822-829.

Liu, H.N., Chang, Y.T., Wong, C.K. (1994). Enrichment of unlabelled human Langerhans cells by modified discontinuous Ficoll-metrizoate density medium and following Langerhans cell culture. *Clin. Exp. Dermatol.* 19: 113-117.

Liu, Y.J., Kanzler, H., Soumelis, V., Gilliet, M. (2001). Dendritic cell lineage, plasticity and cross-regulation. *Nat. Immunol.* 2: 585-589.

Lore, K., Sonnerborg, A., Spetz, A.L., Andersson, U., Andersson, J. (1998). Immunocytochemical detection of cytokines and chemokines in Langerhans cells and in vitro derived dendritic cells. *J. Immunol. Methods* 218: 173-187 (Erratum).

Ma, X., Trinchieri, G. (2001). Regulation of interleukin-12 production in antigen-presenting cells. *Adv. Immunol.* 79: 55-92.

Mackey, M.F., Wang, Z., Eichelberg, K., Germain R.N. (2003). Distinct contributions of different CD40 TRAF binding sites to CD154-induced dendritic cell maturation and IL-12 secretion. *Eur. J. Immunol.* 33: 779-789.

MacPherson, G.G., Liu, L.M. (1999). Intestinal dendritic cells. In: *Dendritic cells*. Eds. Lotze, M.T. und Thomson, A.W., Academic Press, San Diego, pp. 141-151.

Manz, M.G., Traver, D., Miyamoto, T., Weissman, I.L., Akashi, K. (2001). Dendritic cell potentials of early lymphoid and myeloid progenitors. *Blood* 97: 3333-3341.

Matzinger, P. (1994). Tolerance, danger, and the extended family. *Annu. Rev. Immunol.* 12: 991-1045.

McCurdy, J.D., Olynch, T.J., Maher, L.H., Marshall, J.S. (2003). Cutting edge: distinct Toll-like receptor 2 activators selectively induce different classes of mediator production from human mast cells. *J. Immunol.* 170: 1625-1629.

McDermott, R., Bausinger, H., Fricker, D., Spehner, D., Proamer, F., Lipsker, D., Cazenave, J.P., Goud, B., De La Salle, H., Salamelo, J., Hanau, D. (2004). Reproduction of Langerin/CD207 traffic and Birbeck granule formation in a human cell line model. *J. Invest. Dermatol.* 123: 72-77.

McLellan, A.D., Heiser, A., Sorg, R.V., Fearnley, D.B., Hart D.N. (1998). Dermal dendritic cells associated with T lymphocytes in normal human skin display an activated phenotype. *J. Invest. Dermatol.* 111: 841-849.

McMenamin, P.G., Forrester, J.V. (1999). Dendritic cells in the central nervous system and eye and their associated supporting tissues. In: *Dendritic cells*. Eds. Lotze, M.T. und Thomson, A.W., Academic Press, San Diego, pp. 205-248.

Merad, M., Manz, M.G., Karsunky, H., Wagers, A., Peters, W., Charo, I., Weissman, I.L., Cyster, J.G., Engleman, E.G. (2002). Langerhans cells renew in the skin throughout life under steady-state conditions. *Nat. Immunol.* 3: 1135-1141.

Merad, M., Fong, L., Bogenberger, J., Engleman, E.G. (2000). Differentiation of myeloid dendritic cells into CD8alpha-positive dendritic cells in vivo. *Blood* 96: 1865-1872.

Miltenyi, S., Pflüger, E. (1992). High gradient magnetic cell sorting. In: *Flow Cytometry and Cell Sorting*. Ed. Radbruch, A., Springer-Verlag, Berlin, pp. 133-140.

Mitsui, H., Watanabe, T., Saeki, H., Mori, K., Fujita, H., Tada, Y., Asahina, A., Nakamura, K., Tamaki, K. (2004). Differential expression and function of toll-like receptors in Langerhans cells: comparison with splenic dendritic cells. *J. Invest. Dermatol.* 122: 95-102.

Mohamadzadeh, M., Berard, F., Essert, G., Chalouni, C., Pulendran, B., Davoust, J., Bridges, G., Palucka, A.K., Banchereau, J. (2001). Interleukin 15 skews monocyte differentiation into dendritic cells with features of Langerhans cells. *J. Exp. Med.* 194: 1013-1020.

Monks, C.R., Freiberg, B.A., Kupfer, H., Sciaky, N., Kupfer, A. (1998). Three-dimensional segregation of supramolecular activation clusters in T cells. *Nature* 395: 82-86.

Moody, D.B., Porcelli, S.A. (2003). Intracellular pathways of CD1 antigen presentation. *Nat. Rev. Immunol.* 3: 11-22.

Moody, D.B., Briken, V., Cheng, T.Y., Roura-Mir, C., Guy, M.R., Geho, D.H., Tykocinski, M.L., Besra, G.S., Porcelli, S.A. (2002). Lipid length controls antigen entry into endosomal and nonendosomal pathways for CD1b presentation. *Nat. Immunol.* 3: 435-442.

Morhenn, V.B., Benike, C.J., Charron, D.J., Cox, A., Mahrle, G., Wood, G., Engleman, E.G. (1982). Use of the Fluorescence-Activated Cell Sorter to quantitate and enrich for subpopulations of human skin cells. *J. Invest. Dermatol.* 79: 277-281.

Morris, J., Alaibac, M., Jia, M.H., Chu, T. (1992). Purification of functional active epidermal Langerhans cells: a simple and efficient new technique. *J. Invest. Dermatol.* 99: 237-240.

Morse, M.A., Coleman R.E., Akabani, G., Niehaus, N., Coleman, D., Lyerly, H.K (1999). Migration of human dendritic cells after injection in patients with metastatic malignancies. *Cancer Res.* 59: 56-58.

Moschella, F., Maffei, A., Catanzaro, R.P., Papadopoulos, K.P., Skerrett, D., Hesdorffer, C.S., Harris P.E. (2001). Transcript profiling of human dendritic cells maturation-induced under defined culture conditions: comparison of the effects of tumour necrosis factor alpha, soluble CD40 ligand trimer and interferon gamma. *Br. J. Haematol.* 114: 444-457.

Munn, D.H., Sharma, M.D., Lee, J.R., Jhaver, K.G., Johnson, T.S., Keskin, D.B., Marshall, B., Chandler, P., Antonia, S.J., Burgess, R., Slingluff Jr., C.L., Mellor, A.L. (2002). Potential regulatory function of human dendritic cells expresssing indoleamine 2,3-dioxygenase. *Science* 297: 1867-1869.

Murphy, K.M., Reiner, S.L. (2002). The lineage decisions of helper T cells. *Nat. Rev. Immunol.* 2: 933-944.

Muzio, M., Natoli, G., Saccani, S., Levrero, M., Mantovani, A. (1998). The human toll signaling pathway: divergence of nuclear factor kappaB and JNK/SAPK activation upstream of tumor necrosis factor receptor-associated factor 6 (TRAF6). *J. Exp. Med.* 187: 2097-2101.

Nakagawa, S., Koomen, C.W., Bos, J.D., Teunissen, M.B. (1999). Differential modulation of human epidermal Langerhans cell maturation by ultraviolet B radiation. *J. Immunol.* 163: 5192-5200.

Nilsson, H., Johansson, C., Scheynius, A. (1987). Removal of Langerhans cells from human epidermal cell suspensions by immunomagnetic particles. *J. Immunol. Meth.* 105: 165-169.

Oppmann, B., Lesley, R., Blom, B., Timans, J.C., Xu, Y., Hunte, B., Vega, F., Yu, N., Wang, J., Singh, K., Zonin, F., Vaisberg, E., Churakova, T., Liu, M., Gorman, D., Wagner, J., Zurawski, S., Liu, Y., Abrams, J.S., Moore, K.W., Rennick, D., de Waal-Malefyt, R., Hannum, C., Bazan, J.F., Kastelein, R.A. (2000). Novel p19 protein engages IL-12p40 to form a cytokine, IL-23, with biological activities similar as well as distinct from IL-12. *Immunity* 13: 715-725.

Ortner, U., Inaba, K., Koch, F., Heine, M., Miwa, M., Schuler, G., Romani, N. (1996). An improved isolation method for murine migratory cutaneous dendritic cells. *J. Immunol. Meth.* 193: 71-79.

Osada, A., Nakashima, H., Furue, M., Tamaki, K. (1995). Up-regulation of CD44 expression by tumor necrosis factor- α is neutralized by Interleukin-19 in Langerhans cells. *J. Invest. Dermatol.* 105: 124-127.

Osugi, Y., Vuckovic, S., Hart, D.N. (2002). Myeloid blood CD11c $^{+}$ dendritic cells and monocyte-derived dendritic cells differ in their ability to stimulate T lymphocytes. *Blood* 100: 2858-2866.

Peguet-Navarro, J., Dalbiez-Gauthier, C., Rattis, F.M., Van Kooten, C., Banchereau, J., Schmitt, D. (1995). Functional expression of CD40 antigen on human epidermal Langerhans cells. *J. Immunol.* 155: 4241-4247.

Pena-Cruz, V., Ito, S., Oukka, M., Yoneda, K., Dascher, C.C., Von Lichtenberg, F. (2001). Extraction of human Langerhans cells: a method for isolation of epidermis-resident dendritic cells. *J. Immunol. Meth.* 255: 83-91.

Pflanz, S., Timans, J.C., Cheung, J., Rosales, R., Kanzler, H., Gilbert, J., Hibbert, L., Churakova, T., Travis, M., Vaisberg, E., Blumenschein, W.M., Mattson, J.D., Wagner, J.L., To, W., Zurawski, S., McClanahan, T.K., Gorman, D.M., Bazan, J.F., de Waal Malefyt, R., Rennik, D., Kastelein, R.A. (2002). IL-27, a heterodimeric cytokine composed of EBI3 and p28 Protein, induces Proliferation of naïve CD4 $^{+}$ T cells. *Immunity* 16: 779-790.

Piccioli, D., Sbrana, S., Melandri, E., Valiante, N.M. (2002). Contact-dependent stimulation and inhibition of dendritic cells by natural killer cells. *J. Exp. Med.* 195: 335-341.

Picut, C.A., Lee, C. S., Dougherty, E.P., Andersen, K.L., Lewis, R.M. (1987). Purification and identification of murine epidermal dendritic cells: flow cytometry and immunogold labeling. *J. Histochem. Cytochem.* 35: 745-753.

Pope, M., Betjes, M.G., Hirmand, H., Hoffman, L., Steinman, R.M. (1995). Both dendritic cells and memory T lymphocytes emigrate from organ cultures of human skin and form distinctive dendritic-T-cell conjugates. *J. Invest. Dermatol.* 104: 11-17.

Porcelli, S.A., Modlin, R.T. (1999). The CD1 system: antigen-presenting molecules for T cell recognition of lipids and glycolipids. *Annu. Rev. Immunol.* 17: 297-329.

Quezada, S.A., Jarvinen, L.Z., Lind, E.F., Noelle, R.J. (2004). CD40/CD154 interactions at the interface of tolerance and immunity. *Annu. Rev. Immunol.* 22: 307-328.

Rattis, F.M., Concha, M., Dalbiez-Gauthier, C., Courtellemont, P., Schmitt, D., Peguet-Navarro, J. (1998). Effects of ultraviolet B radiation on human Langerhans cells: functional alteration of CD86 upregulation and induction of apoptotic cell death. *J. Invest. Dermatol.* 111: 373-379.

Ratzinger, G., Baggers, J., de Cos, M.A., Yuan, J., Dao, T., Reagan, J.L., Münz, C., Heller, G., Young, J.W. (2004). Mature human Langerhans cells derived from CD34⁺ Hematopoietic progenitors stimulate greater cytolytic T lymphocyte activity in the absence of bioactive IL-12p70, by either single peptide presentation or cross-priming, than do dermal-interstitial or monocyte-derived dendritic cells. *J. Immunol.* 173: 2780-2791.

Reddy, A., Sapp, M., Feldman, M., Subklewe, M., Bhardwaj, N. (1997). A monocyte conditioned medium is more effective than defined cytokines in mediating the terminal maturation of human dendritic cells. *Blood* 90: 3640-3646.

Reis e Sousa, C., Sher, A., Kaye, P. (1999). The role of dendritic cells in the induction and regulation of immunity to microbial infection. *Curr. Opin. Immunol.* 11: 392-399.

Riedl, E., Stockl, J., Majdic, O., Scheinecker, C., Rappersberger, K., Knapp, W., Strobl, H. (2000). Functional involvement of E-cadherin in TGF-β1-induced cell cluster formation of in vitro developing human Langerhans-type dendritic cells. *J. Immunol.* 165: 1381-1386.

Rodriguez, A., Regnault, A., Kleijmeer, M., Ricciardi-Castagnoli, P., Amigorena, S. (1999). Selective transport of internalized antigens to the cytosol for MHC class I presentation in dendritic cells. *Nat. Cell. Biol.* 1: 362-368.

Romani, N., Holzmann, S., Tripp, C.H., Koch, F., Stroitzner, P. (2003). Langerhans cells-dendritic cell of the epidermis. *APMIS* 111: 725-740.

Romani, N., Gruner, S., Brang, D., Kämpgen, E., Lenz, A., Trockenbacher, B., Konwalinka, G., Fritsch, P.O., Steinman, R.M., Schuler, G. (1994). Proliferating dendritic cell progenitors in human blood. *J. Exp. Med.* 180: 83-93.

Romani, N., Koide, S., Crowley, M., Witmer-Pack, M., Livingstone, A.M., Fathman, C.G., Inaba, K., Steinman, R.M. (1989). Presentation of exogenous protein antigens by dendritic cells to T cell clones. Intact protein is presented best by immature, epidermal Langerhans cells. *J. Exp. Med.* 169: 1169-1178.

Safarik, I., Safarikova, M. (1999). Use of magnetic techniques for the isolation of cells. *J. Chromatogr. B. Biomed. Sci. Appl.* 722: 33-53.

Salamero, J., Bausinger, H., Mommaas, A.M., Lipsker, D., Proamer, F., Cazenave, J.P., Goud, B., de la Salle, H., Hanau, D. (2001). CD1a molecules traffic through the early recycling endosomal pathway in human Langerhans cells. *J. Invest. Dermatol.* 116: 401-408.

Salgado, C.G., Nakamura, K., Sugaya, M., Tada, Y., Asahina, A., Koyama, Y., Irie, S., Tamaki, K. (1999). Functional CD40 ligand is expressed on epidermal Langerhans cells. *J. Leukoc. Biol.* 66: 281-285.

Sallusto, F., Lenig, D., Forster, R., Lipp, M., Lanzavecchia, A. (1999). Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature* 401: 659-660.

Sallusto, F., Cella, M., Danieli, C., Lanzavecchia, A., (1995). Dendritic cells use macropinocytosis and the mannose receptor to concentrate antigen to the MHC class II compartment. Downregulation by cytokines and bacterial products. *J. Exp. Med.* 182: 389-400.

Sallusto, F., Lanzavecchia, A. (1994). Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor plus interleukin 4 and downregulated by tumor necrosis factor α . *J. Exp. Med.* 179: 1109-1118.

Sauder, D.N., Dinarello, C.A., Morhenn, V.B. (1984). Langerhans cell production of interleukin-1. *J. Invest. Dermatol.* 82: 605-607.

Schmitt, D., Dezutter-Dambuyant, C., Brochier, J., Thivolet, J. (1986). Subclustering of CD1 monoclonal antibodies based on the reactivity on human Langerhans cells. *Immunol. Lett.* 12: 231-235.

Schreiber, S., Kilgus, O., Payer, E., Kutil, R., Elbe, A., Mueller, C., Stingl, G. (1992). Cytokine pattern of Langerhans cells isolated from murine epidermal cell cultures. *J. Immunol.* 149: 3524-3534.

Schuler, G., Steinman, R.M. (1985). Murine epidermal Langerhans cells mature into potent immunostimulatory dendritic cells in vitro. *J. Exp. Med.* 161: 526-546.

Schulz, O., Edwards, A.D., Schito, M., Aliberti, J., Manickasingham, S., Sher, A., Reis e Sousa, C. (2000). CD40 triggering of heterodimeric IL-12p70 production by dendritic cells in vivo requires a microbial priming signal. *Immunity* 13: 453-462.

Schwandner, R., Dziarski, R., Wesche, H., Rothe, M., Kirschning, C.J. (1999). Peptidoglycan- and lipoteichoic acid-induced cell activation is mediated by toll-like receptor 2. *J. Biol. Chem.* 274: 17406-17409.

Shibaki, A., Katz, S.I. (2001). Activation through CD40 ligation induces functional Fas ligand expression by Langerhans cells. *Eur. J. Immunol.* 31: 3006-3015.

Shortman, K., Liu, Y.J. (2002). Mouse and human dendritic cell subtypes. *Nat. Rev. Immunol.* 2: 151-161.

Siegal, F.P., Kadowaki, N., Shodell, M., Fitzgerald-Bocarsly, P.A., Shah, K., Ho, S., Antonenko, S., Liu, Y.J. (1999). The nature of the principal type 1 interferon-producing cells in human blood. *Science* 284: 1835-1837.

Simon, J.C., Dittmar, H.C., de-Roche, R., Wilting, J., Christ, B., Schopf, E. (1995). Rapid purification of human Langerhans cells using paramagnetic microbeads. *Exp. Dermatol.* 4: 155-161.

Steinman, R.M. (2003). Some interfaces of dendritic cell biology. *APMIS* 111: 675-697.

Steinman, R.M., Turley, S., Mellman, I., Inaba, K. (2000). The induction of tolerance by dendritic cells that have captured apoptotic cells. *J. Exp. Med.* 191: 411-416.

Steinman, R.M., Cohn, Z.A. (1973). Identification of a novel cell type in peripheral lymphoid organs of mice. I. Morphology, quantitation, tissue distribution. *J. Exp. Med.* 137: 1142-1162.

Stoitzner, P., Holzmann, S., McLellan, A.D., Ivarsson, L., Stossel, H., Kapp, M., Kammerer, U., Douillard, P., Kampgen, E., Koch, F., Saeland, S., Romani, N. (2003). Visualization and characterization of migratory Langerhans cells in murine skin and lymph nodes by antibodies against Langerin/CD207. *J. Invest. Dermatol.* 120: 266-274.

Stoll, S., Jonuleit, H., Schmitt, E., Muller, G., Yamauchi, H., Kurimoto, M., Knop, J., Enk, A.H. (1998). Production of functional IL-18 by different subtypes of murine and human dendritic cells (DC): DC-derived IL-18 enhances IL-12-dependent Th1 development. *Eur. J. Immunol.* 28: 3231-3239.

Streptoe, R.J., Thomson, A.W. (1999). Dendritic cells in the liver, kidney, heart and pancreas. In: *Dendritic cells*. Eds. Lotze, M.T. und Thomson, A.W., Academic Press, San Diego, pp. 153-177.

Strobl, H., Riedl, E., Scheinecker, C., Bello-Fernandez, C., Pickl, W.F., Rappersberger, K., Majdic, O., Knapp, W. (1996). TGF- β 1 promotes in vitro development of dendritic cells from CD34 $^{+}$ hemopoietic progenitors. *J. Immunol.* 157: 1499-1507.

Strunk, D., Egger, C., Leitner, G., Hanau, D., Stingl, G. (1997). A skin homing molecule defines the Langerhans cell progenitor in human peripheral blood. *J. Exp. Med.* 185: 1131-1136.

Sugita, M., van Der Wel, N., Rogers, R.A., Peters, P.J., Brenner, M.B. (2000). CD1c molecules broadly survey the endocytic system. *Proc. Natl. Acad. Sci. USA* 97: 8445-8450.

Sugita, M., Grant, E.P., van Donselaar, E., Hsu, V.W., Rogers, R.A., Peters, P.J., Brenner, M.B. (1999). Separate pathways for antigen presentation by CD1 molecules. *Immunity* 11: 743-752.

Suss, G., Shortman, K. (1996). A subclass of dendritic cells kills CD4 T cells via Fas/Fas-ligand induced apoptosis. *J. Exp. Med.* 183: 1789-1796.

Symington, F.W., Brady, W., Linsley, P.S. (1993). Expression and function of B7 on human epidermal Langerhans cells. *J. Immunol.* 150: 1286-1295.

Tada, Y., Asahina, A., Fujita, H., Sugaya, M., Tamaki, K. (2003). Langerhans cells do not produce IFN- γ . *J. Invest. Dermatol.* 120: 891-892.

Tada, Y., Asahina, A., Nakamura, K., Miyazono, K., Tomura, M., Fujiwara, H., Tamaki, K. (2001). Transforming growth factor- β up-regulates CD40-engaged IL-12 production of mouse Langerhans cells. *Eur. J. Immunol.* 31: 294-300.

Tada, Y., Asahina, A., Nakamura, K., Tomura, M., Fujiwara, H., Tamaki, K. (2000). Granulocyte/macrophage colony-stimulating factor inhibits IL-12 production of mouse Langerhans cells. *J. Immunol.* 164: 5113-5119.

Takayama, K., Yokozeki, H., Ghoreishi, M., Satoh, T., Katayama, I., Umeda, T., Nishioka, K.J. (1999). IL-4 inhibits the migration of human Langerhans cells through the downregulation of TNF receptor II expression. *J. Invest. Dermatol.* 113: 541-546.

Tang, A., Amagai, M., Granger, L.G., Stanley, J.R., Udey, M.C. (1993). Adhesion of epidermal Langerhans cells to keratinocytes mediated by E-cadherin. *Nature* 361: 82-85.

Tchou, I., Misery, L., Sabido, O., Dezutter-Dambuyant, C., Bourlet, T., Moja, P., Hamzeh, H., Peguet-Navarro, J., Schmitt, D., Genin, C. (2001). Functional HIV CXCR4 coreceptor on human epithelial Langerhans cells and infection by HIV strain X4. *J. Leukoc. Biol.* 70: 313-321.

Teunissen, M.B., Koomen, C.W., Jansen, J., de Waal Malefyt, R., Schmitt, E., Van den Wijngaard, R.M., Das, P.K., Bos, J.D. (1997A). In contrast to their murine counterparts, normal human keratinocytes and human epidermoid cell lines A431 and HaCaT fail to express IL-10 mRNA and protein. *Clin. Exp. Immunol.* 107: 213-223.

Teunissen, M.B., Kapsenberg, M.L., Bos, J.D. (1997B). Langerhans cells and related skin dendritic cells. In: *Skin immune system*. Ed. Bos, J.D., CRC Press, Boca Raton, pp. 59-83.

Teunissen, M.B., Wormmeester, J., Krieg, S.R., Peters, P.J., Vogels, I.M., Kapsenberg, M.L., Bos, J.D. (1990). Human epidermal Langerhans cells undergo profound morphologic and phenotypical changes during in vitro culture. *J. Invest. Dermatol.* 94: 166-173.

Thery, C., Regnault, A., Garin, J., Wolfers, J., Zitvogel, L., Ricciardi-Castagnoli, P., Raposo, G., Amigorena, S. (1999). Molecular characterization of dendritic cell-derived exosomes. Selective accumulation of the heat shock protein hsc73. *J. Cell. Biol.* 147: 599-610.

Trinchieri, G. (2003). Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nat. Rev. Immunol.* 3: 133-146.

Türeci, Ö., Bian, H., Nestle, F.O., Raddrizzani, L., Rosinski, J.A., Tassis, A., Hilton, H., Walstead, M., Sahin, U., Hammer, J. (2003). Cascades of transcriptional induction during dendritic cell maturation revealed by genome-wide expression analysis. *FASEB J.* 17: 836-847.

Turley, S.J., Inaba, K., Garrett, W.S., Ebersold, M., Unternaehrer, J., Steinman, R.M., Mellman, I. (2000). Transport of peptide-MHC class II complexes in developing dendritic cells. *Science* 288: 522-527.

Valladeau, J., Ravel, O., Dezutter-Dambuyant, C., Moore, K., Kleijmeer, M., Liu, Y., Duvert-Frances, V., Vincent, C., Schmitt, D., Davoust, J., Caux, C., Lebecque, S., Saeland, S. (2000). Langerin, a novel C-type lectin specific to Langerhans cells, is an endocytic receptor that induces the formation of Birbeck granules. *Immunity* 12: 71-81.

Vincent, M.S., Leslie, D.S., Gumperz, J.E., Xiong, X., Grant, E.P., Brenner, M.B. (2002). CD1-dependent dendritic cell instruction. *Nat. Immunol.* 3: 1163-1168.

Wang, B., Amerio, P., Sander, D.M. (1999). Role of cytokines in epidermal Langerhans cell migration. *J. Leukoc. Biol.* 66: 33-39.

Weiss, J.M., Sleeman, J., Renkl, A.C., Dittmar, H., Termeer, C.C., Taxis, S., Howells, N., Hofmann, M., Köhler, G., Schöpf, E., Ponta, H., Herrlich, P., Simon, J.C. (1997). An essential role for CD44 variant isoforms in epidermal Langerhans cell and blood dendritic cell function. *J. Cell Biol.* 137: 1137-1147.

Winzler, C., Rovere, P., Rescigno, M., Granucci, F., Penna, G., Adorini, L., Zimmermann, V.S., Davoust, J., Ricciardi-Castagnoli, P. (1997). Maturation stages of mouse dendritic cells in growth factor-dependent long-term cultures. *J. Exp. Med.* 185: 317-328.

Witmer-Pack, M. D., Olivier, W., Valinsky, J., Schuler, G., Steinman, R.M. (1987). Granulocyte/macrophage colony-stimulating factor is essential for the viability and function of cultured murine epidermal Langerhans cells. *J. Exp. Med.* 166: 1484-1498.

Witsch, E.J., Peiser, M., Hutloff, A., Büchner, K., Dorner, B.G., Jonuleit, H., Mages, H.W., Kroczek, R.A. (2002). ICOS and CD28 reversely regulate IL-10 on re-activation of human effector T cells with mature dendritic cells. *Eur. J. Immunol.* 32: 2680-2686.

Wolff, K. (1972). The Langerhans cell. *Curr. Probl. Dermatol.* 4: 79.

Wykes, M., MacPherson, G. (2000). Dendritic cell-B-cell interaction: dendritic cells provide B cells with CD40-independent proliferation signals and CD40-dependent survival signals. *Immunology* 100: 1-3.

Yanagihara, S., Komura, E., Nagafune, J., Watarai, H., Yamaguchi, Y. (1998). EBI1/CCR7 is a new member of dendritic cell chemokine receptor that is up-regulated upon maturation. *J. Immunol.* 161: 3096-3102.

Yang, D., Biragyn, A., Hoover, D.M., Lubkowski, J., Oppenheim, J.J. (2004). Multiple roles of antimicrobial defensins, cathelicidins, and eosinophil-derived neurotoxin in host defense. *Ann. Rev. Immunol.* 22: 181-215.

Zaitseva, M., Blauvelt, A., Lee, S., Lapham, C.K., Klaus-Kovtun, V., Mostowski, H., Manischewitz, J., Golding, H. (1997). Expression and function of CCR5 and CXCR4 on human Langerhans cells and macrophages: Implications for HIV primary infection. *Nat. Med.* 3: 1369-1375.

Zhang, D., Zhang, G., Hayden, M.S., Greenblatt, M.B., Bussey, C., Flavell, R.A., Ghosh, S. (2004). A toll-like receptor that prevents infection by uropathogenic bacteria. *Science* 303: 1522-1526.

Zhou, L.J., Tedder, T.F. (1995). Human blood dendritic cells selectively express CD83, a member of the immunoglobulin superfamily. *J. Immunol.* 154: 3821-3835.

Zoeteweij, J.P., Golding, H., Mostowski, H., Blauvelt, A. (1998). Cytokines regulate expression and function of the HIV coreceptor CXCR4 on human mature dendritic cells. *J. Immunol.* 161: 3219-3223.

PUBLIKATIONEN

Teile dieser Dissertation sind in folgenden Originalartikeln publiziert:

Peiser, M., Wanner, R., Kolde, G. (2004). Human epidermal Langerhans cells differ from monocyte-derived Langerhans cells in CD80 expression and in secretion of IL-12 after CD40 crosslinking. *J. Leukoc. Biol.* 76: 616-622.

Peiser, M., Grützkau, A., Wanner, R., Kolde, G. (2003). CD1a and CD1c cell sorting yields a homogeneous population of immature enriched human Langerhans cells. *J. Immunol. Meth.* 279: 41-53.

Weitere Veröffentlichungen:

Wanner, R., Peiser, M., Wittig, B. (2004). Keratinocytes rapidly readjust ceramide-sphingomyelin homeostasis and contain a phosphatidylcholine-sphingomyelin transacylase. *J. Invest. Dermatol.* 122: 773-782.

Witsch, E.J., Peiser, M., Hutzloff, A., Büchner, K., Dorner, B.G., Jonuleit, H., Mages, H.W., Kroczeck, R.A. (2002). ICOS and CD28 reversely regulate IL-10 on re-activation of human effector T cells with mature dendritic cells. *Eur. J. Immunol.* 32: 2680-2686.

KONGRESSBEITRÄGE

Peiser, M., Wanner, R. (2003). Negative feedback control of neutral sphingomyelinase by ceramide in keratinocytes. *Eur. J. Cell Biol.* 82, Suppl. 53: 34.

Peiser, M., Grützkau, A., Kolde, G. (2001). CD40 crosslinking induces phenotypical, but not functional maturation of human Langerhans cells. Scientific program and abstracts book of *7th international symposium on dendritic cells*: 44.

Peiser, M., Artuc, M., Grützkau, A., Kolde, G. (2001). Isolated human Langerhans cells require keratinocytes for their cellular and functional properties: a coculturing concept. *Arch. Dermatol. Res.* 293: 55.

Peiser, M., Kolde, G. (2000). Stimulus-dependent activation of epidermal human Langerhans cells isolated by Magnetic-activated cell-sorting. *J. Invest. Dermatol.* 115: 57.

Peiser, M., Grützkau, A., Kolde, G. (2000). Immature Langerhans cells from human epidermis can be eliminated via Fas-mediated apoptosis. *Arch. Dermatol. Res.* 292: 109.

Peiser, M., Grützkau, A., Kolde, G. (1999). Magnetic-activated cell sorting of epidermal Langerhans cells yields highly purified cells with different expression of costimulatory molecules. *J. Invest. Dermatol.* 113: 456.