

10. Literaturverzeichnis

ABRAMS, G.D., BAUM, H., SPRINZ, H. (1963).

Influence of the normal flora on mucosal morphology and cellular renewal in the ileum
Lab.Invest. **12**: 355-364

ALM, E.W. and STAHL, D.A. (2000).

Critical factors influencing the recovery and integrity of rRNA extracted from environmental samples: use of an optimized protocol to measure depth-related biomass distribution in freshwater sediments

J.Microb.Meth. **40** (2000): 153-162

ALM, E.W., ZHENG, D., RASKIN, L. (2000).

The presence of humic substances and DNA in RNA extracts affects hybridization results

Appl.Environ.Microbiol. **66** (10): 4547-4554

ALMIRALL, M. and ESTEVE-GARCIA, E. (1994).

Rate of passage of barley diets with chromium oxide: Influence of age and poultry strain and effect of β -glucanase supplementation

Poult.Sci. **73**: 1433-1440

ALMIRALL, M., FRANCESH, M., PEREZ-VENDRELL, A.M., BRUFAU, J., ESTEVE-GARCIA, E. (1995).

The differences in intestinal viscosity produced by barley and β -glucanase alter digesta enzyme activities and ileal nutrient digestibilities more in broiler chicks than in cocks

J.Nutr. **125**: 947-955

ALODAN, M.A. and MASHALY, M.M. (1999).

Effect of induced molting in laying hens on production and immune parameters

Poult.Sci. **78** (2): 171-177

AMANN, R.I., KRUMHOLZ, L., STAHL, D.A. (1990).

Fluorescent-oligonucleotide probing of whole cells for determinative, phylogenetic, and environmental studies in microbiology

J.Bacteriol. **172**: 762-770

AMANN, R., LUDWIG, W., SCHLEIFER, K.H. (1995).

Phylogenetic identification and In Situ Detection of individual microbial cells without cultivation

Microbiol.Rev. **59** (1): 143-159

AMANN, R. and LUDWIG, W. (2000).

Ribosomal RNA-targeted nucleic acid probes for studies in microbial ecology

FEMS Microbiol.Rev. **24** (5): 555-565

ANNISON, E.F., HILL, K.J., KENWORTHY, R. (1968).

Volatile fatty acids in the digestive tract of the fowl

Br.J.Nutr. **22**: 207-216

ANTONIOU, T. and MARQUARDT, R.R. (1981).

Influence of rye pentosans on the growth of chicks

Poult.Sci. **60**: 1898-1904

ANTONIOU, T., MARQUARDT, R.R., CANSFIELD, E. (1981).

Isolation, partial characterization, and antinutritional activity of a factor (pentosans) in rye grain

J.Agr.Food.Chem. **29**: 1240-1247

ANTONIOU, T. and MARQUARDT, R.R. (1982).

Utilization of rye diets by chicks as affected by lipid type and level and penicillin supplementation

Poult.Sci. **61**: 107-116

AUDISIO, M.C., OLIVER, G., APELLA, M.C. (1999).

Antagonistic effect of *Enterococcus faecium* J96 against human and poultry pathogenic *Salmonella* spp.

J.FoodProtec. **62** (7): 751-755

AXELSSON, L. (1998).

In: Lactic acid bacteria. Microbiology and functional aspects. 2nd Edition.

Salminen, S., von Wright, A. (Ed.)

Marcel Dekker Inc., New York

ISBN 0-8247-0133-X

BAILEY, J.S., COX, N.A., BERRANG, M.E. (1994).

Hatchery-acquired salmonellae in broiler chicks

Poult.Sci. **73** (7): 1153-1157

BARNES, E.M. and IMPEY, C.S. (1970).

The isolation and properties of the predominant anaerobic bacteria in the caecum of chickens and turkeys

Br.Poult.Sci. **11**: 467-481

BARNES, E.M., MEAD, G.C., BARNUM, D.A., HARRY, E.G. (1972).

The intestinal flora of the chicken in the period 2 to 6 weeks of age, with particular reference to the anaerobic bacteria

Poult.Sci. **13**: 311-326

BARNES, E.M. (1979).

The intestinal microflora of poultry and game birds during life and after storage

J.Appl.Bacteriol. **46**: 407-419

BARRY, T., COLLERAN, G., GLENNON, M., DUNICAN, L.K., GANNON, F. (1991).

The 16S/23S ribosomal spacer region as a target for DNA probes to identify eubacteria

PCRMethodsApplic. **1** (1): 51-56

BATT, R.M., RUTGERS, H.C., SANCAK, A.A. (1996).

Enteric bacteria: friend or foe?

J.Small.Anim.Pract. **37** (6): 261-267

BECKER, S., BOGER, P., OEHLMANN, R., ERNST, A. (2000).

PCR bias in ecological analysis: a case study for quantitative Taq nuclease assays in analyses of microbial communities

Appl.Environ.Microbiol. **66** (11): 4945-4953

BECKMANN, L. (1999).

Persönliche Mitteilung

BECKMANN, L., VAHJEN, W., SIMON, O. (2000a).

Isolation of an 1,3-1,4-beta-glucan degrading *Enterococcus faecium* strain from the intestinal tract of chicken and partial characterization of its novel 1,3-1,4-beta-glucanase

J.BasicMicrobiol. **40** (5-6): 303-310

BECKMANN, L., VAHJEN, W., SIMON, O. (2000b).

Isolation of 1,3-1,4- β -glucan degrading bacteria from the intestinal tract of chicken

Microbiology 2000, München, März 2000, Poster 15.P.11.24

BEDFORD, M.R. and CLASSEN, H.L. (1992).

Reduction of intestinal viscosity through manipulation of dietary rye and pentosanase concentration is effected through changes in the carbohydrate composition of the intestinal aqueous phase and results in improved growth rate and feed conversion efficiency of broiler chicks

J.Nutr. **122**: 560-569

BEDFORD, M.R. and CLASSEN, H.L. (1993).

An *in vitro* assay for prediction of broiler intestinal viscosity and growth when fed rye-based diets in the presence of exogenous enzymes

Poult.Sci. **72**: 137-143

BEDFORD, M.R. (1995).

Mechanisms of actions and potential environmental benefits from the use of feed enzymes

Anim.Feed.Sci.Tech. **53**: 145-155

BEDFORD, M.R. and MORGAN, A.J. (1996).

The use of enzymes in poultry diets

WorldPoult.Sci.J. **52**: 61-68

BEST, C., SIMON, O., WEYRAUCH, K.D. (1999).

Einfluss der Digestivviskosität auf die Morphologie des Dünndarms von Broilerküken

Proc.Soc.Nutr.Physiol. **8**: 118

BRENES, A., TREVINO, J., CENTENO, C., YUSTE, P. (1989).

Influence of peas (*Pisum sativum*) as a dietary ingredient and flavomycin supplementation on the performance and intestinal microflora of broiler chicks

Br.Poult.Sci. **30**: 81-89

BRENES, A., SMITH, M., GÜNTHER, R.W., MARQUARDT, R.R. (1993).

Effect of enzyme supplementation on the performance and digestive tract size of broiler chickens fed wheat- and barley-based diets

Poult.Sci. **72**: 1731-1739

CAMPBELL, G.L., CAMPBELL, L.D., CLASSEN, H.L. (1983).

Utilisation of rye by chickens: effect of microbial status, diet gamma irradiation and sodium taurocholate supplementation

Br.Poult.Sci. **24**: 191-203

CAMPBELL, G.L., TEITGE, D.A., CLASSEN, H.L. (1991).

Genotypic and environmental differences in rye fed broiler chicks with dietary pentosanase supplementation

Can.J.Anim.Sci. **71**: 1241-1247

CARRE, B., DEROUET, L., LECLERCQ, B. (1990).

The digestibility of cell-wall polysaccharides from wheat (bran or whole grain), soybean meal and white lupin meal in cockerels, muscovy ducks and rats
Poult.Sci. **69**: 623-633

CHOCT, M. and ANNISON, G. (1990).

Anti-nutritive activity of wheat pentosans in broiler diets
Br.Poult.Sci. **31**: 811-821

CHOCT, M. and ANNISON, G. (1992a).

The inhibition of nutrient digestion by wheat pentosans
Br.J.Nutr. **67**: 123-132

CHOCT, M. and ANNISON, G. (1992b).

Anti-nutritive effect of wheat pentosans in broiler chickens: Role of viscosity and gut microflora
Br.Poult.Sci. **33**: 821-834

CHOCT, M., HUGHES, R.J., TRIMBLE, R.P., ANGKANAPORN, K., ANNISON, G. (1995).

Non-starch polysaccharide-degrading enzymes increase the performance of broiler chickens fed wheat of low apparent metabolizable energy
J.Nutr. **125** (3): 485-492

CHOCT, M., HUGHES, R.J., WANG, J., BEDFORD, M.R., MORGAN, A.J., ANNISON, G. (1996).

Increased small intestine fermentation is partly responsible for the anti-nutritive activity of non-starch polysaccharides in chickens
Br.Poult.Sci. **37**: 609-621

CLAUSEN, M.R. and MORTENSEN, P.B. (1994).

Kinetic studies on the metabolism of short-chain fatty acids and glucose by isolated rat colonocytes
Gastroentology **106** (2): 423-432

COATES, M.E., FULLER, R., HARRISON, G.F., LEV, M., SUFFOLK, S.F. (1963).

A comparison of the growth of chicks in the Gustafsson germ-free apparatus and in a conventional environment, with and without dietary supplements of penicillin
Br.J.Nutr. **17**: 141-150

COATES, M.E., COLE, C.B., FULLER, R., HOUGHTON, S.B., YOKOTA, H. (1981).

The gut microflora and the uptake of glucose from the small intestine of the chick
Br.Poult.Sci. **22**: 289-294

COLE, J.R. and BOYD, F.M. (1967).

Fat absorption from the small intestine of gnotobiotic chicks
Appl.Microbiol. **15** (5): 1229-1234

COLE, C.B. and FULLER, R. (1984).

Bile acid deconjugation and attachment of chicken gut bacteria: their possible role in growth depression
Br.Poult.Sci. **25** (2): 227-231

COX, N.A., BAILEY, J.S., MAULDIN, J.M., BLANKENSHIP, L.C. (1990).

Presence and impact of salmonella contamination in commercial broiler hatcheries
Poult.Sci. **69** (9): 1606-1609

- COX, N.A., BAILEY, J.S., BERRANG, M.E., BUHR, R.J., MAULDIN, J.M. (1994).**
Automated spray sanitizing of broiler hatching eggs. 3. Total bacteria and coliform recovery after using an egg spraying machine
J.Appl.Poult.Res. **3**: 234-237
- DÄNICKE, S., SIMON, O., JEROCH, H., BEDFORD, M. (1997a).**
Interactions between dietary fat type and xylanase supplementation when rye-based diets are fed to broiler chickens. 1. Physico-chemical chyme features
Br.Poult.Sci. **38**: 537-545
- DÄNICKE, S., SIMON, O., JEROCH, H., BEDFORD, M. (1997b).**
Interactions between dietary fat type and xylanase supplementation when rye-based diets are fed to broiler chickens. 2. Performance, nutrient digestibility and the fat soluble vitamin status of livers
Br.Poult.Sci. **38**: 546-556
- DÄNICKE, S., JEROCH, H., BÖTTCHER, W., BEDFORD, M., SIMON, O. (1999a).**
Effects of dietary fat type, pentosan level, and xylanases on digestibility of fatty acids, liver lipids, and of vitamin E in broilers
Fett/Lipid **101**: 90-100
- DÄNICKE, S., JEROCH, H., DUSEL, G., SIMON, O., BEDFORD, M. (1999b).**
Effect of dietary fat type, pentosan level and xylanase supplementation on performance and carcass characteristics of male broilers
Arch.Geflügelk. **63**: 194-203
- DÄNICKE, S., JEROCH, H., DUSEL, G., SIMON, O., BEDFORD, M. (1999c).**
Interactions between dietary fat type and exogenous enzyme supplementation of broiler diets based on maize, wheat, triticale or barley
J.Anim.FeedSci. **8**: 467-483
- DÄNICKE, S., SIMON, O., JEROCH, H., KELLER, K., GLÄSER, K., KLUGE, H., BEDFORD, M.R. (1999d).**
Effects of dietary fat type, pentosan level and xylanase supplementation on digestibility of nutrients and metabolizability of energy in male broilers
Arch.Anim.Nutr. **52**: 245-261
- DÄNICKE, S., VAHJEN, W., SIMON, O., JEROCH, H. (1999e).**
Effects of dietary fat type and xylanase supplementation to rye-based broiler diets on selected bacterial groups adhering to the intestinal epithelium, on transit time of feed, and on nutrient digestibility
Poult.Sci. **78**: 1292-1299
- DÄNICKE, S., BÖTTCHER, W., JEROCH, H., THIELEBEIN, J., SIMON, O. (2000a).**
Replacement of soybean oil with tallow in rye-based diets without xylanase increases protein synthesis in small intestine of broilers
J.Nutr. **130**: 827-834
- DÄNICKE, S., JEROCH, H., SIMON, O. (2000b).**
Endogenous N-losses in broilers estimated by a [¹⁵N]-isotope dilution technique: Effect of dietary fat type and xylanase addition
Arch.Anim.Nutr. **53**: 75-97

DEHORTY, B.A. (1991).

Effects of microbial synergism on fibre digestion in the rumen
Proc.Nutr.Soc. **50** (2): 149-159

DEVRIESE, L.A., HOMMEZ, J., WIJFELS, R., HAESEBROUCK, F. (1991).

Composition of the enterococcal and streptococcal intestinal flora of poultry
J.Appl.Bacteriol. **71**: 46-50

DHO-MOULIN, M. and FAIRBROTHER, J.M. (1999).

Avian pathogenic *Escherichia coli* (APEC)
Vet.Res. **30** (2-3): 299-316

DORE, J. (1995).

Molecular ecology of the human colonic microflora based on the utilization of RNA-targeted hybridization probes
In: DIFE Workshop on molecular approaches to the study of interactions between dietary polysaccharides, the gut flora and its tissues. Potsdam-Rehbrügge (1995)

DUSEL, G., KLUGE, H., JEROCH, H., SIMON, O. (1998).

Xylanase supplementation of wheat-based broiler rations for broilers: influence of wheat characteristics
J.Appl.Poult.Res. **7**: 119-131

EATON, T.J. and GASSON, M.J. (2001).

Molecular screening of *Enterococcus* virulence determinants and potential for genetic exchange between food and medical isolates
Appl.Environ.Microbiol. **67** (4): 1628-1635

EKINCI, M.S., MCCRAE, S.I., FLINT, H.J. (1997).

Isolation and overexpression of a gene encoding an extracellular β -(1,3-1,4)-glucanase from *Streptococcus bovis* JB1
Appl.Environ.Microbiol. **63** (10): 3752-3756

ELKIN, R.G., WOOD, K.V., HAGEY, L.R. (1990).

Biliary bile acid profiles of domestic fowl as determined by high performance liquid chromatography and fast atom bombardment mass spectrometry
Comp.Biochem.Physiol.B. **96**: 157-161

ELWINGER, K. and TEGLÖF, B. (1991).

Performance of broiler chickens as influenced by a dietary enzyme complex with and without antibiotic supplementation
Arch.Geflügelk. **55** (2): 69-73

EL-ZINEY, M.G., VAN DEN TEMPEL, T., DEBEVERE, J., JAKOBSEN, M. (1999).

Application of reuterin produced by *Lactobacillus reuteri* 12002 for meat decontamination and preservation
J.FoodProtec. **62** (3): 257-261

ERNÄHRUNGS- UND AGRARPOLITISCHER BERICHT 2002 DER BUNDESREGIERUNG
Bundesministerium für Verbraucherschutz, Ernährung und Landwirtschaft (Hrsg.)

FEIGHNER, S.D. and DASHKEVICZ, M.P. (1987).

Subtherapeutic levels of antibiotics in poultry feeds and their effects on weight gain, feed efficiency, and bacterial cholytaurine hydrolase activity
Appl.Environ.Microbiol. **53** (2): 331-336

FEIGHNER, S.D. and DASHKEVICZ, M.P. (1988).

Effect of dietary carbohydrates on bacterial cholytaurine hydrolase in poultry intestinal homogenates

Appl.Environ.Microbiol. **54** (2): 337-342

FENGLER, A.I., PAWLICK, J.R., MARQUARDT, R.R. (1988).

Improvement in nutrient retention and changes in excreta viscosity in chicks fed rye-containing diets supplemented with fungal enzymes, sodium taurocholate and penicillin

Can.J.Anim.Sci. **68**: 483-491

FLINT, H.J., MARTIN, J., MCPHERSON, C.A., DANIEL, A.S., ZHANG, J.X. (1993).

A bifunctional enzyme with separate xylanase and β -(1,3-1,4)-glucanase domains, encoded by the xynD gene of *Ruminococcus flavefaciens*

J.Bacteriol. **175**: 2943-2951

FOGEL, G.B., COLLINS, C.R., BRUNK, C.F. (1999).

Prokaryotic genome size and SSU rDNA copy number: Estimation of microbial relative abundance from a mixed population

Microb.Ecol. **38**: 93-113

FORBES, M. and PARK, J.T. (1959).

Growth of germ-free and conventional chicks: effect of diet, dietary penicillin and bacterial environment

J.Nutr. **67**: 69-84

FOX, G.E., WISOTZKEY, J.D., JURTSUK, P. JR (1992).

How close is close: 16S rRNA sequence identity may not be sufficient to guarantee species identity

Int.J.Syst.Bacteriol. **42**: 166-170

FRAHM, E., HEIBER, I., HOFFMANN, S., KOOB, C., MEIER, H., LUDWIG, W., AMANN, R., SCHLEIFER, K.H., OBST, U. (1998).

Application of 23S rDNA-targeted oligonucleotide probes specific for Enterococci to water hygiene control

System.Appl.Microbiol. **21**: 450-453

FRANKS, A.H., HARMSSEN, H.J.M., RAANGS, G.C., JANSEN, G.J., SCHUT, F., WELLING, G.W. (1998).

Variation of bacterial populations in human faeces measured by Fluorescent In Situ Hybridization with group-specific 16S rRNA-targeted oligonucleotide probes

Appl.Env.Microbiol. **64** (9): 3336-3345

FRY, R.E., ALLRED, J.B., JENSEN, L.S., MCGINNIS, J. (1957a).

Influence of water-treatment on nutritional value of barley

Proc.Soc.Exp.Biol.Med. **95**: 249-251

FRY, R.E., ALLRED, J.B., JENSEN, L.S., MCGINNIS, J. (1957b).

Influence of cereal grain components of the diet on the response of chicks and poult to dietary enzyme supplements

Poult.Sci. **36**: 1120

FULLER, R. and TURVEY, A. (1971).

Bacteria associated with the intestinal wall of the fowl (*Gallus domesticus*)

J.Appl.Bact. **34**: 617-622

FULLER, R. (1977).

The importance of Lactobacilli in maintaining normal microbial balance in the crop
Br.Poult.Sci. **18**: 85-94

FULLER, R. (1984).

Microbial activity in the alimentary tract of birds
Proc.Nutr.Soc. **43**: 55-61

FURUSE, M., YANG, S.I., NIWA, N., OKUMERA, J. (1991).

Effect of short chain fatty acids on the performance and intestinal weight in germ free and conventional chicks
Br.Poult.Sci. **32**: 159-165

GARRET, R.L. and YOUNG, J. (1975).

Effect of micelle formation on the absorption of neutral fat and fatty acids by the chicken
J.Nutr. **105**: 827-838

GERLACH, H. (1994).

Funktion und Entwicklung der autochtonen Darmflora beim Wirtschaftsgeflügel
Tierärztl.Praxis **22**: 141-145

GÖBEL, S., VAHJEN, W., JADAMUS, A., SIMON, O. (2000).

PCR-Assay zum Nachweis schweinepathogener *Escherichia coli* Virulenzfaktoren aus dem Verdauungstrakt von Ferkeln bei Einsatz eines sporenbildenden Probiotikums
Proc.Soc.Nutr.Physiol. **9**: 64

GOLDBERG, M. (2002).

Entwicklung und Einsatz von 16S rRNA Sonden zur Identifizierung biotechnologisch genutzter Laktobazillus-Stämme der *L.acidophilus*- und *L.casei*-Gruppe
Dissertation, Freie Universität Berlin

GOLLNICH, K. (1998).

Einfluß eines antimikrobiellen Zusatzstoffes (Avilamycin) und eines Nicht-Stärke-Polysaccharid (NSP)-hydrolysierenden Enzyms allein und in Kombination auf die mikrobielle Darmbesiedlung beim Schwein
Dissertation, TiHo, Hannover

GRAHAM, H. (1996).

Feed enzymes influence intestinal microflora
Zootechnica International **10**: 88-92

GREEN, J. and KELLOG, T.F. (1987).

Bile acid concentration in serum, bile, jejunal contents, and excreta of male broiler chicks during the first six weeks posthatch
Poult.Sci. **66**: 535-540

GREENBERG, E.P. and CANALE-PAROLA, E. (1977).

Motility of flagellated bacteria in viscous environments
J.Bacteriol. **132**: 356-358

HEDGE, S.N., ROLLS, B.A., COATES, M.E. (1982).

The effects of gut microflora and dietary fibre on energy utilization by the chick
Br.J.Nutr. **48**: 73-80

HESELMANN, K. and ÅMAN, P. (1986).

The effect of β -glucanase on the utilization of starch and nitrogen by broiler chickens fed on barley of low- or high-viscosity

Anim.FeedSci.Techol. **15**: 83-93

HOCK, E., HALLE, I., MATTHES, S., JEROCH, H. (1997a).

Effekte eines NSP-hydrolysierenden Enzymes und Zinkbacitracin auf die mikrobielle Besiedlung im Ileum und Caecum von Broilern

In: Vitamine und Zusatzstoffe in der Ernährung von Mensch und Tier

6. Symposium 24./25.1997 in Jena/Thüringen

Schubert, R., Flachowsky, G., Bitsch, R., Jahreis, G. (Hrsg.)

Buch- und Kunstdruckerei Keßler GmbH, Weimar

ISBN 3-00-002381-X

HOCK, E., HALLE, I., MATTHES, S., JEROCH, H. (1997b).

Investigations on the composition of the ileal and caecal microflora of broiler chicks in consideration to dietary enzyme preparation and zinc bacitracin in wheat-based diets

Agribiol.Res. **50**: 85-95

HÜBENER, K., VAHJEN, W., SIMON, O. (1998).

Einfluß des Gehaltes von Nicht-Stärke-Polysacchariden im Futter auf die Mikroflora im Broiler

Proc.Soc.Nutr.Physiol. **7**: 77

HÜBENER, K. (2001).

Untersuchungen zur Wirkungsweise einer Xylanase im Verdauungstrakt von Masthühnern

Dissertation, Freie Universität Berlin

HÜBENER, K., VAHJEN, W., SIMON, O. (2002).

Bacterial responses to different dietary cereal types and xylanase supplementation in the intestine of broiler chicken

Arch.Anim.Nutr. **56**: 167-187

HUHTANEN, C.N. and PENSACK, J.M. (1965).

The development of the intestinal flora of the young chick

Poult.Sci. **44**: 825-830

HURWITZ, S., BAR, A., KATZ, M., SKLAN, D., BUDOWSKI, P. (1973).

Absorption and secretion of fatty acids and bile acids in the intestine of laying fowl

J.Nutr. **103**: 543-547

IJI, P.A., SAKI, A.A., TIVEY, D.R. (2001).

Intestinal development and body growth of broiler chicks on diets supplemented with non-starch-polysaccharides

Anim.Feed.Sci.Technol. **89** (3-4): 175-188

ISAKSSON, G., LILJA, P., LUNDQUIST, I., IHSE, I. (1983).

Influence of dietary fibre on exocrine pancreatic function in the rat

Digestion **27**: 57-62

JACKISCH, B. and JEROCH, H. (1990).

Untersuchungen zur Verbesserung des Futterwertes von roggenreichem Broilermastfutter durch Enzymeinsatz

Arch.Anim.Nutr. **40**: 1109-1118

- JAMROZ, D., EDER, K., WILICZKIEWICZ, A., KIRCHGESSNER, M. (1998a).**
Verdaulichkeit der NSP-gebundenen Zucker bei Verfütterung von Triticale und Enzymen an Hähnchen, Enten und Gänse
J.Anim.Physiol.a.Anim.Nutr. **79**: 113-122
- JAMROZ, D., WILICZKIEWICZ, A., SKORUPINSKA, J., ORDA, J. (1998b).**
Fermentation und scheinbare Verdaulichkeit der Gerüstkohlenhydrate bei Verfütterung von Triticale und Enzymen an Hähnchen, Enten und Gänse
J.Anim.Physiol.a.Anim.Nutr. **79**: 1-17
- JARONI, D., SCHEIDELER, E., BECK, M.M., WYATT, C. (1999).**
The effect of wheat middlings and enzyme supplementation II: Apparent nutrient digestibility, digestive tract size, gut viscosity and gut morphology in two strains of leghorn hens
Poult.Sci. **78**: 1664-1674
- JASKARI, J., KONTULA, P., SIITONEN, A., JOUSIMIES-SOMER, H., MATTILA-SANDHOLM, T., POUTANEN, K. (1998).**
Oat beta glucan and xylan hydrolysates as selective substrates for *Bifidobacterium* and *Lactobacillus* strains
Appl.Microbiol.Biotechnol. **49** (2): 175-181
- JAYNE-WILLIAMS, D.J. and FULLER, R. (1971).**
The influence of the intestinal microflora on nutrition
In: Physiology and biochemistry of the domestic fowl
D.J. Bell and B.M. Freeman (Ed.)
Vol.1; Academic Press, London, New York
- JEROCH, H. (1998).**
In: Jahrbuch für die Geflügelwirtschaft, 1998
Verlag Eugen Ulmer, Stuttgart
- JEROCH, H., DROCHNER, W., SIMON, O. (1999).**
Ernährung landwirtschaftlicher Nutztiere
Verlag Eugen Ulmer, Stuttgart, ISBN 3-8252-8189-9
- JIN, L.Z., HO, Y.W., ALI, M.A., ABDULLAH, N., JALALUDIN, S. (1996).**
Effect of adherent *Lactobacillus spp.* on *in vitro* adherence of salmonellae to the intestinal epithelial cells of chicken
J.Appl.Bacteriol. **81** (2): 201-206
- JOHNSON, I.T. and GEE, J.M. (1981).**
Effect of gel-forming gums on the intestinal unstirred water layer and sugar transport *in vitro*
Gut **22** (5): 398-403
- JONES, D.M., SUTCLIFFE, E.M., CURRY, A. (1991).**
Recovery of viable but non-culturable *Campylobacter jejuni*
J.Gen.Microbiol. **137**: 2477-2482
- JØRGENSEN, H., ZHAO, X.Q., BACH KNUDSEN, K.E., EGGUM, B.O. (1996).**
The influence of fibre source and level on the development of the gastrointestinal tract, digestibility and energy metabolism in broiler chicken
Br.J.Nutr. **75**: 379-395

- JOYA, J.E., TSUJI, T., JACALNE, A.V., ARITA, M., TSUKAMOTO, T., HONDA, T., MIWATANI, T. (1990).**
Demonstration of enterotoxigenic *Escherichia coli* in diarrheic broiler chicks
Eur.J.Epidemiol. **6** (1): 88-90
- KAUFMANN, P., PFEFFERKORN, A., TEUBER, M., MEILE, L. (1997).**
Identification and quantification of *Bifidobacterium species* isolated from food with genus specific 16S rRNA-targeted probes by colony hybridization and PCR
Appl.Env.Microbiol. **63** (4): 1268-1273
- KAUKAS, A., HINTON, M., LINTON, A.H. (1988).**
The effect of growth-promoting antibiotics on the faecal enterococci of healthy young chicks
J.Appl.Bacteriol. **64**: 57-64
- KOSUGI, A., MURASHIMA, K., DOI, R.H. (2001).**
Characterization of xylanolytic enzymes in *Clostridium cellulovorans*: expression of xylanase activity dependent on growth substrates
J.Bacteriol. **183** (24): 7037-7043
- KRITCHEVSKY, D. (1978).**
Influence of dietary fibre on bile acid metabolism
Lipids **13** (12): 982-985
- KROGDAHL, Å. and SELL, J.L. (1989).**
Influence of age on lipase, amylase, and protease activities in pancreatic tissues and intestinal contents of young turkeys
Poult.Sci. **68**: 1561-1568
- KUSSAIBATI, R., GUILLAUME, J., LECLERCQ, B., LAFONT, J.P. (1982).**
Effects of the intestinal microflora and added bile salts on the metabolizable energy and digestibility of saturated fats in the chicken
Arch.Geflügelk. **46**: 42-46
- LABIER, M. and LECLERCQ, B. (1994).**
Nutrition and feeding of poultry
Nottingham University Press
- LANGHOUT, D.J., SCHUTTE, J.B., GEERSE, C., KIES, A.K., DE JONG, J., VERSTEGEN, M.W.A. (1997).**
Effects on chick performance and nutrient digestibility of an endo-xylanase added to a wheat- and rye-based diet in relation to fat source
Br.Poult.Sci. **38** (5): 557-563
- LANGHOUT, D.J., SCHUTTE, J.B., VAN LEEUWEN, P., WIEBENGA, J., TAMMINGA, S. (1999).**
Effect of dietary high- and low-methylated citrus pectin on the activity of the ileal microflora and morphology of the small intestinal wall of broiler chicks
Br.Poult.Sci. **40** (4): 340-347
- LEE, S.S., HA, J.K., CHENG, K. (2000).**
Relative contributions of bacteria, protozoa, and fungi to in vitro degradation of orchard grass cell walls and their interactions
Appl.Enviro.Microbiol. **66** (9): 3807-3813

LINDSAY, O.B. and MARCH, B.E. (1967).

Intestinal absorption of bile salts in cockerel
Poult.Sci. **46**: 164-168

LISBONA, F., JIMENEZ, M., ESTELLER, A., LOPEZ, M.A. (1981).

Basal biliary secretion in conscious chicken and role of enterohepatic circulation
Comp.Biochem.Physiol. **69**: 341-344

LONGSTAFF, M. and McNAB, J.M. (1986).

Influence of site and variety on starch, hemicellulose and cellulose composition of wheats and their digestibilities by adult cockerels
Br.Poult.Sci. **27**: 435-449

MACKIE, R.I., SGHIR, A., GASKINS, H.R. (1999).

Developmental microbial ecology of the neonatal gastrointestinal tract
Am.J.Nutr. **69** (5): 1035S-1045S

MADIGAN, M.T., MARTINKO, J.L., PARKER, J. (1997).

Microbial Growth
In: Biology of microorganisms
Madigan, M.T., Martinko, J.L., Parker, J. (Ed.)
Prentice-Hall International, London
ISBN 0-13-571225-4

MANZ, W., AMANN, R., LUDWIG, W., VANCANNEYT, M., SCHLEIFER, K.H. (1996).

Application of a suite of 16S rRNA specific oligonucleotide probes designed to investigate bacteria of the phylum cytophaga-flavobacter-bacteroides in the natural environment
Microbiol. **142**: 1097-1106

MARQUARDT, R.R., WARD, A.T., MISIR, R. (1979).

The retention of nutrients by chicks fed rye diets supplemented with amino acids and penicillin
Poult.Sci. **58**: 631-640

MARTINEZ-DIEZ, M.C., SERRANO, M.A., MONTE, M.J., MARIN, J.J. (2000).

Comparison of the effects of bile acids on cell viability and DNA synthesis by rat hepatocytes in primary culture
Biochem.Biophys.Acta **1500** (2): 153-160

MEAD, G.C. and ADAMS, B.W. (1975).

Some observations on the caecal microflora of the chicks during the first weeks of life
Br.Poult.Sci. **16**: 169-176

MIWA, N., NISHINA, T., KUBO, S., HONDA, H. (1997).

Most probable numbers of enterotoxigenic *Clostridium perfringens* in intestinal contents of domestic livestock detected by nested PCR
J.Vet.Med.Sci. **59** (7): 557-560

MOHAND-OUSSAID, O., PAYOT, S., GUEDON, E., GELHAYE, E., YOUYOU, A., PETITDEMANGE, H. (1999).

The extracellular xylan degradative system in *Clostridium cellulolyticum* cultivated on xylan: evidence for cell-free cellulosome production
J.Bacteriol. **181** (13): 4035-4040

MORAN, E.T. (1985).

Digestion and absorption of carbohydrates in fowl and events through perinatal development
J.Nutr. **115**: 665-674

MORISHITA, Y., MITSUOKA, T., KANEUCHI, C., YAMAMOTO, S., OGATA, M. (1971).

Specific establishment of lactobacilli in the digestive tract of germ-free chickens
Jpn.J.Microbiol. **15** (6): 531-538

MOSER, S.A. and SAVAGE, D.C. (2001).

Bile salt hydrolase activity and resistance to toxicity of conjugated bile salts are unrelated properties in Lactobacilli

Appl.Environ.Microbiol. **67** (8): 3476-3480

NAGANO, I., KUNISHIMA, M., ITOH, Y., WU, Z., TAKAHASHI, Y. (1998).

Detection of verotoxin-producing *Escherichia coli* O157:H7 by multiplex polymerase chain reaction
Microbiol.Immunol. **42** (5): 371-376

NITSAN, Z., BEN-AHARRAM, G., ZOREF, Z., NIR, I. (1991).

Growth and development of the digestive organs and some enzymes in broiler and egg type chicks after hatching

Br.Poult.Sci. **32**: 515-523

NOY, Y. and SKLAN, D. (1995).

Digestion and absorption in the young chick

Poult.Sci. **74**: 366-373

OCHI, Y., MITSOUKA, T., SEGA, T. (1964).

The intestinal flora of the chicken. 3. Development of the intestinal flora from chick to hen

Zentralbl.Bakteriol. **193**: 80-95

OGRAM, A., SUN, W., BROCKMAN, F.J., FREDERICKSON, J.K. (1995).

Isolation and characterization of RNA from low-biomass deep-surface sediments

Appl.Environ.Microbiol. **61** (2): 763-768

OLSEN, G.J., LANE, D.J., GIOVANNI, S.J., PACE, N.R., STAHL, D.A. (1986).

Microbial ecology and evolution: A ribosomal RNA approach

Ann.Rev.Microbiol. **40**: 337-365

OLSEN, G.J. and WOESE, C.R. (1993).

Ribosomal RNA: a key to phylogeny

FASEB J. **7**: 113-123

ORVOS, D.R. (1992).

Assessing environmental risk from genetically engineered microorganisms and products containing recombinant DNA

Adv.Mod.Environ.Toxicol. **20**: 215-235

O'SULLIVAN, D.J. (1999).

Methods of analysis of the intestinal microflora

In: Probiotics. A critical review

Tannock, G.W. (Ed.)

Horizon Scientific Press

OYOFO, B.A., ABD EL SALAM, S.M., CHURILLA, A.M., WASFY, M.O. (1997).

Rapid and sensitive detection of *Campylobacter spp.* from chicken using the polymerase chain reaction

Zentralbl.Bakteriol. **285** (4): 480-485

PATEL, R., PIPER, K.E., ROUSE, M.S., STECKELBERG, J.M., UHL, J.R., KOHNER, P., HOPKINS, M.K., COCKERILL III, F.R., KLINE, B.C. (1998).

Determination of 16S rRNA sequences of *Enterococci* and application to species identification of nonmotile *Enterococcus gallinarum* isolates

J.Clin.Microbiol. **36** (11): 3399-3407

PETTERSON, D. and ÅMAN, P. (1989).

Enzyme supplementation of a poultry diet containing rye and wheat

Br.J.Nutr. **62**: 139-149

PETTERSON, D., GRAHAM, H., ÅMAN, P. (1991).

The nutritive value for broiler chickens of pelleting and enzyme supplementation of a diet containing barley, wheat and rye.

Anim.FeedSci.Technol. **33**: 1-14

PILLAI, S.D., RICKE, S.C., NISBET, D.J., CORRIER, D.E., DELOACH, J.R. (1994).

A rapid method for screening for *Salmonella typhimurium* in a chicken cecal microbial consortium using gene amplification

AvianDis. **38** (3): 598-604

QUIST, R.G., TON-NU, H.T., LILLIENAU, J., HOFMANN, A.F., BARRET, K.E. (1991).

Activation of mast cells by bile acids

Gastroenterology **101** (2): 446-456

RICKE, S.C. and PILLAI, S.D. (1999).

Conventional and molecular methods for understanding probiotic bacteria functionality in gastrointestinal tracts

Crit.Rev.Microbiol. **25**: 19-38

ROLLS, B.A., TURVEY, A., COATES, M.E. (1978).

The influence of the gut microflora and of dietary fibre on epithelial migration in the chicks intestine

Br.J.Nutr. **39**: 91-98

RUBIO, L.A., BRENES, A., SETIEN, I., DE LA ASUNCION, G., DURAN, N., CUTULI, M.T. (1998).

Lactobacilli counts in crop, ileum and caecum of growing broiler chickens fed on practical diets containing whole or dehulled sweet lupins (*Lupinus angustifolius*) seed meal

Br.Poult.Sci. **39** (3): 354-359

SAKATA, T. (1987).

Stimulatory effect of short-chain fatty acids on epithelial cell proliferation in the rat intestine: a possible explanation for trophic effects of fermentable fibre, gut microbes and luminal trophic factors

Br.J.Nutr. **58**: 95-103

SALANITRO, J.P., BLAKE, I.G., MUIRHEAD, P.A., MAGLIO, M., GOODMAN, J.R. (1978).

Bacteria isolated from the duodenum, ileum and caecum of young chicks

Appl.Environ.Microbiol. **35**: 782-790

- SALMINEN, S., BOULEY, C., BOUTRON-RUAULT, M.C., CUMMINGS, J.H., FRANCK, A., GIBSON, G.R., ISOLAURI, E., MOREAU, M.C., ROBERFROID, M., ROWLAND, I. (1998).**
Functional food science and gastrointestinal physiology and function
Br.J.Nutr. **80** Suppl. 1: S147-S171
- SAVAGE, D.C. (1977).**
Microbial ecology of the gastrointestinal tract
Ann.Rev.Microbiol. **31**: 107-133
- SAVORY, C.J. (1992).**
Gastrointestinal morphology and absorption of monosaccharides in fowls conditioned to different types and levels of dietary fibre
Br.J.Nutr. **67**: 77-89
- SCHAECHTER, M.O., MAALOE, O., KJELGAARD, N.O. (1957).**
Dependency on medium and temperature of cell size and chemical composition during balanced growth of *Salmonella typhimurium*
J.Gen.Microbiol. **19**: 592-606
- SCHUTTE, J.B., DE JONG, J., VAN WEERDEN, E.J., VAN BAAK, M.J. (1992).**
Nutritional value of D-xylose and L-arabinose for broiler chicks
Br.Poult.Sci. **33**: 89-100
- SHACKELFORD, A.D. (1988).**
Modifications of processing methods to control salmonella in poultry
Poult.Sci. **67** (6): 933-935
- SILVERSIDE, F.G. and BEDFORD, M.R. (1999).**
Soluble non-starch polysaccharides, enzymes, and gut viscosity – is there a connection
WorldPoult. **15**: 17-18
- SIMON, O. (1998).**
Auf dem Weg zu neuen Erkenntnissen über die Wirkungsweise NSP-hydrolysierender Enzyme
Lohmann Information **1/98**: 9-14
- SIMON, O. and JEROCH, H. (1999).**
Feeding raw materials – a balancing act
FeedTech. **3** (1): 41-43
- SIMON, O., HÜBENER, K., HIRSCH, K., BECKMANN, L., VAHJEN, W. (2002).**
Einfluß von Xylanasen auf die Darmflora
Lohmann Information Januar-März 2002 1/2002
- SKLAN, D., SCHACHAF, B., BARON, J., HURWITZ, S. (1978).**
Retrograde movement of digesta in the duodenum of the chick: extent frequency and nutritional implantations
J.Nutr. **108**: 1485-1490
- SMITH, H.W. (1965).**
The development of the flora of the alimentary tract in young animals
J.Pathol.Bacteriol. **90**: 495-513

SMITS, C.H., VELDMAN, A., VERSTEGEN, M.W.A., BEYNEN, A.C. (1997).

Dietary carboxymethylcellulose with high instead of low viscosity reduces macronutrient digestion in broiler chickens

J.Nutr. **127**: 483-487

SMITS, C.H., VELDMAN, A., VERKADE, H.J., BEYNEN, A.C. (1998).

The inhibitory effect of carboxymethylcellulose with high viscosity on lipid absorption in broiler chickens coincides with reduced bile salt concentration and raised microbial numbers in small intestine

Poult.Sci. **77** (10): 1534-1539

SMULIKOWSKA, S. (1998).

Relationship between the stage of digestive tract development in chicks and the effect of viscosity reducing enzymes on fat digestion

J.Anim.FeedSci. **7**: 125-134

SPRING, P. (1996).

Effects of mannanoligosaccharides on different cecal parameters and on cecal concentrations of enteric pathogens in poultry

Dissertation, ETH, Zürich

STAHL, D.A., FLESHER, B., MANSFIELD, H.R., MONTGOMERY, L. (1988).

Use of phylogenetically based hybridization probes for studies of ruminal microbial ecology

Appl.Environ.Microbiol. **54** (5): 1079-1084

STALEY, J.T. and KONOPKA, A. (1985).

Measurement of in situ activities of nonphotosynthetic microorganisms in aquatic and terrestrial habitats

Rev.Microbiol. **49**: 321-346

TEATHER, R.M. and ERFLE, J.D. (1990).

DNA sequence of a *Fibrobacter succinogenes* mixed-linkage beta-glucanase (1,3-1,4-beta-D-glucan-4-glucanohydrolase) gene

J.Bacteriol. **127**: 3837-3841

TEBBE, C.C. and VAHJEN, W. (1993).

Interference of humic acids and DNA extracted directly from soil in detection and transformation of recombinant DNA from bacteria and yeast

Appl.Environ.Microbiol. **59**: 2657-2665

TIMMS, L. (1968).

Observation on the bacterial flora of the alimentary tract in three age groups of normal chickens

Br.Vet.J. **124**: 470-477

TSEN, H.Y., LIN, C.K., CHI, W.R. (1998).

Development and use of 16S rRNA gene targeted PCR primers for the identification of *Escherichia coli* cells in water

J.Appl.Microbiol. **85** (3): 554-560

UMESAKI, Y. and SETOYAMA, H. (2000).

Structure of the intestinal flora responsible for development of the gut immune system in a rodent model

Microbes.Infect. **21** (11): 1343-1351

UNTAWALE, G.G. and MCGINNIS, J. (1979).

Effect of rye and levels of raw and autoclaved beans on adhesion of microflora to the intestinal flora

Poult.Sci. **58**: 928-933

VAHJEN, W. and SIMON, O. (1997).

Mögliche Wirkungsebenen NSP-hydrolysierender Enzyme auf intestinale Mikroorganismenpopulationen bei Monogastriden

In: Vitamine und Zusatzstoffe in der Ernährung von Mensch und Tier

6. Symposium 24./25.1997 in Jena/Thüringen

Schubert, R., Flachowsky, G., Bitsch, R., Jahreis, G. (Hrsg.)

Buch- und Kunstdruckerei Keßler GmbH, Weimar

ISBN 3-00-002381-X

VAHJEN, W., GLÄSER, K., SCHÄFER, K., SIMON, O. (1998).

Influence of xylanase-supplemented feed on the development of selected bacterial groups in the intestinal tract of broiler chicks

J.Agricult.Sci. **130**: 489-500

VAHJEN, W. and SIMON, O. (1999).

Biochemical characteristics of non starch polysaccharide hydrolyzing enzyme preparations designed as feed additives for poultry and piglet nutrition

Arch.Anim.Nutr. **52**: 1-14

VAN DER KLIES, J.D. and VAN VOORST, A. (1993).

The effect of carboxy methyl cellulose (a soluble polysaccharide) on the rate of marker excretion from the gastrointestinal tract of broilers

Poult.Sci. **72**: 503-512

VELDMANN, A. and VAHL, H.A. (1994).

Xylanase in broiler diets with differences in characteristics and content of wheat

Br.Poult.Sci. **35**: 537-550

VISEK, W.J. (1978).

The mode of growth promotion by antibiotics

J.Anim.Sci. **46**: 1447-1469

VIVEROS, A., BRENES, A., PIZARRO, M., CASTAÑO, M. (1994).

Effect of enzyme supplementation of a diet based on barley, and autoclave treatment, on apparent digestibility, growth performance and gut morphology of broilers

Anim.FeedSci.Tech. **48** (3-4): 237-251

WAGNER, D.D. and THOMAS, O.P. (1978).

Influence of diets containing rye or pectin on the intestinal flora of chicks

Poult.Sci. **57**: 971-975

WAGNER, M.P., AMANN, R., LEMMER, H., SCHLEIFER, K.H. (1993).

Probing activated sludge with oligonucleotides specific for proteobacteria: inadequacy of culture dependent methods for describing microbial community structures

Appl.Environ.Microbiol. **59**: 1520-1525

WARD, A.T. and MARQUARDT, R.R. (1983).

The effect of saturation, chain length of pure triglycerides, and age of bird on the utilization of rye diets

Poult.Sci. **62**: 1054-1062

WARD, D.M., WELLER, R., BATESON, M.M. (1990).

16S rRNA sequences reveal numerous uncultured microorganisms in a natural community

Nature (London) **345**: 63-65

ZOETENDAL, E.G., VON WRIGHT, A., VLIPPONEN-SALMELA, T., BEN-ARMOR, K., AKKERMANS, A.D., DEVOS, W.M. (2002).

Mucosa-associated bacteria in the human intestinal tract are uniformly distributed along the colon and differ from the community recovered from faeces

Appl.Environ.Microbiol. **68** (7): 3401-3407