

8. LITERATURVERZEICHNIS

- Aberer, E., C. Brunner, G. Suchanek, H. Klade, A. Barbour, G. Stanek and H. Lassmann** (1989): Molecular mimicry and Lyme borreliosis: a shared antigenic determinant between *B. burgdorferi* and human tissue. Ann. Neurol. **26**, 732-737.
- Ackermann, R., J. Kabatzki, H.P. Boisten, A.C. Steere, R.C. Grodzicky, S. Hartung und U. Runne** (1984): Spirochaeten-Ätiologie der Erythema-chronicum-migrans-Krankheit. Dtsch. Med. Wochenschr. **109**, 92-97.
- Adam, T., G.S. Gassmann, C. Rasiah and U.B. Göbel** (1991): Phenotypic and genotypic analysis of *Borrelia burgdorferi* isolates from various sources. Infect. Immun. **59**, 2579-2585.
- Afzelius, A.** (1909): Verhandlung der Deutschen Gesellschaft zu Stockholm. Seminar vom 28. Oktober 1909. Arch. Derm. Syph. (Berlin) **101**, 404.
- Alekseev, A.N., H.V. Dubinina, L.P. Antykova, T.I. Dzhivanyan, S.G.T. Rijpkema, N. Verbeek- de Kruif and M. Cinco** (1998): Tick-borne borreliosis pathogen identification in *Ixodes* ticks (Acarina, Ixodidae) collected in St. Petersburg and Kalingrad baltic regions of Russia. J. Med. Entomol. **35**, 136-142.
- Anderson, J.F.** (1991): Epizootiology of Lyme Borreliosis. Scand. J. Infect. Dis. – Suppl. **77**, 23-34.
- Anonymous** (1988): Lyme disease in British Columbia. Can. Vet. J. **29**, 393-394.
- Appel, M.J.G.** (1990): Lyme disease in dogs and cats. Compend. Contin. Educ. Prac. Vet. **12**, 617-626.
- Appel, M.J.G., S. Allan, R.H. Jacobson, T.L. Lauderdale, Y.F. Chang, S.J. Shin, J.W. Thomford, R.J. Todhunter and B.A. Summers** (1993): Experimental Lyme disease in dogs produces arthritis and persistent infection. J. Infect. Dis. **167**, 651-664.
- Austin, F.E.** (1993): Maintenance of infective *Borrelia burgdorferi* Sh-2-82 in 4% oxygen - 5% carbon dioxide in vitro. Can. J. Microbiol. **39**, 1103-1110.

Bäverstedt, B. (1943): Über Lymphadenosis benigna cutis. Acta Derm. Venereol. Suppl. (Stockh.) **11**, (Suppl. zu **24**), 1-202.

Balmelli, T. and J.C. Piffaretti (1995): Association between different clinical manifestations of Lyme disease and different species of *B. burgdorferi* sensu lato. Res. Microbiol. **146** (4), 329-340.

Bannwarth, A. (1941): Chronische Lymphozytäre Meningitis, entzündliche Polyneuritis und Rheumatismus. Arch. Psychiatr. Nervenkr. **113**, 284-376.

Baranton, G., D. Postic, I. Saint-Girons, P. Boerlin, J.C. Piffaretti, M. Assous and P.A.D. Grimont (1992): Delineation of *Borrelia burgdorferi* sensu stricto, *Borrelia garinii* sp. Nov., and Group VS461 associated with Lyme borreliosis. Int. J. Syst. Bacteriol. **42**, 378-383.

Barbour, A.G. (1984): Isolation and cultivation of Lyme disease spirochetes. Yale J. Biol. Med. **57**, 521-525.

Barbour, A.G. and S.F. Hayes (1986): Biology of *Borrelia* species. Microbiol. Rev. **50**, 381-400.

Barbour, A.G., W. Burgdorfer, S.F. Hayes, O. Peter and A. Aeschlimann (1983): Isolation of a cultivable spirochete from *Ixodes ricinus* ticks in Switzerland. Curr. Microbiol. **8**, 123-126.

Bauerfeind, R., U. Kreis, R. Weiß, L.H. Wieler and G. Baljer (1998): Detection of *Borrelia burgdorferi* in urine specimens from dogs by a nested polymerase chain reaction. Zentr.bl. Bakt. **287**, 347-361.

Baumeister, K. (1999): Die Lyme-Borreliose des Hundes – Epidemiologie, Klinik, Probleme der Diagnostik. Tierärztl. Umschau **54**, 239-242.

Bayer M.E., L. Zhang and M.H. Bayer (1996): *Borrelia burgdorferi* DNA in the urine of treated patients with chronic Lyme disease symptoms. A PCR study of 97 cases. Infection **24**, 347-353.

- Becker, W.** (1996): Zoonose-Fibel. Zwischen Tier und Mensch übertragbare Krankheiten. Berlin: H. Hoffmann Verlag.
- Beder, G.** (1988): Rasterelektronenmikroskopische Untersuchung von Zecken der Gattung *Ixodes* an einheimischen Säugetieren. Hannover, Univ., Dipl. Biol..
- Belfaiza, J., D. Postic, E. Bellenger, G. Baranton and I. Saint Girons** (1993): Genomic fingerprinting of *Borrelia burgdorferi* sensu lato by pulsed-field gel electrophoresis. *J. Clin. Microbiol.* **31**, 2873-2877.
- Benach, J.L., B. Fernandez Villar, A. Szczepanski and J.C. Garcia-Monco** (1991): Lyme borreliosis: non-specific interactions of the organism with the host. *Scand. J. Infect. Dis. – Suppl.* **77**, 130-135.
- Berger, B.W., R.C. Johnson, C. Kodner and L. Coleman** (1992): Cultivation of *Borrelia burgdorferi* from erythema migrans lesions and perilesional skin. *J. Clin. Microbiol.* **30**, 359-361.
- Bergmann, J., A. Liebisch und K. Pohlmeyer** (1992): Zum Vorkommen der einheimischen Borreliose bei Zecken, Wild- und Haustieren in einem niedersächsischen Moor. *Vet.* **7**, 12-15.
- Bergström, S.** (1999): Variation in antigen expression. VIII Int. Conf. on Lyme Borreliosis and other Emerging Tick-Borne Diseases, Munich, Germany, June 20-24, 1999, Abstr. **39**.
- Bergström, S., L. Noppa, Å. Gylfe and Y. Östberg** (2002): Molecular and Cellular Biology of *Borrelia burgdorferi* sensu lato. CAB Int. Lyme Borreliosis: Biology, Epidemiology and Control (eds J. Gray, O. Kahl, R.S. Lane and G. Stanek), **47-90**.
- Bernard, W.V., D. Cohen, E. Bosler and D. Zamos** (1990): Serologic survey for *Borrelia burgdorferi* antibody in horses referred to a mid-Atlantic veterinary teaching hospital. *J. Am. Vet. Med. Assoc.* **196**, 1255-1258.
- Binder, E., R. Döpfner und O. Hornstein** (1955): Experimentelle Übertragung des Erythema chronicum migrans von Mensch zu Mensch. *Hautarzt* **6**, 494-496.

Boerner, J. (1994): Empfindlichkeitsprüfung von *Borrelia burgdorferi* gegen Antibiotika und Chemotherapeutika in vitro. Vet.med. Diss., Giessen.

Bosler, E.M., D.P. Cohen, T.L. Schulze, C. Olsen, W. Bernard and B. Lissman (1988): Host Responses to *Borrelia burgdorferi* in Dogs and Horses. Ann. N.Y. Acad. Sci. **539**, 221-228.

Brand, A. (1990): Vergleichende seroepidemiologische Untersuchung von Rindern auf Zeckenborreliose in der Südheide und im Weserbergland. Vet.med. Diss., TiHo, Hannover.

Breitschwerdt, E.B., W.L. Nicholson, A.R. Kiehl, D.J. Steers and J.F. Levine (1994): Natural infections with *Borrelia* spirochetes in two dogs from Florida. J. Clin. Microbiol. **32**, 352-357.

Brettschneider, S., H. Bruckbauer, N. Klugbauer and H. Hofmann (1998): Diagnostic value of PCR for detection of *Borrelia burgdorferi* in skin biopsy and urine samples from patients with skin borreliosis. J. Clin. Microbiol. **36**, 2658-2665.

Browning, A., S.D. Carter, A. Barnes, C. May and D. Bennett (1993): Lameness associated with *Borrelia burgdorferi* infection in the horse. Vet. Rec. **132**, 610-611.

Bruckbauer, H.R., V. Preac-Mursic, R. Fuchs and B. Wilske (1992): Cross-reactive proteins of *Borrelia burgdorferi*. Eur. J. Clin. Microbiol. Infect. Dis. **11**, 224-232.

Buchwald, A. (1883): Ein Fall von diffuser idiopathischer Hautatrophie. Arch. Derm. Syph. (Berlin) **15**, 553-556.

Bühl, A. und P. Zöfel (2002): SPSS 11 Einführung in die moderne Datenanalyse unter Windows 8. Aufl. Pearson Education Deutschl. GmbH.

Burgdorfer, W., A.G. Barbour, S.F. Hayes, J.L. Benach, E. Grunwald and J.P. Davis (1982): Lyme disease – a tick borne spirochaetosis? Science **216**, 1317-1319.

Burgess, E.C. (1986): Experimental inoculation of dogs with *Borrelia burgdorferi*. Zbl. Bakt. Hyg. A **263**, 49-54.

- Burgess, E.C.** (1988): *Borrelia burgdorferi* infection in Wisconsin horses and cows. Ann. N.Y. Acad. Sci. **539**, 235-243.
- Burgess, E.C.** (1992): Experimentally induced infection of cats with *Borrelia burgdorferi*. Am. J. Vet. Res. **53** (9), 1507-1511.
- Burgess, E.C. and M. Mattison** (1987): Encephalitis associated with *Borrelia burgdorferi* infection in a horse. JAVMA **191**, 1457-1458.
- Burgess, E.C. and A. Gendron-Fitzpatrick** (1990): Experimental infection of equines with *Borrelia burgdorferi*. IV International Conf. on Lyme Borreliosis, Stockholm.
- Burgess, E.C., D. Gillette and J.P. Pickett** (1986): Arthritis and panuveitis as manifestations of *Borrelia burgdorferi* infection in a Wisconsin pony. JAVMA **189**, 1340-1342.
- Burgess, E.C., A. Gendron-Fitzpatrick and W.O. Wright** (1987): Arthritis and systemic disease caused by *Borrelia burgdorferi* infection in a cow. J. Am. Vet. Med. Assoc. **191** (11), 1468-1470.
- Burgess, E.C., A. Gendron-Fitzpatrick and M. Mattison** (1990): Foal mortality associated with natural infection of pregnant mares with *Borrelia burgdorferi*. Eq. Inf. Diseases V. (Proc. 5th Intern. Disease Conf.) Powell, D.G. (Ed). Univ. Press of Kentucky. 217-220.
- Burmester, G.R.** (1999): Internistische Manifestationen bei der Lyme Borreliose. In: A. Krause, G.R. Burmester (Hrsg.): Lyme Borreliose. Stuttgart [u.a.]: Georg Thieme Verlag, 25-32.
- Bushmich, S.L. and J.E. Post** (1989): Lyme disease update for veterinarians. Cooperative Extension Publication **89-31**, Spring 1989; The University of Connecticut, Storrs, Ct.
- Canica, M.M., F. Nato, L. du Merle, J.C. Mazie, G. Baraton and D. Postic** (1993): Monoclonal antibodies for identification of *Borrelia afzelii* sp. nov. associated with late cutaneous manifestations of Lyme borreliosis. Scand. J. Infect. Dis. **25**, 441-448.
- Carter, S.D., C. May, A. Barnes and D. Bennett** (1994): *Borrelia burgdorferi* infection in UK horses. Equine Vet. J. **26**, 187-190.

Cerri, D., F. Farina, E. Andreani, R. Nuvoloni and A. Pedrini (1994): Experimental infection of dogs with *Borrelia burgdorferi*. Res. Vet. Sci. **57**, 256-258.

Champion, C.I., D.R. Blanco, J.T. Skare, D.A. Haake, M. Giladi, D. Foley, J.N. Miller and M.A. Lovett (1994): A 9.0-kilobase-pair circular plasmid of *Borrelia burgdorferi* encodes an exported protein: evidence for expression only during infection. Infect. Immun. **62**, 2653-2661.

Chang, Y.-F. (1996): Dissemination of *Borrelia burgdorferi* after experimental infection in dogs. Journal of Spirochetal and Tick-Borne Disease **3** (1), 80-86.

Chang, Y., V. Novosol, S.P. McDonough, C.F. Chang, R.H. Jacobson, T. Divers, F.W. Quimby, K.S. Shin and D.H. Lein (1999): Vaccination against Lyme disease with recombinant *Borrelia burgdorferi* outer-surface protein A (rOspA) in horses. Vaccine **18** (5-6), 540-548.

Chang, Y., S.P. McDonough, C.F. Chang, K.S. Shin, W. Yen and T. Divers (2000): Human granulocytic ehrlichiosis agent infection in a pony vaccinated with a *Borrelia burgdorferi* recombinant OspA vaccine and challenged by exposure to naturally infected ticks. Clin. Diag. Lab. Immunol. **7** (1), 68-71.

Cimmino, M., M. Granström, J.S. Gray, E.C. Guy, S. O'Connell and G. Stanek (1998): European Lyme borreliosis clinical spectrum. Zentr.bl. Bakteriol. **287**, 248-252.

Cohen, N.D. and D. Cohen (1990): Borreliosis in horses: a comparative review. Equine Comp. **12**, 1449-1458.

Cohen, D., E.M. Bosler, W. Bernard, D. Meirs II, R. Eisner and T.L. Schulze (1988): Epidemiologic Studies of Lyme Disease in Horses and Their Public Health Significance. Ann. N.Y. Acad. Sci. **539**, 244-251.

Cohen, N.D. C.N. Carter, M.A. Thomas Jr., A.B. Angulo and A.K. Engster (1990): Clinical and epizootiologic characteristics of dogs seropositive for *Borrelia burgdorferi* in Texas: 110 cases (1988). J. Am. Vet. Med. Assoc. **197**, 893-898.

- Cohen, N.D., F.C. Heck, B. Heim, D.M. Flad, E.M. Bosler and D. Cohen** (1992): Seroprevalence of antibodies to *Borrelia burgdorferi* in a population of horses in central Texas. J. Am. Vet. Med. Assoc. **201**, 1030-1034.
- Coons, A.H., H.J. Creech and R.N. Jones** (1941): Immunological properties of an antibody containing a fluorescent group. Proc. Soc. Exp. Biol. Med. **47**, 200-202.
- Cutler, S.J. and M.J. Woodward** (2001): Lyme borreliosis in the UK – ecology and risks to domestic animals and Man. Rev. Med. Microbiol. **12** (4), 199-209.
- Dambach, D.M., C.A. Smith, R.M. Lewis and T.J. van Winkle** (1997): Morphologic, immunhistochemical and ultrastructural characterization of a distinctive renal lesion in dogs putatively associated with *Borrelia burgdorferi* infection: 49 cases (1987-1992). Vet. Pathol. **34**, 85-96.
- Dejmekova, H., D. Hulinska, D. Tegzova, K. Pavelka, J. Gatterova and P. Vavrik** (2002): Seronegative Lyme arthritis caused by *Borrelia garinii*. Clin. Rheumatol. **21** (4), 330-334.
- Del Rio, C., S.R. Granter and P.H. Duray** (1997): Lyme Borreliosis. In: C.R. Horsburgh, Jr., A.M. Nelson (ed.): Pathology of Emerging Infections. Washington, DC.: ASM Press. 269-283.
- Demaerschalck, I., A. Ben Messaoud, M. De Kesel, B. Hoyois, Y. Lobet, P. Hoet, G. Bigaignon, A. Bollen and E. Godfroid** (1995): Simultaneous presence of different *Borrelia burgdorferi* genospecies in biological fluid of Lyme disease Patients. J. Clin. Microbiol. **33**, 602-608.
- Dennis, D.T.** (1997): Epidemiology of Lyme Borreliosis. Proc. 2nd Int. Symp. Lyme Disease, 27.-28. Okt. 1997 in Japan, 121-164.
- Deutsches Grünes Kreuz und Bundesinstitut für gesundheitlichen Verbraucherschutz und Veterinärmedizin** (Hrsg.) (1997): Krankheitsüberträger Zecke. Lyme Borreliose und FSME. Marburg: Verlag im KILIAN.

- Divers, T.J., Y.F. Chang and P.L. McDonough** (2003): Equine Lyme Disease: A Review of Experimental Disease Production, Treatment Efficacy and Vaccine Protection. In: 49th Ann. Am. Assoc. Equine Pract. Conv.
- Doby, J.M., S. Chevrier et A. Couatarmanach** (1987): Spirochétose à tiques par *Borrelia burgdorferi* chez le cheval en Bretagne. Résultats d' une enquête sérologique portant sur 400 chevaux. Bulletin de la Société Française de Parasitologie **5**, 285-298.
- Dorn, W. and U. Sünder** (1997): Lyme borreliosis in Thuringia (Germany): Prevalence of the causative agent in field-collected ticks. In: J. Süss, O. Kahl (eds.): 4th International Potsdam Symposium on Tick-Borne Diseases: Tick-Borne Encephalitis and Lyme Borreliosis, 21.-22. Februar 1997. Lengerich: PABST, 175-196.
- Dzierzecka, M. and J. Kita** (2002a): The use of chosen serological diagnostic methods in Lyme disease in horses. Part I. Indirect immunofluorescence and enzyme-linked immunosorbent assay (ELISA). J. Vet. Sci. **5** (2), 71-77.
- Dzierzecka, M. and J. Kita** (2002b): The use of chosen serological diagnostic methods in Lyme disease in horses. Part II. Western Blot. J. Vet. Sci. **5** (2), 79-84.
- Egenvall, A., P. Franzén, A. Gunnarsson, E.O. Engvall, I. Vågsholm, U.-B. Wikström and K. Artursson** (2001): Cross-sectional study of the seroprevalence to *Borrelia burgdorferi* sensu lato and granulocytic *Ehrlichia* ssp. and demographic, clinical and tick-exposure factors in Swedish horses. Prev. Vet. Med. **49** (3-4), 191-208.
- Eiffert, H., A. Ohlenbusch, H.J. Christen, R. Thomssen, A. Spielmann and F.R. Matuschka** (1995): Nondifferentiation between Lyme disease spirochetes from vector ticks and human cerebrospinal fluid. J. Infect. Dis. **171**, 476-479.
- Eng, T.R., M.L. Wilson, A. Spielman and C.C. Lastavica** (1988): Greater risk of *Borrelia burgdorferi* infection in dogs than in people. J. Infect. Dis. **158**, 1410-1411.
- Fischer, A. und G. Leuterer** (1992): *Borrelia-burgdorferi*-assoziierte Lymphadenitis purulenta bei einem Hund. Kleintierpr. **37**, 13-16.
- Fraenkel, C.J., U. Garpmo and J. Berglund** (2002): Determination of novel *Borrelia* genospecies in Swedish *Ixodes ricinus* ticks. J. Clin. Microbiol. **40** (9), 3308-3312.

Frese, S. (1997): Untersuchungen zum klinischen Verlauf der Lyme-Borreliose beim Hund nach experimenteller Infektion durch den Holzbock *Ixodes ricinus*. Vet. med. Diss., FU-Berlin.

Fridriksdóttir, V., G. Overnes and S. Stuen (1992): Suspected Lyme borreliosis in sheep. Vet. Rec. **130** (15), 323-324.

Fukunaga, M., Y. Takahashi, Y. Tsuruta, O. Matsushita, D. Ralph, M. Mc Clelland and M. Nakao (1998): Genetic and phenotypic analysis of *Borrelia miyamotoi* sp. nov., isolated from the ixodid tick *Ixodes persulcatus*, the vector for Lyme disease in Japan. Int. J. Syst. Bacteriol. **45**, 804-810.

Garcia-Monco, J.C. (1995): Pathomechanismen der Neuroborreliose. Wien Med. Wochenschr. **7**, 174-177.

Garcia-Monco, J.C. and J.L. Benach (1989): The pathogenesis of Lyme disease. Rheum. Dis. Clin. North Am. **15**, 711-726.

Garin, C. et C. Bujadoux (1922): Paralysis par les tiques. J. Med. Lyon **71**, 765-767.

Gerhards, H. und B. Wollanke (1996): Antikörpertiter gegen Borrelien bei Pferden im Serum und im Auge und Vorkommen der equinen rezidivierenden Uveitis (ERU). Berl. Münch. Tierärztl. Wschr. **109**, 273-278.

Gern, L., P.F. Humair, C.M. Hu and S. Leuba-Garcia (1997): Ecology of *Borrelia burgdorferi* sensu lato in Europe. In: J. Süss, O. Kahl (eds.): 4th International Potsdam Symposium on Tick-Borne Diseases: Tick-Borne Encephalitis and Lyme Borreliosis, 21.-22. Februar 1997. Lengerich: PABST, 271-280.

Gern, L., A. Estrada-Peña, F. Frandsen, J.S. Gray, T.G.T. Jaenson, F. Jongejan, O. Kahl, E. Korenberg, R. Mehl and P.A. Nuttall (1998): European reservoir hosts of *Borrelia burgdorferi* sensu lato. Zentr.bl. Bakteriol. **287**, 196-204.

Goebel, P. (1995): Vergleich serologischer Nachweismethoden der Lyme-Borreliose bei experimentell und natürlich infizierten Hunden. Vet.med. Diss., FU Berlin, Berlin.

Golubic, D., Rijkema, S., Tkalec-Makovec and Ruzic, E. (1998): Epidemiologic, ecologic and clinical characteristics of Lyme borreliosis in northwest Croatica. *Acta med. Croat.* **52**, 7-13.

Grauer, G.F., E.C. Burgess, A.C. Cooley and J.H. Hagee (1988): Renal lesions associated with *Borrelia burgdorferi* infection in a dog. *J. Am. Vet. Med. Assoc.* **193**, 237-239.

Gray, J.S. (1991): The development and seasonal activity of the tick *Ixodes ricinus*: a vector of Lyme Borreliosis. *Rev. Med. Vet. Entomol.* **79**, 323-333.

Gray, J.S., O. Kahl, C. Janetzki-Mittmann, J. Stein and E. Guy (1994): Acquisition of *Borrelia burgdorferi* by *Ixodes ricinus* ticks fed on the European hedgehog, *Erinaceus europaeus* L. *Exp. Appl. Acarol.* **18** (8), 485-491.

Gray, J.S., A. Schönberg, D. Postic, J. Belfaiza and I. Saint-Girons (1996): First isolation and characterisation of *Borrelia garinii*, agent of Lyme borreliosis, from Irish ticks. *Irish J. of Medical Science* **165**, 24-26.

Gupta, S.K., A. Schönberg and T. Hiepe (1995): Prevalence of ticks in relation to their role as vector of *Borrelia burgdorferi* under autochthonous conditions. *Appl. Parasitol.* **36**, 97-106.

Guy, E.C., J.N. Robertson, M. Cimmino, L. Gern, Y. Moosmann, S.G.T. Rijkema, V. Sambri and G. Stanek (1998): European interlaboratory comparison of Lyme borreliosis serology. *Zent.bl. Bakteriol.* **287**, 241-247.

Hahn, C.N., I.G. Maayhew, K.E. Whitwell, D. Carey, S.D. Carter and R.A. Read (1996): A possible case of Lyme borreliosis in a horse in the UK. *Eq. Vet. J.* **28**, 84-88.

Halouzka, J., D. Postic and Z. Hubalek (1998): Isolation of the spirochaete *Borrelia afzelii* from the mosquito *Aedes vexans* in the Czech Republic. *Med. Vet. Entomol.* **12**, 103-105.

Hartmann, K. (2002): Interpretation criteria for standardized Western blots for three European species of *Borrelia burgdorferi* sensu lato. *J. Clin. Microbiol.* **35**, 1433-1444.

Hauser, U., G. Lehnert and B. Wilske (1998): Diagnostic value of proteins of three *Borrelia* species (*Borrelia burgdorferi* sensu lato) and implications for development and use of recombinant antigens for serodiagnosis of Lyme borreliosis in Europe. *Clin. Diagn. Labor Immunol.* **5**, 456-462.

Hauser, U., G. Lehnert, R. Lobentanzer and B. Wilske (1997): Interpretation criteria for standardized Western blots for three European species of *Borrelia burgdorferi* sensu lato. *J. Clin. Microbiol.* **35**, 1433-1444.

Hayes, S.F. and W. Burgdorfer (1993): Ultrastructure of *Borrelia burgdorferi*. In: Weber, K. & Burgdorfer, W. (Hrsg.), *Aspects of Lyme Borreliosis*, Springer, Berlin, Heidelberg, 29-43.

Heid, C.A., J. Stevens, K.J. Livak and P.M. Williams (1996): Real time quantitative PCR. *Genome Res.* **6**, 686-694.

Heidrich, J., A. Schönberg, S. Steuber, K. Nöckler, P. Schulze, W.-P. Voigt and E. Schein (1999): Investigation of Skin Samples from Red Foxes (*Vulpes vulpes*) in Eastern Brandenburg (Germany) for the Detection of *Borrelia burgdorferi* s.l.. *Zentr.bl. Bakteriol.* **289**, 666-672.

Heidrich, J. (2000): Untersuchungen zur Prävalenz von *Borrelia burgdorferi* sensu lato beim Rotfuchs (*Vulpes vulpes*) in Ostbrandenburg. *Vet.med. Diss.*, FU Berlin, Berlin.

Herxheimer, K. und K. Hartmann (1902): Über Acrodermatitis chronica atrophicans. *Arch. Dermatol. (Berlin)* **61**, 57-76.

Herzer, P. (1990): Lyme Borreliose: Epidemiologie, Ätiologie, Diagnostik, Klinik und Therapie. Darmstadt: Steinkopff Verlag. 2. Aufl.

Hollström, E. (1951): Succesful treatment of Erythema migrans Afzelius. *Acta Derm. Venereol. (Stockh.)* **31**, 235-243.

Horst, H. (1997): Einheimische Zeckenborreliose (Lyme-Krankheit) bei Mensch und Tier. Balingen: Demeter-Verl. im Spitta Verl., 3. überarbeitete Auflage.

- Hovind-Hougen, K.** (1984): Ultrastructure of spirochetes isolated from *Ixodes ricinus* and *Ixodes dammini*. In: Steere, A.C., Malawista, S.E., Craft, J.E., Fischer, D.K., Garcia-Blanco, M. (Hrsg.), Lyme disease, First Int. Symp., Yale J. Biol. Med., Connecticut, 93-98.
- Hovmark, A., E. Åsbrink, O. Schwan, B. Hederstedt and D. Christensson** (1986): Antibodies to *Borrelia* spirochetes in sera from Swedish cattle and sheep. Acta Vet. Scand. **27**, 479-485.
- Hubbard, M.J., A.S. Baker and K.J. Cann** (1998): Distribution of *Borrelia burgdorferi* s.l. spirochete DNA in British ticks (Argasidae and Ixodidae) since the 19th century, assessed by PCR. Med. Vet. Entomol. **12**, 89-97.
- Hunfeld, K.P., G. Stanek, E. Straube, H.J. Hagedorn, C. Schorner, F. Muhschlegel and V. Brade** (2002): Quality of Lyme disease serology. Lessons from the German Proficiency Testing Program 1999-2001. A preliminary report. Wien. Klin. Wochenschr. **114** (13-14), 591-600.
- Hyde, F. and R.C. Johnson** (1984): Relationship of the Lyme disease spirochetes to *Leptospira*, *Treponema* and *Borrelia*. Based on DNA Guanine-plus-cytosine Content and DNA reassociation studies. In: A. Schönberg (ed.), 4th Meeting of European Leptospira Workers Berlin (West), 12-14 October 1983, Selected Papers and Abstracts. Zbl.Bakt.Hyg. A **257**, 479.
- Jenal, K.D.** (2002): Studie zur Pathogenese und klinischen Symptomatik der Borreliose beim Hund nach experimenteller Infektion. Vet.med. Diss., Universität Zürich, Zürich.
- Johnson, R.C., G.P. Schmid, F.W. Hyde, A.G. Teigerwald and D.J. Brenner** (1984): *Borrelia burgdorferi*, sp. nov.: Etiologic Agent of Lyme disease. Int. J. Syst. Bacteriol. **34**, 496-497.
- Kahl, O.** (1991): Lyme Borreliosis – an ecological perspective of a tick-borne human disease. Anz. Schädlingskd. Pflanzenschutz Umweltschutz **64**, 45-55.
- Kahl, O.** (1994): Die Zecke als Vektor. In J. Süss (Hrsg.): Durch Zecken übertragbare Erkrankungen: FSME und Lyme-Borreliose. 2. Potsdamer Symposium, 13. März 1993. Schriesheim: Weller Verlag, 1-19.

Kahl, O. (1996): Fatal Attraction or How We Get Tick Bites? Infect. **24** (5), 394-395.

Kahl, O., L. Gern, J.S. Gray, E.C. Guy, F. Jongejan, F. Kirstein, K. Kurtenbach, S.G.T. Rijpkema and G. Stanek (1998): Detection of *Borrelia burgdorferi* sensu lato in ticks: immunofluorescence assay versus polymerase chain reaction. Zentr.bl. Bakteriol. **287**, 205-210.

Käsbohrer, A. und A. Schönberg (1990): Serologische Untersuchungen zum Vorkommen von *Borrelia burgdorferi* bei Haustieren in Berlin (West). Berl. Münch. Tierärztl. Wschr. **103**, 374-378.

Käsbohrer, A. and A. Schönberg (1990): Crossreactions between *Borrelia burgdorferi* and other spirochetes. IV Int. Conf. on Lyme Borreliosis, Stockholm 1990 (Abstr.).

Kawabata, H., H. Tashibu, K. Yamada, T. Masuzawa and Y. Yanagihara (1994): Polymerase chain reaction analysis of *Borrelia species* isolated in Japan. Microbiol. Immunol. **38**, 591-598.

Kelly, R.T. (1984): The Spirochetes: Genus IV. *Borrelia*. In: Krieg, N.R. and J.G. Holt (Hrsg.), Burgey's Manual of Systematic Bacteriology, Vol 1. William and Wilkins, Baltimore, London 57-62.

Kimsey, R.B. and A. Spielman (1990): Motility of Lyme disease spirochetes in fluids as viscous as the extracellular matrix. J. Infect. Dis. **162**, 1205-1208.

Kirstein, F., S. Rijpkema, M. Molkenboer and J.S. Gray (1997): The distribution and prevalence of *Borrelia burgdorferi* genomospecies in *Ixodes ricinus* ticks in Ireland. Eur. J. Epidemiol. **13**, 67-72.

Klapper, B.-M. (1999): Untersuchungen zur Epidemiologie der Borreliose im Freistaat Sachsen – Verbreitung von *Borrelia burgdorferi* in Zecken und Antikörperbestimmung in Humanseren-. Med. Diss., Medizinische Fakultät, Universität Leipzig, Leipzig.

Klempner, M.S., R. Noring and R.A. Rogers (1993): Invasion of human skin fibroblast by the Lyme disease spirochete *Borrelia burgdorferi*. J. Infect. Dis. **167**, 1074-1081.

Klich, M., M.W. Lankester, K. Wu and K.W. Wu (1996): Spring migratory birds (Aves) extend the northern occurrence of blacklegged tick (Acari: Ixodidae). *J. Med. Entomol.* **33**, 581-585.

Kopp, A. (1990): Natürliche und experimentelle Infektion bei Haustieren mit durch Zecken übertragene Borrelien. Vet.med. Diss., TiHo, Hannover.

Kornblatt, A.N., P.H. Urband and A.C. Steere (1985): Arthritis caused by *Borrelia burgdorferi* in dogs. *J. Am. Vet. Med. Assoc.* **186** (9), 960-964.

Kraiczy, P., G. Acker und V. Brade (1998): Erregereigenschaften. In: Oschmann, P., Kraiczy P. (Hrsg.), Lyme-Borreliose und Frühsommer-Meningoenzephalitis, UNIMED, Bremen, 16-25.

Kramer, M.D., R. Wallich and M.M. Simon (1996): The outer surface protein A (OspA) of *Borrelia burgdorferi*: A vaccine candidate and bioactive mediator. *Infection* **24**, 190-193.

Kurtenbach, K., H.S. Sewell, S.S. Schäfer and S. de Michelis (1999): Biology, ecology and systematics of the genus *Borrelia*. *Zentr.bl. Bakteriol.* **289**, 639-642.

Kurtti, T.J., U.G. Munderloh, R.C. Johnson and G.G. Ahlstrand (1987): Colony formation and morphology in *Borrelia burgdorferi*. *J. Clin. Microbiol.* **25**, 2054-2058.

Layfield, D. and P. Guilfoile (2002): The prevalence of *Borrelia burgdorferi* (Spirochetales: spirochaetaceae) and the agent of human granulocytic ehrlichiosis (Rickettsiaceae: Ehrlichieae) in *Ixodes scapularis* (Acari: Ixodidae) collected during 1998 and 1999 from Minnesota. *J. Med. Entomol.* **39** (1), 218-220.

Le Fleche, A., D. Postic, K. Girardet, O. Peter and G. Baraton (1997): Characterization of *Borrelia lusitaniae* sp. nov. by 16S ribosomal DNS sequence analysis. *Int. J. Syst. Bacteriol.* **47**, 921-925.

Leutenegger, C.M., N. Pusterla, C.N. Mislin, R. Weber and H. Lutz (1999): Molecular evidence of coinfection of ticks with *Borrelia burgdorferi* sensu lato and the human granulocytic ehrlichiosis agent in Switzerland. *J. Clin. Microbiol.* **37**, 3390-3391.

Leutenegger, C.M., N. Pusterla, R. Wicki und H. Lutz (2002): Neue molekularbiologische Nachweismethoden am Beispiel zeckengebundener Infektionserreger. Schweiz. Arch. Tierheilk. **144** (8), 395-404.

Levy, S.A. and P.H. Duray (1988): Complete heart block in a dog seropositive for *Borrelia burgdorferi*. J. Vet. Intern. Med. **2**, 138-144.

Liebisch, G. (1993): Zeckenborreliose bei Haustieren. In: Horst, H. (Hrsg.), Einheimische Zeckenborreliose (Lyme-Krankheit) bei Mensch und Tier, PERIMED-spitta, Nürnberg, 164-187.

Liebisch, A. und G. Liebisch (1993): Lyme-Borreliose bei Hunden – Literaturübersicht und Ergebnisse aus Norddeutschland. Mh. Vet. Med. **48**, 479-487.

Liebisch, G., B. Sohns and W. Bautsch (1998): Detection and typing of *Borrelia burgdorferi* sensu lato in *Ixodes ricinus* ticks attached to human skin by PCR. J. Clin. Microbiol. **36**, 3355-3358.

Liebisch, G., G. Assmann und A. Liebisch (1999): Infektion mit *Borrelia burgdorferi* s.l. als Krankheitsursache der Lyme-Borreliose bei Pferden in Deutschland. Prakt. Tierarzt **80:6**, 498-516.

Liebisch, G., A. Schlotzhauer, B. Sohns, G. Assmann and A. Liebisch (2002): Isolation of *Borrelia afzelii* from a horse with Lyme Borreliosis. IX Int. Conf. on Lyme Borreliosis and other tick-borne diseases 2002, New York, Poster, P-109.

Lindenmeyer, J.L., M. Weber, A. Ondertonk and J. Plain (1989): *Borrelia burgdorferi* infection in horses. JAVMA **194**, 1384.

Lindner, A. und K. Böckel (1995): Vorkommen von Antikörpern gegen *Borrelia burgdorferi* bei Hunden und Katzen. Prakt. Tierarzt **76** (3), 177-185.

Lischer, C.J., C.M. Leutenegger, U. Braun and H. Lutz (2000): Diagnosis of Lyme disease in two cows by the detection of *Borrelia burgdorferi* DNA. Vet. Rec. **146**, 497-499.

- Lissmann, B.A., E.M. Bosler, H. Camay, B.G. Ormiston and J.L. Benach** (1984): Spirochete-associated arthritis (Lyme disease) in a dog. *J. Am. Vet. Med. Assoc.* **185** (2), 219-220.
- Livesley, M.A., I.P. Thompson, M.J. Bailey and P.A. Nuttall** (1993): Comparison of the fatty acid profiles of *Borrelia*, *Serpulina* and *Leptospira* species. *J. Gen. Microbiol.* **139**, 889-895.
- Livesley, M.A., D. Carey, L. Gern and P.A. Nuttall** (1994): Problems of isolating *Borrelia burgdorferi* from ticks collected in United Kingdom foci of Lyme disease. *Med. Vet. Entomol.* **8**, 172-178.
- Macedo-Aguirre, S., A. Schönberg, V. Fingerle, L. Pradel and B. Wilske** (1996): Immunological characterisation of *Borrelia burgdorferi* sensu lato isolated from Eastern Germany. VII International Congress on Lyme Borreliosis, 1996, San Francisco. Abstracts, p. 353.
- Madigan, J.** (1993): Lyme disease (Lyme borreliosis) in horses. *Vet. Clin. of North Am. Eq. Pract.* **9**, 429-434.
- Magnarelli, L.A. and J.F. Anderson** (1989): Class specific and polyvalent enzyme-linked immunosorbent assays for detection of antibodies to *Borrelia burgdorferi* in equids. *J. Am. Vet. Med. Assoc.* **195**, 1365-1368.
- Magnarelli, L.A., J.F. Anderson and R.C. Johnson** (1987a): Cross-reactivity in serological tests for Lyme disease and other spirochetal infections. *J. Infect. Dis.* **156**, 183-188.
- Magnarelli, L.A., J.F. Anderson, A.B. Schreier and C.M. Ficke** (1987b): Clinical and serological studies of canine borreliosis. *J. Am. Vet. Med. Assoc.* **191** (9), 1089-1094.
- Magnarelli, L.A., J.F. Anderson, E. Shaw, J.E. Post and F.C. Palka** (1988): Borreliosis in equids in northeastern United States. *Am. J. Vet. Res.* **49**, 359-362.
- Magnarelli, L.A., J.F. Anderson, H.R. Levine and S.A. Levy** (1990a): Tick parasitism and antibodies to *Borrelia burgdorferi* in cats. *J. Am. Vet. Med. Assoc.* **197** (1), 63-66.

Magnarelli, L.A., J.F. Anderson and A.B. Schreier (1990b): Persistence of antibodies to *Borrelia burgdorferi* in dogs of New York and Connecticut. *J. Am. Vet. Med. Assoc.* **196** (7), 1064-1068.

Magnarelli, L.A., J.F. Anderson, R.C. Johnson, R.B. Nadelman and G.P. Wormser (1994): Comparison of different strains of *Borrelia burgdorferi* sensu lato used as antigens in enzym-linked immunosorbent assays. *J. Clin. Microbiol.* **32**, 1154-1158.

Magnarelli, L.A., J.W. Ijdo, A.E. van Andel, C. Wu, S.J. Padula and E. Frikg (2000): Serologic confirmation of *Ehrlichia equi* and *Borrelia burgdorferi* infections in horses from northeastern United States. *J. Am. Vet. Med. Assoc.* **217**, 1045-1050.

Magnarelli, L.A., S.A. Levy, J.W. Ijdo, C. Wu, S.J. Padula and E. Frikg (2001): Reactivity of dog sera to whole-cell or recombinant of *Borrelia burgdorferi* by ELISA and immunoblot analysis. *J. Med. Microbiol.* **50**, 889-895.

Maloney, E.M. and J.L. Lindenmayer (1992): Seroprevalence and clinical signs of Lyme disease in Cape Code horses. *Eq. Pract.* **14**, 15-19.

Manion, T.B., M.I. Khan, J. Dinger and S.L. Bushmich (1998): Viable *Borrelia burgdorferi* in the urine of two clinically normal horses. *J. Vet. Diag. Invest.* **10**, 196-199.

Marconi, R.T. and C.F. Garon (1992): Development of polymerase chain reaction primer sets for diagnosis of Lyme disease and for species-specific identification of Lyme disease isolates by 16S rRNA signature nucleotide analysis. *J. Clin. Microbiol.* **30**, 2830-2834.

Marcus, L.C., M.M. Patterson, R.E. Gilfillan and P.H. Urband (1985): Antibodies to *B. burgdorferi* in New England horses. *Am. J. Vet. Res.* **46**, 2570-2571.

Masuzawa, T. und Y. Yanagihara (1998): Die Erreger-Klassifikation, Vorkommen und klinische Relevanz. In: Talaska, T. (Hrsg.), Für die Praxis: Lyme Borreliose, Chromik Offsetdruck, Brieskow-Finkenheerd, ISBN 3-00-002363-1, 11-18.

Matuschka, F.-R., H. Eiffert, A. Ohlenbusch and A. Spielman (1994): Amplifying role of edible dormice in Lyme disease transmission in Central Europe. *J. Infect. Dis.* **170**, 122-127.

- Matuschka, F.-R., A. Ohlenbusch, H. Eiffert, D. Richter and A. Spielman** (1996): Characteristics of Lyme disease spirochetes in archived European ticks. *J. Infect. Dis.* **174** (2), 424-426.
- Matuschka, F.R., T.W. Schinkel, B. Klug, A. Spielman and D. Richter** (1998): Failure of Ixodes ticks to inherit *Borrelia afzelii* infection. *Appl. Environ. Microbiol.* **64**, 3089-3091.
- Mitchell, P.D., K.D. Reed and J.M. Hofkes** (1996): Immunoserologic evidence of coinfection with *Borrelia burgdorferi*, *Babesia microti* and human granulocytic Ehrlichia species in residents of Wisconsin and Minnesota. *J. Clin. Microbiol.* **34**, 724-727.
- Mönchenberg, J.S.** (1998): Polymerase-Kettenreaktion und Restriktionsenzym-Analyse zum Nachweis und zur Differenzierung von *Borrelia burgdorferi* sensu lato in Schildzecken. *Vet.med. Diss.*, Tierärztliche Fakultät der LMU München, München.
- Müller, E.** (1994): Borreliose bei Hunden – klinische und serologische Befunde. *Kleintierpraxis* **39**, 375-380.
- Müller, I.** (1999): Immunoblot-Muster von Pferdeseren mit fünf verschiedenen Borrelien-Arten. *Vet. med. Diss.*, Vet.med. Univers. Wien.
- Munderloh, U.G., T.J. Kurtti, R.C. Johnson and G.G. Ahlstrand** (1988): Colony formation by Lyme disease spirochetes. *Ann. N.Y. Acad. Sci.* **539**, 404-406.
- Murgia, R., C. Piazetta and M. Cinco** (2002): Cystic forms of *Borrelia burgdorferi* sensu lato: induction, development and the role of RpoS. *Wien. Klin. Wochenschr.* **114** (13-14), 574-579.
- National Committee for Clinical Laboratory Standards (NCCLS)** (1998): Western blot assay for antibodies to *Borrelia burgdorferi*; proposed guideline. M34-9, Vol. **18** No. **12**, 1-30.
- Niesenbaum, K.** (1991): Die Borreliose (Lyme disease) des Hundes. *Waltham Internat. Focus* **1**, 6-9.

Oehme, R. und P. Kimmig (1999): Epidemiologie der FSME und der Lyme-Borreliose in Baden-Württemberg. PCR-Workshop FSMEV, Berlin, 17.-18.12.1999 (unveröffentlicht).

Oheim, S. und H. Herrmann (1994): Untersuchungen zur Verbreitung von *Borrelia burgdorferi* im Raum Greifswald in der Region Vorpommern des Landes Mecklenburg-Vorpommern. Z. f. Ä. Fb. **88**, 807-810.

Oschmann, P., P. Kraiczký, J. Halperin and V. Brade (1999): Lyme Borreliosis and Tick-Borne Encephalitis. In: Oschmann, Kraiczký (Hrsg.), Bremen: UNI-MED Verlag.

Ott, A. und A. Schönberg (1988): Erste kulturelle Anzüchtung von *Borrelia burgdorferi* aus der Hautprobe einer Patientin mit Erythema migrans in Berlin. Dt. Ärztebl. **85**, Heft 50 (27), 2187-2188.

Panelius, J., P. Lahdenne, T. Heikkila, M. Peltomaa, J. Oksi and I. Seppala (2002): Recombinant OspC from *Borrelia burgdorferi* sensu stricto, *B. afzelii* and *B. garinii* in the serodiagnosis of Lyme borreliosis. J. Med. Microbiol. **51** (9), 731-739.

Parker, J.L. and K.K. White (1992): Lyme borreliosis in cattle and horses: A review of the literature. Cornell Vet. **82**, 253-274.

Pfister, K., B. Bigler, J. Neswadba, L. Gern and A. Aeschlimann (1989): *Borrelia burgdorferi* infections of dogs in Switzerland. Zbl. Bakt. **18**, 26-31.

Philipp, M.T. and B.J. Johnson (1994): Animal models of Lyme disease: pathogenesis and immunoprophylaxis. Trends in Microbiology **2** (11), 431-437.

Pichon, B., L. Mousson, C. Figureau, F. Rodhain and C. Perez-Eid (1999): Density of deer in relation to the prevalence of *Borrelia burgdorferi* s.l. in *Ixodes ricinus* nymphs in Rambouillet Forest, France. Expt. Appl. Acarol. **23**, 267-275.

Picken, R.N. (1992): Polymerase chain reaction primers and probes derived from flagellin gene sequences for specific detection of the agents of Lyme disease and North American Relapsing fever. J. Clin. Microbiol. **30**, 99-114.

Pollack, R.J., S.R. Telford III and A. Spielman (1993): Standardization of medium for culturing Lyme disease spirochetes. *J. Clin. Microbiol.* **31**, 1251-1255.

Post, J.E., E.E. Shaw and F.C. Palka (1987): Lyme disease in horses. *Vet. Cl. Of North Am. Eq. Pract.* **32**, 415-420.

Post, J.E., E.E. Shaw and S.D. Wright (1988): Suspected borreliosis in cattle. *Ann. N.Y. Acad. Sci.* **539**, 488.

Preac-Mursic, V. und B. Wilske (1992): Flexible Schraubenbakterien – Gattung *Borrelia*. In: Burghardt F (ed) *Mikrobiologische Diagnostik*. Thieme, Stuttgart, New York, 629.

Preac-Mursic, V. and B. Wilske (1993): Biology of *Borrelia burgdorferi*. In: Weber, K. & Burgdorfer, W. (Hrsg.), *Aspects of Lyme Borreliosis*, Springer, Berlin, Heidelberg, 44-58.

Preac-Mursic, V., B. Wilske, G. Schierz, H.-W. Pfister and K. Einhäupl (1984): Repeated isolation of spirochetes from cerebrospinal fluid of a patient with meningoradiculitis Bannwarth. *Eur. J. Clin. Microbiol.* **3**, 564-565.

Preac-Mursic, V., B. Wilske and G. Schierz (1986): European *Borrelia burgdorferi* isolated from humans and ticks – Culture conditions and antibiotic susceptibility. *Zbl. Bakt. Hyg. A* **263**, 112-118.

Preac-Mursic, V., B. Wilske and S. Reinhardt (1991): Culture of *Borrelia burgdorferi* on six solid media. *Eur. J. Clin. Microbiol. Infect. Dis.* **10**, 1076-1079.

Priem, S. und A. Krause (1999): Labordiagnostik der Lyme Borreliose. In: A. Krause, G.R. Burmester (Hrsg.): *Lyme Borreliose*. Stuttgart: Georg Thieme Verlag, 17-24.

Reichl, U. (1996): Application of molecular-biology-based methods to the diagnosis of infectious diseases. *Front Biosc.* **1**, 72-77.

- Rijpkema, S.G.T., D.J. Tazelaar, M.J.C.H. Molkenboer, G.T. Noordhoek, G. Plantinga, L.M. Schouls and J.F.P. Schellekens** (1997): Detection of *Borrelia afzelii*, *Borrelia burgdorferi* sensu stricto, *Borrelia garinii* and group VS116 by PCR in skin biopsies of patients with erythema migrans and acrodermatitis chronica atrophicans. *Clin. Microbiol.* **3**, 109-116.
- Roessler, D., H. Eiffert, S. Jauris-Heipke, G. Lehnert, V. Preac-Mursic, J. Teepe, T. Schlott, E. Soutschek and B. Wilske** (1995): Molecular and immunological characterization of the p 83/100 protein of various *Borrelia burgdorferi* sensu lato strains. *Med. Microbiol. Immunol.* **184**, 23-32.
- Roessler, D., V. Vasiliu and B. Wilske** (1996): Development of OspA type specific PCR for characterization of *Borrelia burgdorferi* sensu lato. *J. Microbiol. Meth.* **27**, 102, Abstr. 19.
- Rolle, M. und A. Mayr** (2002): Medizinische Mikrobiologie, Infektions- und Seuchenlehre. 7. Aufl., Enke Verlag, Stuttgart.
- Rosa, P.A., D. Hogan and T.G. Schwan** (1991): Polymerase chain reaction analysis identify two distinct classes of *Borrelia burgdorferi*. *J. Clin. Microbiol.* **29**, 524-532.
- Russel, H., J.S. Sampson, G.P. Schmid, H.W. Wilkinson and B. Plikaytis** (1984): Enzyme-linked immunosorbent assay and indirect immunofluorescence assay for Lyme disease. *J. Infect. Dis.* **149**, 465-470.
- Ruzic-Sabljic, E., V. Maraspin, J. Cimperman, S. Lotric-Furlan and F. Strle** (2002): Evaluation of immunofluorescence test (IFT) and immuno (western) blot (WB) test in patients with erythema migrans. *Wien. Klin. Wochenschr.* **114** (13-14), 586-590.
- Saint Girons, I., I.G. Old and B.E. Davidson** (1994): Molecular biology of the *Borrelia*, bacteria with linear replicons. *Microbiol.* **140** (8), 1803-1816.
- Saint Girons, I., L. Gern, J.S. Gray, E.C. Guy, E. Korenberg, P.A. Nuttall, S.G.T. Rijpkema, A. Schönberg, G. Stanek and D. Postic** (1998): Identification of *Borrelia burgdorferi* sensu lato species in Europe. *Zentr.bl. Bakteriol.* **287**, 190-195.
- Satz, N.** (1993): Klinik der Lyme-Borreliose. Verlag Hans Huber, Bern.

Salyers, A. and D. Whitt (1994): Lyme disease and Syphilis. Bacterial pathogenesis: a molecular approach. Washington D.C., American Society for Microbiology, 290-299.

Schmid, G.P. (1985): The global distribution of Lyme disease. Rev. Infect. Dis. **7** (1), 41-50.

Schneider, P. (2002): Prof. Enderleins Forschung aus heutiger Sicht – Lassen sich die Forschungsergebnisse mit modernen Untersuchungsmethoden bestätigen? Ganzheitl. Tiermedizin **16**, 23-31.

Schönberg, A. (1997): Epidemiology of Lyme Borreliosis in Europe. Proc. 2nd Int. Symp. Lyme Disease, 27-28 October 1997 in Japan, 165-188.

Schönberg, A., C. Camey, O. Kahl, B. Wilske, V. Preac-Mursic and K. Hovind-Hougen (1988): First isolation of *Borrelia burgdorferi*, the agent of Lyme Borreliosis, from *Ixodes ricinus* (Acari: Ixodidae) in Berlin (West). Zbl. Bakt. Hyg. A **268**, 487-494.

Schönberg, A., K. Schmidt und A. Käsbohrer (1989): Eine durch Zecken übertragbare Zoonose: Lyme-Borreliose. Bundesgesundhbl. **32**, Sonderdruck, 190-193.

Schönberg, A., C. Loser and S. Gupta (1995): *Borrelia-burgdorferi*-Isolate aus Zecken der neuen Bundesländer und ihre Reaktionen mit monoklonalen Antikörpern. In: Süss, J. (Hrsg.), Durch Zecken übertragbare Erkrankungen/FSME und Lyme-Borreliose, 3. Potsdamer Symp., 11. März 1995, Weller, Schriesheim, 155-161.

Schulze, P., S. Macedo, A. Schönberg und M. Schwebs (1995): Morphologische Unterschiede von *Borrelia-burgdorferi*-Stämmen bei elektronenmikroskopischen Untersuchungen. In: Süss, J. (Hrsg.), Durch Zecken übertragbare Erkrankungen/FSME und Lyme-Borreliose, 3. Potsdamer Symp., 11. März 1995, Weller, Schriesheim, 175-181.

Scott Hefty, P., S.E. Jolliff, M.J. Caimano, S.K. Wikle and D.R. Akins (2002): Changes in Temporal and Spatial Patterns of Outer Surface Lipoprotein Expression Generate Population Heterogeneity and Antigenic Diversity in the Lyme Disease Spirochete, *Borrelia burgdorferi*. Infect. and Immunol. **70** (7), 3468-3478.

Sigal, L.H. (1993): Cross-reactivity between *B. burgdorferi* flagellin and a human axonal 64.000 molecular weight protein. J. Infect. Dis. **167**, 1372-1378.

Simon, M.M., U. Altenschmidt, E. Böggemeyer, L. Gern, H. Fuchs, N. Honarvar, U. Hurtenbach, K. Kurtenbach, M. Modolell, S. Moter, C. Museteau, U.E. Schaible, R. Wallich and M.D. Kramer (1994): Pathogenesis of Lyme borreliosis: lessons from the mouse model. Proc. VI Int. Conf. on Lyme Borreliosis, Bologna, Italy, June 19-22, 1994, 69-75.

Simon, M.M., Y. Bauer, W. Zhong, H. Hofman and R. Wallich (1999): Lyme disease: Pathogenesis and vaccine development. Zentr.bl. Bakteriol. **289**, 690-695.

Simpson, W.J., W. Burgdorfer, M.E. Schrumpf, R.H. Karstens and T.G. Schwan (1991): Antibody to a 39-kilodalton *Borrelia burgdorferi* antigen (P39) as a marker for infection in experimentally and naturally inoculated animals. J. Clin. Microbiol. **29**, 236-243.

Skotarczak, B., B. Wodecka and A. Stachow (1998): Detection of Lyme disease spirochetes, *Borrelia burgdorferi* sensu lato DNA by polymerase chain reaction in ticks in the recreative areas of Szczecin. 18. Tagung der DVG, 24.-28. März 1998 in Dresden, Abstr. P 122.

Speck, S., K. Failing, B. Reiner and M.M. Wittenbrink (2002): Evaluation of different media and a BGM cell culture assay for isolation of *Borrelia burgdorferi* sensu lato from ticks and dogs. Vet. Microbiol. **89** (4), 291-302.

Stanek, G. (1997a): Biology of *Borrelia burgdorferi*, risk of infection of Lyme borreliosis. In: Süss, J., Kahl, O. (Hrsg.), Tick-borne Encephalitis and Lyme borreliosis, 4th Int. Potsdam Symp. On Tick-Borne Dis., 21.-22. Februar 1997, Pabst Science Publishers, Lengerich, 230-237.

Stanek, G. (1997b): Neues in der Behandlung der Lyme-Borreliose. Antibiotika Monitor XIII, 65-80.

Stanek, G. (1998): Klinische Falldefinitionen der Lyme-Borreliose für Europa. In: Talaska, T. (Hrsg.), Für die Praxis: Lyme-Borreliose, Chromik Offsetdruck, Brieskow-Finkenheerd, ISBN 3-00-002363-1, 114-117.

Stanek, G., H. Flamm, A.G. Barbour and W. Burgdorfer (1987): Lyme borreliosis. Proceedings of the second international symposium on Lyme disease and related disorders, Vienna 1985. Gustav Fischer, Stuttgart, New York.

- Stanek, G., M. Pletschette, H. Flamm, A.M. Hirschl, E. Aberer, W. Kristoferitsch and E. Schmutzhard** (1988): European Lyme borreliosis. Ann. N.Y. Acad. Sci. **539**, 274-282.
- Steere, A.C.** (1989): Lyme disease. New Engl. J. Med. **321**, 586-596.
- Steere, A.C., S.E. Malavista, D.R. Snydman, D.E. Shope, W.A. Andiman, M.R. Ross and F.M. Steele** (1977a): Lyme arthritis: An endemic of oligoarticular arthritis in children and adults in three Connecticut communities. Arthritis Rheum. **20**, 7-17.
- Steere, A.C., S.E. Malavista, J.A. Hardin, S. Ruddy, P.W. Askenase and W.A. Andiman** (1977b): Erythema chronicum migrans and Lyme arthritis the enlarging clinical spectrum. Ann. Intern. Med. **86**, 685-698.
- Steere, A.C., T.F. Broderick and S.E. Malavista** (1978): Erythema chronicum migrans and Lyme arthritis: Epidemiologic evidence for a tick vector. Am. J. Epidemiol. **108**, 312-321.
- Steere, A.C., N.H. Bartenhagen, J.E. Craft, G.J. Hutchinson, J.H. Newman, D.W. Rahn, L.H. Sigal, P.N. Spieler, K.S. Stenn and S.E. Malavista** (1983): The early clinical manifestations of Lyme disease. Ann. Intern. Med. **99**, 76-82.
- Steere, A.C., R.L. Grodzicky, J.E. Craft, M. Shrestha, A.N. Kornblatt and S.E. Malavista** (1984): Recovery of Lyme disease spirochetes from patients. Yale J. Biol. Med. **57**, 557-560.
- Stiernstedt, G.T.** (1985): Tick borne *Borrelia* infection in Sweden. Scand. J. Infect. Dis. (suppl **45**), 1-70.
- Straubinger, R., A. Straubinger, B. Summers, R. Jacobson and H. Erb** (1998): Clinical manifestations, pathogenesis and effect of antibiotic treatment on Lyme borreliosis in dogs. Wien. Klin. Woch. **110** (24), 874-881.
- Suter, P.** (1994): Infektionskrankheiten. In: Praktikum der Hundeklinik. Niemand, G. und P. Suter (Hrsg.) 8. Auflage, 231, Blackwell Wissenschaftsverlag Berlin.

- Svartz, N.** (1946): Penicillin Behandlung vid dermatitis atrophicans Herxheimer. Nord. Med. **32**, 2783.
- Talaska, T.** (1998a): Diagnostische Methoden bei Borrelien-Infektionen – Übersicht. In: T. Talaska (Hrsg.): Für die Praxis: Lyme-Borreliose. 48-59.
- Talaska, T.** (1998b): Komplementbindungsreaktion in der Borreliose-Diagnostik? In: T. Talaska (Hrsg.): Für die Praxis: Lyme-Borreliose. 62-67.
- Tasai, M., I. Takashima, H. Kariwa, N. Hashimoto, T. Kondo, T. Sugiura and M. Kamada** (1993): Seroimmunological survey of Lyme borreliosis in horses in Japan by immunofluorescent enzyme-linked immunosorbent assay. Bull. Eq. Res. Inst. **30**, 37-42.
- Taylor, F.G.R. and M.H. Hillyer** (2001): Klinische Diagnostik in der Pferdepraxis. Hannover, Schlüterscher Verlag, 19-20.
- Terekhova, D., M.L. Sartakova, G.P. Wormser, I. Schwartz and F.C. Cabello** (2002): Erythromycin resistance in *B. burgdorferi*. Antimicrobial agents and chemotherapy. Vol. **46** (11), 3637-3640.
- Thomas, L.** (1992): Labor und Diagnose. 4. Aufl. In: Med. Verlagsgesellsch. Marburg 1992.
- Toutoungi, L.N. and L. Gern** (1993): Ability of transovarially and subsequent transstadially infected *Ixodes hexagonus* ticks to maintain and transmit *Borrelia burgdorferi* in the laboratory. Exp. Appl. Acarol. **17**, 581-586.
- Tylewska-Wierzbanowska, S. and T. Chmielewski** (2002): Limitation of serological testing for Lyme borreliosis: evaluation of ELISA and western blot in comparison with PCR and culture methods. Wien. Klin. Wochenschr. **114** (13-14), 601-605.
- van Dam, A.P., H. Kuiper, K. Vos et al.** (1993): Different genospecies of *Borrelia burgdorferi* are associated with distinct clinical manifestations of Lyme borreliosis. Clin. Infect. Dis. **17**, 708-717.

Venner M. und E. Deegen (1996): Interpretation von *Borrelia burgdorferi* Antikörpertitern beim Pferd unter Berücksichtigung der Kenntnisse zur Borreliose beim Menschen – eine Literaturübersicht. Pferdeheilkunde **12**, 865-873.

Wasmoen, T.L., R.W. Sebring, B.M. Blumer, L.G. Chavez, H.-J. Chu and W.M. Acree (1992): Examination of Koch's postulates for *Borrelia burgdorferi* as the causative agent of limb/joint dysfunction in dogs with borreliosis. J. Am. Vet. Med. Assoc. **201** (3), 412-418.

Weber, A., U. Heim und R. Schäfer (1991): Zum Vorkommen von Antikörpern gegen *Borrelia burgdorferi* bei Hunden einer Kleintierpraxis in Nordbayern. Berl. Münch. Tierärztl. Wschr. **104**, 384-386.

Weber, R., N. Pusterla, M. Loy and H. Lutz (1998): Fever, leukopenia and thrombocytopenia in a patient with acute Lyme borreliosis were due to human granulocytic ehrlichiosis. Clin. Inf. Dis. **26**, 253-254.

Wheeler, C.M., J.C.G. Monco, J.L. Benach, M.G. Golightly, G.S. Habicht and A.C. Steere (1993): Nonprotein antigens of *Borrelia burgdorferi*. J. Infect. Dis. **167**, 665-674.

Wicki, R., P. Sauter, C. Mettler, S. Natsch, T. Enzler, N. Pusterla, P. Kuhnert, G. Egli, M. Bernasconi, R. Lienhard, H. Lutz and C.M. Leutenegger (2000): A swiss army survey in Switzerland to determine the prevalence of *Francisella tularensis*, members of the *Ehrlichia phagocytophila* genogroup, *Borrelia burgdorferi* sensu lato and tick-borne encephalitis virus in ticks. Europ. J. Clin. Microbiol. Inf. Dis. **20**, 427-432.

Wilske, B. and V. Preac-Mursic (1993): Microbiological diagnosis of Lyme Borreliosis. In: Weber, K. & Burgdorfer, W. (Hrsg.), Aspects of Lyme Borreliosis, Springer, Berlin, Heidelberg, 44-58.

Wilske, B., V. Preac-Mursic, G. Schierz, R. Kuhbeck, A.G. Barbour and M. Kramer (1988): Antigenic variability of *Borrelia burgdorferi*. Ann. N.Y. Acad. Sci. **539**, 126-143.

Wilske, B., V. Fingerle, P. Herzer, A. Hoffmann, G. Lehnert, H. Peters, H.-W. Pfister, V. Preac-Mursic, E. Soutschek and K. Weber (1993): Recombinant immunoblot in the serodiagnosis of Lyme borreliosis. Med. Microbiol. Immunol. **182**, 255-270.

Wilske, B., V. Fingerle, U. Hauser und D. Rössler (1997): Diagnostische Bibliothek – Borrelien. Lab. Med. **48** (6), 1-12.

Wilske, B., L. Zöller, V. Brade, H. Eiffert, U.B. Göbel, G. Stanek und H.W. Pfister (2000): Lyme-Borreliose. MiQ 12/2000 Qualitätsstandards in der mikrobiologisch-infektiologischen Diagnostik. In: H. Mauch und R. Lütticken (Hrsg.), Urban & Fischer.

Wolf, S. (1996): Elektronenmikroskopische Untersuchungen zur Ultrastruktur von *Borrelia burgdorferi*. Vet.med. Diss., FU Berlin.

Zeman, P. (1998): *Borrelia*-infection rates in tick and insect vectors accompanying human risk of acquiring Lyme borreliosis in a highly endemic region in Central Europe. Folia Parasitol. **45**, 319-325.

Zhioua, E., A. Aeschlimann and L. Gern (1994): Infection of field-collected *Ixodes ricinus* (Acari: *Ixodidae*) larvae with *Borrelia burgdorferi* in Switzerland. J. Med. Entomol. **31**, 735-766.

Zhioua, E., D. Postic, F. Rodhain and C. Perez-Eid (1996): Infection of *Ixodes ricinus* (Acari: *Ixodidae*) by *Borrelia burgdorferi* in Ile de France. J. Med. Entomol. **33**, 694-697.

Zipfel, P.F., P. Kraiczy und J. Hellwage (2002): Das tägliche Versteckspiel: Wie Mikroorganismen der Immunabwehr entgehen. In: Biologie in unserer Zeit, Weinheim: WILEY-VCH Verl. GmbH, Vol. 32, **6**, 371-379.