

## 6. Literaturverzeichnis

1. Klemperer G. Untersuchungen über Stoffwechsel und Ernährung in Krankheiten. In: Leyden E, ed. Arbeiten aus der ersten medicinischen Klinik der Charité zu Berlin. Berlin: August Hirschwald 1891:118-73.
2. Ezzati M, Lopez AD, Rodgers A, Vander HS, Murray CJ. Selected major risk factors and global and regional burden of disease. Lancet 2002;360:1347-60.
3. Dramaix M, Hennart P, Brasseur D et al. Serum albumin concentration, arm circumference, and oedema and subsequent risk of dying in children in central Africa. BMJ 1993;307:710-3.
4. Gomez-Almaguer D, Ruiz-Arguelles GJ, Ponce-de-Leon S. Nutritional status and socio-economic conditions as prognostic factors in the outcome of therapy in childhood acute lymphoblastic leukemia. Int.J.Cancer Suppl 1998;11:52-5.
5. Muller MJ, Mast M, Langnase K. [WHO warns of obesity epidemic. Are we becoming a society of obese persons?]. MMW.Fortschr.Med. 2001;143:28-32.
6. WHO. Report of a WHO Expert Committee. Physical status: the use and interpretation of anthropometry. WHO. 854. 1995. Genf.
7. Langnase K, Muller MJ. Nutrition and health in an adult urban homeless population in Germany. Public Health Nutr. 2001;4:805-11.
8. Reuben DB, Greendale GA, Harrison GG. Nutrition screening in older persons. J.Am.Geriatr.Soc. 1995;43:415-25.
9. Pirlich M, Luhmann N, Schutz T, Plauth M, Lochs H. Mangelernährung bei Klinikpatienten: Diagnostik und klinische Bedeutung. Akt.Ernähr.-Med. 1999;24:260-6.
10. Committee of Experts on Nutrition FSaCP. Food and nutritional care in hospitals: how to prevent undernutrition. Strasburg: Council of Europe Publishing, 2002.
11. Pirlich M, Schwenk A, Müller MJ et al. DGEM-Leitlinie Enterale Ernährung: Ernährungsstatus. Akt.Ernähr.-Med. 2003;28:10-25.
12. Klein S, Kinney J, Jeejeebhoy K et al. Nutrition support in clinical practice: review of published data and recommendations for future research directions. Summary of a conference sponsored by the National Institutes of Health, American Society for Parenteral and Enteral Nutrition, and American Society for Clinical Nutrition. Am.J Clin Nutr 1997;66:683-706.
13. Kondrup J, Allison SP, Elia M, Vellas B, Plauth M. ESPEN guidelines for nutrition screening 2002. Clin.Nutr. 2003;22:415-21.
14. Stratton RJ, Green CJ, Elia M. Disease-related Malnutrition: An evidence-based approach to treatment. Oxon, UK: CABI Publishing, 2003.
15. White H, Pieper C, Schmader K. The association of weight change in Alzheimer's disease with severity of disease and mortality: a longitudinal analysis. J.Am.Geriatr.Soc. 1998;46:1223-7.
16. Pirlich M, Lochs H. Nutrition in the elderly. Best.Pract.Res.Clin Gastroenterol 2001;15:869-84.
17. Tierney AJ. Undernutrition and elderly hospital patients: a review. J Adv.Nurs. 1996;23:228-36.
18. Volkert D, Kruse W, Oster P, Schlierf G. Malnutrition in geriatric patients: diagnostic and prognostic significance of nutritional parameters. Ann.Nutr.Metab 1992;36:97-112.

19. Muller MJ, Lautz HU, Plogmann B, Burger M, Korber J, Schmidt FW. Energy expenditure and substrate oxidation in patients with cirrhosis: the impact of cause, clinical staging and nutritional state. *Hepatology* 1992;15:782-94.
20. Suttmann U, Ockenga J, Hoogestraat L et al. Resting energy expenditure and weight loss in human immunodeficiency virus-infected patients. *Metabolism* 1993;42:1173-9.
21. Anker SD, Coats AJ. Cardiac cachexia: a syndrome with impaired survival and immune and neuroendocrine activation. *Chest* 1999;115:836-47.
22. Anker SD, Chua TP, Ponikowski P et al. Hormonal changes and catabolic/anabolic imbalance in chronic heart failure and their importance for cardiac cachexia. *Circulation* 1997;96:526-34.
23. Argiles JM, Busquets S, Lopez-Soriano FJ. Cytokines in the pathogenesis of cancer cachexia. *Curr.Opin.Clin.Nutr.Metab Care* 2003;6:401-6.
24. Mitch WE, Price SR. Mechanisms activated by kidney disease and the loss of muscle mass. *Am.J.Kidney Dis.* 2001;38:1337-42.
25. Tisdale MJ. Cachexia in cancer patients. *Nat.Rev.Cancer* 2002;2:862-71.
26. Ott M, Lembcke B, Fischer H et al. Early changes of body composition in human immunodeficiency virus-infected patients: tetrapolar body impedance analysis indicates significant malnutrition. *Am J Clin Nutr* 1993;57:15-9.
27. Suttmann U, Ockenga J, Selberg O, Hoogestraat L, Deicher H, Muller MJ. Incidence and prognostic value of malnutrition and wasting in human immunodeficiency virus-infected outpatients. *J.Acquir.Immune Defic.Syndr.Hum.Retrovirol.* 1995;8:239-46.
28. Andreyev HJ, Norman AR, Oates J, Cunningham D. Why do patients with weight loss have a worse outcome when undergoing chemotherapy for gastrointestinal malignancies? *Eur.J Cancer* 1998;34:503-9.
29. Selberg O, Bottcher J, Tusch G, Pichlmayr R, Henkel E, Muller MJ. Identification of high- and low-risk patients before liver transplantation: a prospective cohort study of nutritional and metabolic parameters in 150 patients. *Hepatology* 1997;25:652-7.
30. Caregaro L, Alberino F, Amodio P et al. Malnutrition in alcoholic and virus-related cirrhosis. *Am.J Clin.Nutr.* 1996;63:602-9.
31. Selberg O, Bottcher J, Pirlich M, Henkel E, Manns MP, Muller MJ. Clinical significance and correlates of whole body potassium status in patients with liver cirrhosis. *Hepatol.Res.* 1999;1999:36-48.
32. Schols AM, Slangen J, Volovics L, Wouters EF. Weight loss is a reversible factor in the prognosis of chronic obstructive pulmonary disease. *Am.J Respir.Crit Care Med.* 1998;157:1791-7.
33. Anker SD, Ponikowski P, Varney S et al. Wasting as independent risk factor for mortality in chronic heart failure. *Lancet* 1997;349:1050-3.
34. Ciciora M, Davos CH, Florea V et al. Chronic heart failure in the very elderly: clinical status, survival, and prognostic factors in 188 patients more than 70 years old. *Am Heart J* 2001;142:174-80.
35. Anker SD, Negassa A, Coats AJ et al. Prognostic importance of weight loss in chronic heart failure and the effect of treatment with angiotensin-converting-enzyme inhibitors: an observational study. *Lancet* 2003;361:1077-83.
36. Nixon DW, Heymsfield SB, Cohen AE et al. Protein-calorie undernutrition in hospitalized cancer patients. *Am J Med* 1980;68:683-90.

37. Aviles A, Yanez J, Lopez T, Garcia EL, Guzman R, Diaz-Maqueo JC. Malnutrition as an adverse prognostic factor in patients with diffuse large cell lymphoma. *Arch.Med.Res.* 1995;26:31-4.
38. Rey-Ferro M, Castano R, Orozco O, Serna A, Moreno A. Nutritional and immunologic evaluation of patients with gastric cancer before and after surgery. *Nutrition* 1997;13:878-81.
39. Viana MB, Murao M, Ramos G et al. Malnutrition as a prognostic factor in lymphoblastic leukaemia: a multivariate analysis. *Arch.Dis.Child* 1994;71:304-10.
40. Kotler DP, Tierney AR, Wang J, Pierson RN, Jr. Magnitude of body-cell-mass depletion and the timing of death from wasting in AIDS. *Am J Clin Nutr* 1989;50:444-7.
41. Ott M, Fischer H, Polat H et al. Bioelectrical impedance analysis as a predictor of survival in patients with human immunodeficiency virus infection. *J Acquir Immune Defic Syndr Hum Retrovirol* 1995;9:20-5.
42. Mendenhall CL, Tosch T, Weesner RE et al. VA cooperative study on alcoholic hepatitis. II: Prognostic significance of protein-calorie malnutrition. *Am.J Clin.Nutr.* 1986;43:213-8.
43. Hedlund J. Community-acquired pneumonia requiring hospitalisation. Factors of importance for the short- and long term prognosis. *Scand.J.Infect.Dis.Supp* 1995;97:1-60.
44. Gariballa SE, Parker SG, Taub N, Castleden CM. Influence of nutritional status on clinical outcome after acute stroke. *Am J Clin Nutr* 1998;68:275-81.
45. Giner M, Laviano A, Meguid MM, Gleason JR. In 1995 a correlation between malnutrition and poor outcome in critically ill patients still exists. *Nutrition* 1996;12:23-9.
46. Fiatarone MA, O'Neill EF, Ryan ND et al. Exercise training and nutritional supplementation for physical frailty in very elderly people. *N Engl J Med* 1994;330:1769-75.
47. Delmi M, Rapin CH, Bengoa JM, Delmas PD, Vasey H, Bonjour JP. Dietary supplementation in elderly patients with fractured neck of the femur. *Lancet* 1990;335:1013-6.
48. Dempsey DT, Mullen JL, Buzby GP. The link between nutritional status and clinical outcome: can nutritional intervention modify it? *Am J Clin Nutr* 1988;47:352-6.
49. Müller J, Thul P, B. A. Perioperative Ernährungstherapie und ihre Relevanz für postoperative Ergebnisse. *Chirurg* 1997;574-82.
50. Vellas B, Baumgartner RN, Wayne SJ et al. Relationship between malnutrition and falls in the elderly. *Nutrition* 1992;8:105-8.
51. Vetta F, Ronzoni S, Taglieri G, Bollea MR. The impact of malnutrition on the quality of life in the elderly. *Clin Nutr* 1999;18:259-67.
52. Coats KG, Morgan SL, Bartolucci AA, Weinsier RL. Hospital-associated malnutrition: a reevaluation 12 years later. *J Am Diet Assoc* 1993;93:27-33.
53. McWhirter JP, Pennington CR. Incidence and recognition of malnutrition in hospital. *BMJ* 1994;308:945-8.
54. Naber TH, Schermer T, de Bree A et al. Prevalence of malnutrition in nonsurgical hospitalized patients and its association with disease complications. *Am.J Clin Nutr* 1997;66:1232-9.
55. Pirlich M, Schutz T, Kemps M et al. Prevalence of malnutrition in hospitalized medical patients: impact of underlying disease. *Dig Dis* 2003;21:245-51.
56. Robinson G, Goldstein M, Levine GM. Impact of nutritional status on DRG length of stay. *JPEN J Parenter.Enteral Nutr* 1987;11:49-51.

57. Reilly JJ, Jr., Hull SF, Albert N, Waller A, Bringardener S. Economic impact of malnutrition: a model system for hospitalized patients. *JPEN J Parenter. Enteral Nutr* 1988;12:371-6.
58. Tucker HN, Miguel SG. Cost containment through nutrition intervention. *Nutr Rev* 1996;54:111-21.
59. Weinsier RL, Hunker EM, Krumdieck CL, Butterworth CE, Jr. Hospital malnutrition. A prospective evaluation of general medical patients during the course of hospitalization. *Am.J Clin Nutr* 1979;32:418-26.
60. Bruun LI, Bosaeus I, Bergstad I, Nygaard K. Prevalence of malnutrition in surgical patients: evaluation of nutritional support and documentation. *Clin.Nutr.* 1999;18:141-7.
61. Barton AD, Beigg CL, Macdonald IA, Allison SP. High food wastage and low nutritional intakes in hospital patients. *Clin Nutr* 2000;19:445-9.
62. Barton AD, Beigg CL, Macdonald IA, Allison SP. A recipe for improving food intakes in elderly hospitalized patients. *Clin Nutr* 2000;19:451-4.
63. Blackburn GL, Bistrian BR, Maini BS, Schlamm HT, Smith MF. Nutritional and metabolic assessment of the hospitalized patient. *JPEN J Parenter. Enteral Nutr.* 1977;1:11-22.
64. DeJong PCM, Wesdorp RICO, Volovics A, et al. The value of objective measurements to select patients who are malnourished. *Clin.Nutr.* 1985;4:61-6.
65. Bergmann KE, Mensink GB. [Anthropometric data and obesity]. *Gesundheitswesen* 1999;61 Spec No:S115-S120.
66. Cunningham D, Pyrhonen S, James RD et al. Randomised trial of irinotecan plus supportive care versus supportive care alone after fluorouracil failure for patients with metastatic colorectal cancer. *Lancet* 1998;352:1413-8.
67. Herndon JE, Fleishman S, Kornblith AB, Kosty M, Green MR, Holland J. Is quality of life predictive of the survival of patients with advanced nonsmall cell lung carcinoma? *Cancer* 1999;85:333-40.
68. Detsky AS, McLaughlin JR, Baker JP et al. What is subjective global assessment of nutritional status? *JPEN J Parenter. Enteral Nutr* 1987;11:8-13.
69. Lochs H, Lübke H, Weimann A (Hrsg.). Leitlinie Enterale Ernährung. *Akt.Ernähr.-Med.* 2003; 28(S1): 1-120.
70. Waitzberg DL, Caiaffa WT, Correia MI. Hospital malnutrition: the Brazilian national survey (IBRANUTRI): a study of 4000 patients. *Nutrition* 2001;17:573-80.
71. Baker JP, Detsky AS, Wesson DE et al. Nutritional assessment: a comparison of clinical judgement and objective measurements. *N Engl J Med* 1982;306:969-72.
72. Lawson JA, Lazarus R, Kelly JJ. Prevalence and prognostic significance of malnutrition in chronic renal insufficiency. *J Ren Nutr.* 2001;11:16-22.
73. Persson C, Sjoden PO, Glimelius B. The Swedish version of the patient-generated subjective global assessment of nutritional status: gastrointestinal vs urological cancers. *Clin Nutr* 1999;18:71-7.
74. Sacks GS, Dearman K, Replogle WH, Cora VL, Meeks M, Canada T. Use of subjective global assessment to identify nutrition-associated complications and death in geriatric long-term care facility residents. *J.Am.Coll.Nutr.* 2000;19:570-7.
75. Buzby GP, Knox LS, Crosby LO et al. Study protocol: a randomized clinical trial of total parenteral nutrition in malnourished surgical patients. *Am J Clin Nutr* 1988;47:366-81.

76. Kyle UG, Pirlich M, Schütz T, Lochs H, Pichard C. Is nutritional depletion by Nutritional Risk Index (NRI) associated with increased length of hospital stay (LOS)? JPEN J Parenter Enteral Nutr 2004 (in press).
77. Kondrup J, Rasmussen HH, Hamberg O, Stanga Z. Nutritional risk screening (NRS 2002): a new method based on an analysis of controlled clinical trials. Clin.Nutr. 2003;22:321-36.
78. Lautz HU, Selberg O, Korber J, Burger M, Muller MJ. Protein-calorie malnutrition in liver cirrhosis. Clin Investig 1992;70:478-86.
79. Klein S. The myth of serum albumin as a measure of nutritional status. Gastroenterology 1990;99:1845-6.
80. Selberg O. The adjunctive value of routine biochemistry in nutritional assessment of hospitalized patients. Clin.Nutr. 2001;20:477-85.
81. Pirlich M, Schutz T, Spachos T et al. Bioelectrical impedance analysis is a useful bedside technique to assess malnutrition in cirrhotic patients with and without ascites. Hepatology 2000;32:1208-15.
82. Schwenk A, Beisenherz A, Kremer G, Diehl V, Salzberger B, Fatkenheuer G. Bioelectrical impedance analysis in HIV-infected patients treated with triple antiretroviral treatment. Am J Clin Nutr 1999;70:867-73.
83. Heymsfield SB, Wang Z, Visser M, Gallagher D, Pierson RN, Jr. Techniques used in the measurement of body composition: an overview with emphasis on bioelectrical impedance analysis. Am J Clin Nutr 1996;64:478S-84S.
84. Lukaski HC. Methods for the assessment of human body composition: traditional and new. Am J Clin Nutr 1987;46:537-56.
85. Heymsfield SB, Arteaga C, McManus C, Smith J, Moffitt S. Measurement of muscle mass in humans: validity of the 24-hour urinary creatinine method. Am.J Clin.Nutr. 1983;37:478-94.
86. Bistrian BR, Blackburn GL, Sherman M, Scrimshaw NS. Therapeutic index of nutritional depletion in hospitalized patients. Surg Gynecol.Obstet. 1975;141:512-6.
87. Bistrian BR. Nutritional assessment and therapy of protein--calorie malnutrition in the hospital. J.Am.Diet.Assoc. 1977;71:393-7.
88. Heymsfield SB, Waki M, Reinus J. Are patients with chronic liver disease hypermetabolic? Hepatology 1990;11:502-5.
89. Gurney JM, Jelliffe DB. Arm anthropometry in nutritional assessment: nomogram for rapid calculation of muscle circumference and cross-sectional muscle and fat areas. Am.J Clin Nutr 1973;26:912-5.
90. Heymsfield SB, McManus C, Smith J, Stevens V, Nixon DW. Anthropometric measurement of muscle mass: revised equations for calculating bone-free arm muscle area. Am.J Clin.Nutr. 1982;36:680-90.
91. Frisancho AR. New norms of upper limb fat and muscle areas for assessment of nutritional status. Am.J Clin Nutr 1981;34:2540-5.
92. Pirlich M, Plauth M, Lochs H. Bioelektrische Impedanzanalyse: Fehlerquellen und methodische Grenzen bei der klinischen Anwendung zur Analyse der Körperzusammensetzung. Akt.Ernähr.-Med. 1999;24:81-90.
93. Hermann L. Über eine Wirkung galvanischer Ströme auf Muskeln und Nerven. Pflügers Archiv für die gesamte Physiologie 1871;5:223-75.
94. Foster KR, Lukaski HC. Whole-body impedance--what does it measure? Am.J.Clin.Nutr. 1996;64:388S-96S.
95. Kushner RF. Bioelectrical impedance analysis: a review of principles and applications. J Am Coll Nutr 1992;11:199-209.

96. Selberg O, Selberg D. Norms and correlates of bioimpedance phase angle in healthy human subjects, hospitalized patients, and patients with liver cirrhosis. *Eur.J.Appl.Physiol* 2002;86:509-16.
97. Deurenberg P, Schouten FJ. Loss of total body water and extracellular water assessed by multifrequency impedance. *Eur J Clin Nutr* 1992;46:247-55.
98. Heath EM, Adams TD, Daines MM, Hunt SC. Bioelectric impedance and hydrostatic weighing with and without head submersion in persons who are morbidly obese. *J Am.Diet.Assoc.* 1998;98:869-75.
99. Bioelectrical Impedance Analysis in Body Composition Measurement. Proceedings of a National Institutes of Health Technology Assessment Conference. Bethesda, Maryland, December 12-14, 1994. *Am J Clin Nutr* 1996;64:387S-532S.
100. Plauth M, Weimann A, Holm E, Muller MJ. [Guidelines of GASL for nutrition in liver diseases and liver transplantation]. *Z.Gastroenterol.* 1999;37:301-12.
101. Plauth M, Merli M, Kondrup J, Ferenci P, Weimann A, Muller MJ. ESPEN guidelines for nutrition in liver disease and transplantation. *Clin.Nutr.* 1997;16:43-55.
102. Child CG, Turcotte JG. Surgery and portal hypertension. *Major.Probl.Clin.Surg.* 1964;1:1-85.
103. Pugh RN, Murray-Lyon IM, Dawson JL, Pietroni MC, Williams R. Transection of the oesophagus for bleeding oesophageal varices. *Br.J.Surg.* 1973;60:646-9.
104. Durazo-Arvizu RA, Goldbourt U, McGee DL. Body-mass index and mortality. *N Engl J Med.* 2000;342:287-9.
105. Waaler HT. Height, weight and mortality. The Norwegian experience. *Acta Med.Scand.Suppl* 1984;679:1-56.
106. Kyle UG, Pirlich M, Schuetz T, Luebke HJ, Lochs H, Pichard C. Prevalence of malnutrition in 1760 patients at hospital admission: a controlled population study of body composition. *Clin.Nutr.* 2003;22:473-81.
107. Pirlich M, Biering H, Gerl H et al. Loss of body cell mass in Cushing's syndrome: effect of treatment. *J Clin Endocrinol.Metab* 2002;87:1078-84.
108. Pirlich M, Schutz T, Gastell S, Lochs H. Malnutrition affects long-term prognosis in hospitalized patients. *Gastroenterology* 2003;122:A363.
109. Pirlich M, Schutz T, Ockenga J et al. Improved assessment of body cell mass by segmental bioimpedance analysis in malnourished subjects and acromegaly. *Clin Nutr* 2003;22:167-74.
110. Pichard C, Jeejeebhoy KN. Muscle dysfunction in malnourished patients. *Q J Med* 1988;69:1021-45.
111. Muller JM, Brenner U, Dienst C, Pichlmaier H. Preoperative parenteral feeding in patients with gastrointestinal carcinoma. *Lancet* 1982;1:68-71.
112. Muller MJ. Malnutrition in cirrhosis. *J.Hepatol.* 1995;23 Suppl 1:31-5.
113. Pirlich M, Selberg O, Boker K, Schwarze M, Muller MJ. The creatinine approach to estimate skeletal muscle mass in patients with cirrhosis. *Hepatology* 1996;24:1422-7.
114. Mitch WE. Robert H Herman Memorial Award in Clinical Nutrition Lecture, 1997. Mechanisms causing loss of lean body mass in kidney disease. *Am.J.Clin.Nutr.* 1998;67:359-66.
115. Plauth M, Schutz T, Buckendahl DP et al. Weight gain after transjugular intrahepatic portosystemic shunt is associated with improvement in body composition in malnourished patients with cirrhosis and hypermetabolism. *J.Hepatol.* 2004;40:228-33.

116. Lukaski HC. Applications of bioelectrical impedance analysis: a critical review. *Basic Life Sci* 1990;55:365-74.
117. NIH Conference. Bioelectrical impedance analysis in body composition measurement: National Institute of Health Technology Assessment Conference Statement. *Am.J Clin.Nutr.* 1996;64:524S-32S.
118. Guglielmi FW, Contento F, Laddaga L, Panella C, Francavilla A. Bioelectric impedance analysis: experience with male patients with cirrhosis. *Hepatology* 1991;13:892-5.
119. McCullough AJ, Mullen KD, Kalhan SC. Measurements of total body and extracellular water in cirrhotic patients with and without ascites. *Hepatology* 1991;14:1102-11.
120. Schloerb PR, Forster J, Delcore R, Kindscher JD. Bioelectrical impedance in the clinical evaluation of liver disease. *Am.J.Clin.Nutr.* 1996;64:510S-4S.
121. Zillikens MC, van den Berg JW, Wilson JH, Rietveld T, Swart GR. The validity of bioelectrical impedance analysis in estimating total body water in patients with cirrhosis. *J.Hepatol.* 1992;16:59-65.
122. Prijatmoko D, Strauss BJ, Lambert JR et al. Early detection of protein depletion in alcoholic cirrhosis: role of body composition analysis. *Gastroenterology* 1993;105:1839-45.
123. Moore FD, Boyden CM. Body cell mass and limits of hydration of the fat-free body: their relation to estimated skeletal weight. *Ann.N.Y.Acad.Sci.* 1963;110:62-71.
124. Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1986;1:307-10.
125. Rallison LR, Kushner RF, Penn D, Schoeller DA. Errors in estimating peritoneal fluid by bioelectrical impedance analysis and total body electrical conductivity. *J Am Coll Nutr* 1993;12:66-72.
126. Bistrian BR, Blackburn GL, Hallowell E, Heddle R. Protein status of general surgical patients. *JAMA* 1974;230:858-60.
127. Bistrian BR, Blackburn GL, Vitale J, Cochran D, Naylor J. Prevalence of malnutrition in general medical patients. *JAMA* 1976;235:1567-70.
128. Blackburn GL, Benotti PN, Bistrian BR et al. Nutritional assessment and treatment of hospital malnutrition. *Infusionsther.Klin.Ernahr.* 1979;6:238-50.
129. Roubenoff R, Roubenoff RA, Preto J, Balke CW. Malnutrition among hospitalized patients. A problem of physician awareness. *Arch Intern Med* 1987;147:1462-5.
130. Cederholm T, Jagren C, Hellstrom K. Outcome of protein-energy malnutrition in elderly medical patients. *Am.J Med* 1995;98:67-74.
131. Ferguson ML, Bauer J, Gallagher B, Capra S, Christie DR, Mason BR. Validation of a malnutrition screening tool for patients receiving radiotherapy. *Australas.Radiol.* 1999;43:325-7.
132. Edington J, Boorman J, Durrant ER et al. Prevalence of malnutrition on admission to four hospitals in England. The Malnutrition Prevalence Group. *Clin Nutr* 2000;19:191-5.
133. Pirlich M, Schütz T, Kemps M et al. Social risk factors for hospital malnutrition. *Nutrition* 2004 (in press).
134. Alberino F, Gatta A, Amodio P et al. Nutrition and survival in patients with liver cirrhosis. *Nutrition* 2001;17:445-50.
135. Merli M, Riggio O, Dally L. Does malnutrition affect survival in cirrhosis? PINC (Policentrica Italiana Nutrizione Cirrosi). *Hepatology* 1996;23:1041-6.

136. Moukarzel AA, Najm I, Vargas J, McDiarmid SV, Busuttil RW, Ament ME. Effect of nutritional status on outcome of orthotopic liver transplantation in pediatric patients. *Transplant Proc.* 1990;22:1560-3.
137. Pikul J, Sharpe MD, Lowndes R, Ghent CN. Degree of preoperative malnutrition is predictive of postoperative morbidity and mortality in liver transplant recipients. *Transplantation* 1994;57:469-72.
138. Ockenga J, Pirlisch M, Gastell S, Lochs H. [Tumour anorexia--tumour cachexia in case of gastrointestinal tumours: standards and visions]. *Z.Gastroenterol* 2002;40:929-36.
139. ASPEN Board of Directors and The Clinical Guidelines Task Force. Guidelines for the use of parenteral and enteral nutrition in adult and pediatric patients. *JPEN J Parenter. Enteral Nutr.* 2002;26:9SA-12SA.
140. DeWys WD, Begg C, Lavin PT, et al. Prognostic effect of weight loss prior to chemotherapy in cancer patients. *Am J Med* 1980;69:491-7.
141. Barber MD, Ross JA, Voss AC, Tisdale MJ, Fearon KC. The effect of an oral nutritional supplement enriched with fish oil on weight-loss in patients with pancreatic cancer. *Br.J.Cancer* 1999;81:80-6.
142. Barber MD, Fearon KC, Tisdale MJ, McMillan DC, Ross JA. Effect of a fish oil-enriched nutritional supplement on metabolic mediators in patients with pancreatic cancer cachexia. *Nutr.Cancer* 2001;40:118-24.
143. Tjonneland A, Gronbaek M, Stripp C, Overvad K. Wine intake and diet in a random sample of 48763 Danish men and women. *Am.J.Clin.Nutr.* 1999;69:49-54.