

## 5. Literature

- Alba, B.M. and Gross, C.A. (2004) Regulation of the E.coli sigma-dependent envelope stress response. *Mol Microbiol*, **3**, 613-619.
- Andersson, F.I., Blakytyn, R., Kirstein, J., Turgay, K., Bukau, B., Mogk, A. and Clarke, A.K. (2006) Cyanobacterial ClpC/HSP100 Protein Displays Intrinsic Chaperone Activity. *J Biol Chem*, **281**, 5468-5475.
- Becker, G., Klauck, E. and Hengge-Aronis, R. (1999) Regulation of RpoS proteolysis in Escherichia coli: the response regulator RssB is a recognition factor that interacts with the turnover element in RpoS. *Proc Natl Acad Sci U S A*, **96**, 6439-6444.
- Bochtler, M., Hartmann, C., Song, H.K., Bourenkov, G.P., Bartunik, H.D. and Huber, R. (2000) The structures of HsiU and the ATP-dependent protease HsiU-HsIV. *Nature*, **403**, 800-805.
- Bouche, S., Klauck, E., Fischer, D., Lucassen, M., Jung, K. and Hengge-Aronis, R. (1998) Regulation of RssB-dependent proteolysis in E. coli: a role for acetyl phosphate in a response regulator-controlled process. *Mol Microbiol*, **4**, 787-795.
- Bougdour, A., Wickner, S. and Gottesman, S. (2006) Modulating RssB activity: IraP, a novel regulator of sigma(S) stability in Escherichia coli. *Genes Dev*, **20**, 884-897.
- Brotz-Oesterhelt, H., Beyer, D., Kroll, H.P., Endermann, R., Ladel, C., Schroeder, W., Hinzen, B., Raddatz, S., Paulsen, H., Henninger, K., Bandow, J.E., Sahl, H.G. and Labischinski, H. (2005) Dysregulation of bacterial proteolytic machinery by a new class of antibiotics. *Nat Med*, **11**, 1082-1087.
- Derre, I., Rapoport, G. and Msadek, T. (1999) CtsR, a novel regulator of stress and heat shock response, controls clp and molecular chaperone gene expression in gram-positive bacteria. *Mol Microbiol*, **31**, 117-131.
- Deuerling, E., Patzelt, H., Vorderwulbecke, S., Rauch, T., Kramer, G., Schaffitzel, E., Mogk, A., Schulze-Specking, A., Langen, H. and Bukau, B. (2003) Trigger Factor and DnaK possess overlapping substrate pools and binding specificities. *Mol Microbiol*, **47**, 1317-1328.
- Dougan, D.A., Mogk, A., Zeth, K., Turgay, K. and Bukau, B. (2002a) AAA+ proteins and substrate recognition, it all depends on their partner in crime. *FEBS Lett*, **529**, 6-10.
- Dougan, D.A., Reid, B.G., Horwich, A.L. and Bukau, B. (2002b) ClpS, a substrate modulator of the ClpAP machine. *Mol Cell*, **9**, 673-683.
- Dougan, D.A., Weber-Ban, E. and Bukau, B. (2003) Targeted Delivery of an ssrA-Tagged Substrate by its Adaptor Protein SspB to Its Cognate AAA+ Protein ClpX. *Mol Cell*, **12**, 373-380.
- Dreveny, I., Kondo, H., Uchiyama, K., Shaw, A., Zhang, X. and Freemont, P.S. (2004) Structural basis of the interaction between the AAA ATPase p97/VCP and its adaptor protein p47. *Embo J*, **23**, 1030-1039.
- Dubnau, D. and Roggiani, M. (1990) Growth medium-independent genetic competence mutants of B. subtilis. *J Bacteriol*, **7**, 4048-4055.
- Erbse, A., Schmidt, R., Bornemann, T., Schneider-Mergener, J., Mogk, A., Zahn, R., Dougan, D.A. and Bukau, B. (2006) ClpS is an essential component of the N-end rule pathway in E. coli. *Nature*, **439**, 753-756.

- Flynn, J.M., Neher, S.B., Kim, Y.I., Sauer, R.T. and Baker, T.A. (2003) Proteomic discovery of cellular substrates of the ClpXP protease reveals five classes of ClpX-recognition signals. *Mol Cell*, **11**, 671-683.
- Gerth, U., Krüger, E., Derré, I., Msadek, T. and Hecker, M. (1998) Stress induction of the *Bacillus subtilis* *clpP* gene encoding a homologue of the proteolytic component of the Clp protease and the involvement of ClpP and ClpX in stress tolerance. *Mol. Microbiol.*, **28**, 787-802.
- Hattendorf, D.A. and Lindquist, S.L. (2002) Cooperative kinetics of both HSP104 ATPase domains and interdomain communication revealed by AAA sensor-1 mutants. *Embo J*, **1-2**, 12-21.
- Heinrich, J. and Wiegert, T. (2006) YpdC determines site-1 degradation in regulated intramembrane proteolysis of the RsiW anti-sigma factor of *Bacillus subtilis*. *Mol Microbiol*, **2**, 566-579.
- Hinnerwisch, J., Fenton, W.A., Furtak, K.J., Farr, G.W. and Horwich, A.L. (2005) Loops in the Central Channel of ClpA Chaperone Mediate Protein Binding, Unfolding, and Translocation. *Cell*, **121**, 1029-1041.
- Jürgen, B., Hanschke, R., Sarvas, M., Hecker, M. and Schweder, T. (2001) Proteome and transcriptome based analysis of *B. subtilis* cells overproducing an insoluble heterologous protein. *Appl Microbiol Biotechnol*, **3**, 326-332.
- Kang, S.G., Dimitrova, M.N., Ortega, J., Ginsburg, A. and Maurizi, M.R. (2005) Human mitochondrial ClpP is a stable heptamer that assembles into a tetradecamer in the presence of ClpX. *J Biol Chem*.
- Karata, K., Inagawa, T., Wilkinson, A.J., Tatsuta, T. and Ogura, T. (1999) Dissecting the role of a conserved motif (the second region of homology) in the AAA family of ATPases. Site directed mutagenesis of the ATP-dependent protease FtsH. *J Biol Chem*, **37**, 26225-26232.
- Kenniston, J.A., Baker, T.A., Fernandez, J.M. and Sauer, R.T. (2003) Linkage between ATP Consumption and Mechanical Unfolding during the Protein Processing Reactions of an AAA(+) Degradation Machine. *Cell*, **114**, 511-520.
- Kim, Y.I., Levchenko, I., Fraczkowska, K., Woodruff, R.V., Sauer, R.T. and Baker, T.A. (2001) Molecular determinants of complex formation between Clp/Hsp100 ATPases and the ClpP peptidase. *Nat Struct Biol*, **8**, 230-233.
- Kirstein, J., Dougan, D.A., Gerth, U., Hecker, M. and Turgay, K. (2007) The tyrosine kinase McsB is a regulated adaptor protein for ClpCP. *Embo J*, **8**, 2061-2070.
- Kirstein, J., Schlothauer, T., Dougan, D.A., Lilie, H., Tischendorf, G., Mogk, A., Bukau, B. and Turgay, K. (2006) Adaptor protein controlled oligomerization activates the AAA+ protein ClpC. *EMBO J*, **25**, 1481-1491.
- Kirstein, J., Zuhlke, D., Gerth, U., Turgay, K. and Hecker, M. (2005) A tyrosine kinase and its activator control the activity of the CtsR heat shock repressor in *B. subtilis*. *EMBO J*, **24**, 3435-3445.
- Kock, H., Gerth, U. and Hecker, M. (2004a) The ClpP peptidase is the major determinant of bulk protein turnover in *Bacillus subtilis*. *J Bacteriol*, **186**, 5856-5864.
- Kock, H., Gerth, U. and Hecker, M. (2004b) MurAA, catalysing the first committed step in peptidoglycan biosynthesis, is a target of Clp-dependent proteolysis in *Bacillus subtilis*. *Mol Microbiol*, **51**, 1087-1102.
- Krüger, E. and Hecker, M. (1998) The first gene of the *Bacillus subtilis* *clpC* operon, *ctsR*, encodes a negative regulator of its own operon and other class III heat shock genes. *J Bacteriol*, **180**, 6681-6688.

- Krüger, E., Witt, E., Ohlmeier, S., Hanschke, R. and Hecker, M. (2000) The Clp proteases in *B. subtilis* are directly involved in degradation of misfolded proteins. *J Bacteriol*, **11**, 3259-3265.
- Krüger, E., Zühlke, D., Witt, E., Ludwig, H. and Hecker, M. (2001) Clp-mediated proteolysis in Gram-positive bacteria is autoregulated by the stability of a repressor. *EMBO J*, **20**, 852-863.
- Levchenko, I., Grant, R.A., Flynn, J.M., Sauer, R.T. and Baker, T.A. (2005) Versatile modes of peptide recognition by the AAA+ adaptor protein SspB. *Nat Struct Mol Biol*, **12**, 520-525.
- Levchenko, I., Grant, R.A., Wah, D.A., Sauer, R.T. and Baker, T.A. (2003) Structure of a delivery protein for an AAA+ protease in complex with a peptide degradation tag. *Mol Cell*, **12**, 365-372.
- Levchenko, I., Luo, L. and Baker, T.A. (1995) Disassembly of the Mu transposase tetramer by the ClpX chaperone. *Genes Dev*, **9**, 2399-2408.
- Martin, A., Baker, T.A. and Sauer, R.T. (2005) Rebuilt AAA + motors reveal operating principles for ATP-fuelled machines. *Nature*, **437**, 1115-1120.
- Martin, A., Baker, T.A. and Sauer, R.T. (2007) Distinct static and dynamic interactions control ATPase-peptidase communication in an AAA+ protease. *Mol Cell*, **1**, 41-52.
- McGinness, K.E., Baker, T.A. and Sauer, R.T. (2006) Engineering controllable protein degradation. *Mol Cell*, **5**, 701-707.
- McGrath, P.T., Iniesta, A.A., Ryan, K.R., Shapiro, L. and McAdams, H.H. (2006) A dynamically localized protease complex and a polar specificity factor control a cell cycle master regulator. *Cell*, **124**, 535-547.
- Miethke, M., Hecker, M. and Gerth, U. (2006) Involvement of *Bacillus subtilis* ClpE in CtsR Degradation and Protein Quality Control. *J Bacteriol*, **188**, 4610-4619.
- Mika, F. and Hengge, R. (2005) A two-component phosphotransfer network involving ArcB, ArcA and RssB coordinates synthesis and proteolysis of RpoS in *E. coli*. *Genes Dev*, **22**, 2770-2781.
- Mogk, A., Schlieker, C., Strub, C., Rist, W., Weibezahn, J. and Bukau, B. (2003) Roles of individual domains and conserved motifs of the AAA+ chaperone ClpB in oligomerization, ATP hydrolysis, and chaperone activity. *J Biol Chem*, **20**, 17615-17624.
- Msadek, T., Dartois, V., Kunst, F., Herbaud, M.L., Denizot, F. and Rapoport, G. (1998) ClpP of *Bacillus subtilis* is required for competence development, motility, degradative enzyme synthesis, growth at high temperature and sporulation. *Mol Microbiol*, **27**, 899-914.
- Neher, S.B., Flynn, J.M., Sauer, R.T. and Baker, T.A. (2003a) Latent ClpX-recognition signals ensure LexA destruction after DNA damage. *Genes Dev*, **17**, 1084-1089.
- Neher, S.B., Sauer, R.T. and Baker, T.A. (2003b) Distinct peptide signals in the UmuD and UmuD' subunits of UmuD/D' mediate tethering and substrate processing by the ClpXP protease. *Proc Natl Acad Sci U S A*, **100**, 13219-13224.
- Neuwald, A.F., Aravind, L., Spouge, J.L. and Koonin, E.V. (1999) AAA+: A class of chaperone-like ATPases associated with the assembly, operation, and disassembly of protein complexes. *Genome Res*, **1**, 27-43.
- Ortega, J., Lee, H.S., Maurizi, M.R. and Steven, A.C. (2004) ClpA and ClpX ATPases bind simultaneously to opposite ends of ClpP peptidase to form active hybrid complexes. *J Struct Biol*, **146**, 217-226.

- Ortega, J., Singh, S.K., Ishikawa, T., Maurizi, M.R. and Steven, A.C. (2000) Visualization of Substrate Binding and Translocation by the ATP- Dependent Protease, ClpXP. *Mol Cell*, **6**, 1515-1521.
- Pan, Q., Garsin, D.A. and Losick, R. (2001) Self-Reinforcing Activation of a Cell-Specific Transcription Factor by Proteolysis of an Anti-sigma Factor in *B. subtilis*. *Mol Cell*, **8**, 873-883.
- Patel, S. and Latterich, M. (1998) The AAA team: related ATPases with diverse functions. *Trends Cell Biol*, **2**, 65-71.
- Persuh, M., Mandic-Mulec, I. and Dubnau, D. (2002) A MecA paralog, YpbH, binds ClpC, affecting both competence and sporulation. *J Bacteriol*, **184**, 2310-2313.
- Persuh, M., Turgay, K., Mandic-Mulec, I. and Dubnau, D. (1999) The N- and C-terminal domains of MecA recognize different partners in the competence molecular switch. *Mol Microbiol*, **33**, 886-894.
- Prepiak, P. and Dubnau, D. (2007) A peptide signal for adaptor protein-mediated degradation by the AAA+ protease ClpCP. *Mol Cell*, **5**, 639-647.
- Pruteanu, M. and Hengge-Aronis, R. (2002) The cellular level of the recognition factor RssB is rate-limiting for sigmaS proteolysis: implications for RssB regulation and signal transduction in sigmaS turnover in *Escherichia coli*. *Mol Microbiol*, **45**, 1701-1713.
- Reid, B.G., Fenton, W.A., Horwich, A.L. and Weber-Ban, E. (2001) ClpA mediates directional translocation of substrate proteins into the ClpP protease. *Proc Natl Acad Sci U S A*, **7**, 3768-3772.
- Ruvolo, M.V., Mach, K.E. and Burkholder, W.F. (2006) Proteolysis of the replication checkpoint protein Sda is necessary for the efficient initiation of sporulation after transient replication stress in *Bacillus subtilis*. *Molecular Microbiology*, **60**, 1490-1508.
- Sauer, R.T., Bolon, D.N., Burton, B.M., Burton, R.E., Flynn, J.M., Grant, R.A., Hersch, G.L., Joshi, S.A., Kenniston, J.A., Levchenko, I., Neher, S.B., Oakes, E.S., Siddiqui, S.M., Wah, D.A. and Baker, T.A. (2004) Sculpting the proteome with AAA(+) proteases and disassembly machines. *Cell*, **119**, 9-18.
- Schlieker, C., Weibezahn, J., Patzelt, H., Tessarz, P., Strub, C., Zeth, K., Erbse, A., Schneider-Mergener, J., Chin, J.W., Schultz, P.G., Bukau, B. and Mogk, A. (2004) Substrate recognition by the AAA+ chaperone ClpB. *Nat Struct Mol Biol*, **7**, 607-615.
- Schlothauer, T., Mogk, A., Dougan, D.A., Bukau, B. and Turgay, K. (2003) MecA, an adaptor protein necessary for ClpC chaperone activity. *Proc Natl Acad Sci U S A*, **100**, 2306.
- Schöbel, S., Zellmeier, S., Schumann, W. and Wiegert, T. (2004) The *Bacillus subtilis* sigmaW anti-sigma factor RsiW is degraded by intramembrane proteolysis through YluC. *Mol Microbiol*, **52**, 1091-1105.
- Siddiqui, S.M., Sauer, R.T. and Baker, T.A. (2004) Role of the processing pore of the ClpX AAA+ ATPase in the recognition and engagement of specific protein substrates. *Genes Dev*, **4**, 369-374.
- Singh, S.K. and Maurizi, M.R. (1994) Mutational analysis demonstrates different functional roles for the two ATP-binding sites in ClpAP protease from *E. coli*. *J Biol Chem*, **47**, 29537-29545.
- Stüdemann, A., Noirclerc-Savoye, M., Klauck, E., Becker, G., Schneider, D. and Hengge, R. (2003) Sequential recognition of two distinct sites in sigma(S) by the proteolytic targeting factor RssB and ClpX. *EMBO J*, **22**, 4111-4120.

- Thibault, G., Yudin, J., Wong, P., Tsitrin, V., Sprangers, R., Zhao, R. and Houry, W.A. (2006) Specificity in substrate and cofactor recognition by the N-terminal domain of the chaperone ClpX. *Proc Natl Acad Sci U S A*, **47**, 17724-17729.
- Thompson, M.W., Singh, S.K. and Maurizi, M.R. (1994) Processive degradation of proteins by the ATP-dependent Clp protease from *Escherichia coli*. Requirement for the multiple array of active sites in ClpP but not ATP hydrolysis. *J Biol Chem*, **269**, 18209-18215.
- Tobias, J.W., Shrader, T.E., Rocap, G. and Varshavsky, A. (1991) The N-end rule in bacteria. *Science*, **254**, 1374-1377.
- Turgay, K., Hahn, J., Burghoorn, J. and Dubnau, D. (1998) Competence in *Bacillus subtilis* is controlled by regulated proteolysis of a transcription factor. *EMBO J*, **17**, 6730-6738.
- Ullers, R.S., Luirink, J., Harms, N., Schwager, F., Georgopolulos, C. and Genevaux, P. (2004) SecB is a bona fide generalized chaperone in *E. coli*. *Proc Natl Acad Sci U S A*, **20**, 7583-7588.
- Wah, D.A., Levchenko, I., Rieckhof, G.E., Bolon, D.N., Baker, T.A. and Sauer, R.T. (2003) Flexible linkers leash the substrate binding domain of SspB to a peptide module that stabilizes delivery complexes with the AAA+ ClpXP protease. *Mol Cell*, **12**, 355-363.
- Wang, J., Hartling, J.A. and Flanagan, J.M. (1998) Crystal structure determination of *E. coli* ClpP starting from an EM-derived mask. *J Struct Biol*, **2-3**, 151-163.
- Wang, K.H., Sauer, R.T. and Baker, T.A. (2007) ClpS modulates but is not essential for bacterial N-end rule degradation. *Genes Dev*, **4**, 403-408.
- Watanabe, Y.H., Takano, M. and Yoshida, M. (2005) ATP binding to nucleotide binding domain (NBD)1 of the ClpB chaperone induces motion of the long coiled-coil, stabilizes the hexamer, and activates NBD2. *J Biol Chem*, **26**, 24562-24567.
- Weibezahn, J., Bukau, B. and Mogk, A. (2004) Unscrambling an egg: protein disaggregation by AAA+ proteins. *Microb Cell Fact*, **3**, 1.
- Wickner, S., Maurizi, M.R. and Gottesman, S. (1999) Posttranslational quality control: folding, refolding, and degrading proteins. *Science*, **286**, 1888-1893.
- Yousef, M.S., Fabiola, F., Gattis, J.L., Somasundaram, T. and Chapman, M.S. (2002) Refinement of the arginine kinase transition-state analogue complex at 1.2 Å resolution: mechanistic insights. *Acta Crystallogr D Biol Crystallogr*, **12**, 2009-2017.
- Yu, A.Y. and Houry, W.A. (2007) ClpP: A distinctive family of cylindrical energy-dependent serine proteases. *FEBS Lett*.
- Zeth, K., Ravelli, R.B., Paal, K., Cusack, S., Bukau, B. and Dougan, D.A. (2002) Structural analysis of the adaptor protein ClpS in complex with the N-terminal domain of ClpA. *Nat Struct Biol*, **9**, 906-911.
- Zhou, G., Somasundaram, T., Blanc, E., Parthasarathy, G., Ellington, W.R. and Chapman, M.S. (1998) Transition state structure of arginine kinase: implications for catalysis of bimolecular reactions. *Proc Natl Acad Sci U S A*, **15**, 8449-8454.
- Zhou, Y. and Gottesmann, S. (1998) Regulation of the proteolysis of the stationary-phase sigma factor RpoS. *J Bacteriol*, **5**, 1154-1158.

