

7. References

- Albert, A. P. & Large, W. A. (2001). The effect of external divalent cations on spontaneous non-selective cation channel currents in rabbit portal vein myocytes. *J Physiol* **536**, 409-420.
- Albert, A. P. & Large, W. A. (2002). A Ca^{2+} -permeable non-selective cation channel activated by depletion of internal Ca^{2+} stores in single rabbit portal vein myocytes. *J Physiol* **538**, 717-728.
- Aromolaran, A. S. & Large, W. A. (1999). Comparison of the effects of divalent cations on the noradrenaline-evoked cation current in rabbit portal vein smooth muscle cells. *J Physiol* **520 Pt 3**, 771-782.
- Asano, M., Nakajima, T., Hazama, H., Iwasawa, K., Tomaru, T., Omata, M., Soma, M., Asakura, Y., Mizutani, M., Suzuki, S., Yamashita, K. & Okuda, Y. (1998). Influence of cellular incorporation of n-3 eicosapentaenoic acid on intracellular Ca^{2+} concentration and membrane potential in vascular smooth muscle cells. *Atherosclerosis* **138**, 117-127.
- Asano, M., Nakajima, T., Iwasawa, K., Hazama, H., Omata, M., Soma, M., Yamashita, K. & Okuda, Y. (1997). Inhibitory effects of ω -3 polyunsaturated fatty acids on receptor-mediated non-selective cation currents in rat A7r5 vascular smooth muscle cells. *Br J Pharmacol* **120**, 1367-1375.
- Barritt, G. J. (1999). Receptor-activated Ca^{2+} inflow in animal cells: a variety of pathways tailored to meet different intracellular Ca^{2+} signalling requirements. *Biochem. J.* **337**, 153-169.
- Benham, C. D., Hess, P. & Tsien, R. W. (1987). Two types of calcium channels in single smooth muscle cells from rabbit ear artery studied with whole-cell and single-channel recordings. *Circ Res* **61**, 110-16.
- Berchtold, M. W., Brinkmeier, H. & Muntener, M. (2000). Calcium ion in skeletal muscle: its crucial role for muscle function, plasticity, and disease. *Physiol Rev* **80**, 1215-1265.
- Berridge, M. J. (1993). Inositol trisphosphate and calcium signalling. *Nature* **361**, 315-325.
- Berridge, M. J., Lipp, P. & Bootman, M. D. (2000). The versatility and universality of calcium signalling. *Nat Rev Mol Cell Biol* **1**, 11-21.
- Bigay, J., Deterre, P., Pfister, C. & Chabre, M. (1985). Fluoroaluminates activate transducin-GDP by mimicking the γ -phosphate of GTP in its binding site. *FEBS Lett* **191**, 181-185.
- Bigay, J., Deterre, P., Pfister, C. & Chabre, M. (1987). Fluoride complexes of aluminium or beryllium act on G-proteins as reversibly bound analogues of the γ phosphate of GTP. *EMBO J* **6**, 2907-2913.
- Blaustein, M. P. & Lederer, W. J. (1999). Sodium/calcium exchange: its physiological implications. *Physiol Rev* **79**, 763-854.
- Bolton, T. B., Prestwich, S. A., Zholos, A. V. & Gordienko, D. V. (1999). Excitation-contraction coupling in gastrointestinal and other smooth muscles. *Annu Rev Physiol* **61**, 85-115.
- Boulay, G., Zhu, X., Peyton, M., Jiang, M., Hurst, R., Stefani, E. & Birnbaumer, L. (1997). Cloning and expression of a novel mammalian homologue of Drosophila transient receptor potential (TRP) involved in calcium entry secondary to activation of receptors coupled by the G_q class of G protein. *J. Biol. Chem.* **272**, 29672-29680.
- Broad, L. M., Cannon, T. R. & Taylor, C. W. (1999). A non-capacitative pathway activated by arachidonic acid is the major Ca^{2+} entry mechanism in rat A7r5 smooth muscle cells stimulated with low concentrations of vasopressin. *J Physiol* **517**, 121-134.
- Brock, J. A., McLachlan, E. M. & Rayner, S. E. (1997). Contribution of α -adrenoceptors to depolarization and contraction evoked by continuous asynchronous sympathetic nerve activity in rat tail artery. *Br J Pharmacol* **120**, 1513-1521.

- Byron, K. & Taylor, C. W. (1995). Vasopressin stimulation of Ca^{2+} mobilization, two bivalent cation entry pathways and Ca^{2+} efflux in A7r5 rat smooth muscle cells. *J Physiol* **485**, 455-468.
- Byron, K. L. (1996). Vasopressin stimulates Ca^{2+} spiking activity in A7r5 vascular smooth muscle cells via activation of phospholipase A_2 . *Circ Res* **78**, 813-820.
- Byron, K. L. & Lucchesi, P. A. (2002). Signal transduction of physiological concentrations of vasopressin in A7r5 vascular smooth muscle cells. A role for PYK2 and tyrosine phosphorylation of K^+ channels in the stimulation of Ca^{2+} spiking. *J Biol Chem* **277**, 7298-7307.
- Byron, K. L. & Taylor, C. W. (1993). Spontaneous Ca^{2+} spiking in a vascular smooth muscle cell line is independent of the release of intracellular Ca^{2+} stores. *J Biol Chem* **268**, 6945-6952.
- Carafoli, E. (1994). Biogenesis: plasma membrane calcium ATPase: 15 years of work on the purified enzyme. *Faseb J* **8**, 993-1002.
- Caterina, M., Schumacher, M. A., Tominaga, M., Rosen, T. A., Levine, J. D. & Julius, D. (1997). The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* **389**, 816-824.
- Caterina, M. J., Rosen, T. A., Tominaga, M., Brake, A. J. & Julius, D. (1999). A capsaicin receptor homologue with a high threshold for noxious heat. *Nature* **398**, 436-441.
- Chang, A. S., Chang, S. M., Garcia, R. L. & Schilling, W. P. (1997). Concomitant and hormonally regulated expression of *trp* genes in bovine aortic endothelial cells. *FEBS Letters* **415**, 335-340.
- Chen, S., Inoue, R. & Ito, Y. (1993). Pharmacological characterization of muscarinic receptor-activated cation channels in guinea-pig ileum. *Br J Pharmacol* **109**, 793-801.
- Chen, X. L. & Rembold, C. M. (1995). Phenylephrine contracts rat tail artery by one electromechanical and three pharmacomechanical mechanisms. *Am J Physiol* **268**, H74-81.
- Chin, D. & Means, A. R. (2000). Calmodulin: a prototypical calcium sensor. *Trends Cell Biol* **10**, 322-328.
- Chyb, S., Raghunath, P. & Hardie, R. C. (1999). Polyunsaturated fatty acids activate light-sensitive channels TRP and TRPL. *Nature* **397**, 255-259.
- Clapham, D. E. (1995). Calcium signaling. *Cell* **80**, 259-268.
- Clapham, D. E., Runnels, L. W. & Strübing, C. (2001). The TRP ion channel family. *Nat Rev Neurosci* **2**, 387-396.
- Clementi, E. & Meldolesi, J. (1996). Pharmacological and functional properties of voltage-independent Ca^{2+} channels. *Cell Calcium* **19**, 269-279.
- Clementi, E., Scheer, H., Zacchetti, D., Fasolato, C., Pozzan, T. & Meldolesi, J. (1992). Receptor-activated Ca^{2+} influx. Two independently regulated mechanisms of influx stimulation coexist in neurosecretory PC12 cells. *J Biol Chem* **267**, 2164-2172.
- Coronado, R., Morrissette, J., Sukhareva, M. & Vaughan, D. M. (1994). Structure and function of ryanodine receptors. *Am J Physiol* **266**, C1485-1504.
- Cosens, D. J. & Manning, A. (1969). Abnormal electroretinogram from a *Drosophila* mutant. *Nature* **224**, 285-287.
- Davis, M. J. & Hill, M. A. (1999). Signaling mechanisms underlying the vascular myogenic response. *Physiol Rev* **79**, 387-423.
- Delmas, P., Wanaverbecq, N., Abogadie, F. C., Mistry, M. & Brown, D. A. (2002). Signaling microdomains define the specificity of receptor-mediated InsP_3 pathways in neurons. *Neuron* **34**, 209-220.
- Duncan, L. M., Deeds, J., Hunter, J., Shao, J., Holmgren, L. M., Woolf, E. A., Tepper, R. L. & Shyjan, A. W. (1998). Down-regulation of the novel gene melastatin correlates with potential for melanoma metastasis. *Cancer Res.* **58**, 1515-1520.
- Ebashi, S. (1972). Calcium ions and muscle contraction. *Nature* **240**, 217-218.
- Elliott, A. C. (2001). Recent developments in non-excitable cell calcium entry. *Cell Calcium* **30**, 73-93.

- Enklaar, T., Esswein, M., Oswald, M., Hilbert, K., Winterpacht, A., Higgins, M., Zabel, B. & Prawitt, D. (2000). Mtr1, a novel biallelically expressed gene in the center of the mouse distal chromosome 7 imprinting cluster, is a member of the Trp gene family. *Genomics* **67**, 179-187.
- Fan, J. & Byron, K. L. (2000). Ca²⁺ signalling in rat vascular smooth muscle cells: a role for protein kinase C at physiological vasoconstrictor concentrations of vasopressin. *J Physiol* **524 Pt 3**, 821-831.
- Fang, D. & Setaluri, V. (2000). Expression and Up-regulation of alternatively spliced transcripts of melastatin, a melanoma metastasis-related gene, in human melanoma cells. *Biochem Biophys Res Commun* **279**, 53-61.
- Fasolato, C., Innocenti, B. & Pozzan, T. (1994). Receptor-activated Ca²⁺ influx: How many mechanisms for how many channels. *Trends Pharmacol. Sci.* **15**, 77-83.
- Fasolato, C. & Nilius, B. (1998). Store depletion triggers the calcium release-activated calcium current (I_{CRAC}) in macrovascular endothelial cells: a comparison with Jurkat and embryonic kidney cell lines. *Pflugers Arch* **436**, 69-74.
- Felder, C. C., Poulter, M. O. & Wess, J. (1992). Muscarinic receptor-operated Ca²⁺ influx in transfected fibroblast cells is independent of inositol phosphates and release of intracellular Ca²⁺. *Proc Natl Acad Sci U S A* **89**, 509-513.
- Fenwick, E. M., Marty, A. & Neher, E. (1982). Sodium and calcium channels in bovine chromaffin cells. *J Physiol* **331**, 599-635.
- Franke, W. W., Schmid, E., Kartenbeck, J., Meyer, D., Hacker, H.-J., Bannasch, P., Osborn, M., Weber, K., Denk, H., Wanson, J. C. & Drochmans, P. (1979). Characterization of the intermediate sized filaments in liver cells by immunofluorescence and electron microscopy. *Biol Cell* **34**, 99-110.
- Franke, W. W., Schmid, E., Vandekerckhove, J. & Weber, K. (1980). Permanently proliferating rat vascular smooth muscle cell with maintained expression of smooth muscle characteristics, including actin of the vascular smooth muscle type. *J Cell Biol* **87**, 594-600.
- Franzini-Armstrong, C. & Protasi, F. (1997). Ryanodine receptors of striated muscles: a complex channel capable of multiple interactions. *Physiol Rev* **77**, 699-729.
- Freichel, M., Suh, S. H., Pfeifer, A., Schweig, U., Trost, C., Weissgerber, P., Biel, M., Philipp, S., Freise, D., Droogmans, G., Hofmann, F., Flockerzi, V. & Nilius, B. (2001). Lack of an endothelial store-operated Ca²⁺ current impairs agonist-dependent vasorelaxation in TRP4^{-/-} mice. *Nat Cell Biol* **3**, 121-127.
- Funayama, M., Goto, K. & Kondo, H. (1996). Cloning and expression localization of cDNA for rat homolog of TRP protein, a possible store-operated calcium (Ca²⁺) channel. *Brain Res Mol Brain Res* **43**, 259-266.
- Furuichi, T., Kohda, K., Miyawaki, A. & Mikoshiba, K. (1994). Intracellular channels. *Curr Opin Neurobiol* **4**, 294-303.
- Galione, A., Lee, H. C. & Busa, W. B. (1991). Ca²⁺-induced Ca²⁺ release in sea urchin egg homogenates: modulation by cyclic ADP-ribose. *Science* **253**, 1143-1146.
- Galvan, D. L., Borrego-Diaz, E., Perez, P. J. & Mignery, G. A. (1999). Subunit oligomerization, and topology of the inositol 1,4,5-trisphosphate receptor. *J Biol Chem* **274**, 29483-29492.
- Ganitkevich, V. & Isenberg, G. (1990). Isolated guinea pig coronary smooth muscle cells. Acetylcholine induces hyperpolarization due to sarcoplasmic reticulum calcium release activating potassium channels. *Circ Res* **67**, 525-528.
- Garcia, R. L. & Schilling, W. P. (1997). Differential expression of mammalian TRP homologues across tissues and cell lines. *Biochemical and Biophysical Research Communications* **239**, 279-283.
- Gardner, J. D. & Benoît, J. N. (2000). Effects of capacitative calcium entry on agonist-induced calcium transients in A7r5 vascular smooth muscle cells. *J Biomed Sci* **7**, 304-310.
- Genazzani, A. A. & Billington, R. A. (2002). NAADP: an atypical Ca²⁺-release messenger? *Trends Pharmacol Sci* **23**, 165-167.

- Gibson, A., McFadzean, I., Wallace, P. & Wayman, C. P. (1998). Capacitative Ca^{2+} entry and the regulation of smooth muscle tone. *Trends Pharmacol Sci* **19**, 266-269.
- Gillo, B., Chorna, I., Cohen, H., Cook, B., Manistersky, I., Chorev, M., Arnon, A., Pollock, J. A., Selinger, Z. & Minke, B. (1996). Coexpression of *Drosophila* TRP and TRP-like proteins in *Xenopus* oocytes reconstitutes capacitative Ca^{2+} entry. *Proc Natl Acad Sci U S A* **93**, 14146-14151.
- Golovina, V. A., Platoshyn, O., Bailey, C. L., Wang, J., Limsuwan, A., Sweeney, M., Rubin, L. J. & Yuan, J. X. (2001). Upregulated TRP and enhanced capacitative Ca^{2+} entry in human pulmonary artery myocytes during proliferation. *Am J Physiol Heart Circ Physiol* **280**, H746-755.
- Goto, K., Kasuya, Y., Matsuki, N., Takuwa, Y., Kurihara, H., Ishikawa, T., Kimura, S., Yanagisawa, M. & Masaki, T. (1989). Endothelin activates the dihydropyridine-sensitive, voltage-dependent Ca^{2+} channel in vascular smooth muscle. *Proc Natl Acad Sci U S A* **86**, 3915-3918.
- Groschner, K., Hingel, S., Lintschinger, B., Balzer, M., Romanin, C., Zhu, X. & Schreibmayer, W. (1998). Trp proteins form store-operated channels in human vascular endothelial cells. *FEBS Lett.* **437**, 101-106.
- Grynkiewicz, G., Poenie, M. & Tsien, R. Y. (1985). A new generation of Ca^{2+} indicators with greatly improved fluorescence properties. *J Biol Chem* **260**, 3440-3450.
- Guibert, C. & Beech, D. J. (1999). Positive and negative coupling of the endothelin ET_A receptor to Ca^{2+} -permeable channels in rabbit cerebral cortex arterioles. *J Physiol* **514**, 843-856.
- Gunthorpe, M. J., Benham, C. D., Randall, A. & Davis, J. B. (2002). The diversity in the vanilloid (TRPV) receptor family of ion channels. *Trends Pharmacol Sci* **23**, 183-191.
- Halaszovich, C. R., Zitt, C., Jüngling, E. & Lückhoff, A. (2000). Inhibition of TRP3 channels by lanthanides. Block from the cytosolic side of the plasma membrane. *J Biol Chem* **275**, 37423-37428.
- Hamill, O. P., Marty, A., Neher, E., Sakmann, B. & Sigworth, F. J. (1981). Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflugers Arch* **391**, 85-100.
- Hara, Y., Wakamori, M., Ishii, M., Maeno, E., Nishida, M., Yoshida, T., Yamada, H., Shimizu, S., Mori, E., Kudoh, J., Shimizu, N., Kurose, H., Okada, Y., Imoto, K. & Mori, Y. (2002). LTRPC2 Ca^{2+} -permeable channel activated by changes in redox status confers susceptibility to cell death. *Mol Cell* **9**, 163-173.
- Hardie, R. C. & Minke, B. (1992). The *trp* gene is essential for a light-activated Ca^{2+} channel in *Drosophila* photoreceptors. *Neuron* **8**, 643-651.
- Hardie, R. C. & Minke, B. (1995). Phosphoinositide-mediated phototransduction in *Drosophila* photoreceptors: the role of Ca^{2+} and *trp*. *Cell Calcium* **18**, 256-274.
- Harteneck, C., Plant, T. D. & Schultz, G. (2000). From worm to man: three subfamilies of TRP channels. *Trends Neurosci* **23**, 159-166.
- Helliwell, R. M. & Large, W. A. (1996). Dual effect of external Ca^{2+} on noradrenaline-activated cation current in rabbit portal vein smooth muscle cells. *J Physiol* **492**, 75-88.
- Helliwell, R. M. & Large, W. A. (1997). α_1 -adrenoceptor activation of a non-selective cation current in rabbit portal vein by 1,2-diacyl-*sn*-glycerol. *J Physiol* **499**, 417-428.
- Helliwell, R. M. & Large, W. A. (1998). Facilitatory effect of Ca^{2+} on the noradrenaline-evoked cation current in rabbit portal vein smooth muscle cells. *J Physiol* **512**, 731-741.
- Hill, M. A., Zou, H., Potocnik, S. J., Meininger, G. A. & Davis, M. J. (2001). Invited review: arteriolar smooth muscle mechanotransduction: Ca^{2+} signaling pathways underlying myogenic reactivity. *J Appl Physiol* **91**, 973-983.
- Hoenderop, J. G. J., van der Kemp, A. W. C. M., Hartog, A., van de Graaf, S. F. J., van Os, C. H., Willems, P. H. G. M. & Bindels, R. J. M. (1999). Molecular identification of the apical

- Ca²⁺ channel in 1,25-dihydroxyvitamin D₃-responsive epithelia. *J. Biol. Chem.* **274**, 8375-8378.
- Hofmann, T., Obukhov, A. G., Schaefer, M., Harteneck, C., Gudermann, T. & Schultz, G. (1999). Direct activation of human TRPC6 and TRPC3 channels by diacylglycerol. *Nature* **397**, 259-263.
- Hofmann, T., Schaefer, M., Schultz, G. & Gudermann, T. (2000). Cloning, expression and subcellular localization of two novel splice variants of mouse transient receptor potential channel 2. *Biochem J* **351**, 115-122.
- Hofmann, T., Schaefer, M., Schultz, G. & Gudermann, T. (2002). Subunit composition of mammalian transient receptor potential channels in living cells. *Proc Natl Acad Sci U S A* **99**, 7461-7466.
- Hoth, M. & Penner, R. (1992). Depletion of intracellular calcium stores activates a calcium current in mast cells. *Nature* **355**, 353-355.
- Hu, Y., Vaca, L., Zhu, X., Birnbaumer, L., Kunze, D. L. & Schilling, W. P. (1994). Appearance of a novel Ca²⁺ influx pathway in Sf9 insect cells following expression of the transient receptor potential-like (*trpl*) protein of *Drosophila*. *Biochem Biophys Res Commun* **201**, 1050-1056.
- Hurst, R. S., Zhu, X., Boulay, G., Birnbaumer, L. & Stefani, E. (1998). Ionic currents underlying TRP3 mediated agonist-dependent Ca²⁺ influx in stably transfected HEK293 cells. *FEBS Lett.* **422**, 333-338.
- Hussain, A. & Inesi, G. (1999). Involvement of Sarco/endoplasmic reticulum Ca²⁺ ATPases in cell function and the cellular consequences of their inhibition. *J Membr Biol* **172**, 91-99.
- Hwang, S. W., Cho, H., Kwak, J., Lee, S. Y., Kang, C. J., Jung, J., Cho, S., Min, K. H., Suh, Y. G., Kim, D. & Oh, U. (2000). Direct activation of capsaicin receptors by products of lipoxygenases: endogenous capsaicin-like substances. *Proc Natl Acad Sci U S A* **97**, 6155-6160.
- Inoue, R., Morita, H., Yanagida, H. & Ito, Y. (1998). Potentiating actions of lanthanum on ACh-induced cation current in guinea-pig ileal smooth muscle cells. *J Smooth Muscle Res* **34**, 69-81.
- Inoue, R., Okada, T., Onoue, H., Hara, Y., Shimizu, S., Naitoh, S., Ito, Y. & Mori, Y. (2001). The transient receptor potential protein homologue TRP6 is the essential component of vascular α_1 -adrenoceptor-activated Ca²⁺-permeable cation channel. *Circ Res* **88**, 325-332.
- Irvine, R. F. (1990). 'Quantal' Ca²⁺ release and the control of Ca²⁺ entry by inositol phosphates--a possible mechanism. *FEBS Lett* **263**, 5-9.
- Ives, H. E., Schultz, G. S., Galaray, R. E. & Jamieson, J. D. (1978). Preparation of functional smooth muscle cells from the rabbit aorta. *J Exp Med* **148**, 1400-1413.
- Iwamuro, Y., Miwa, S., Minowa, T., Enoki, T., Zhang, X. F., Ishikawa, M., Hashimoto, N. & Masaki, T. (1998). Activation of two types of Ca²⁺-permeable nonselective cation channel by endothelin-1 in A7r5 cells. *Br J Pharmacol* **124**, 1541-1549.
- Iwasawa, K., Nakajima, T., Hazama, H., Goto, A., Shin, W. S., Toyo-oka, T. & Omata, M. (1997). Effects of extracellular pH on receptor-mediated Ca²⁺ influx in A7r5 rat smooth muscle cells: involvement of two different types of channel. *J Physiol* **503**, 237-251.
- Jackson, W. F. (2000). Ion channels and vascular tone. *Hypertension* **35**, 173-178.
- Johnson, S., Michalak, M., Opas, M. & Eggleton, P. (2001). The ins and outs of calreticulin: from the ER lumen to the extracellular space. *Trends Cell Biol* **11**, 122-129.
- Jordt, S. E. & Julius, D. (2002). Molecular basis for species-specific sensitivity to "hot" chili peppers. *Cell* **108**, 421-430.
- Jordt, S. E., Tominaga, M. & Julius, D. (2000). Acid potentiation of the capsaicin receptor determined by a key extracellular site. *Proc Natl Acad Sci U S A* **97**, 8134-8139.
- Julius, D. & Basbaum, A. I. (2001). Molecular mechanisms of nociception. *Nature* **413**, 203-210.

- Jungnickel, M. K., Marrero, H., Birnbaumer, L., Lemos, J. R. & Florman, H. M. (2001). Trp2 regulates entry of Ca^{2+} into mouse sperm triggered by egg ZP3. *Nat Cell Biol* **3**, 499-502.
- Kahn, R. A. (1991). Fluoride is not an activator of the smaller (20-25 kDa) GTP-binding proteins. *J Biol Chem* **266**, 15595-15597.
- Kamouchi, M., Phillip, S., Flockerzi, V., Wissenbach, U., Mamin, A., Raeymaekers, L., Eggermont, J., Droogmans, G. & Nilius, B. (1999). Properties of heterologously expressed hTRP3 channels in bovine pulmonary artery endothelial cells. *J. Physiol.* **518**, 345-358.
- Kanki, H., Kinoshita, M., Akaike, A., Satoh, M., Mori, Y. & Kaneko, S. (2001). Activation of inositol 1,4,5-trisphosphate receptor is essential for the opening of mouse TRP5 channels. *Mol Pharmacol* **60**, 989-998.
- Kanzaki, M., Zhang, Y. Q., Mashima, H., Li, L., Shibata, H. & Kojima, I. (1999). Translocation of a calcium-permeable cation channel induced by insulin-like growth factor-I. *Nat Cell Biol* **1**, 165-170.
- Kaplan-Albuquerque, N. & Di Salvo, J. (1998). Protein kinase C: modulation of vasopressin-induced calcium influx and release in A7r5 vascular smooth muscle cells. *Arch Biochem Biophys* **359**, 209-214.
- Karaki, H., Ozaki, H., Hori, M., Mitsui-Saito, M., Amano, K., Harada, K., Miyamoto, S., Nakazawa, H., Won, K. J. & Sato, K. (1997). Calcium movements, distribution, and functions in smooth muscle. *Pharmacol Rev* **49**, 157-230.
- Kawanabe, Y., Okamoto, Y., Hashimoto, N. & Masaki, T. (2001). Characterization of Ca^{2+} channels involved in endothelin-1-induced mitogenic responses in vascular smooth muscle cells. *Eur J Pharmacol* **422**, 15-21.
- Kedei, N., Szabo, T., Lile, J. D., Treanor, J. J., Olah, Z., Iadarola, M. J. & Blumberg, P. M. (2001). Analysis of the native quaternary structure of vanilloid receptor 1. *J Biol Chem* **276**, 28613-28619.
- Kimes, B. W. & Brandt, B. L. (1976). Properties of a clonal muscle cell line from rat heart. *Exp Cell Res* **98**, 367-381.
- Kinoshita, M., Akaike, A., Satoh, M. & Kaneko, S. (2000). Positive regulation of capacitative Ca^{2+} entry by intracellular Ca^{2+} in *Xenopus* oocytes expressing rat TRP4. *Cell Calcium* **28**, 151-159.
- Kiselyov, K., Mignery, G. A., Zhu, M. X. & Muallem, S. (1999). The N-terminal domain of the IP_3 receptor gates store-operated hTrp3 channels. *Mol Cell* **4**, 423-429.
- Kiselyov, K. & Muallem, S. (1999). Fatty acids, diacylglycerol, $\text{Ins}(1, 4, 5)\text{P}_3$ receptors and Ca^{2+} influx. *Trends Neurosci.* **22**, 334-337.
- Kiselyov, K., Xu, X., Mozhayeva, G., Kuo, T., Pessah, I., Mignery, G., Zhu, X., Birnbaumer, L. & Muallem, S. (1998). Functional interaction between InsP_3 receptors and store-operated channels. *Nature* **396**, 478-482.
- Kiselyov, K. I., Shin, D. M., Wang, Y., Pessah, I. N., Allen, P. D. & Muallem, S. (2000). Gating of store-operated channels by conformational coupling to ryanodine receptors. *Mol Cell* **6**, 421-431.
- Koch, J. & Korbmayer, C. (1999). Osmotic shrinkage activates nonselective cation (NSC) channels in various cell types. *J Membr Biol* **168**, 131-139.
- Krautwurst, D., Degtiar, V. E., Schultz, G. & Hescheler, J. (1994). The isoquinoline derivative LOE 908 selectively blocks vasopressin-activated nonselective cation currents in A7r5 aortic smooth muscle cells. *Naunyn Schmiedebergs Arch Pharmacol* **349**, 301-307.
- Kuriyama, H., Kitamura, K., Itoh, T. & Inoue, R. (1998). Physiological features of visceral smooth muscle cells, with special reference to receptors and ion channels. *Physiol Rev* **78**, 811-920.
- Lapidot, S. A., Huang, B. K., Fayazi, A., Russek, L. N., Strickberger, S. A., Brooks, A. E. & Phair, R. D. (1996). Mechanisms for Ca signaling in vascular smooth muscle: resolved from ^{45}Ca uptake and efflux experiments. *Cell Calcium* **19**, 167-184.

- Leung, H. T., Geng, C. & Pak, W. L. (2000). Phenotypes of *trpl* mutants and interactions between the transient receptor potential (TRP) and TRP-like channels in *Drosophila*. *J Neurosci* **20**, 6797-6803.
- Li, H. S., Xu, X. Z. & Montell, C. (1999). Activation of a TRPC3-dependent cation current through the neurotrophin BDNF. *Neuron* **24**, 261-273.
- Li, Y., Shiels, A. J., Maszak, G. & Byron, K. L. (2001). Vasopressin-stimulated Ca^{2+} spiking in vascular smooth muscle cells involves phospholipase D. *Am J Physiol Heart Circ Physiol* **280**, H2658-2664.
- Liedtke, W., Choe, Y., Marti-Renom, M. A., Bell, A. M., Denis, C. S., Sali, A., Hudspeth, A. J., Friedman, J. M. & Heller, S. (2000). Vanilloid receptor-related osmotically activated channel (VR-OAC), a candidate vertebrate osmoreceptor. *Cell* **103**, 525-535.
- Liman, E. R., Corey, D. P. & Dulac, C. (1999). TRP2: A candidate transduction channel for mammalian pheromone sensory signaling. *Proc. Natl. Acad. Sci. USA* **96**, 5791-5796.
- Lintschinger, B., Balzer-Geldsetzer, M., Baskaran, T., Graier, W. F., Romanin, C., Zhu, M. X. & Groschner, K. (2000). Coassembly of Trp1 and Trp3 proteins generates diacylglycerol- and Ca^{2+} -sensitive cation channels. *J Biol Chem* **275**, 27799-27805.
- Liu, X., Wang, W., Singh, B. B., Lockwich, T., Jadowiec, J., O'Connell, B., Wellner, R., Zhu, M. X. & Ambudkar, I. S. (2000). Trp1, a candidate protein for the store-operated Ca^{2+} influx mechanism in salivary gland cells. *J Biol Chem* **275**, 3403-3411.
- Llinás, R. R. (1982). Calcium in synaptic transmission. *Sci Am* **247**, 56-65.
- Lockwich, T. P., Liu, X., Singh, B. B., Jadowiec, J., Weiland, S. & Ambudkar, I. S. (2000). Assembly of Trp1 in a signaling complex associated with caveolin-scaffolding lipid raft domains. *J Biol Chem* **275**, 11934-11942.
- Loirand, G., Pacaud, P., Baron, A., Mironneau, C. & Mironneau, J. (1991). Large conductance calcium-activated non-selective cation channel in smooth muscle cells isolated from rat portal vein. *J Physiol* **437**, 461-475.
- Lückhoff, A. & Clapham, D. E. (1992). Inositol 1,3,4,5-tetrakisphosphate activates an endothelial Ca^{2+} -permeable channel. *Nature* **355**, 356-358.
- Ma, H. T., Patterson, R. L., van Rossum, D. B., Birnbaumer, L., Mikoshiba, K. & Gill, D. L. (2000). Requirement of the inositol trisphosphate receptor for activation of store-operated Ca^{2+} channels. *Science* **287**, 1647-1651.
- Margaroli, A., Milani, D., Meldolesi, J. & Pozzan, T. (1987). Fura-2 measurement of cytosolic free Ca^{2+} in monolayers and suspensions of various types of animal cells. *J Cell Biol* **105**, 2145-2155.
- Marchenko, S. M. & Sage, S. O. (1993). Electrical properties of resting and acetylcholine-stimulated endothelium in intact rat aorta. *J Physiol* **462**, 735-751.
- Marks, T. N., Dubyak, G. R. & Jones, S. W. (1990). Calcium currents in the A7r5 smooth muscle-derived cell line. *Pflugers Arch* **417**, 433-439.
- Mathias, R. S., Zhang, S. J., Wilson, E., Gardner, P. & Ives, H. E. (1997). Non-capacitative calcium entry in Chinese hamster ovary cells expressing the platelet-derived growth factor receptor. *J Biol Chem* **272**, 29076-29082.
- Matsuda, J. J., Volk, K. A. & Shibata, E. F. (1990). Calcium currents in isolated rabbit coronary arterial smooth muscle myocytes. *J Physiol* **427**, 657-680.
- McDaniel, S. S., Platoshyn, O., Wang, J., Yu, Y., Sweeney, M., Krick, S., Rubin, L. J. & Yuan, J. X. (2001). Capacitative Ca^{2+} entry in agonist-induced pulmonary vasoconstriction. *Am J Physiol Lung Cell Mol Physiol* **280**, L870-880.
- McFadzean, I. & Gibson, A. (2002). The developing relationship between receptor-operated and store-operated calcium channels in smooth muscle. *Br J Pharmacol* **135**, 1-13.
- McKay, R. R., Szymeczek-Seay, C. L., Lievreumont, J. P., Bird, G. S., Zitt, C., Jüngling, E., Lückhoff, A. & Putney, J. W. (2000). Cloning and expression of the human transient receptor potential 4 (TRP4) gene: localization and functional expression of human TRP4 and TRP3. *Biochem J* **351 Pt 3**, 735-746.

- McKemy, D. D., Neuhauser, W. M. & Julius, D. (2002). Identification of a cold receptor reveals a general role for TRP channels in thermosensation. *Nature* **416**, 52-58.
- Meissner, G. (1994). Ryanodine receptor/ Ca^{2+} release channels and their regulation by endogenous effectors. *Annu Rev Physiol* **56**, 485-508.
- Meldolesi, J. & Pozzan, T. (1998). The endoplasmic reticulum Ca^{2+} store: a view from the lumen. *Trends Biochem Sci* **23**, 10-14.
- Mery, L., Magnino, F., Schmidt, K., Krause, K. H. & Dufour, J. F. (2001). Alternative splice variants of hTrp4 differentially interact with the C-terminal portion of the inositol 1,4,5-trisphosphate receptors. *FEBS Lett* **487**, 377-383.
- Michalak, M., Corbett, E. F., Mesaali, N., Nakamura, K. & Opas, M. (1999). Calreticulin: one protein, one gene, many functions. *Biochem J* **344 Pt 2**, 281-292.
- Mignen, O. & Shuttleworth, T. J. (2000). I_{ARC} , a novel arachidonate-regulated, noncapacitative Ca^{2+} entry channel. *J Biol Chem* **275**, 9114-9119.
- Mikoshiha, K. & Hattori, M. (2000). IP_3 receptor-operated calcium entry. *Sci STKE* **2000**, E1.
- Minke, B., Wu, C. & Pak, W. L. (1975). Induction of photoreceptor voltage noise in the dark in *Drosophila* mutant. *Nature* **258**, 84-87.
- Misquitta, C. M., Mack, D. P. & Grover, A. K. (1999). Sarco/endoplasmic reticulum Ca^{2+} (SERCA)-pumps: link to heart beats and calcium waves. *Cell Calcium* **25**, 277-290.
- Missiaen, L., Declerck, I., Droogmans, G., Plessers, L., De Smedt, H., Raeymaekers, L. & Casteels, R. (1990). Agonist-dependent Ca^{2+} and Mn^{2+} entry dependent on state of filling of Ca^{2+} stores in aortic smooth muscle cells of the rat. *J Physiol* **427**, 171-186.
- Mizuno, M., Kitayama, S., Saishin, Y., Shimada, S., Morita, K., Mitsuhata, C., Kurihara, H. & Dohi, T. (1999). Molecular cloning and characterization of rat *trp* homologues from brain. *Mol. Brain Res.* **64**, 41-51.
- Moneer, Z. & Taylor, C. W. (2002). Reciprocal regulation of capacitative and non-capacitative Ca^{2+} entry in A7r5 vascular smooth muscle cells: only the latter operates during receptor activation. *Biochem J* **362**, 13-21.
- Montell, C. (2001). Physiology, phylogeny, and functions of the TRP superfamily of cation channels. *Sci STKE* **2001**, RE1.
- Montell, C., Birnbaumer, L. & Flockerzi, V. (2002a). The TRP channels, a remarkably functional family. *Cell* **108**, 595-598.
- Montell, C., Birnbaumer, L., Flockerzi, V., Bindels, R. J., Bruford, E. A., Caterina, M. J., Clapham, D. E., Harteneck, C., Heller, S., Julius, D., Kojima, I., Mori, Y., Penner, R., Prawitt, D., Scharenberg, A. M., Schultz, G., Shimizu, N. & Zhu, M. X. (2002b). A unified nomenclature for the superfamily of TRP cation channels. *Mol Cell* **9**, 229-231.
- Montell, C. & Rubin, G. M. (1989). Molecular characterization of the *Drosophila trp* locus: a putative integral membrane protein required for phototransduction. *Neuron* **2**, 1313-1323.
- Mori, Y., Inoue, R., Ishii, M., Hara, Y. & Imoto, K. (2001). Dissecting receptor-mediated Ca^{2+} influx pathways: TRP channels and their native counterparts. *Jpn J Pharmacol* **87**, 245-252.
- Nadler, M. J., Hermosura, M. C., Inabe, K., Perraud, A. L., Zhu, Q., Stokes, A. J., Kurosaki, T., Kinet, J. P., Penner, R., Scharenberg, A. M. & Fleig, A. (2001). LTRPC7 is a Mg-ATP-regulated divalent cation channel required for cell viability. *Nature* **411**, 590-595.
- Nagamine, K., Kudoh, J., Minoshima, S., Kawasaki, K., Asakawa, S., Ito, F. & Shimizu, N. (1998). Molecular cloning of a novel putative Ca^{2+} channel protein (TRPC7) highly expressed in brain. *Genomics* **54**, 124-131.
- Nakajima, T., Hazama, H., Hamada, E., Wu, S. N., Igarashi, K., Yamashita, T., Seyama, Y., Omata, M. & Kurachi, Y. (1996). Endothelin-1 and vasopressin activate Ca^{2+} -permeable non-selective cation channels in aortic smooth muscle cells: mechanism of receptor-mediated Ca^{2+} influx. *J Mol Cell Cardiol* **28**, 707-722.
- Nakajima, T., Iwasawa, K., Hazama, H., Asano, M., Okuda, Y. & Omata, M. (1997). Extracellular Mg^{2+} inhibits receptor-mediated Ca^{2+} -permeable non-selective cation currents in aortic smooth muscle cells. *Eur J Pharmacol* **320**, 81-86.

- Neher, E. & Sakmann, B. (1976). Single-channel currents recorded from membrane of denervated frog muscle fibres. *Nature* **260**, 799-802.
- Nelson, M. T., Patlak, J. B., Worley, J. F. & Standen, N. B. (1990). Calcium channels, potassium channels, and voltage dependence of arterial smooth muscle tone. *Am J Physiol* **259**, C3-18.
- Nelson, M. T. & Worley, J. F. (1989). Dihydropyridine inhibition of single calcium channels and contraction in rabbit mesenteric artery depends on voltage. *J Physiol* **412**, 65-91.
- Ng, L. C. & Gurney, A. M. (2001). Store-operated channels mediate Ca^{2+} influx and contraction in rat pulmonary artery. *Circ Res* **89**, 923-929.
- Niemeyer, B. A., Suzuki, E., Scott, K., Jalink, K. & Zuker, C. S. (1996). The *Drosophila* light-activated conductance is composed of the two channels TRP and TRPL. *Cell* **85**, 651-659.
- Nilius, B., Vennekens, R., Prenen, J., Hoenderop, J. G., Droogmans, G. & Bindels, R. J. (2001). The single pore residue Asp542 determines Ca^{2+} permeation and Mg^{2+} block of the epithelial Ca^{2+} channel. *J Biol Chem* **276**, 1020-1025.
- Noguera, M. A., Madrero, Y., Ivorra, M. D. & D'Ocon, P. (1998). Characterization of two different Ca^{2+} entry pathways dependent on depletion of internal Ca^{2+} pools in rat aorta. *Naunyn Schmiedebergs Arch Pharmacol* **357**, 92-99.
- Numberger, M. & Draguhn, A. (1996). *Patch-Clamp-Technik*. Heidelberg: Spektrum Akademischer Verlag GmbH.
- Obejero-Paz, C. A., Lakshmanan, M., Jones, S. W. & Scarpa, A. (1993). Effects of dexamethasone on L-type calcium currents in the A7r5 smooth muscle-derived cell line. *FEBS Lett* **333**, 73-77.
- Ogden, D. C. & Stanfield, P. R. (1994). *Patch clamp techniques for single channel and whole-cell recording*. In *Microelectrode Techniques: The Plymouth Workshop Handbook*, Ogden, D. (ed.), (pp. 53-78). The Company of Biologists Ltd., Cambridge.
- Okada, T., Inoue, R., Yamazaki, K., Maeda, A., Kurosaki, T., Yamakuni, T., Tanaka, I., Shimizu, S., Ikenaka, K., Imoto, K. & Mori, Y. (1999). Molecular and functional characterization of a novel mouse transient receptor potential protein homologue TRP7. Ca^{2+} -permeable cation channel that is constitutively activated and enhanced by stimulation of G protein-coupled receptor. *J Biol Chem* **274**, 27359-27370.
- Okada, T., Shimizu, S., Wakamori, M., Maeda, A., Kurosaki, T., Takada, N., Imoto, K. & Mori, Y. (1998). Molecular cloning and functional characterization of a novel receptor-activated TRP Ca^{2+} channel from mouse brain. *J. Biol. Chem.* **273**, 10279-10287.
- Osol, G., Laher, I. & Kelley, M. (1993). Myogenic tone is coupled to phospholipase C and G protein activation in small cerebral arteries. *Am J Physiol* **265**, H415-420.
- Otsuka, Y., Sakagami, H., Owada, Y. & Kondo, H. (1998). Differential localization of mRNAs for mammalian trps, presumptive capacitative calcium entry channels, in the adult mouse brain. *Tohoku J Exp Med* **185**, 139-146.
- Palmer, C. P., Zhou, X. L., Lin, J., Loukin, S. H., Kung, C. & Saimi, Y. (2001). A TRP homolog in *Saccharomyces cerevisiae* forms an intracellular Ca^{2+} -permeable channel in the yeast vacuolar membrane. *Proc Natl Acad Sci U S A* **98**, 7801-7805.
- Parekh, A. B. & Penner, R. (1997). Store depletion and calcium influx. *Physiological Reviews* **77**, 901-930.
- Patel, S., Churchill, G. C. & Galione, A. (2001). Coordination of Ca^{2+} signalling by NAADP. *Trends Biochem Sci* **26**, 482-489.
- Peier, A. M., Moqrich, A., Hergarden, A. C., Reeve, A. J., Andersson, D. A., Story, G. M., Earley, T. J., Dragoni, I., McIntyre, P., Bevan, S. & Patapoutian, A. (2002). A TRP channel that senses cold stimuli and menthol. *Cell* **108**, 705-715.
- Peier, A. M., Reeve, A. J., Andersson, D. A., Moqrich, A., Earley, T. J., Hergarden, A. C., Story, G. M., Colley, S., Hogenesch, J. B., McIntyre, P., Bevan, S. & Patapoutian, A. (2002). A heat-sensitive TRP channel expressed in keratinocytes. *Science, in press*.

- Peng, J. B., Chen, X. Z., Berger, U. V., Vassilev, P. M., Tsukaguchi, H., Brown, E. M. & Hediger, M. A. (1999). Molecular cloning and characterization of a channel-like transporter mediating intestinal calcium absorption. *J Biol Chem* **274**, 22739-22746.
- Peng, J. B., Chen, X. Z., Berger, U. V., Weremowicz, S., Morton, C. C., Vassilev, P. M., Brown, E. M. & Hediger, M. A. (2000). Human calcium transport protein CaT1. *Biochem Biophys Res Commun* **278**, 326-332.
- Penniston, J. T. & Enyedi, A. (1998). Modulation of the plasma membrane Ca^{2+} pump. *J Membr Biol* **165**, 101-109.
- Perraud, A. L., Fleig, A., Dunn, C. A., Bagley, L. A., Launay, P., Schmitz, C., Stokes, A. J., Zhu, Q., Bessman, M. J., Penner, R., Kinet, J. P. & Scharenberg, A. M. (2001). ADP-ribose gating of the calcium-permeable LTRPC2 channel revealed by Nudix motif homology. *Nature* **411**, 595-599.
- Petersen, O. H., Tepikin, A. & Park, M. K. (2001). The endoplasmic reticulum: one continuous or several separate Ca^{2+} stores? *Trends Neurosci* **24**, 271-276.
- Petr, M. J. & Wurster, R. D. (1997). Determination of in situ dissociation constant for Fura-2 and quantitation of background fluorescence in astrocyte cell line U373-MG. *Cell Calcium* **21**, 233-240.
- Pfützer, G. (2001). Invited review: regulation of myosin phosphorylation in smooth muscle. *J Appl Physiol* **91**, 497-503.
- Philipp, S., Cavalié, A., Freichel, M., Wissenbach, U., Zimmer, S., Trost, C., Marquart, A., Murakami, M. & Flockerzi, V. (1996). A mammalian capacitative calcium entry channel homologous to *Drosophila* TRP and TRPL. *EMBO Journal* **15**, 6166-6171.
- Philipp, S., Hambrecht, J., Braslavski, L., Schroth, G., Freichel, M., Murakami, M., Cavalié, A. & Flockerzi, V. (1998). A novel capacitative calcium entry channel expressed in excitable cells. *EMBO J.* **17**, 4274-4282.
- Philipp, S., Trost, C., Warnat, J., Rautmann, J., Himmerkus, N., Schroth, G., Kretz, O., Nastainczyk, W., Cavalié, A., Hoth, M. & Flockerzi, V. (2000). TRP4 (CCE1) protein is part of native calcium release-activated Ca^{2+} -like channels in adrenal cells. *J Biol Chem* **275**, 23965-23972.
- Phillips, A. M., Bull, A. & Kelly, L. E. (1992). Identification of a *Drosophila* gene encoding a calmodulin-binding protein with homology to the trp phototransduction gene. *Neuron* **8**, 631-642.
- Popp, R., Englert, H. C., Lang, H. J. & Gögelein, H. (1993). Inhibitors of nonselective cation channels in cells of the blood-brain barrier. *Exs* **66**, 213-218.
- Pozzan, T., Rizzuto, R., Volpe, P. & Meldolesi, J. (1994). Molecular and cellular physiology of intracellular calcium stores. *Physiol Rev* **74**, 595-636.
- Putney, J. W., Jr. (1986). A model for receptor-regulated calcium entry. *Cell Calcium* **7**, 1-12.
- Putney, J. W., Jr. (1987). Formation and actions of calcium-mobilizing messenger, inositol 1,4,5-trisphosphate. *Am J Physiol* **252**, G149-157.
- Putney, J. W., Jr., Broad, L. M., Braun, F. J., Lievreumont, J. P. & Bird, G. S. (2001). Mechanisms of capacitative calcium entry. *J Cell Sci* **114**, 2223-2229.
- Putney, J. W. J. (1997). *Capacitative Calcium Entry*. Georgetown, TX, U.S.A.: R. G. Landes Company.
- Ramanan, S. V., Fan, S. F. & Brink, P. R. (1992). Model invariant method for extracting single-channel mean open and closed times from heterogeneous multichannel records. *J Neurosci Methods* **42**, 91-103.
- Ranganathan, R., Malicki, D. M. & Zuker, C. S. (1995). Signal transduction in *Drosophila* photoreceptors. *Annu Rev Neurosci* **18**, 283-317.
- Reuss, H., Mojet, M. H., Chyb, S. & Hardie, R. C. (1997). In vivo analysis of the *drosophila* light-sensitive channels, TRP and TRPL. *Neuron* **19**, 1249-1259.
- Riccio, A., Mattei, C., Kelsell, R. E., Medhurst, A. D., Calver, A. R., Randall, A. D., Davis, J. B., Benham, C. D. & Pangalos, M. N. (2002). Cloning and functional expression of human

- short TRP7, a candidate protein for store-operated Ca^{2+} influx. *J Biol Chem* **277**, 12302-12309.
- Rosado, J. A. & Sage, S. O. (2001). Activation of store-mediated calcium entry by secretion-like coupling between the inositol 1,4,5-trisphosphate receptor type II and human transient receptor potential (hTrp1) channels in human platelets. *Biochem J* **356**, 191-198.
- Runnels, L. W., Yue, L. & Clapham, D. E. (2001). TRP-PLIK, a bifunctional protein with kinase and ion channel activities. *Science* **291**, 1043-1047.
- Runnels, L. W., Yue, L. & Clapham, D. E. (2002). The TRPM7 channel is inactivated by PIP_2 hydrolysis. *Nat Cell Biol* **4**, 329-336.
- Ryazanov, A. G. (2002). Elongation factor-2 kinase and its newly discovered relatives. *FEBS Lett* **514**, 26-29.
- Sakmann, B. & Neher, E. (Ed.). (1983). *Single-Channel Recording* (first edition ed.). New York: Plenum Press.
- Sanders, K. M. (2001). Invited review: mechanisms of calcium handling in smooth muscles. *J Appl Physiol* **91**, 1438-1449.
- Schaefer, M., Plant, T. D., Obukhov, A. G., Hofmann, T., Gudermann, T. & Schultz, G. (2000). Receptor-mediated regulation of the nonselective cation channels TRPC4 and TRPC5. *J Biol Chem* **275**, 17517-17526.
- Schaefer, M., Plant, T. D., Stresow, N., Albrecht, N. & Schultz, G. (2002). Functional differences between TRPC4 splice variants. *J Biol Chem* **277**, 3752-3759.
- Shuttleworth, T. J. & Thompson, J. L. (1998). Muscarinic receptor activation of arachidonate-mediated Ca^{2+} entry in HEK293 cells is independent of phospholipase C. *J Biol Chem* **273**, 32636-32643.
- Silberberg, S. D., Poder, T. C. & Lacerda, A. E. (1989). Endothelin increases single-channel calcium currents in coronary arterial smooth muscle cells. *FEBS Lett* **247**, 68-72.
- Singh, B. B., Liu, X. & Ambudkar, I. S. (2000). Expression of truncated transient receptor potential protein 1α (Trp1 α): evidence that the Trp1 C terminus modulates store-operated Ca^{2+} entry. *J Biol Chem* **275**, 36483-36486.
- Singh, B. B., Zheng, C., Liu, X., Lockwich, T., Liao, D., Zhu, M. X., Birnbaumer, L. & Ambudkar, I. S. (2001). Trp1-dependent enhancement of salivary gland fluid secretion: role of store-operated calcium entry. *Faseb J* **15**, 1652-1654.
- Sinkins, W. G., Estacion, M. & Schilling, W. P. (1998). Functional expression of TrpC1: a human homologue of the *Drosophila* Trp channel. *Biochem J* **331**, 331-339.
- Smart, D., Gunthorpe, M. J., Jerman, J. C., Nasir, S., Gray, J., Muir, A. I., Chambers, J. K., Randall, A. D. & Davis, J. B. (2000). The endogenous lipid anandamide is a full agonist at the human vanilloid receptor (hVR1). *Br J Pharmacol* **129**, 227-230.
- Sterner, O. & Szallasi, A. (1999). Novel natural vanilloid receptor agonists: new therapeutic targets for drug development. *Trends Pharmacol Sci* **20**, 459-465.
- Sternweis, P. C. & Gilman, A. G. (1982). Aluminum: a requirement for activation of the regulatory component of adenylate cyclase by fluoride. *Proc Natl Acad Sci U S A* **79**, 4888-4891.
- Stevens, T. (2001). Is there a role for store-operated calcium entry in vasoconstriction? *Am J Physiol Lung Cell Mol Physiol* **280**, L866-869.
- Striggow, F. & Ehrlich, B. E. (1996). Ligand-gated calcium channels inside and out. *Curr Opin Cell Biol* **8**, 490-495.
- Strotmann, R., Harteneck, C., Nunnenmacher, K., Schultz, G. & Plant, T. D. (2000). OTRPC4, a nonselective cation channel that confers sensitivity to extracellular osmolarity. *Nat Cell Biol* **2**, 695-702.
- Strübing, C., Krapivinsky, G., Krapivinsky, L. & Clapham, D. E. (2001). Trpc1 and trpc5 form a novel cation channel in mammalian brain. *Neuron* **29**, 645-655.
- Sutko, J. L. & Airey, J. A. (1996). Ryanodine receptor Ca^{2+} release channels: does diversity in form equal diversity in function? *Physiol Rev* **76**, 1027-1071.

- Sylvester, J. B., Mwanjewe, J. & Grover, A. K. (2001). Transient receptor potential protein mRNA expression in rat substantia nigra. *Neurosci Lett* **300**, 83-86.
- Szallasi, A. & Blumberg, P. M. (1999). Vanilloid (Capsaicin) receptors and mechanisms. *Pharmacol Rev* **51**, 159-212.
- Takahashi, A., Camacho, P., Lechleiter, J. D. & Herman, B. (1999). Measurement of intracellular calcium. *Physiol Rev* **79**, 1089-1125.
- Taylor, C. W. (1998). Inositol trisphosphate receptors: Ca^{2+} -modulated intracellular Ca^{2+} channels. *Biochim Biophys Acta* **1436**, 19-33.
- Tesfai, Y., Brereton, H. M. & Barritt, G. J. (2001). A diacylglycerol-activated Ca^{2+} channel in PC12 cells (an adrenal chromaffin cell line) correlates with expression of the TRP-6 (transient receptor potential) protein. *Biochem J* **358**, 717-726.
- Thibonnier, M., Bayer, A. L., Simonson, M. S. & Kester, M. (1991). Multiple signaling pathways of V_1 -vascular vasopressin receptors of A7r5 cells. *Endocrinology* **129**, 2845-2856.
- Tominaga, M., Caterina, M., Malmberg, A. B., Rosen, T. A., Gilbert, H., Skinner, K., Raumann, B. E., Basbaum, A. I. & Julius, D. (1998). The cloned capsaicin receptor integrates multiple pain-producing stimuli. *Neuron* **21**, 531-543.
- Tomita, Y., Kaneko, S., Funayama, M., Kondo, H., Satoh, M. & Akaike, A. (1998). Intracellular Ca^{2+} store-operated influx of Ca^{2+} through TRP-R, a rat homolog of TRP, expressed in *Xenopus* oocytes. *Neurosci. Lett.* **248**, 195-198.
- Trepakova, E. S., Gericke, M., Hirakawa, Y., Weisbrod, R. M., Cohen, R. A. & Bolotina, V. M. (2001). Properties of a native cation channel activated by Ca^{2+} store depletion in vascular smooth muscle cells. *J Biol Chem* **276**, 7782-7790.
- Trump, B. F. & Berezsky, I. K. (1995). Calcium-mediated cell injury and cell death. *Faseb J* **9**, 219-228.
- Tsavalier, L., Shaper, M. H., Morkowski, S. & Laus, R. (2001). *Trp-p8*, a novel prostate-specific gene, is up-regulated in prostate cancer and other malignancies and shares high homology with transient receptor potential calcium channel proteins. *Cancer Res* **61**, 3760-3769.
- Tsien, R. W. & Tsien, R. Y. (1990). Calcium channels, stores, and oscillations. *Annu Rev Cell Biol* **6**, 715-760.
- Vaca, L. & Kunze, D. L. (1995). IP_3 -activated Ca^{2+} channels in the plasma membrane of cultured vascular endothelial cells. *American Journal of Physiology: Cell Physiology* **269**, C733-C738.
- Vaca, L., Sinkins, W. G., Hu, Y., Kunze, D. L. & Schilling, W. P. (1994). Activation of recombinant trp by thapsigargin in Sf9 insect cells. *Am J Physiol* **267**, C1501-1505.
- van Breemen, C. & Saida, K. (1989). Cellular mechanisms regulating $[Ca^{2+}]_i$ smooth muscle. *Annu Rev Physiol* **51**, 315-329.
- van Haasteren, G., Li, S., Muda, M., Susini, S. & Schlegel, W. (1999). Calcium signalling and gene expression. *J Recept Signal Transduct Res* **19**, 481-492.
- Van Renterghem, C., Romey, G. & Lazdunski, M. (1988). Vasopressin modulates the spontaneous electrical activity in aortic cells (line A7r5) by acting on three different types of ionic channels. *Proc Natl Acad Sci U S A* **85**, 9365-9369.
- van Rossum, D. B., Patterson, R. L., Ma, H. T. & Gill, D. L. (2000). Ca^{2+} entry mediated by store depletion, S-nitrosylation, and TRP3 channels. Comparison of coupling and function. *J Biol Chem* **275**, 28562-28568.
- Vannier, B., Peyton, M., Boulay, G., Brown, D., Qin, N., Jiang, M., Zhu, X. & Birnbaumer, L. (1999). Mouse trp2, the homologue of the human trpc2 pseudogene, encodes mTrp2, a store depletion-activated capacitative Ca^{2+} entry channel. *Proc. Natl. Acad. Sci. USA* **96**, 2060-2064.
- Vannier, B., Zhu, X., Brown, D. & Birnbaumer, L. (1998). The membrane topology of human transient receptor potential 3 as inferred from glycosylation-scanning mutagenesis and epitope immunocytochemistry. *J. Biol. Chem.* **273**, 8675-8679.

- Venkatachalam, K., Ma, H. T., Ford, D. L. & Gill, D. L. (2001). Expression of functional receptor-coupled TRPC3 channels in DT40 triple receptor InsP₃ knockout cells. *J Biol Chem* **276**, 33980-33985.
- Vennekens, R., Hoenderop, J. G., Prenen, J., Stuiver, M., Willems, P. H., Droogmans, G., Nilius, B. & Bindels, R. J. (2000). Permeation and gating properties of the novel epithelial Ca²⁺ channel. *J Biol Chem* **275**, 3963-3969.
- Vennekens, R., Prenen, J., Hoenderop, J. G., Bindels, R. J., Droogmans, G. & Nilius, B. (2001). Pore properties and ionic block of the rabbit epithelial calcium channel expressed in HEK 293 cells. *J Physiol* **530**, 183-191.
- Voets, T., Prenen, J., Fleig, A., Vennekens, R., Watanabe, H., Hoenderop, J. G., Bindels, R. J., Droogmans, G., Penner, R. & Nilius, B. (2001). CaT1 and the calcium release-activated calcium channel manifest distinct pore properties. *J Biol Chem* **276**, 47767-47770.
- Volpe, P., Pozzan, T. & Meldolesi, J. (1990). Rapidly exchanging Ca²⁺ stores of non-muscle cells. *Semin Cell Biol* **1**, 297-304.
- Walker, R. L., Hume, J. R. & Horowitz, B. (2001). Differential expression and alternative splicing of TRP channel genes in smooth muscles. *Am J Physiol Cell Physiol* **280**, C1184-1192.
- Warnat, J., Philipp, S., Zimmer, S., Flockerzi, V. & Cavalié, A. (1999). Phenotype of a recombinant store-operated channel: highly selective permeation of Ca²⁺. *J. Physiol.* **518**, 631-638.
- Watanabe, H., Davis, J. B., Smart, D., Jerman, J. C., Smith, G. D., Hayes, P., Vriens, J., Cairns, W., Wissenbach, U., Prenen, J., Flockerzi, V., Droogmans, G., Benham, C. D. & Nilius, B. (2002). Activation of TRPV4 channels (hVRL-2/mTRP12) by phorbol derivatives. *J Biol Chem* **277**, 13569-13577.
- Welsh, D. G., Morielli, A. D., Nelson, M. T. & Brayden, J. E. (2002). Transient receptor potential channels regulate myogenic tone of resistance arteries. *Circ Res* **90**, 248-250.
- Wes, P. D., Chevesich, J., Jeromin, A., Rosenberg, C., Stetten, G. & Montell, C. (1995). TRPC1, a human homolog of a *Drosophila* store-operated channel. *PNAS* **92**, 9652-9656.
- Williams, D. A. & Fay, F. S. (1990). Intracellular calibration of the fluorescent calcium indicator Fura-2. *Cell Calcium* **11**, 75-83.
- Wilson, E., Mai, Q., Sudhir, K., Weiss, R. H. & Ives, H. E. (1993). Mechanical strain induces growth of vascular smooth muscle cells via autocrine action of PDGF. *J Cell Biol* **123**, 741-747.
- Wissenbach, U., Boddling, M., Freichel, M. & Flockerzi, V. (2000). Trp12, a novel Trp related protein from kidney. *FEBS Lett* **485**, 127-134.
- Worley, J. F., Deitmer, J. W. & Nelson, M. T. (1986). Single nisoldipine-sensitive calcium channels in smooth muscle cells isolated from rabbit mesenteric artery. *Proc Natl Acad Sci U S A* **83**, 5746-5750.
- Wu, X., Babnigg, G. & Villereal, M. L. (2000). Functional significance of human trp1 and trp3 in store-operated Ca²⁺ entry in HEK-293 cells. *Am J Physiol Cell Physiol* **278**, C526-536.
- Xu, L., Tripathy, A., Pasek, D. A. & Meissner, G. (1998). Potential for pharmacology of ryanodine receptor/calcium release channels. *Ann N Y Acad Sci* **853**, 130-148.
- Xu, S. Z. & Beech, D. J. (2001). TrpC1 is a membrane-spanning subunit of store-operated Ca²⁺ channels in native vascular smooth muscle cells. *Circ Res* **88**, 84-87.
- Xu, X. Z., Chien, F., Butler, A., Salkoff, L. & Montell, C. (2000). TRP γ , a *Drosophila* TRP-related subunit, forms a regulated cation channel with TRPL. *Neuron* **26**, 647-657.
- Xu, X. Z., Li, H. S., Guggino, W. B. & Montell, C. (1997). Coassembly of TRP and TRPL produces a distinct store-operated conductance. *Cell* **89**, 1155-1164.
- Yamada, H., Wakamori, M., Hara, Y., Takahashi, Y., Konishi, K., Imoto, K. & Mori, Y. (2000). Spontaneous single-channel activity of neuronal TRP5 channel recombinantly expressed in HEK293 cells. *Neurosci Lett* **285**, 111-114.

- Yue, L., Peng, J. B., Hediger, M. A. & Clapham, D. E. (2001). CaT1 manifests the pore properties of the calcium-release-activated calcium channel. *Nature* **410**, 705-709.
- Zhang, L. & Saffen, D. (2001). Muscarinic acetylcholine receptor regulation of TRP6 Ca²⁺ channel isoforms. Molecular structures and functional characterization. *J Biol Chem* **276**, 13331-13339.
- Zhang, Z., Tang, J., Tikunova, S., Johnson, J. D., Chen, Z., Qin, N., Dietrich, A., Stefani, E., Birnbaumer, L. & Zhu, M. X. (2001). Activation of Trp3 by inositol 1,4,5-trisphosphate receptors through displacement of inhibitory calmodulin from a common binding domain. *Proc Natl Acad Sci U S A* **98**, 3168-3173.
- Zhu, X., Chu, P.B., Peyton, M., Birnbaumer, L. (1995). Molecular cloning of a widely expressed human homologue for the *Drosophila* trp gene. *FEBS Lett.* **373**, 193-198.
- Zhu, X., Jiang, M. & Birnbaumer, L. (1998). Receptor-activated Ca²⁺ influx via human Trp3 stably expressed in human embryonic kidney (HEK)293 cells. *J. Biol. Chem.* **273**, 133-142.
- Zhu, X., Jiang, M., Peyton, M., Boulay, G., Hurst, R., Stefani, E. & Birnbaumer, L. (1996). trp, a novel mammalian gene family essential for agonist-activated capacitative Ca²⁺ entry. *Cell* **85**, 661-671.
- Zitt, C., Halaszovich, C. R. & Lückhoff, A. (2002). The TRP family of cation channels: probing and advancing the concepts on receptor-activated calcium entry. *Prog Neurobiol* **66**, 243-264.
- Zitt, C., Obukhov, A. G., Strübing, C., Zobel, A., Kalkbrenner, F., Lückhoff, A. & Schultz, G. (1997). Expression of TRPC3 in Chinese hamster ovary cells results in calcium-activated cation currents not related to store depletion. *Journal of Cell Biology* **138**, 1333-1341.
- Zitt, C., Zobel, A., Obukhov, A. G., Harteneck, C., Kalkbrenner, F., Lückhoff, A. & Schultz, G. (1996). Cloning and functional expression of a human Ca²⁺-permeable cation channel activated by calcium store depletion. *Neuron* **16**, 1189-1196.
- Zucchi, R. & Ronca-Testoni, S. (1997). The sarcoplasmic reticulum Ca²⁺ channel/ryanodine receptor: modulation by endogenous effectors, drugs and disease states. *Pharmacol Rev* **49**, 1-51.
- Zygmunt, P. M., Petersson, J., Andersson, D. A., Chuang, H., Sorgard, M., Di Marzo, V., Julius, D. & Hogestatt, E. D. (1999). Vanilloid receptors on sensory nerves mediate the vasodilator action of anandamide. *Nature* **400**, 452-457.