

## 7 Literaturverzeichnis

- [1] Deutsche Gesellschaft für Kardiologie - Herz- und Kreislaufforschung e.V. (2005): Leistungsbilanz der Herzchirurgie, DGTHG Hamburg 2005.
- [2] Green, G. E.; Stertzer, S. H.; Gordon, R. B. und Tice, D. A. (1970): Anastomosis of the internal mammary artery to the distal left anterior descending coronary artery, *Circulation* 41 [5 Suppl], Seite II79-II85. URL: PM:4952697
- [3] Moccetti, T.; Lichtlen, P.; Minale, C.; Halter, J.; Schonbeck, M.; Meier, W. und Senning, A. (1972): [Aorto-coronary bypass with graft of the saphenous vein (108 cases). Clinical, angiographic and functional long-term results], *G.Ital.Cardiol.* 2 [5], Seite 644-654. URL: PM:4538714
- [4] Shander, A.; Knight, K.; Thurer, R.; Adamson, J. und Spence, R. (2004): Prevalence and outcomes of anemia in surgery: a systematic review of the literature, *Am.J.Med.* 116 Suppl 7A, Seite 58S-69S. URL: PM:15050887
- [5] Bundesärztekammer (2003): Leitlinien zur Therapie mit Blutkomponenten und Plasmaderivaten, 3.überarb. Aufl., Deutscher Ärzte Verlag-Köln 2003 16, Seite 259-278.
- [6] Blumberg, N. (1997): Allogeneic transfusion and infection: economic and clinical implications, *Semin.Hematol.* 34 [3 Suppl 2], Seite 34-40. URL: PM:9253782
- [7] Innerhofer, P.; Tilz, G.; Fuchs, D.; Luz, G.; Hobisch-Hagen, P.; Schobersberger, W.; Nussbaumer, W.; Lochs, A. und Irschick, E. (2000): Immunologic changes after transfusion of autologous or allogeneic buffy coat-poor versus WBC-reduced blood transfusions in patients undergoing arthroplasty. II. Activation of T cells, macrophages, and cell-mediated lympholysis, *Transfusion* 40 [7], Seite 821-827. URL: PM:10924610
- [8] von, Heymann C.; Pruss, A.; Foer, A.; Volk, T.; Braun, J.; Rohrs, A.; Kiesewetter, H. und Spies, C. (2005): [Quality assurance of haemotherapy in anaesthesia. Auswertung eines 3-jährigen Erfahrungszeitraums], *Anaesthesist* 54 [2], Seite 107-116. URL: PM:15599688
- [9] Carson, J. L.; Altman, D. G.; Duff, A.; Noveck, H.; Weinstein, M. P.; Sonnenberg, F. A.; Hudson, J. I. und Provenzano, G. (1999): Risk of bacterial infection associated with allogeneic blood transfusion among patients undergoing hip fracture repair, *Transfusion* 39 [7], Seite 694-700. URL: PM:10413276
- [10] Marchetti, M. und Barosi, G. (2000): Cost-effectiveness of epoetin and autologous blood donation in reducing allogeneic blood transfusions in coronary artery bypass graft surgery, *Transfusion* 40 [6], Seite 673-681. URL: PM:10864987

- [11] Shander, A. (2004): Emerging risks and outcomes of blood transfusion in surgery, *Semin.Hematol.* 41 [1 Suppl 1], Seite 117-124. URL: PM:14872432
- [12] Vamvakas, E. C. und Carven, J. H. (2000): Length of storage of transfused red cells and postoperative morbidity in patients undergoing coronary artery bypass graft surgery, *Transfusion* 40 [1], Seite 101-109. URL: PM:10644819
- [13] Habler, O.; Kleen, M.; Kemming, G. und Zwissler, B. (2002): Hyperoxia in extreme hemodilution, *Eur.Surg.Res.* 34 [1-2], Seite 181-187. URL: PM:11867921
- [14] Habler, O. und Messmer, K. (1997): [Methods for reduction of homologous blood transfusions in operative medicine], *Anaesthesist* 46 [11], Seite 915-926. URL: PM:9490578
- [15] Bryson, G. L.; Laupacis, A. und Wells, G. A. (1998): Does acute normovolemic hemodilution reduce perioperative allogeneic transfusion? A meta-analysis. The International Study of Perioperative Transfusion, *Anesth.Analg.* 86 [1], Seite 9-15. URL: PM:9428843
- [16] Feldman, J. M.; Roth, J. V. und Bjoraker, D. G. (1995): Maximum blood savings by acute normovolemic hemodilution, *Anesth.Analg.* 80 [1], Seite 108-113. URL: PM:7802266
- [17] Matot, I.; Scheinin, O.; Jurim, O. und Eid, A. (2002): Effectiveness of acute normovolemic hemodilution to minimize allogeneic blood transfusion in major liver resections, *Anesthesiology* 97 [4], Seite 794-800. URL: PM:12357142
- [18] Monk, T. G. (2005): Acute normovolemic hemodilution, *Anesthesiol.Clin.North America.* 23 [2], Seite 271-81, vi. URL: PM:15922899
- [19] Habler, O. P. und Messmer, K. F. (1997): The physiology of oxygen transport, *Transfus.Sci.* 18 [3], Seite 425-435. URL: PM:10175156
- [20] Spahn, D. R.; Leone, B. J.; Reves, J. G. und Pasch, T. (1994): Cardiovascular and coronary physiology of acute isovolemic hemodilution: a review of nonoxygen-carrying and oxygen-carrying solutions, *Anesth.Analg.* 78 [5], Seite 1000-1021. URL: PM:8160966
- [21] Kreimeier, U. und Messmer, K. (1996): Hemodilution in clinical surgery: state of the art 1996, *World J Surg.* 20 [9], Seite 1208-1217. URL: PM:8864083
- [22] Bruckner, U. B. und Messmer, K. (1990): Blood rheology and systemic oxygen transport, *Biorheology* 27 [6], Seite 903-912. URL: PM:2093398

- [23] Klovekorn, W. P.; Pichlmaier, H.; Ott, E.; Bauer, H.; Sunder-Plassmann, L. und Messmer, K. (1974): [Acute preoperative hemodilution--possibility for autologous blood transfusion], *Chirurg* 45 [10], Seite 452-458. URL: PM:4462985
- [24] Lauth, W. W. (1977): Control of hepatic and intestinal blood flow: effect of isovolaemic haemodilution on blood flow and oxygen uptake in the intact liver and intestines, *J.Physiol* 265 [2], Seite 313-326. URL: PM:15108
- [25] Chamorro, G.; Rodriguez, J. A.; Dzindzio, B. und Rapaport, E. (1973): Effect of acute isovolemic anemia on cardiac output and estimated hepatic blood flow in the conscious dog, *Circ.Res.* 32 [4], Seite 530-535. URL: PM:4573829
- [26] Noldge, G. F.; Priebe, H. J. und Geiger, K. (1992): Splanchnic hemodynamics and oxygen supply during acute normovolemic hemodilution alone and with isoflurane-induced hypotension in the anesthetized pig, *Anesth.Analg.* 75 [5], Seite 660-674. URL: PM:1416116
- [27] Kleen, M.; Habler, O.; Hutter, J.; Podtschaske, A.; Tiede, M.; Kemming, G.; Corso, C.; Batra, S.; Keipert, P.; Faithfull, S. und Messmer, K. (1997): Effects of hemodilution on splanchnic perfusion and hepatorenal function. I. Splanchnic perfusion, *Eur.J.Med.Res.* 2 [10], Seite 413-418. URL: PM:9348267
- [28] Kleen, M.; Habler, O.; Hutter, J.; Podtschaske, A.; Tiede, M.; Kemming, G.; Welte, M.; Corso, C. und Messmer, K. (1996): Effects of hemodilution on gastric regional perfusion and intramucosal pH, *Am.J.Physiol* 271 [5 Pt 2], Seite H1849-H1855. URL: PM:8945900
- [29] Laks, H.; O'Connor, N. E.; Pilon, R. N.; Anderson, W.; MacCallum, J. R.; Klovekorn, W. P. und Moore, F. D. (1973): Acute normovolemic hemodilution: effects on hemodynamics, oxygen transport, and lung water in anesthetized man, *Surg.Forum* 24, Seite 201-202. URL: PM:4805985
- [30] Weiskopf, R. B.; Viele, M. K.; Feiner, J.; Kelley, S.; Lieberman, J.; Noorani, M.; Leung, J. M.; Fisher, D. M.; Murray, W. R.; Toy, P. und Moore, M. A. (1998): Human cardiovascular and metabolic response to acute, severe isovolemic anemia, *JAMA* 279 [3], Seite 217-221. URL: PM:9438742
- [31] Bracey, A. W.; Radovancevic, R.; Riggs, S. A.; Houston, S.; Cozart, H.; Vaughn, W. K.; Radovancevic, B.; McAllister, H. A., Jr. und Cooley, D. A. (1999): Lowering the hemoglobin threshold for transfusion in coronary artery bypass procedures: effect on patient outcome, *Transfusion* 39 [10], Seite 1070-1077. URL: PM:10532600
- [32] DeFoe, G. R.; Ross, C. S.; Olmstead, E. M.; Surgenor, S. D.; Fillinger, M. P.; Groom, R. C.; Forest, R. J.; Pieroni, J. W.; Warren, C. S.; Bogosian, M. E.; Krumholz, C. F.; Clark, C.; Clough, R. A.; Weldner, P. W.; Lahey, S. J.; Leavitt, B. J.; Marrin, C. A.; Charlesworth, D. C.; Marshall, P. und O'Connor, G. T. (2001):

- Lowest hematocrit on bypass and adverse outcomes associated with coronary artery bypass grafting. Northern New England Cardiovascular Disease Study Group, *Ann.Thorac.Surg.* 71 [3], Seite 769-776. URL: PM:11269449
- [33] Fang, W. C.; Helm, R. E.; Krieger, K. H.; Rosengart, T. K.; DuBois, W. J.; Sason, C.; Lesser, M. L.; Isom, O. W. und Gold, J. P. (1997): Impact of minimum hematocrit during cardiopulmonary bypass on mortality in patients undergoing coronary artery surgery, *Circulation* 96 [9 Suppl], Seite II-9. URL: PM:9386097
- [34] Hardy, J. F.; Martineau, R.; Couturier, A.; Belisle, S.; Cartier, R. und Carrier, M. (1998): Influence of haemoglobin concentration after extracorporeal circulation on mortality and morbidity in patients undergoing cardiac surgery, *Br.J.Anaesth.* 81 Suppl 1, Seite 38-45. URL: PM:10318987
- [35] Fromes, Y.; Gaillard, D.; Ponzio, O.; Chauffert, M.; Gerhardt, M. F.; Deleuze, P. und Bical, O. M. (2002): Reduction of the inflammatory response following coronary bypass grafting with total minimal extracorporeal circulation, *Eur.J.Cardiothorac.Surg.* 22 [4], Seite 527-533. URL: PM:12297167
- [36] Hirai, S. (2003): Systemic inflammatory response syndrome after cardiac surgery under cardiopulmonary bypass, *Ann.Thorac.Cardiovasc.Surg.* 9 [6], Seite 365-370. URL: PM:15003097
- [37] Kumle, B.; Boldt, J.; Suttner, S. W.; Piper, S. N.; Lehmann, A. und Blome, M. (2003): Influence of prolonged cardiopulmonary bypass times on splanchnic perfusion and markers of splanchnic organ function, *Ann.Thorac.Surg.* 75 [5], Seite 1558-1564. URL: PM:12735579
- [38] Orime, Y.; Shiono, M.; Hata, H.; Yagi, S.; Tsukamoto, S.; Okumura, H.; Nakata, K.; Kimura, S.; Hata, M.; Sezai, A. und Sezai, Y. (1999): Cytokine and endothelial damage in pulsatile and nonpulsatile cardiopulmonary bypass, *Artif.Organs* 23 [6], Seite 508-512. URL: PM:10392275
- [39] Rao, S. V.; O'Grady, K.; Pieper, K. S.; Granger, C. B.; Newby, L. K.; Van de Werf F.; Mahaffey, K. W.; Califf, R. M. und Harrington, R. A. (2005): Impact of bleeding severity on clinical outcomes among patients with acute coronary syndromes, *Am.J.Cardiol.* 96 [9], Seite 1200-1206. URL: PM:16253582
- [40] Landow, L. (1993): Splanchnic lactate production in cardiac surgery patients, *Crit Care Med.* 21 [2 Suppl], Seite S84-S91. URL: PM:8428503
- [41] Ott, M. J.; Buchman, T. G. und Baumgartner, W. A. (1995): Postoperative abdominal complications in cardiopulmonary bypass patients: a case-controlled study, *Ann.Thorac.Surg.* 59 [5], Seite 1210-1213. URL: PM:7733723
- [42] Braun, J. P.; Schroeder, T.; Buehner, S.; Dohmen, P.; Moshirzadeh, M.; Grosse, J.; Streit, F.; Schlaefke, A.; Armstrong, V. W.; Oellerich, M.; Lochs, H.; Konertz,

- W.; Kox, W. J. und Spies, C. (2004): Splanchnic oxygen transport, hepatic function and gastrointestinal barrier after normothermic cardiopulmonary bypass, *Acta Anaesthesiol.Scand.* 48 [6], Seite 697-703. URL: PM:15196101
- [43] Fiddian-Green, R. G. (1993): Associations between intramucosal acidosis in the gut and organ failure, *Crit Care Med.* 21 [2 Suppl], Seite S103-S107. URL: PM:8428490
- [44] Quigley, R. L.; Caplan, M. S.; Perkins, J. A.; Arentzon, C. E.; Alexander, J. C.; Kuehn, B. E.; Hoff, W. J. und Wallock, M. E. (1995): Cardiopulmonary bypass with adequate flow and perfusion pressures prevents endotoxaemia and pathologic cytokine production, *Perfusion* 10 [1], Seite 27-31. URL: PM:7795310
- [45] Riddington, D. W.; Venkatesh, B.; Boivin, C. M.; Bonser, R. S.; Elliott, T. S.; Marshall, T.; Mountford, P. J. und Bion, J. F. (1996): Intestinal permeability, gastric intramucosal pH, and systemic endotoxemia in patients undergoing cardiopulmonary bypass, *JAMA* 275 [13], Seite 1007-1012. URL: PM:8596232
- [46] Boyle, E. M., Jr.; Pohlman, T. H.; Cornejo, C. J. und Verrier, E. D. (1996): Endothelial cell injury in cardiovascular surgery: ischemia-reperfusion, *Ann.Thorac.Surg.* 62 [6], Seite 1868-1875. URL: PM:8957415
- [47] Boyle, E. M., Jr.; Pohlman, T. H.; Johnson, M. C. und Verrier, E. D. (1997): Endothelial cell injury in cardiovascular surgery: the systemic inflammatory response, *Ann.Thorac.Surg.* 63 [1], Seite 277-284. URL: PM:8993292
- [48] Landow, L. und Andersen, L. W. (1994): Splanchnic ischaemia and its role in multiple organ failure, *Acta Anaesthesiol.Scand.* 38 [7], Seite 626-639. URL: PM:7839769
- [49] Li, S.; Price, R.; Phiroz, D.; Swan, K. und Crane, T. A. (2005): Systemic inflammatory response during cardiopulmonary bypass and strategies, *J.Extra.Corpor.Technol.* 37 [2], Seite 180-188. URL: PM:16117456
- [50] Chen, Y. F.; Tsai, W. C.; Lin, C. C.; Tsai, L. Y.; Lee, C. S.; Huang, C. H.; Pan, P. C. und Chen, M. L. (2004): Effect of leukocyte depletion on endothelial cell activation and transendothelial migration of leukocytes during cardiopulmonary bypass, *Ann.Thorac.Surg.* 78 [2], Seite 634-642. URL: PM:15276536
- [51] Biffi, W. L. und Moore, E. E. (1996): Splanchnic ischaemia/reperfusion and multiple organ failure, *Br.J.Anaesth.* 77 [1], Seite 59-70. URL: PM:8703631
- [52] de, Vroege R.; te, Meerman F.; Eijnsman, L.; Wildevuur, W. R.; Wildevuur, ChR und van, Oeveren W. (2004): Induction and detection of disturbed homeostasis in cardiopulmonary bypass, *Perfusion* 19 [5], Seite 267-276. URL: PM:15508198

- [53] Maynard, N. D.; Bihari, D. J.; Dalton, R. N.; Beale, R.; Smithies, M. N. und Mason, R. C. (1997): Liver function and splanchnic ischemia in critically ill patients, *Chest* 111 [1], Seite 180-187. URL: PM:8996014
- [54] Wang, P.; Ba, Z. F. und Chaudry, I. H. (1991): Hepatic extraction of indocyanine green is depressed early in sepsis despite increased hepatic blood flow and cardiac output, *Arch.Surg.* 126 [2], Seite 219-224. URL: PM:1992997
- [55] Benedetti, M.; De, Caterina R.; Bionda, A.; Gardinali, M.; Cicardi, M.; Maffei, S.; Gazzetti, P.; Pistolesi, P.; Vernazza, F.; Michelassi, C. und . (1990): Blood--artificial surface interactions during cardiopulmonary bypass. A comparative study of four oxygenators, *Int.J.Artif.Organs* 13 [8], Seite 488-497. URL: PM:2146229
- [56] Harig, F.; Hohenstein, B.; von der, Emde J. und Weyand, M. (2001): Modulating IL-6 and IL-10 levels by pharmacologic strategies and the impact of different extracorporeal circulation parameters during cardiac surgery, *Shock* 16 Suppl 1, Seite 33-38. URL: PM:11770031
- [57] Birdi, I.; Caputo, M.; Underwood, M.; Bryan, A. J. und Angelini, G. D. (1999): The effects of cardiopulmonary bypass temperature on inflammatory response following cardiopulmonary bypass, *Eur.J.CardiThorac.Surg.* 16 [5], Seite 540-545. URL: PM:10609905
- [58] Tonz, M.; Mihaljevic, T.; von Segesser, L. K.; Schmid, E. R.; Joller-Jemelka, H. I.; Pei, P. und Turina, M. I. (1995): Normothermia versus hypothermia during cardiopulmonary bypass: a randomized, controlled trial, *Ann.Thorac.Surg.* 59 [1], Seite 137-143. URL: PM:7818312
- [59] Nagano, K.; Gelman, S.; Bradley, E. L., Jr. und Parks, D. (1990): Hypothermia, hepatic oxygen supply-demand, and ischemia-reperfusion injury in pigs, *Am.J.Physiol* 258 [6 Pt 1], Seite G910-G918. URL: PM:2360637
- [60] Okano, N.; Hiraoka, H.; Owada, R.; Fujita, N.; Kadoi, Y.; Saito, S.; Goto, F. und Morita, T. (2001): Hepatosplanchnic oxygenation is better preserved during mild hypothermic than during normothermic cardiopulmonary bypass, *Can.J.An aesth.* 48 [10], Seite 1011-1014. URL: PM:11698321
- [61] Caputo, M.; Bays, S.; Rogers, C. A.; Pawade, A.; Parry, A. J.; Suleiman, S. und Angelini, G. D. (2005): Randomized comparison between normothermic and hypothermic cardiopulmonary bypass in pediatric open-heart surgery, *Ann.Thorac.Surg.* 80 [3], Seite 982-988. URL: PM:16122470
- [62] Grunenfelder, J.; Zund, G.; Schoeberlein, A.; Schmid, E. R.; Schurr, U.; Frisullo, R.; Maly, F. und Turina, M. (2000): Expression of adhesion molecules and cytokines after coronary artery bypass grafting during normothermic and

- hypothermic cardiac arrest, *Eur.J.Cardiothorac.Surg.* 17 [6], Seite 723-728. URL: PM:10856867
- [63] Speziale, G.; Ferroni, P.; Ruvolo, G.; Fattouch, K.; Pulcinelli, F. M.; Lenti, L.; Gazzaniga, P. P. und Marino, B. (2000): Effect of normothermic versus hypothermic cardiopulmonary bypass on cytokine production and platelet function, *J.Cardiovasc.Surg.(Torino)* 41 [6], Seite 819-827. URL: PM:11232964
- [64] Hampton, W. W.; Townsend, M. C.; Schirmer, W. J.; Haybron, D. M. und Fry, D. E. (1989): Effective hepatic blood flow during cardiopulmonary bypass, *Arch.Surg.* 124 [4], Seite 458-459. URL: PM:2930354
- [65] Mathie, R. T.; Ohri, S. K.; Batten, J. J.; Peters, A. M. und Keogh, B. E. (1997): Hepatic blood flow during cardiopulmonary bypass operations: the effect of temperature and pulsatility, *J.Thorac.Cardiovasc.Surg.* 114 [2], Seite 292-293. URL: PM:9270653
- [66] DeFoe, G. R.; Krumholz, C. F.; DioDato, C. P.; Ross, C. S.; Olmstead, E. M.; Groom, R. C.; Pieroni, J. W.; Forest, R. J.; O'Connor, B. R.; Bogosian, M. E.; Warren, C. S. und O'Connor, G. T. (2003): Lowest core body temperature and adverse outcomes associated with coronary artery bypass surgery, *Perfusion* 18 [2], Seite 127-133. URL: PM:12868791
- [67] Neuhof, C.; Wendling, J.; Dapper, F.; Bauer, J.; Zickmann, B.; Jochum, M.; Tillmanns, H. und Neuhoft, H. (2001): Endotoxemia and cytokine generation in cardiac surgery in relation to flow mode and duration of cardiopulmonary bypass, *Shock* 16 Suppl 1, Seite 39-43. URL: PM:11770032
- [68] Whitten, C. W.; Hill, G. E.; Ivy, R.; Greilich, P. E. und Lipton, J. M. (1998): Does the duration of cardiopulmonary bypass or aortic cross-clamp, in the absence of blood and/or blood product administration, influence the IL-6 response to cardiac surgery?, *Anesth.Analg.* 86 [1], Seite 28-33. URL: PM:9428846
- [69] Kennergren, C.; Mantovani, V.; Lonroth, P.; Nystrom, B.; Berglin, E. und Hamberger, A. (1999): Monitoring of extracellular aspartate aminotransferase and troponin T by microdialysis during and after cardioplegic heart arrest, *Cardiology* 92 [3], Seite 162-170. URL: PM:10754346
- [70] Redick, J. A.; Jakoby, W. B. und Baron, J. (1982): Immunohistochemical localization of glutathione S-transferases in livers of untreated rats, *J.Biol.Chem.* 257 [24], Seite 15200-15203. URL: PM:7174691
- [71] Beckett, G. J.; Dyson, E. H.; Chapman, B. J.; Templeton, A. J. und Hayes, J. D. (1985): Plasma glutathione S-transferase measurements by radioimmunoassay: a sensitive index of hepatocellular damage in man, *Clin.Chim.Acta* 146 [1], Seite 11-19. URL: PM:3987035

- [72] Beckett, G. J. und Hayes, J. D. (1993): Glutathione S-transferases: biomedical applications, *Adv.Clin.Chem.* 30, Seite 281-380. URL: PM:8237562
- [73] Jakob, S. M.; Ruokonen, E. und Takala, J. (2000): Assessment of the adequacy of systemic and regional perfusion after cardiac surgery, *Br.J.Anaesth.* 84 [5], Seite 571-577. URL: PM:10844831
- [74] Autschbach, R.; Falk, V.; Lange, H.; Oellerich, M.; Walther, T.; Mohr, F. W. und Dalichau, H. (1996): Assessment of metabolic liver function and hepatic blood flow during cardiopulmonary bypass, *Thorac.Cardiovasc.Surg.* 44 [2], Seite 76-80. URL: PM:8782332
- [75] Bradley, S. E.; CHILDS, A. W.; COMBES, B.; COURNAND, A.; WADE, O. L. und Wheeler, H. O. (1956): The effect of exercise on the splanchnic blood flow and splanchnic blood volume in normal man, *Clin.Sci.(Lond)* 15 [3], Seite 457-463. URL: PM:13356576
- [76] Bradley, S. E. (1960): Estimation of hepatic blood flow, *Methods Med.Res.* 8, Seite 275-283. URL: PM:13803722
- [77] Kramer, W. G. und Romagnoli, A. (1986): Effect of surgery and cardiopulmonary bypass on indocyanine green pharmacokinetics, *Tex.Heart Inst.J.* 13 [1], Seite 77-82. URL: PM:15226835
- [78] LEEVY, C. M.; MENDENHALL, C. L.; LESKO, W. und HOWARD, M. M. (1962): Estimation of hepatic blood flow with indocyanine green, *J.Clin.Invest* 41, Seite 1169-1179. URL: PM:14463639
- [79] Uusaro, A.; Ruokonen, E. und Takala, J. (1995): Estimation of splanchnic blood flow by the Fick principle in man and problems in the use of indocyanine green, *Cardiovasc.Res.* 30 [1], Seite 106-112. URL: PM:7553712
- [80] Lehmann, U.; Armstrong, V. W.; Schutz, E.; Regel, G.; Pape, D. und Oellerich, M. (1995): Monoethylglycinexylidide as an early predictor of posttraumatic multiple organ failure, *Ther.Drug Monit.* 17 [2], Seite 125-132. URL: PM:7624899
- [81] Oellerich, M. und Armstrong, V. W. (2001): The MEGX test: a tool for the real-time assessment of hepatic function, *Ther.Drug Monit.* 23 [2], Seite 81-92. URL: PM:11294522
- [82] Schroter, J.; Wandel, C.; Bohrer, H.; Schmidt, H.; Bottiger, B. W. und Martin, E. (1995): Lignocaine metabolite formation: an indicator for liver dysfunction and predictor of survival in surgical intensive care patients, *Anaesthesia* 50 [10], Seite 850-854. URL: PM:7485872
- [83] Demers, P.; Elkouri, S.; Martineau, R.; Couturier, A. und Cartier, R. (2000): Outcome with high blood lactate levels during cardiopulmonary bypass in adult



- cardiac operation, *Ann.Thorac.Surg.* 70 [6], Seite 2082-2086. URL: PM:11156124
- [84] Tao, W.; Zwischenberger, J. B.; Nguyen, T. T.; Vertrees, R. A.; McDaniel, L. B.; Nutt, L. K.; Herndon, D. N. und Kramer, G. C. (1995): Gut mucosal ischemia during normothermic cardiopulmonary bypass results from blood flow redistribution and increased oxygen demand, *J.Thorac.Cardiovasc.Surg.* 110 [3], Seite 819-828. URL: PM:7564451
- [85] Toraman, F.; Evrenkaya, S.; Yuce, M.; Aksoy, N.; Karabulut, H.; Bozkulak, Y. und Alhan, C. (2004): Lactic acidosis after cardiac surgery is associated with adverse outcome, *Heart Surg.Forum* 7 [2], Seite E155-E159. URL: PM:15138095
- [86] Kox, W. J. und C.D.Spies (2005): *Check-up Anästhesiologie, 2., erw. und aktual. Aufl. Auflage*, Springer, ISBN: 3-540-23093-9.
- [87] Branthwaite, M. A. und Bradley, R. D. (1968): Measurement of cardiac output by thermal dilution in man, *J Appl.Physiol* 24 [3], Seite 434-438. URL: PM:5640735
- [88] Ganz, W.; Tamura, K.; Marcus, H. S.; Donoso, R.; Yoshida, S. und Swan, H. J. (1971): Measurement of coronary sinus blood flow by continuous thermodilution in man, *Circulation* 44 [2], Seite 181-195. URL: PM:4935053
- [89] Ganz, W.; Donoso, R.; Marcus, H. S.; Forrester, J. S. und Swan, H. J. (1971): A new technique for measurement of cardiac output by thermodilution in man, *Am.J Cardiol.* 27 [4], Seite 392-396. URL: PM:4929422
- [90] Hoeft, A (1995): *Dilutionstechniken und Ficksches Prinzip*, Springer Verlag 1995.
- [91] Chang, G. W. und Kam, P. C. (1999): The physiological and pharmacological roles of cytochrome P450 isoenzymes, *Anaesthesia* 54 [1], Seite 42-50. URL: PM:10209369
- [92] Reichel, C.; Skodra, T.; Nacke, A.; Spengler, U. und Sauerbruch, T. (1998): The lignocaine metabolite (MEGX) liver function test and P-450 induction in humans, *Br.J.Clin.Pharmacol.* 46 [6], Seite 535-539. URL: PM:9862241
- [93] Thomson, A. H.; Elliott, H. L.; Kelman, A. W.; Meredith, P. A. und Whiting, B. (1987): The pharmacokinetics and pharmacodynamics of lignocaine and MEGX in healthy subjects, *J.Pharmacokinet.Biopharm.* 15 [2], Seite 101-115. URL: PM:3612496
- [94] Bennett, P. N.; Aarons, L. J.; Bending, M. R.; Steiner, J. A. und Rowland, M. (1982): Pharmacokinetics of lidocaine and its deethylated metabolite: dose and time dependency studies in man, *J Pharmacokinet.Biopharm.* 10 [3], Seite 265-281. URL: PM:7175699

- [95] Buchta, M.; Hoffmann, A.; Sönnichsen, A. und Wolff, A. (2006): Pharmakologie, Hammerexamen , 1. Auflage, Seite 1536-1727, Elsevier GmbH, Urban & Fischer Verlag, München, ISBN: 987-3-437-43800-4.
- [96] Scharpf, O. und Sibold, M. (2004): Modellieren mit Mathematik, Bericht zur 1. Arbeitsphase "Medikamente" des gleichnamigen Seminars an der TU Berlin im SS 2004, 1. Auflage , Seminar, TU-Berlin. URL: <http://www.math.tu-berlin.de>
- [97] Thomson, P. D.; Melmon, K. L.; Richardson, J. A.; Cohn, K.; Steinbrunn, W.; Cudihee, R. und Rowland, M. (1973): Lidocaine pharmacokinetics in advanced heart failure, liver disease, and renal failure in humans, *Ann.Intern.Med.* 78 [4], Seite 499-508. URL: PM:4694036
- [98] Habler, O.; Kleen, M.; Kemming, G. und Zwissler, B. (2002): Hyperoxia in extreme hemodilution, *Eur.Surg.Res.* 34 [1-2], Seite 181-187. URL: PM:11867921
- [99] Johnson, L. B.; Plotkin, J. S. und Kuo, P. C. (1998): Reduced transfusion requirements during major hepatic resection with use of intraoperative isovolemic hemodilution, *Am.J.Surg.* 176 [6], Seite 608-611. URL: PM:9926799
- [100] Nollert, G.; Sperling, J.; Sakamoto, T.; Jaeger, B. R. und Jonas, R. A. (2001): Higher hematocrit improves liver blood flow and metabolism during cardiopulmonary bypass in piglets, *Thorac.Cardiovasc.Surg.* 49 [4], Seite 226-230. URL: PM:11505319
- [101] Noldge, G. F.; Priebe, H. J.; Bohle, W.; Buttler, K. J. und Geiger, K. (1991): Effects of acute normovolemic hemodilution on splanchnic oxygenation and on hepatic histology and metabolism in anesthetized pigs, *Anesthesiology* 74 [5], Seite 908-918. URL: PM:1708653
- [102] Chouker, A.; Martignoni, A.; Schauer, R. J.; Dugas, M.; Schachtner, T.; Kaufmann, I.; Setzer, F.; Rau, H. G.; Lohe, F.; Jauch, K. W.; Peter, K. und Thiel, M. (2005): Alpha-gluthathione S-transferase as an early marker of hepatic ischemia/reperfusion injury after liver resection, *World J.Surg.* 29 [4], Seite 528-534. URL: PM:15776301
- [103] Strom, S.; Bendz, R.; Olin, C. und Lundberg, S. (1979): Serum enzymes with special reference to CK-MB following coronary bypass surgery, *Scand.J Thorac.Cardiovasc.Surg.* 13 [1], Seite 53-59. URL: PM:311943
- [104] Olsson, R.; Hermodsson, S.; Roberts, D. und Waldenstrom, J. (1984): Hepatic dysfunction after open-heart surgery, *Scand.J Thorac.Cardiovasc.Surg.* 18 [3], Seite 217-222. URL: PM:6152078
- [105] randa-Michel, J. und Sherman, K. E. (1998): Tests of the liver: use and misuse, *Gastroenterologist.* 6 [1], Seite 34-43. URL: PM:9531115

- [106] Habler, O.; Kleen, M.; Hutter, J.; Podtschaske, A.; Tiede, M.; Kemming, G.; Corso, C.; Batra, S.; Keipert, P.; Faithfull, S. und Messmer, K. (1997): Effects of hemodilution on splanchnic perfusion and hepatorenal function. II. Renal perfusion and hepatorenal function, *Eur.J Med Res.* 2 [10], Seite 419-424. URL: PM:9348268
- [107] Strom, S.; Bendz, R.; Olin, C. und Lundberg, S. (1979): Serum enzymes with special reference to CK-MB following coronary bypass surgery, *Scand.J Thorac.Cardiovasc.Surg.* 13 [1], Seite 53-59. URL: PM:311943
- [108] Tiainen, P.; Lindgren, L. und Rosenberg, P. H. (1995): Disturbance of hepatocellular integrity associated with propofol anaesthesia in surgical patients, *Acta Anaesthesiol.Scand.* 39 [6], Seite 840-844. URL: PM:7484045
- [109] Murray, J. M.; Phillips, A. S. und Fee, J. P. (1994): Comparison of the effects of isoflurane and propofol on hepatic glutathione-S-transferase concentrations during and after prolonged anaesthesia, *Br.J.Anaesth.* 72 [5], Seite 599-601. URL: PM:8198917
- [110] Tiainen, P.; Lindgren, L. und Rosenberg, P. H. (1995): Disturbance of hepatocellular integrity associated with propofol anaesthesia in surgical patients, *Acta Anaesthesiol.Scand.* 39 [6], Seite 840-844. URL: PM:7484045
- [111] Tiainen, P.; Lindgren, L. und Rosenberg, P. H. (1998): Changes in hepatocellular integrity during and after desflurane or isoflurane anaesthesia in patients undergoing breast surgery, *Br.J Anaesth.* 80 [1], Seite 87-89. URL: PM:9505785
- [112] Koo, D. J.; Zhou, M.; Chaudry, I. H. und Wang, P. (2000): Plasma alpha-glutathione S-transferase: a sensitive indicator of hepatocellular damage during polymicrobial sepsis, *Arch.Surg.* 135 [2], Seite 198-203. URL: PM:10668881
- [113] Beckett, G. J.; Chapman, B. J.; Dyson, E. H. und Hayes, J. D. (1985): Plasma glutathione S-transferase measurements after paracetamol overdose: evidence for early hepatocellular damage, *Gut* 26 [1], Seite 26-31. URL: PM:3965363
- [114] Trull, A. K.; Facey, S. P.; Rees, G. W.; Wight, D. G.; Noble-Jamieson, G.; Joughin, C.; Friend, P. J. und Alexander, G. J. (1994): Serum alpha-glutathione S-transferase--a sensitive marker of hepatocellular damage associated with acute liver allograft rejection, *Transplantation* 58 [12], Seite 1345-1351. URL: PM:7809927
- [115] Mathie, R. T. (1993): Hepatic blood flow during cardiopulmonary bypass, *Crit Care Med.* 21 [2 Suppl], Seite S72-S76. URL: PM:8428501
- [116] Tiainen, P.; Lindgren, L. und Rosenberg, P. H. (1995): Disturbance of hepatocellular integrity associated with propofol anaesthesia in surgical patients, *Acta Anaesthesiol.Scand.* 39 [6], Seite 840-844. URL: PM:7484045

- [117] Schmidt, C. C.; Suttner, S. W.; Piper, S. N.; Nagel, D. und Boldt, J. (1999): Comparison of the effects of desflurane and isoflurane anaesthesia on hepatocellular function assessed by alpha glutathione S-transferase, *Anaesthesia* 54 [12], Seite 1207-1211. URL: PM:10594421
- [118] Granados Llamas, L. M.; Navas, Rivera E.; Marengo de la Fuente ML; Balanza, Ortiz R.; Suarez, Collazos F. und Puente Egido, J. J. (1999): [Mobilization of alpha-glutathione S-transferase in the anesthetized patient], *Rev.Esp.Anesthesiol.Reanim.* 46 [8], Seite 350-353. URL: PM:10563141
- [119] Tiainen, P. und Rosenberg, P. H. (2000): Glutathione transferase alpha and inhalation anaesthetics, *Anaesthesia* 55 [9], Seite 931-932. URL: PM:10947785
- [120] FOX, I. J.; BROOKER, L. G.; HESELTINE, D. W.; ESSEX, H. E. und WOOD, E. H. (1957): A tricarbo-cyanine dye for continuous recording of dilution curves in whole blood independent of variations in blood oxygen saturation, *Mayo Clin.Proc.* 32 [18], Seite 478-484. URL: PM:13465828
- [121] WIEGAND, B. D.; KETTERER, S. G. und Rapaport, E. (1960): The use of indocyanine green for the evaluation of hepatic function and blood flow in man, *Am.J.Dig.Dis.* 5, Seite 427-436. URL: PM:13844597
- [122] Pollack, D. S.; Sufian, S. und Matsumoto, T. (1979): Indocyanine green clearance in critically ill patients, *Surg.Gynecol.Obstet.* 149 [6], Seite 852-854. URL: PM:388703
- [123] Kimura, S.; Yoshioka, T.; Shibuya, M.; Sakano, T.; Tanaka, R. und Matsuyama, S. (2001): Indocyanine green elimination rate detects hepatocellular dysfunction early in septic shock and correlates with survival, *Crit Care Med.* 29 [6], Seite 1159-1163. URL: PM:11395594
- [124] Maynard, N. D.; Bihari, D. J.; Dalton, R. N.; Smithies, M. N. und Mason, R. C. (1995): Increasing splanchnic blood flow in the critically ill, *Chest* 108 [6], Seite 1648-1654. URL: PM:7497776
- [125] Sakka, SG und Meier-Hellmann, A (2001): Indocyanin Green for Assessment of liver function in Critically Ill Patients, *Yearbook of Intensive Care and Emergency Medicine* 2001 , Berlin-Heidelberg-New York: Springer Verlag.
- [126] Sakka, S. G.; Reinhart, K. und Meier-Hellmann, A. (2002): Prognostic value of the indocyanine green plasma disappearance rate in critically ill patients, *Chest* 122 [5], Seite 1715-1720. URL: PM:12426276
- [127] Kholoussy, A. M.; Pollack, D. und Matsumoto, T. (1984): Prognostic significance of indocyanine green clearance in critically ill surgical patients, *Crit Care Med.* 12 [2], Seite 115-116. URL: PM:6697728

- [128] Wynne, H. A.; Cope, L. H.; Mutch, E.; Rawlins, M. D.; Woodhouse, K. W. und James, O. F. (1989): The effect of age upon liver volume and apparent liver blood flow in healthy man, *Hepatology* 9 [2], Seite 297-301. URL: PM:2643548
- [129] Wynne, H. A. und James, O. F. (1990): The ageing liver, *Age Ageing* 19 [1], Seite 1-3. URL: PM:2316418
- [130] Zotz, R. B.; von, Schonfeld J.; Erhard, J.; Breuer, N.; Lange, R.; Beste, M.; Goebell, H. und Eigler, F. W. (1997): Value of an extended monoethylglycinexylidide formation test and other dynamic liver function tests in liver transplant donors, *Transplantation* 63 [4], Seite 538-541. URL: PM:9047147
- [131] Haisjackl, M.; Birnbaum, J.; Redlin, M.; Schmutzler, M.; Waldenberger, F.; Lochs, H.; Konertz, W. und Kox, W. (1998): Splanchnic oxygen transport and lactate metabolism during normothermic cardiopulmonary bypass in humans, *Anesth.Analg.* 86 [1], Seite 22-27. URL: PM:9428845
- [132] Koo, D. J.; Chaudry, I. H. und Wang, P. (1999): Kupffer cells are responsible for producing inflammatory cytokines and hepatocellular dysfunction during early sepsis, *J.Surg.Res.* 83 [2], Seite 151-157. URL: PM:10329110
- [133] Chetty, G.; Sharpe, D. A.; Nandi, J.; Butler, S. J. und Mitchell, I. M. (2004): Liver blood flow during cardiac surgery, *Perfusion* 19 [3], Seite 153-156. URL: PM:15298422
- [134] Potter, J. M.; Hickman, P. E.; Henderson, A.; Balderson, G. A.; Lynch, S. V. und Strong, R. W. (1996): The use of the lidocaine-monoethylglycinexylidide test in the liver transplant recipient, *Ther.Drug Monit.* 18 [4], Seite 383-387. URL: PM:8857555
- [135] Conti, F.; Dousset, B.; Cherruau, B.; Guerin, C.; Soubrane, O.; Houssin, D. und Calmus, Y. (2004): Use of lidocaine metabolism to test liver function during the long-term follow-up of liver transplant recipients, *Clin.Transplant.* 18 [3], Seite 235-241. URL: PM:15142042
- [136] Nies, A. S.; Shand, D. G. und Wilkinson, G. R. (1976): Altered hepatic blood flow and drug disposition, *Clin.Pharmacokinet.* 1 [2], Seite 135-155. URL: PM:13954
- [137] Berkenstadt, H.; Segal, E.; Mayan, H.; Almog, S.; Rotenberg, M.; Perel, A. und Ezra, D. (1999): The pharmacokinetics of morphine and lidocaine in critically ill patients, *Intensive Care Med.* 25 [1], Seite 110-112. URL: PM:10051088
- [138] Hein, O. V.; Ohring, R.; Schilling, A.; Oellerich, M.; Armstrong, V. W.; Kox, W. J. und Spies, C. (2004): N-acetylcysteine decreases lactate signal intensities in liver tissue and improves liver function in septic shock patients, as shown by magnetic resonance spectroscopy: extended case report, *Crit Care* 8 [2], Seite R66-R71. URL: PM:15025780

- [139] Jakob, S. M.; Ruokonen, E.; Rosenberg, P. H. und Takala, J. (2002): Effect of dopamine-induced changes in splanchnic blood flow on MEGX production from lidocaine in septic and cardiac surgery patients, *Shock* 18 [1], Seite 1-7. URL: PM:12095126
- [140] Andersen, L. W.; Landow, L.; Baek, L.; Jansen, E. und Baker, S. (1993): Association between gastric intramucosal pH and splanchnic endotoxin, antibody to endotoxin, and tumor necrosis factor-alpha concentrations in patients undergoing cardiopulmonary bypass, *Crit Care Med.* 21 [2], Seite 210-217. URL: PM:8428471
- [141] Mustafa, I.; Roth, H.; Hanafiah, A.; Hakim, T.; Anwar, M.; Siregar, E. und Leverage, X. M. (2003): Effect of cardiopulmonary bypass on lactate metabolism, *Intensive Care Med.* 29 [8], Seite 1279-1285. URL: PM:12845428
- [142] Mizock, B. A. und Falk, J. L. (1992): Lactic acidosis in critical illness, *Crit Care Med* 20 [1], Seite 80-93. URL: PM:1309494
- [143] Ariza, M.; Gothard, J. W.; Macnaughton, P.; Hooper, J.; Morgan, C. J. und Evans, T. W. (1991): Blood lactate and mixed venous-arterial PCO<sub>2</sub> gradient as indices of poor peripheral perfusion following cardiopulmonary bypass surgery, *Intensive Care Med* 17 [6], Seite 320-324. URL: PM:1755880
- [144] Koyama, K.; Okuyama, S.; Fukuyama, T.; Okamoto, Y.; Kawasaki, J. und Miyao, H. (2003): [Blood lactate levels during fast-track cardiac anesthesia], *Masui* 52 [11], Seite 1191-1194. URL: PM:14661564
- [145] Cain, S. M. und Curtis, S. E. (1991): Systemic and regional oxygen uptake and delivery and lactate flux in endotoxic dogs infused with dopexamine, *Crit Care Med.* 19 [12], Seite 1552-1560. URL: PM:1959377
- [146] Tsunooka, N.; Hamada, Y.; Imagawa, H.; Nakamura, Y.; Shiozaki, T.; Suzuki, H.; Kikkawa, H.; Miyauchi, K.; Watanabe, Y. und Kawachi, K. (2003): Ischemia of the intestinal mucosa during cardiopulmonary bypass, *J.Artif.Organs* 6 [2], Seite 149-151. URL: PM:14621695
- [147] Hochleitner, M. und Bader, A. (2003): [Gender differences in cardiology], *Acta Med Austriaca* 30 [3], Seite 69-71. URL: PM:14671822