

7. Literaturverzeichnis

- Aebersold R., Leavitt J., Saavedra R. A., Hood L. E., and Kent S. B. (1987): Internal amino acid sequence analysis of proteins separated by one- or two-dimensional gel electrophoresis after in situ protease digestion on nitrocellulose. *Proceedings of the National Academy of Science*; 84: 6970-6974
- Aebersold R., and Mann M. (2003): Mass spectrometry-based proteomics. *Nature*; 422: 198-207
- Akin D. T., Shapira R., and Kinkade Jr. J. M. (1985): The determination of molecular weights of biologically active proteins by cetyltrimethylammonium bromid-polyacrylamide gel electrophoresis. *Analytical Biochemistry*; 145: 170-176
- Akins R., Levin P., and Tuan R. (1992): Cetyltrimethylammonium bromide discontinuous gel electrophoresis: Mr-based separation of proteins with retention of enzymatic activity. *Analytical Biochemistry*; 202: 172-178
- Berton O., and Nestler E. J. (2006): New approaches to antidepressant drug discovery: beyond monoamides. *Nature Reviews Neuroscience*; 7: 137-151
- Bjellquist B., Ek K., Righetti P. G., et al. (1982): Isoelectric focusing in immobilized pH gradients: principle, methodology and some applications. *Journal of Biochemical and Biophysical Methods*; 6: 317-339
- Blomberg A., Blomberg L., Norbeck J., et al. (1995): Interlaboratory reproducibility of yeast protein patterns analyzed by immobilized pH gradient two-dimensional gel electrophoresis. *Electrophoresis*; 16: 1935-1945
- Blondeau F., Ritter B., Allaire P. D., et al. (2004): Tandem MS analysis of brain clathrin-coated vesicles reveals their critical involvement in synaptic vesicle recycling. *Proceedings of the National Academy of Sciences of the United States of America*; 101: 3833-3838
- Bradford M. M. (1979): A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*; 72: 248-254
- Braun J. E. A., and Madison D. V. (2000): A novel SNAP25-caveolin complex correlates with the onset of persistent synaptic potentiation. *The Journal of Neuroscience*; 20: 5997-6006
- Broedel O., Eravci M., Fuxius S., et al. (2003): Effects of hyper- and hypothyroidism on thyroid hormone concentrations in regions of the rat brain. *American Journal of Physiology Endocrinology and Metabolism*; 285: E470-E480
- Buxbaum E. (2003): Cationic electrophoresis and electrotransfer of membrane glycoproteins. *Analytical Biochemistry*; 314: 70-76

- Campostrini N., Areces L. B., Rappsilber J., et al. (2005): Spot overlapping in two-dimensional maps: A serious problem ignored for much too long. *Proteomics*; 5: 2385-2395
- Cao R., Li X., Peng X., et al. (2006): Integration of a two-phase partition method into proteomics research on rat liver membrane proteins. *Journal of proteome research*; 5: 634-642
- Carboni L., Puibelli C., Righetti P. G., Jansson B., and Domenici E. (2002): Proteomic analysis of rat brain tissue: Comparison of protocols for two-dimensional gel electrophoresis analysis based on different solubilizing agents. *Electrophoresis*; 23: 4132-4141
- Carrette O., Burkhard P. R., Hochstrasser D. F., and Sanchez J. C. (2006): Age-related proteome analysis of the mouse brain: A 2-DE study. *Proteomics*; 6: 4940-4949
- Challapalli K. K., Zabel C., Schuchhardt J., et al. (2004): High reproducibility of large-gel two-dimensional electrophoresis. *Electrophoresis*; 25: 3040-3047
- Cole R. N., and Hart G. W. (2001): Cytosolic O-glycosylation is abundant in nerve terminals. *Journal of Neurochemistry*; 79: 1080-1089
- Coughenour H. D., Spaulding R. S., and Thompson A. (2004): The synaptic vesicle proteome: A comparative study in membrane protein identification. *Proteomics*; 4: 3141-3155
- Cramer J. A., Marchesi V. T., and Armitage I. M. (1980): Reversible trifluoroacetic acid-induced conformational changes in glycophorin A as detected by proton nuclear magnetic resonance spectroscopy. *Biochimica et Biophysica Acta*; 595: 235-243
- de Vocht M. L., Scholtmeijer K., van der Vegte E. W., et al. (1998): Structural characterization of the hydrophobin SC3, as a monomer and after self-assembly at hydrophobic/hydrophilic interfaces. *Biophysical Journal*; 74: 2059-2068
- Deshusses J. M., Burgess J. A., Scherl A., et al. (2003): Exploitation of specific properties of trifluoroethanol for extraction and separation of membrane proteins. *Proteomics*; 3: 1418-1424
- Dodd P. R., Hardy J. A., Oakley A. E., and Strong A. J. (1981): Synaptosomes prepared from fresh human cerebral cortex; morphology, respiration and release of transmitter amino acids. *Brain Research*; 224: 419-425
- Domon B., and Aebersold R. (2006): Mass spectrometry and protein analysis. *Science*; 312: 212-217
- Edgar P. F., Douglas J. E., Cooper G. J. S., et al. (2000): Comparative proteome analysis of the hippocampus implicates chromosome 6q in schizophrenia. *Molecular Psychiatry*; 5: 85-90

- Eichacker L. A., Granvogl B., Mirus O., et al. (2004): Hiding behind hydrophobicity. *The Journal of Biological Chemistry*; 279: 50915-50922
- Enard W., Khaitovich P., Klose J., et al. (2002): Intra- and interspecific variation in primate gene expression patterns. *Science*; 296: 340-343
- Everberg H., Sivars U., Emanuelsson C., et al. (2004): Protein pre-fractionation in detergent-polymer aqueous two-phase systems for facilitated proteomic studies of membrane proteins. *Journal of Chromatography A*; 1029: 113-124
- Ferro M., Salvi D., Riviere-Rolland H., et al. (2002): Integral membrane proteins of the chloroplast envelope: Identification and subcellular localization of new transporters. *Proceedings of the National Academy of Science*; 99: 11487-11492
- Fountoulakis M., Juranville J.-F., Dierssen M., and Lubec G. (2002): Proteomic analysis of the fetal brain. *Proteomics*; 2: 1539-1546
- Friso G., and Wikström L. (1999): Analysis of proteins from membrane-enriched cerebellar preparations by two-dimensional gel electrophoresis and mass spectrometry. *Electrophoresis*; 20: 917-927
- Fröhlich T., Helmstetter D., Zobawa M., et al. (2006): Analysis of the HUP0 Brain Proteome reference samples using 2-D DIGE and 2-D LC-MS/MS. *Proteomics*; 6: 4950-4966
- Fujiki Y., Hubbard A. J., Fowler S., and Lazarow P. B. (1982): Isolation of intracellular membranes by means of sodium carbonate treatment: application to endoplasmic reticulum. *The Journal of Cell Biology*; 93: 97-102
- Gauss C., Kalkum M., Löwe M., Lehrach H., and Klose J. (1999): Analysis of the mouse proteome. (I) Brain proteins: Separation by two-dimensional electrophoresis and identification by mass spectrometry and genetic variation. *Electrophoresis*; 20: 575-600
- Gendreau S., Voswinkel S., Torres-Salazar D., et al. (2004): A trimeric quaternary structure is conserved in bacterial and human glutamate transporters. *The Journal of Biological Chemistry*; 279: 39505-39512
- Glowinski J., and Iversen L. L. (1966): Regional studies of catecholamine metabolism in the rat brain, I. The disposition of (3H) dopamin and (3H) dopa in various regions of the brain. *Journal of Neurochemistry*; 13: 655-669
- Görg A., Postel W., Gunther S., et al. (1988): Approach to stationary two-dimensional pattern: Influence of focusing time and immobiline/carrier ampholytes concentrations. *Electrophoresis*; 9: 37-46
- Görg A., Obermaier C., Boguth G., et al. (1997): Very alkaline immobilize pH gradients for two-dimensional electrophoresis of ribosomal and nuclear proteins. *Electrophoresis*; 18: 328-337

- Görg A., Obermaier C., Boguth G., et al. (2000): The current state of two-dimensional electrophoresis with immobilized pH gradients. *Electrophoresis*; 21: 1037-1053
- Görg A., Weiss W., and Dunn M. (2004): Current two-dimensional electrophoresis technology for proteomics. *Proteomics*; 4: 3665-3685
- Grayson S., and Sequeira S. M. (1990): Dispersal of proteolipid macroaggregates with trifluoroacetic acid and analysis by sodium dodecyl sulfate polyacrylamide gel electrophoresis. *Analytical Biochemistry*; 189: 192-196
- Gygi S. P., Rist B., Gerber S. A., et al. (1999a): Quantitative analysis of complex protein mixtures using isotope-coded affinity tags. *Nature Biotechnology*; 17: 994-999
- Gygi S. P., Rochon Y., Franza R., and Aebersold R. (1999b): Correlation between protein and mRNA abundance in yeast. *Molecular and Cell Biology*; 19: 1720-1730
- Gygi S. P., Corthals G. L., Zhang Y., Rochon Y., and Aebersold R. (2000): Evaluation of two-dimensional gel electrophoresis-based proteome analysis technology. *Proceedings of the National Academy of Science*; 97: 9390-9395
- Hartinger J., Stenius K., Hogemann D., and Jahn R. (1996): 16-BAC/SDS-PAGE: a two-dimensional gel electrophoresis system suitable for the separation of integral membrane proteins. *Analytical Biochemistry*; 240: 126-133
- Heinz C., Engelhardt H., and Niederweis M. (2003): The core of the tetrameric mycobacterial porin MspA is an extremely stable β -sheet domain. *The Journal of Biological Chemistry*; 278: 8678-8685
- Hennessey J. P., Jr., and Scarborough G. A. (1989): An optimized procedure for sodium dodecyl sulfate-electrophoresis analysis of hydrophobic peptides from an integral membrane protein. *Analytical Biochemistry*; 176: 284-289
- Herbert B. R., Molloy M. P., Gooley A. A., et al. (1998): Improved protein solubility in two-dimensional electrophoresis using tributyl phosphine as reducing agent. *Electrophoresis*; 19: 845-851
- Ho Y., Gruhler A., Heilbut A., et al. (2002): Systematic identification of protein complexes in *Saccharomyces cerevisiae* by mass spectrometry. *Nature*; 415: 180-183
- Hoving S., Gerrits B., Voshol H., et al. (2002): Preparative two-dimensional gel electrophoresis at alkaline pH using narrow range immobilized pH gradients. *Proteomics*; 2: 127-134
- Ilias Y., Rudaz S., Mathieu P., Christen P., and J.L. V. (2005): Extraction and analysis of different Cannabis samples by headspace solid-phase microextraction combined with gas chromatography-mass spectrometry. *Journal of Sep Science*; 17: 2293-2300

- Jimenez C. R., Eyman M., Lavina Z. S., et al. (2002): Protein synthesis in synaptosomes: a proteomics analysis. *Journal of Neurochemistry*; 81: 735-744
- Julka S., and Regnier F. (2004): Quantification in proteomics through stable isotope coding: A review. *Journal of Proteome Research*; 3: 350-363
- Karp N. A., Kreil D. P., and Lilley K. S. (2004): Determining a significant change in protein expression with DeCyder during a pair-wise comparison using two-dimensional difference gel electrophoresis. *Proteomics*; 4: 1421-1432
- Karp N. A., and Lilley K. S. (2005): Maximising sensitivity for detecting changes in protein expression: Experimental design using minimal CyDyes. *Proteomics*; 5: 3105-3115
- Klein C., Garcia-Rizo C., Bisle B., et al. (2005): The membrane proteome of *Halobacterium salinarium*. *Proteomics*; 5: 180-197
- Klose J. (1975): Protein mapping by combined isoelectric focusing and electrophoresis of mouse tissues. A novel approach to testing for induced point mutations in mammals. *Humangenetik*; 26: 231-243
- Klose J. (1999): Large-gel 2D electrophoresis. *Methods in Molecular Biology*; 112: 147-172
- Klose J., Nock C., Herrmann M., et al. (2002): Genetic analysis of the mouse brain proteome. *Nature Genetics*; 30: 385-393
- Kuhn K., Prinz T., Schäfer J., et al. (2005): Protein sequence tags: A novel solution for comparative proteomics. *Proteomics*; 5: 2364-2368
- Laemmli U. K. (1970): Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*; 227: 680-685
- Li X.-M., Patel B. B., Blagoi E. I., et al. (2004): Analyzing alkaline proteins in human colon crypt proteome. *Journal of Proteome Research*; 3: 821-833
- Liberatori S., Canas B., Tani C., et al. (2004): Proteomic approach to the identification of voltage-dependent anion channel protein isoforms in guinea pig brain synaptosomes. *Proteomics*; 4: 1335-1340
- Llinás M., De Marco A., and Lecomte J. T. J. (1980): Proton magnetic resonance study of crambin, a hyperstable hydrophobic protein. *Biochemistry*; 19: 1140-1145
- Lovtrup-Rein H., and McEwen B. S. (1966): Isolation and fractionation of rat brain nuclei. *Journal of Cell Biology*; 30: 405-415
- Lowry O. H., Rosbrough N. J., Farr A. L., and Randall R. J. (1951): Protein measurement with the folin phenol reagent.

- MacCoss M. J., MacDonald W. H., Saraf A., et al. (2002): Shotgun identification of protein modification from protein complexes and lens tissue. *Proceedings of the National Academy of Science*; 99: 7900-7905
- MacFarlane D. E. (1983): Use of benzyltrimethyl-n-hexadecylammonium chloride ("16-BAC"), a cationic detergent, in an acidic polyacrylamide gel electrophoresis system to detect base labile protein methylation in intact cells. *Analytical Biochemistry*; 132: 231-235
- Manning M., and Colòn W. (2004): Structural basis of protein kinetic stability: Resistance to sodium dodecyl sulfate suggests a central role for rigidity and a bias toward beta-sheet structure. *Biochemistry*; 43: 11248-11254
- Martin P. T. (2002): Glycobiology of the synapse. *Glycobiology*; 12: 1R-7R.
- Mata J., Marguerat S., and Bahler J. (2005): Post-transcriptional control of gene expression: A genome-wide perspective. *Trends in Biochemical Sciences*; 30: 506-514
- Meddows E., Le Bourdellès B., Grimwood S., et al. (2001): Identification of molecular determinants that are important in the assembly of N-methyl-D-aspartate receptors. *The Journal of Biological Chemistry*; 276: 18795-18803
- Meier T., and Dimroth P. (2002): Intersubunit bridging by Na⁺ ions as a rationale for the unusual stability of the c-rings of Na⁺-translocating F1F0 ATP synthases. *EMBO reports*; 3: 1094-1098
- Molloy M., Herbert B. R., Walsh B. J., et al. (1998): Extraction of membrane proteins by differential solubilization for separation using two-dimensional gel electrophoresis. *Electrophoresis*; 19: 837-844
- Molloy M., Herbert B. R., Williams K. L., and Gooley A. A. (1999): Extraktion of Escherichia coli proteins with organic solvents prior to two-dimensional electrophoresis. *Electrophoresis*; 20: 701-704
- Monier S., Dietzen D. J., Hastings W. R., Lublin D. M., and Kurzchalia T. V. (1996): Oligomerization of VIP21-caveolinin vitro is stabilized by long chain fatty acylation or cholesterol. *FEBS Letters*; 388: 143-149
- Moritz B., and Meyer H. E. (2003): Approaches for the quantification of protein concentration ratios. *Proteomics*; 3: 2208-2220
- Navarre C., Degand H., Bennett K. L., et al. (2002): Subproteomics: Identification of plasma membrane proteins from the yeast Saccharomyces cerevisiae. *Proteomics*; 2: 1709-1714
- Neuhoff V., Arold N., Taube D., and Ehrhardt W. (1988): Improved staining of proteins in polyacrylamide gels including isoelectric focusing gels with clear background at nanogram sensitivity using Coomassie Brilliant Blue G-250 and R-250. *Electrophoresis*; 9: 255-262

- Nielsen P. A., Olsen J. V., Podtelejnikov A. V., et al. (2005): Proteomic mapping of brain plasma membrane proteins. *Molecular and Cellular Proteomics*; 4: 402-408
- Nishihara J. C., and Champion K. M. (2002): Quantitative evaluation of proteins in one- and two-dimensional polyacrylamide gels using a fluorescent stain. *Electrophoresis*; 23: 2203-2215
- O'Farrell P. H. (1975): High resolution two-dimensional electrophoresis of proteins. *Journal of Biological Chemistry*; 250: 4007-4021
- Pedersen S. K., Harry J. L., Sebastian L. T., et al. (2003): Unseen proteome: Mining below the tip of the iceberg to find low abundance and membrane proteins. *Journal of Proteome Research*; 2: 303-311.
- Peng J., Kim M. J., Cheng D., et al. (2004): Semiquantitative proteomic analysis of rat forebrain postsynaptic density fractions by mass spectrometry. *Journal of Biological Chemistry*; 279: 21003-21011
- Pinna G., Broedel O., Eravci M., et al. (2003): Thyroid hormones in the rat amygdala as common targets for antidepressant drugs, mood stabilizers, and sleep deprivation. *Biological Psychiatry*; 54: 1049-1059
- Pritchard D. I., Crawford C. R., Duce I. R., and Behnke J. M. (1985): Antigen stripping from the nematode epicuticle using the cationic detergent cethylmethylammonium bromide (CTAB). *Parasite Immunology*; 7: 575-585
- Rabilloud T. (1998): Use of thiourea to increase the solubility of membrane proteins in two-dimensional electrophoresis. *Electrophoresis*; 19: 758-760
- Rabilloud T. (1999): Solubilization of proteins in 2-D electrophoresis. *Methods in Molecular Biology*; 112: 9-19
- Rabilloud T., Blisnick T., Heller M., et al. (1999): Analysis of membrane proteins by two-dimensional electrophoresis: Comparison of the proteins extracted from normal or *Plasmodium falciparum*-infected erythrocyte ghosts. *Electrophoresis*; 20: 3603-3610
- Rabilloud T., Strub J.-M., Luche S., Dorselaer A. v., and Lunardi J. (2001): A comparison between Sypro Ruby and ruthenium II tris (bathophenanthroline disulfonate) as fluorescent stains for protein detection in gels. *Proteomics*; 1: 699-704
- Rais I., Karas M., and Schägger H. (2004): Two-dimensional electrophoresis for the isolation of integral membrane proteins and mass spectrometric identification. *Proteomics*; 4: 2567-2571
- Rauh J. J., Lambert M. P., Cho N. J., Chin H., and Klein W. (1986): Glycoprotein properties of muscarinic acetylcholine receptors from bovine cerebral cortex. *Journal of Neurochemistry*; 46: 23-32

- Raymond S., and Weintraub L. (1959): Acrylamide gel as a supporting medium for zone electrophoresis. *Science*; 18: 711
- Reidegeld K. A., Müller M., Stephan C., et al. (2006): The power of cooperative investigation: Summary and comparison of the HUPO Brain Proteome pilot study results. *Proteomics*; 6: 4997-5014
- Reifschneider N. H., Goto S., Nakamoto H., et al. (2006): Defining the mitochondrial proteomes from five rat organs in a physiologically significant context using 2D blue-native/SDS-PAGE. *Journal of Proteome Research*; 5: 1117-1132
- Righetti P. G., Castagna A., Antonucci F., et al. (2004): Critical survey of quantitative proteomics in two-dimensional electrophoretic approaches. *Journal of Chromatography A* 3-17
- Righetti P. G. (2006): Real and imaginary artefacts in proteome analysis via two-dimensional maps. *Journal of Chromatography B*; 841: 14-22
- Rosengren A. T., Salmi J. M., Aittokallio T., et al. (2003): Comparison of PDQuest and Progenesis software packages in the analysis of two-dimensional electrophoresis gels. *Proteomics*; 3: 1936-1946
- Ross P. L., Huang Y. N., Marchese J. N., et al. (2004): Multiplexed protein quantitation in *Saccharomyces cerevisiae* using amine-reactive isobaric tagging reagents. *Molecular and Cellular Proteomics*; 3: 1154-1169
- Sagné C., Isambert M.-F., Henry J.-P., and Gasnier B. (1996): SDS-resistant aggregation of membrane proteins: Application to the purification of the vesicular monoamine transporter. *Biochemical Journal*; 316: 825-831
- Santoni V., Molloy M., and Rabilloud T. (2000): Membrane proteins and proteomics: un amour impossible? *Electrophoresis* 1054-1070
- Schägger H., and Ohm T. G. (1995): Humal diseases with defects in oxidative phosphorylation 2. F₁F₀ ATP-synthase defects in Alzheimer disease revealed by blue native polyacrylamide gel electrophoresis. *European Journal of Biochemistry*; 227: 916-921
- Schindler J., Jung S., Niedner-Schatteburg G., Friauf E., and Nothwang H. G. (2006): Enrichment of integral membrane proteins from small amounts of brain tissue. *Journal of Neural Transmission*; 113: 995-1013
- Schmidt A., Kellermann J., and Lottspeich F. (2005): A novel strategy for quantitative proteomics using isotope-coded protein labels. *Proteomics*; 5: 4-15
- Schmidt F., Dahlmann B., Janek K., et al. (2006): Comprehensive quantitative proteome analysis of 20S proteasome subtypes from rat liver by isotope coded affinity tag and 2-D gel-based approaches. *Proteomics*; 6: 4622 - 4632

- Schrimpf S. P., Meskenaitel V., Brunner E., et al. (2005): Proteomic analysis of synaptosomes using isotope-coded affinity tags and mass spectrometry. *Proteomics*; 5: 2531-2541
- Schwartz R., Ting C., and King J. (2001): Whole proteome pI values correlate with subcellular localizations of proteins for organisms within the three domains of life. *Genome Research*; 11: 703-709
- Seigneurin-Berny D., Rolland N., Garin J., and Joyard J. (1999): Differential extraction of hydrophobic protein from chloroplast envelope membranes: A subcellular-specific proteomic approach to identify rare intrinsic membrane proteins. *The Plant Journal*; 19: 217-228
- Smolka M., Zhou H., and Aebersold R. (2002): Quantitative protein profiling using two-dimensional gel electrophoresis, isotope-coded affinity tag labeling, and mass spectrometry. *Molecular and Cellular Proteomics*; 1: 19-29
- Stevens S. M. J., Zharikova A. D., and Prokai L. (2003): Proteomic analysis of the synaptic plasma membrane fraction isolated from rat forebrain. *Molecular Brain Research*; 117: 116-128
- Svenningsson P., Chergui K., Rachleff I., et al. (2006): Alterations in 5-HT_{1B} receptor function by p11 in depression-like states. *Science*; 311: 77-80
- Thiede B., Höhenwarter W., Krah A., et al. (2005): Peptide mass fingerprinting. *Methods*; 35: 237-247
- Thompson A., Cronin M. S., and Kirschner D. A. (2002): Myelin protein zero exists as dimers and tetramers in native membranes of *Xenopus laevis* peripheral nerve. *Journal of Neuroscience Research*; 67: 766-771
- Ünlü M., Morgan M. E., and Minden J. S. (1997): Difference gel electrophoresis: A single gel method for detecting changes in protein extracts. *Electrophoresis*; 18: 2071-2077
- van Montfort B. A., Doeven M. K., Canas B., et al. (2002): Combined in-gel tryptic digestion and CNBr cleavage for the generation of peptide maps of an integral membrane protein with MALDI-TOF mass spectrometry. *Biochimica et Biophysica Acta*; 1555: 111-115
- Walikonis R. S., Jensen O. N., Mann M., et al. (2000): Identification of proteins in the postsynaptic density fraction by mass spectrometry. *The Journal of Neuroscience*; 20: 4069-4080
- Wallin E., and von Heijne G. (1998): Genome-wide analysis of integral membrane proteins from eubacterial, archaean, and eukaryotic organisms. *Protein Science*; 7: 1029-1038

- Wang G., Wu W. W., Zeng W., Chou C.-L., and Shen R.-F. (2006): Label-free protein quantification using LC-coupled ion trap or FT mass spectrometry: Reproducibility, linearity, and application with complex proteomes. *Journal of Proteome Research*; 5: 1214-1223
- Washburn M. P., Wolters D., and III J. R. Y. (2001): Large-scale analysis of the yeast proteome by multidimensional protein identification technology. *Nature Biotechnology*; 19: 242-247
- Wheelock A., and Buckpitt A. (2005): Software-induced variances in two-dimensional gel electrophoresis image analysis. *Proteomics*; 26: 4508-4520
- Wilkins M. R., Appel R. D., Van Eyk J. E., et al. (2006): Guidelines for the next 10 years of proteomics. *Proteomics*; 6: 4-8
- Witzmann F. A., Arnold R. J., Bai F., et al. (2005): A proteomic survey of rat cerebral cortical synaptosomes. *Proteomics*; 5: 2177-2201
- Wu C. C., MacCoss M. J., Howell K. L., and Yates J. R. (2003): A method for the comprehensive proteomic analysis of membrane proteins. *Nature Biotechnology*; 21: 532-538
- Wu C. C., and Yates III J. R. (2003): The application of mass spectrometry to membrane proteomics. *Nature Biotechnology*; 21: 262-267
- Yang J.-W., Czech T., and Lubec G. (2004): Proteomic profiling of human hippocampus. *Electrophoresis*; 25: 1169-1174
- Yoshimura Y., Yamauchi Y., Shinkawa T., et al. (2004): Molecular constituents of the postsynaptic density fraction revealed by proteomic analysis using multidimensional liquid chromatography-tandem mass spectrometry. *Journal of Neurochemistry*; 88: 759-768
- Zabel C., Sagi D., Kaindl A. M., et al. (2006): Comparative proteomics in neurodegenerative and non-neurodegenerative diseases suggest nodal point proteins in regulatory networking. *Journal of Proteome Research*; 5: 1948-1958
- Zhou S., Bailey M. J., Dunn M., Preedy V. R., and Emery P. W. (2005): A quantitative investigation into the loss of proteins at different stages of a two-dimensional gel electrophoresis procedure. *Proteomics*; 5: 2739-2747
- Zhu W., Venable J., Giometti C. S., et al. (2005): Large-scale LC-MS/MS for silver- and Coomassie blue-stained polyacrylamide gels. *Electrophoresis*; 26: 4495-4507
- Zimny-Arndt U., Schmid M., Ackermann R., and Jungblut P. R. (2007). Classical proteomics: two-dimensional electrophoresis/MALDI. In *Mass Spectrometry of Proteins and Peptides*. (ed. L. Pasa-Tolic). Humana Press.