

Bibliography

Allen, D.W., and Zamecnik, P.C. (1962). The effect of puromycin on rabbit reticulocyte ribosomes. *Biochim Biophys Acta* 55, 865-874.

Allen, G., Zavialov, A., Gursky, R., Ehrenberg, M., and Frank, J. (2005). The cryo-EM structure of a translation initiation complex from *Escherichia coli*. *Cell* 121, 703-712.

Ban, N., Nissen, P., Hansen, J., Moore, P.B., and Steitz, T.A. (2000). The complete atomic structure of the large ribosomal subunit at 2.4 Å resolution. *Science* 289, 905-920.

Bartetzko, A., and Nierhaus, K.H. (1988). A simple Mg^{2+}/NH_4^+ /polyamine system for poly(U) dependent poly(Phe) synthesis with near *in vivo* characteristics. *Methods Enzymol* 164, 650-658.

Berg, J.M., Tymoczko, J.L., and Stryer, L. (2002). *Biochemistry*, 5th edition edn (New York, W. H. Freeman and Company).

Bergemann, K., and Nierhaus, K.H. (1983). Spontaneous, elongation factor G independent translocation of *Escherichia coli* ribosomes. *J Biol Chem* 258, 15105-15113.

Bijlsma, J.J., Lie, A.L.M., Nootenboom, I.C., Vandenbroucke-Grauls, C.M., and Kusters, J.G. (2000). Identification of loci essential for the growth of *Helicobacter pylori* under acidic conditions. *J Infect Dis* 182, 1566-1569.

Blaha, G., Burkhardt, N., and Nierhaus, K.H. (2002). Formation of 70S ribosomes: Large activation energy is required for the adaptation of exclusively the small ribosomal subunit. *Biophys Chem* 96, 153-161.

Blaha, G., and Nierhaus, K.H. (2001). Features and functions of the ribosomal E site. *Cold Spring Harbor Symposia on Quantitative Biology* 65, 135-145.

Blaha, G., Stelzl, U., Spahn, C.M.T., Agrawal, R.K., Frank, J., and Nierhaus, K.H. (2000). Preparation of functional ribosomal complexes and the effect of buffer conditions on tRNA positions observed by cryoelectron microscopy. *Methods Enzymol* 317, 292-309.

Bommer, U., Burkhardt, N., Jünemann, R., Spahn, C.M.T., Triana-Alonso, F.J., and Nierhaus, K.H. (1996). Ribosomes and polysomes. In *Subcellular fractionation A practical approach*, J. Graham, and D. Rickwoods, eds. (Oxford, IRL Press at Oxford University Press), pp. 271-301.

Bonner, G., Lafer, E.M., and Sousa, R. (1994). Characterization of a set of T7 RNA polymerase active site mutants. *J Biol Chem* 269, 25120-25128.

Bouadloun, F., Donner, D., and Kurland, C.G. (1983). Codon-specific missense errors *in vivo*. *EMBO J* 2, 1351-1356.

Bremer, H., and Dennis, P.P. (1996). Modulation of chemical composition and other parameters of the cell by growth rate. In *Escherichia coli and Salmonella*, F.C. Neidhardt, R.C. III, J.L. Ingraham, E.C.C. Lin, K.B. Low, B. Magasanik, W.S. Reznikow, M. Riley, M. Schaechter, and H.E. Umbarger, eds. (Washington D.C., ASM Press), pp. 1553-1569.

Brock, S., Szkaradkiewicz, K., and Sprinzl, M. (1998). Initiation factors of protein biosynthesis in bacteria and their structural relationship to elongation and termination factors. *Mol Microbiol* 29, 409-417.

Brodersen, D.E., Clemons, W.M., Carter, A.P., Morgan-Warren, R.J., Wimberly, B.T., and Ramakrishnan, V. (2000). The structural basis for the action of the antibiotics tetracycline, pactamycin, and hygromycin B on the 30S ribosomal subunit. *Cell* 103, 1143-1154.

Brodersen, D.E., Clemons, W.M., Jr., Carter, A.P., Wimberly, B.T., and Ramakrishnan, V. (2002). Crystal structure of the 30 S ribosomal subunit from *Thermus thermophilus*: structure of the proteins and their interactions with 16 S RNA. *J Mol Biol* 316, 725-768.

Carpousis, A.J., and Dreyfus, M. (2004). mRNA decay and RNA-degrading machines in prokaryotes and eukaryotes. In *Protein Synthesis and Ribosome Structure*, K.H. Nierhaus, Wilson, D. N., ed. (Weinheim, Wiley-VCH Verlag GmbH & Co.), pp. 185-206.

Carter, A.P., Clemons, W.M., Brodersen, D.E., Morgan-Warren, R.J., Wimberly, B.T., and Ramakrishnan, V. (2000). Functional insights from the structure of the 30S ribosomal subunit and its interactions with antibiotics. *Nature* 407, 340-348.

Carter, A.P., Clemons, W.M., Jr., Brodersen, D.E., Morgan-Warren, R.J., Hartsch, T., Wimberly, B.T., and Ramakrishnan, V. (2001). Crystal structure of an initiation factor bound to the 30S ribosomal subunit. *Science* 291, 498-501.

Cassan, M., Berteaux, V., Angrand, P.O., and Rousset, J.P. (1990). Expression vectors for quantitating *in vivo* translational ambiguity: their potential use to analyse frameshifting at the HIV gag-pol junction. *Res Virol* 141, 597-610.

Cate, J.H., Yusupov, M.M., Yusupova, G.Z., Earnest, T.N., and Noller, H.F. (1999). X-ray crystal structures of 70S ribosome functional complexes. *Science* 285, 2095-2104.

Chamberlin, M., and Ring, J. (1973). Characterization of T7-specific ribonucleic acid polymerase. 1. General properties of the enzymatic reaction and the template specificity of the enzyme. *J Biol Chem* 248, 2235-2244.

Chang, H.C., Kaiser, C.M., Hartl, F.U., and Barral, J.M. (2005). *De novo* folding of GFP fusion proteins: high efficiency in eukaryotes but not in bacteria. *J Mol Biol* 353, 397-409.

Chin, K., Shean, C.S., and Gottesman, M.E. (1993). Resistance of lambda cl translation to antibiotics that inhibit translation initiation. *J Bacteriol* 175, 7471-7473.

Claude, A. (1943). The constitution of protoplasm. *Science* 97, 451-456.

Clemons, W.M., Jr., Brodersen, D.E., McCutcheon, J.P., May, J.L., Carter, A.P., Morgan-Warren, R.J., Wimberly, B.T., and Ramakrishnan, V. (2001). Crystal structure of the 30 S ribosomal subunit from *Thermus thermophilus*: purification, crystallization and structure determination. *J Mol Biol* 310, 827-843.

Cramer, A., Whitehorn, E.A., Tate, E., and Stemmer, W.P. (1996). Improved green fluorescent protein by molecular evolution using DNA shuffling. *Nat Biotechnol* 14, 315-319.

Cukras, A.R., Southworth, D.R., Brunelle, J.L., Culver, G.M., and Green, R. (2003). Ribosomal proteins S12 and S13 function as control elements for translocation of the mRNA:tRNA complex. *Mol Cell* 12, 321-328.

Cummings, H.S., and Hershey, J.W.B. (1994). Translation initiation factor IF1 is essential for cell viability in *Escherichia coli*. *J Bacteriol* 176, 198-205.

Czworkowski, J., Wang, J., Seitz, T.A., and Moore, P.B. (1994). The crystal structure of elongation factor G complexed with GDP, at 2.7 Å resolution. *EMBO J* 13, 3661-3668.

Dahlquist, K.D., and Puglisi, J.D. (2000). Interaction of translation initiation factor IF1 with the *E. coli* ribosomal A site. *J Mol Biol* 299, 1-15.

Dallas, A., and Noller, H.F. (2001). Interaction of translation initiation factor 3 with the 30S ribosomal subunit. *Mol Cell* 8, 855-864.

Dandliker, P., Pratt, S., Nilius, A., Black-Schaefer, C., Ruan, X., Towne, D., Clark, R., Englund, E., Wagner, R., Weitzberg, M., *et al.* (2003). Novel antibacterial class. *Antimicrob Agents Chemother* 47, 3831-3839.

Davies, J., Gilbert, W., and Gorini, L. (1964). Streptomycin, suppression, and the code. *Proc Natl Acad Sci USA* 51, 883-890.

de Smit, M.H., and van Duin, J. (1994). Control of translation by mRNA secondary structure in *Escherichia coli* - A quantitative analysis of literature data. *J Mol Biol* 244, 144-150.

Dennis, P.P., and Bremer, H. (1973). A method for determination of the synthesis rate of stable and unstable ribonucleic acid in *Escherichia coli*. *Analytical biochemistry* 56, 489-501.

Dibb, N.J., and Wolfe, P.B. (1986). *lep* operon proximal gene is not required for growth or secretion by *Escherichia coli*. *J Bacteriol* 166, 83-87.

Dinos, G., Wilson, D.N., Teraoka, Y., Szaflarski, W., Fucini, P., Kalpaxis, D., and Nierhaus, K.H. (2004). Dissecting the ribosomal inhibition mechanisms of edeine and pactamycin: the universally conserved residues G693 and C795 regulate P-site tRNA binding. *Mol Cell* 13, 113-124.

Frank, J. (1989). Image analysis of single macromolecules. *Electron Microsc Rev* 2, 53-74.

Frank, J. (2001). Cryo-electron microscopy as an investigative tool: the ribosome as an example. *Bioessays* 23, 725-732.

Gale, E.F., Cundliffe, E., Reynolds, P.E., Richmond, M.H., and Waring, M.J. (1981). Antibiotic inhibitors of ribosome function. In *The molecular basis of antibiotic action* (Bristol, UK, John Wiley and sons), pp. 278-379.

Gavrilova, L.P., and Spirin, A.S. (1971). Stimulation of 'non-enzymic' translocation in ribosome by p -chloromercuribenzoate. *FEBS Lett* 17, 324-326.

Gnirke, A., and Nierhaus, K.H. (1986). tRNA binding sites on the subunits of *Escherichia coli* ribosomes. *J Biol Chem* 261, 14506-14514.

Gowrishankar, J., and Harinarayanan, R. (2004). Why is transcription coupled to translation in bacteria? *Mol Microbiol* 54, 598-603.

Gualerzi, C., Brandi, L., Caserta, E., La teana, A., Spurio, R., Tomsic, J., and Pon, C. (2000). Translation initiation in bacteria. In *The ribosome structure, function, antibiotics, and cellular interactions*, R.A. Garrett, S.R. Douthwaite, A. Liljas, A.T. Matheson, P.B. Moore, and H.F. Noller, eds. (Washington D.C., ASM Press), pp. 477-494.

Gualerzi, C.O., and Pon, C.L. (1990). Initiation of messenger-RNA translation in prokaryotes. *Biochemistry* 29, 5881-5889.

Guillerez, J., Lopez, P.J., Proux, F., Launay, H., and Dreyfus, M. (2005). A mutation in T7 RNA polymerase that facilitates promoter clearance. *Proc Natl Acad Sci U S A* 102, 5958-5963.

Hamada, M., Hashimoto, T., Takahashi, T., Yokoyama, S., Miyake, M., Takeuchi, T., Okami, Y., and Umezawa, H. (1965). Antimicrobial activity of kasugamycin. *J Antibiot* 18, 104-106.

Hansen, J.L., Ippolito, J.A., Ban, N., Nissen, P., Moore, P.B., and Steitz, T.A. (2002). The structures of four macrolide antibiotics bound to the large ribosomal subunit. *Mol Cell* 10, 117-128.

Hapke, B., and Noll, H. (1976). Structural dynamics of bacterial ribosomes: IV. Classification of ribosomes by subunit interaction. *J Mol Biol* 105, 97-109.

Harms, J., Schluenzen, F., Zarivach, R., Bashan, A., Gat, S., Agmon, I., Bartels, H., Franceschi, F., and Yonath, A. (2001). High resolution structure of the large ribosomal subunit from a mesophilic eubacterium. *Cell* 107, 679-688.

Hartz, D., McPheeters, D.S., Traut, R., and Gold, L. (1988). Extension inhibition analysis of translation initiation complexes. *Methods Enzymol* 164, 419-425.

Hausner, T.P., Geigenmüller, U., and Nierhaus, K.H. (1988). The allosteric three site model for the ribosomal elongation cycle. New insights into the inhibition mechanisms of aminoglycosides, thiostrepton, and viomycin. *J Biol Chem* 263, 13103-13111.

Hirashima, A., Childs, G., and Inouye, M. (1973). Differential inhibitory effects of antibiotics on the biosynthesis of envelope proteins of *Escherichia coli*. *J Mol Biol* 79, 373-389.

Iost, I., and Dreyfus, M. (1995). The stability of *Escherichia coli lacZ* mRNA depends upon the simultaneity of its synthesis and translation. *Embo J* 14, 3252-3261.

Iskakova, M.B., Szaflarski, W., Dreyfus, M., Remme, J., and Nierhaus, K.H. (2006). Troubleshooting coupled *in vitro* transcription–translation system derived from *Escherichia coli* cells: synthesis of high-yield fully active proteins. *Nucl Acids Res*, e135.

Jelenc, P.C., and Kurland, C.G. (1979). Nucleoside triphosphate regeneration decreases the frequency of translation errors. *Proc Natl Acad Sci USA* 76, 3174-3178.

Jewett, M.C., and Swartz, J.R. (2004). Substrate replenishment extends protein synthesis with an *in vitro* translation system designed to mimic the cytoplasm. *Biotechnol Bioeng* 87, 465-472.

Jewett, M.C., Voloshin, A., Swartz, J. R. (2002). Procaryotic systems for *in vitro* expression. In Gene Cloning and Expression Technologies, M.P.a.L. Weiner, Q., ed. (Westborough, MA, BioTechniques Press), pp. 391-411.

Jorgensen, F., and Kurland, C.G. (1990). Processivity errors of gene expression in *Escherichia coli*. *J Mol Biol* 215, 511-521.

Kerner, M.J., Naylor, D.J., Ishihama, Y., Maier, T., Chang, H.C., Stines, A.P., Georgopoulos, C., Frishman, D., Hayer-Hartl, M., Mann, M., *et al.* (2005). Proteome-wide analysis of chaperonin-dependent protein folding in *Escherichia coli*. *Cell* 122, 209-220.

Kim, D.M., and Choi, C.Y. (1996). A semi-continuous prokaryotic coupled transcription/translation system using a dialysis membrane. *Biotechnol Progr* 12, 645-649.

Kim, D.M., and Swartz, J.R. (2000). Prolonging cell-free protein synthesis by selective reagent additions. *Biotechnol Prog* 16, 385-390.

Kozak, M., and Nathans, D. (1972). Differential inhibition of coliphage MS2 protein synthesis by ribosome- directed antibiotics. *J Mol Biol* 70, 41-55.

Kubota, K., Okuyama, A., and Tanaka, N. (1972). Differential effects of antibiotics on peptidyl transferase reactions. *Biochem Biophys Res Comm* 47, 1196-1202.

Kurland, C.G., Jørgensen, F., Richter, A., Ehrenberg, M., Bilgin, N., and Rojas, A.-M. (1990). Through the accuracy window. In *The Ribosome- Structure, Function, and Evolution*, A. Dahlberg, W.E. Hill, R.A. Garrett, P.B. Moore, D. Schlessinger, and J.R. Warner, eds. (Washington, D. C., Amer. Soc. Microbiol.), pp. 513-526.

Laemmli, U.K., and Favre, M. (1973). Maturation of the head of bacteriophage T4. I. DNA packaging events. *J Mol Biol* 80, 575-599.

Lewicki, B.T.U., Margus, T., Remme, J., and Nierhaus, K.H. (1993). Coupling of rRNA transcription and ribosomal assembly *in vivo* - formation of active ribosomal subunits in *Escherichia coli* requires transcription of rRNA genes by host RNA polymerase which cannot be replaced by bacteriophage-T7 RNA polymerase. *J Mol Biol* 231, 581-593.

Liiv, A., and Remme, J. (2004). Importance of transient structures during post-transcriptional refolding of the pre-23S rRNA and ribosomal large subunit assembly. *J Mol Biol* 342, 725-741.

Liiv, A., Tenson, T., and Remme, J. (1996). Analysis of the ribosome large subunit assembly and 23S rRNA stability *in vivo*. *J Mol Biol* 263, 396-410.

Makarova, O.V., Makarov, E.M., Sousa, R., and Dreyfus, M. (1995). Transcribing of *Escherichia coli* genes with mutant T7 RNA polymerases: Stability of *lacZ* mRNA inversely correlates with polymerase speed. *Proc Natl Acad Sci USA* 92, 12250-12254.

Makino, Y., Amada, K., Taguchi, H., and Yoshida, M. (1997). Chaperonin-mediated folding of green fluorescent protein. *J Biol Chem* 272, 12468-12474.

Maniatis, R.B., Fritsch, E.F., and Sambrook, J. (1982). *Molecular Cloning: A Laboratory Manual*. (Cold Spring Harbour Laboratory).

Márquez, V. (2002). Switching off the mechanism for maintaining the ribosomal reading frame: Translational regulation of release factor 2 (Berlin, Free University Berlin).

Masukawa, H., Tanaka, N., and Umezawa, H. (1968). Inhibition by kasugamycin of protein synthesis in *Piricularia oryzae*. *J Antibiot* 21, 73-74.

Menninger, J.R. (1976). Peptidyl transfer RNA dissociates during protein synthesis from ribosomes of *Escherichia coli*. *J Biol Chem* 251, 3392-3398.

Milligan, F., and Uhlenbeck, O.C. (1989). Synthesis of small RNAs using T7 RNA polymerase. *Meth Enzymol* 180, 51-62.

Moazed, D., and Noller, H.F. (1987). Interaction of antibiotics with functional sites in 16S ribosomal RNA. *Nature* 327, 389-394.

Moazed, D., and Noller, H.F. (1989). Interaction of tRNA with 23S rRNA in the ribosomal A, P, and E sites. *Cell* 57, 585-597.

Moazed, D., and Noller, H.F. (1990). Binding of tRNA to the ribosomal A and P sites protects two distinct sets of nucleotides in the 16S rRNA. *J Mol Biol* 211, 135-145.

Mohanty, B.K., and Kushner, S.R. (1999). Analysis of the function of *Escherichia coli* poly(A) polymerase I in RNA metabolism. *Mol Microbiol* 34, 1094-1108.

Moll, I., and Bläsi, U. (2002). Differential inhibition of 30S and 70S translation initiation complexes on leaderless mRNA by kasugamycin. *Biochem Biophys Res Comm* 297, 1021-1026.

Murzin, A. (1993). OB (oligonucleotide/oligosaccharide binding)-fold: common structural and functional solution for non-homologous sequences. *EMBO J* 12, 861-867.

Nierhaus, K.H. (1990). The allosteric three-site model for the ribosomal elongation cycle: features and future. *Biochemistry* 29, 4997-5008.

Nissen, P., Ippolito, J.A., Ban, N., Moore, P.B., and Steitz, T.A. (2001). RNA tertiary interactions in the large ribosomal subunit: the A-minor motif. *Proc Natl Acad Sci USA* 98, 4899-4903.

Ogawa, H., Inouye, S., Tsuji, F.I., Yasuda, K., and Umesono, K. (1995). Localization, trafficking, and temperature-dependence of the *Aequorea* green

fluorescent protein in cultured vertebrate cells. *Proc Natl Acad Sci U S A* 92, 11899-11903.

Ogle, J., Carter, A., and Ramakrishnan, V. (2003). Insights into the decoding mechanism from recent ribosome structures. *TIBS* 28, 259-266.

Ogle, J.M., Brodersen, D.E., Clemons Jr, W.M., Tarry, M.J., Carter, A.P., and Ramakrishnan, V. (2001). Recognition of cognate transfer RNA by the 30S ribosomal subunit. *Science* 292, 897-902.

Ogle, J.M., Murphy, F.V., Tarry, M.J., and Ramakrishnan, V. (2002). Selection of tRNA by the ribosome requires a transition from an open to a closed form. *Cell* 111, 721-732.

Okuyama, A., Machiyama, N., Kinoshita, T., and Tanaka, N. (1971). Inhibition by kasugamycin of initiation complex formation on 30S ribosomes. *Biochem Biophys Res Commun* 43, 196-169.

Okuyama, A., and Tanaka, N. (1972). Differential effects of aminoglycosides on cistron-specific initiation of protein synthesis. *Biochem Biophys Res Commun* 49, 951-957.

Ormo, M., Cubitt, A.B., Kallio, K., Gross, L.A., Tsien, R.Y., and Remington, S.J. (1996). Crystal structure of the *Aequorea victoria* green fluorescent protein. *Science* 273, 1392-1395.

Pestka, S. (1969). Studies on the formation of transfer ribonucleic acid-ribosome complexes. VI. Oligopeptide synthesis and translocation on ribosomes in the presence and absence of soluble transfer factors. *J Biol Chem* 244, 1533-1539.

Pioletti, M., Schlunzen, F., Harms, J., Zarivach, R., Gluhmann, M., Avila, H., Bashan, A., Bartels, H., Auerbach, T., Jacobi, C., *et al.* (2001). Crystal structures of complexes of the small ribosomal subunit with tetracycline, edeine and IF3. *EMBO J* 20, 1829-1839.

Poldermans, B., Goosen, N., and Van Knippenberg, P.H. (1979). Studies on the function of two adjacent N⁶,N⁶-dimethyladenosines near the 3' end of 16 S ribosomal RNA of *Escherichia coli*. I. The effect of kasugamycin on initiation of protein synthesis. *J Biol Chem* 254, 9085-9089.

Potapov, A.P. (1982). A stereospecific mechanism for the aminoacyl-tRNA selection at the ribosome. *FEBS Lett* 146, 28-33.

Potapov, A.P., Triana-Alonso, F.J., and Nierhaus, K.H. (1995). Ribosomal decoding processes at codons in the A or P sites depend differently on 2'-OH groups. *J Biol Chem* 270, 17680-17684.

Prasher, D.C., Eckenrode, V.K., Ward, W.W., Prendergast, F.G., and Cormier, M.J. (1992). Primary structure of the *Aequorea victoria* green-fluorescent protein. *Gene* 111, 229-233.

Qin, Y., Polacek, N., Vesper, O., Staub, E., Einfeldt, E., Wilson, D.N., and Nierhaus, K.H. (2006). The highly conserved LepA is a ribosomal elongation factor that back-translocates the ribosome. *Cell* 127, 721-733.

Remme, J., Margus, T., VILLEMS, R., and Nierhaus, K.H. (1989). The third ribosomal tRNA-binding site, the E site, is occupied in native polysomes. *Eur J Biochem* 183, 281-284.

Rheinberger, H.-J. (2004). A history of protein biosynthesis and ribosome research.

Rheinberger, H.-J., Geigenmüller, U., Wedde, M., and Nierhaus, K.H. (1988). Parameters for the preparation of *Escherichia coli* ribosomes and ribosomal subunits active in tRNA binding. *Methods Enzymol* 164, 658-670.

Rheinberger, H.-J., and Nierhaus, K.H. (1986). Allosteric interactions between the ribosomal transfer RNA-binding sites A and E. *J Biol Chem* 261, 9133-9139.

Rodnina, M.V., Gromadski, K.B., Kothe, U., and Wieden, H.J. (2005). Recognition and selection of tRNA in translation. *FEBS Lett* 579, 938-942.

Ryabova, L.A., Vinokurov, L.M., Shekhovtsova, E.A., Alakhov, Y.B., and Spirin, A.S. (1995). Acetyl phosphate as an energy source for bacterial cell-free translation systems. *Analytical biochemistry* 226, 184-186.

Sacchetti, A., Cappetti, V., Marra, P., Dell'Arciprete, R., El Sewedy, T., Crescenzi, C., and Alberti, S. (2001). Green fluorescent protein variants fold differentially in prokaryotic and eukaryotic cells. *J Cell Biochem Suppl Suppl* 36, 117-128.

Schäfer, M.A., Tastan, A.O., Patzke, S., Blaha, G., Spahn, C.M., Wilson, D.N., and Nierhaus, K.H. (2002). Codon-anticodon interaction at the P Site is a prerequisite for tRNA interaction with the small ribosomal subunit. *J Biol Chem* 277, 19095-19105.

Schilling-Bartetzko, S., Bartetzko, A., and Nierhaus, K.H. (1992). Kinetic and thermodynamic parameters for transfer RNA binding to the ribosome and for the translocation reaction. *J Biol Chem* 267, 4703-4712.

Schlieker, C., Bukau, B., and Mogk, A. (2002). Prevention and reversion of protein aggregation by molecular chaperones in the *E. coli* cytosol: implications for their applicability in biotechnology. *J Biotechnol* 96, 13-21.

Schlunzen, F., Takemoto, C., Wilson, D.N., Kaminishi, T., Harms, J.M., Hanawa-Suetsugu, K., Szaflarski, W., Kawazoe, M., Shirouzu, M., Nierhaus, K.H., *et al.* (2006). The antibiotic kasugamycin mimics mRNA nucleotides to destabilize tRNA binding and inhibit canonical translation initiation. *Nat Struct Mol Biol* 13, 871-878.

Schlunzen, F., Tocilj, A., Zarivach, R., Harms, J., Gluehmann, M., Janell, D., Bashan, A., Bartels, H., Agmon, I., Franceschi, F., *et al.* (2000). Structure of functionally activated small ribosomal subunit at 3.3 Å resolution. *Cell* 102, 615-623.

Schmeing, T.M., Huang, K.S., Kitchen, D.E., Strobel, S.A., and Steitz, T.A. (2005). Structural insights into the roles of water and the 2' hydroxyl of the P site tRNA in the peptidyl transferase reaction. *Mol Cell* 20, 437-448.

Schmitt, E., Mechulam, Y., Fromant, M., Plateau, P., and Blanquet, S. (1997). Crystal structure at 1.2 Å resolution and active site mapping of *Escherichia coli* peptidyl-tRNA hydrolase. *Embo J* 16, 4760-4769.

Selinger, D.W., Saxena, R.M., Cheung, K.J., Church, G.M., and Rosenow, C. (2003). Global RNA half-life analysis in *Escherichia coli* reveals positional patterns of transcript degradation. *Genome Res* 13, 216-223.

Semenkov, Y.P., Rodnina, M.V., and Wintermeyer, W. (1996). The "allosteric three-site model" of elongation cannot be confirmed in a well-defined ribosome system from *Escherichia coli*. *Proc Natl Acad Sci U S A* 93, 12183-12188.

Shen, L.L., Black-Schaefer, C., Cai, Y., Dandliker, P.J., and Beutel, B.A. (2005). Mechanism of action of a novel series of naphthyridine-type ribosome inhibitors: enhancement of tRNA footprinting at the decoding site of 16S rRNA. *Antimicrob Agents Chemother* 49, 1890-1897.

Shimomura, O., Johnson, F.H., and Saiga, Y. (1962). Extraction, purification and properties of aequorin, a bioluminescent protein from the luminous hydromedusan, *Aequorea*. *J Cell Comp Physiol* 59, 223-239.

Shine, J., and Dalgarno, L. (1974). The 3'-terminal sequence of *E. coli* 16S rRNA: Complementarity to nonsense triplets and ribosome binding sites. *Proc Natl Acad Sci USA* 71, 1342-1346.

Shoji, S., Walker, S.E., and Fredrick, K. (2006). Reverse translocation of tRNA in the ribosome. *Mol Cell* 24, 931-942.

Spahn, C.M.T., and Nierhaus, K.H. (1998). Models of the elongation cycle: An evaluation. *Biol Chem* 379, 753-772.

Spirin, A.S., Baranov, V.I., Ryabova, L.A., Ovodov, S.Y., and Alakhov, Y.B. (1988). A continuous cell-free translation system capable of producing polypeptides in high yield. *Science* 242, 1162-1164.

Srivastava, A.K., and Schlessinger, D. (1990). Mechanism and regulation of bacterial ribosomal RNA processing. *Annu Rev Microbiol* 44, 105-129.

Stark, H., Mueller, F., Orlova, E.V., Schatz, M., Dube, P., Erdemir, T., Zemlin, F., Brimacombe, R., and Heel, M.v. (1995). The 70S *Escherichia coli* ribosome at 23 Å resolution: Fitting the ribosomal RNA. *Structure* 3, 815-821.

Stöffler-Meilicke, M., and Stöffler, G. (1990). Topography of the ribosomal proteins from *Escherichia coli* within the intact subunits as determined by immunoelectron microscopy and protein-protein cross-linking. In *The ribosome Structure, function and evolution*, W.E. Hill, A. Dahlberg, R.A. Garrett, P.B. Moore, D. Schlessinger, and J.R. Warner, eds. (Washington D C, American Society for Microbiology), pp. 123-133.

Studer, S., and Joseph, S. (2006). Unfolding of mRNA secondary structure by the bacterial translation initiation complex. *Mol Cell* 22, 105-115.

Suhara, Y., Maeda, K., and Umezawa, H. (1966). Chemical studies on kasugamycin. V. The structure of kasugamycin. *Tetrahedron Lett* 12, 1239-1244.

Sussman, J.K., Simons, E.L., and Simons, R.W. (1996). *Escherichia coli* translation initiation factor 3 discriminates the initiation codon *in vivo*. *Mol Microbiol* 21, 347-360.

Takeuchi, T., Ishizuka, M., Takayama, H., Kureha, K., Hamada, M., and Umezawa, H. (1965). Pharmacology of kasugamycin and the effect on *Pseudomonas* infection. *J Antibiot* 18, 107-110.

Tanaka, N., Yamaguchi, H., and Umezawa, H. (1966a). Mechanism of kasugamycin action on polypeptide synthesis. *J Biochem (Tokyo)* 60, 429-434.

Tanaka, N., Yoshida, Y., Sashikata, K., Yamaguchi, H., and Umezawa, H. (1966b). Inhibition of polypeptide synthesis by kasugamycin, an aminoglycosidic antibiotic. *J Antibiot* 19, 65-68.

Thompson, J., O'Connor, M., Mills, J.A., and Dahlberg, A.E. (2002). The protein synthesis inhibitors, oxazolidinones and chloramphenicol, cause extensive translational inaccuracy *in vivo*. *J Mol Biol* 322, 273-279.

Traut, R.R., and Monro, R.E. (1964). The puromycin reaction and its relation to protein synthesis. *J Mol Biol* 10, 63-72.

Triana-Alonso, F.J., Dabrowski, M., Wadzack, J., and Nierhaus, K.H. (1995). Self-coded 3'-extension of run-off transcripts produces aberrant products during *in vitro* transcription with T7 RNA polymerase. *J Biol Chem* 270, 6298-6307.

Tsien, R.Y. (1998). The green fluorescent protein. *Annu Rev Biochem* 67, 509-544.

Umezawa, H., Hamada, M., Suhara, Y., Hashimoto, T., and Ikekawa, T. (1965). Kasugamycin, a new antibiotic. *Antimicrobial Agents Chemother* 5, 753-757.

Underwood, K.A., Swartz, J.R., and Puglisi, J.D. (2005). Quantitative polysome analysis identifies limitations in bacterial cell-free protein synthesis. *Biotechnol Bioeng* 91, 425-435.

Van Buul, C.P.J.J., Visser, W., and Van Knippenberg, P.H. (1984). Increased translational fidelity caused by the antibiotic kasugamycin and ribosomal ambiguity in mutants harboring the *ksgA* gene. *FEBS Lett* 177, 119-124.

Vila-Sanjurjo, A., Squires, C.L., and Dahlberg, A.E. (1999). Isolation of kasugamycin resistant mutants in the 16 S ribosomal RNA of *Escherichia coli*. *J Mol Biol* 293, 1-8.

Voss, N.R., Gerstein, M., Steitz, T.A., and Moore, P.B. (2006). The geometry of the ribosomal polypeptide exit tunnel. *Journal of molecular biology* 360, 893-906.

Ward, W.W., and Bokman, S.H. (1982). Reversible denaturation of *Aequorea* green-fluorescent protein: physical separation and characterization of the renatured protein. *Biochemistry* 21, 4535-4540.

Watanabe, S. (1972). Interaction of siomycin with the acceptor site of *Escherichia coli* ribosomes. *J Mol Biol* 67, 443-457.

Weber, M.H., Beckering, C.L., and Marahiel, M.A. (2001). Complementation of cold shock proteins by translation initiation factor IF1 in vivo. *J Bacteriol* 183, 7381-7386.

Weijland, A., and Parmeggiani, A. (1993). Toward a model for the interaction between elongation factor Tu and the ribosome. *Science* 259, 1311-1314.

Wilson, D.N., and Nierhaus, K.H. (2003). The ribosome through the looking glass. *Angew Chem Int Ed Engl* 42, 3464-3486.

Wilson, D.N., and Nierhaus, K.H. (2006). The E-site story: the importance of maintaining two tRNAs on the ribosome during protein synthesis. *Cell Mol Life Sci* 63, 2725-2737.

Wimberly, B.T., Brodersen, D.E., Clemons, W.M., Morgan-Warren, R.J., Carter, A.P., Vornrhein, C., Hartsch, T., and Ramakrishnan, V. (2000). Structure of the 30S ribosomal subunit. *Nature* 407, 327-339.

Woodcock, J., Moazed, D., Cannon, M., Davies, J., and Noller, H.F. (1991). Interaction of antibiotics with A- and P-site-specific bases in 16S ribosomal RNA. *EMBO J* 10, 3099-3103.

Yang, F., Moss, L.G., and Phillips, G.N., Jr. (1996). The molecular structure of green fluorescent protein. *Nat Biotechnol* 14, 1246-1251.

Yusupov, M.M., Yusupova, G.Z., Baucom, A., Lieberman, K., Earnest, T.N., Cate, J.H., and Noller, H.F. (2001). Crystal structure of the ribosome at 5.5 Å resolution. *Science* 292, 883-896.

Zaniewski, R., Petkaites, E., and Deutscher, M.P. (1984). A multiple mutant of *Escherichia coli* lacking the exoribonucleases RNase II, RNase D, and RNase BN. *J Biol Chem* 259, 11651-11653.