

---

**6. LITERATURVERZEICHNIS**

- Akamatsu, H., Zouboulis, C.C. and Orfanos, C.E. (1992) Control of human sebocyte proliferation in vitro by testosterone and 5-alpha-dihydrotestosterone is dependent on the localization of the sebaceous glands. *J Invest Dermatol*, **99**, 509-511.
- Akamatsu, H., Zouboulis, C.C. and Orfanos, C.E. (1993) Spironolactone directly inhibits proliferation of cultured human facial sebocytes and acts antagonistically to testosterone and 5 alpha-dihydrotestosterone in vitro. *J Invest Dermatol*, **100**, 660-662.
- Ando, Y., Yamaguchi, Y., Hamada, K., Yoshikawa, K. and Itami, S. (1999) Expression of mRNA for androgen receptor, 5alpha-reductase and 17beta-hydroxysteroid dehydrogenase in human dermal papilla cells. *Br J Dermatol*, **141**, 840-845.
- Anigbogu, A.N., Williams, A.C. and Barry, B.W. (1996) Permeation characteristics of 8-methoxypsoralen through human skin; relevance to clinical treatment. *J Pharm Pharmacol*, **48**, 357-366.
- Anttila, H.S., Reitamo, S. and Saurat, J.H. (1992) Interleukin 1 immunoreactivity in sebaceous glands. *Br J Dermatol*, **127**, 585-588.
- Azziz, R. (2003) The evaluation and management of hirsutism. *Obstet Gynecol*, **101**, 995-1007.
- Bashir, S.J., Chew, A.L., Anigbogu, A., Dreher, F. and Maibach, H.I. (2001) Physical and physiological effects of stratum corneum tape stripping. *Skin Res Technol*, **7**, 40-48.
- Battmann, T., Bonfils, A., Branche, C., Humbert, J., Goubet, F., Teutsch, G. and Philibert, D. (1994) RU 58841, a new specific topical antiandrogen: a candidate of choice for the treatment of acne, androgenetic alopecia and hirsutism. *J Steroid Biochem Mol Biol*, **48**, 55-60.
- Bayne, E.K., Flanagan, J., Einstein, M., Ayala, J., Chang, B., Azzolina, B., Whiting, D.A., Mumford, R.A., Thiboutot, D., Singer, II and Harris, G. (1999) Immunohistochemical localization of types 1 and 2 5alpha-reductase in human scalp. *Br J Dermatol*, **141**, 481-491.
- Birch, M.P., Lalla, S.C. and Messenger, A.G. (2002) Female pattern hair loss. *Clin Exp Dermatol*, **27**, 383-388.
- Blaschke T., Sivaramakrishnan R, Gross M, Kramer KD.(2006) Organ mapping using parelectric spectroscopy. *Phys Med Biol*, **51**, 1623-31
- Blauer, M., Vaalasti, A., Pauli, S.L., Ylikomi, T., Joensuu, T. and Tuohimaa, P. (1991) Location of androgen receptor in human skin. *J Invest Dermatol*, **97**, 264-268.
- Boehm, K.D., Yun, J.K., Strohl, K.P. and Elmetts, C.A. (1995) Messenger RNAs for the multifunctional cytokines interleukin-1 alpha, interleukin-1 beta and tumor necrosis factor-alpha are present in adnexal tissues and in dermis of normal human skin. *Exp Dermatol*, **4**, 335-341.
- Bojar, R.A. and Holland, K.T. (2004) Acne and Propionibacterium acnes. *Clin Dermatol*, **22**, 375-379.
- Breiner, M., Romalo, G. and Schweikert, H.U. (1986) Inhibition of androgen receptor binding by natural and synthetic steroids in cultured human genital skin fibroblasts. *Klin Wochenschr*, **64**, 732-737.
- Brinkmann, A.O., Faber, P.W., van Rooij, H.C., Kuiper, G.G., Ris, C., Klaassen, P., van der Korput, J.A., Voorhorst, M.M., van Laar, J.H., Mulder, E. and et al. (1989) The human androgen receptor: domain structure, genomic organization and regulation of expression. *J Steroid Biochem*, **34**, 307-310.

- Bronaugh, R.L., Stewart, R.F. and Simon, M. (1986) Methods for in vitro percutaneous absorption studies. VII: Use of excised human skin. *J Pharm Sci*, **75**, 1094-1097.
- Brown, T.R., Rothwell, S.W., Sultan, C. and Migeon, C.J. (1981) Inhibition of androgen binding in human foreskin fibroblasts by antiandrogens. *Steroids*, **37**, 635-648.
- Bunjes, H., Drechsler, M., Koch, M.H.J. and Westesen, K. (2001) Incorporation of the model drug ubidecarenone into solid lipid nanoparticles. *Pharm Res*, **18**, 287-293.
- Campisi, G., Giandalia, G., De Caro, V., Di Liberto, C., Arico, P. and Giannola, L.I. (2004) A new delivery system of clobetasol-17-propionate (lipid-loaded microspheres 0.025%) compared with a conventional formulation (lipophilic ointment in a hydrophilic phase 0.025%) in topical treatment of atrophic/erosive oral lichen planus. A Phase IV, randomized, observer-blinded, parallel group clinical trial. *Br J Dermatol*, **150**, 984-990.
- Cannell, G.R., Mortimer, R.H. and Thomas, M.J. (1981) High-performance liquid chromatographic estimation of cyproterone acetate in human plasma. *J Chromatogr*, **226**, 492-497.
- Chase, H.B. (1954) Growth of the hair. *Physiol Rev*, **34**, 113-126.
- Chen, W., Yang, C.C., Sheu, H.M., Seltmann, H. and Zouboulis, C.C. (2003) Expression of peroxisome proliferator-activated receptor and CCAAT/enhancer binding protein transcription factors in cultured human sebocytes. *J Invest Dermatol*, **121**, 441-447.
- Choudhry, R., Hodgins, M.B., Van der Kwast, T.H., Brinkmann, A.O. and Boersma, W.J. (1992) Localization of androgen receptors in human skin by immunohistochemistry: implications for the hormonal regulation of hair growth, sebaceous glands and sweat glands. *J Endocrinol*, **133**, 467-475.
- Crooks, S.W. and Stockley, R.A. (1998) Leukotriene B<sub>4</sub>. *Int J Biochem Cell Biol*, **30**, 173-178.
- Crovato, F., Moretti, G. and Bertamino, R. (1973) 17- $\alpha$ -Hydroxy-steroid-dehydrogenases in hair follicles of normal and bald scalp: a histochemical study. *J Invest Dermatol*, **60**, 126-133.
- Danilenko, D.M., Ring, B.D. and Pierce, G.F. (1996) Growth factors and cytokines in hair follicle development and cycling: recent insights from animal models and the potentials for clinical therapy. *Mol Med Today*, **2**, 460-467.
- de Hassonville, S., Chiap, P., Liegeois, J.F., Evrard, B., Delattre, L., Crommen, J., Piel, G. and Hubert, P. (2004) Development and validation of a high-performance liquid chromatographic method for the determination of cyproterone acetate in human skin. *J Pharm Biomed Anal*, **36**, 133-143.
- de Jalon, E.G., Blanco-Prieto, M.J., Ygartua, P. and Santoyo, S. (2001) Topical application of acyclovir-loaded microparticles: quantification of the drug in porcine skin layers. *J Control Release*, **75**, 191-197.
- Denk, W., Strickler, J.H. and Webb, W.W. (1990) Two-photon laser scanning fluorescence microscopy. *Science*, **248**, 73-76.
- Deplewski, D. and Rosenfield, R.L. (1999) Growth hormone and insulin-like growth factors have different effects on sebaceous cell growth and differentiation. *Endocrinology*, **140**, 4089-4094.
- Deplewski, D. and Rosenfield, R.L. (2000) Role of hormones in pilosebaceous unit development. *Endocr Rev*, **21**, 363-392.
- DeVillez, R.L. (1987) Topical minoxidil for androgenetic alopecia: optimizing the chance for success by appropriate patient selection. *Dermatologica*, **175 Suppl 2**, 50-53.

- Dreher, F., Modjtahedi, B.S., Modjtahedi, S.P. and Maibach, H.I. (2005) Quantification of stratum corneum removal by adhesive tape stripping by total protein assay in 96-well microplates. *Skin Res Technol*, **11**, 97-101.
- du Cros, D.L. (1993) Fibroblast growth factor and epidermal growth factor in hair development. *J Invest Dermatol*, **101**, 106S-113S.
- Dumont, M., Luu-The, V., Dupont, E., Pelletier, G. and Labrie, F. (1992) Characterization, expression, and immunohistochemical localization of 3 beta-hydroxysteroid dehydrogenase/delta 5-delta 4 isomerase in human skin. *J Invest Dermatol*, **99**, 415-421.
- Epstein, E. and Epstein, W. (1966) New cell formation in human sebaceous glands. *J Invest Dermatol*, **46**, 453-458.
- Fiedler, H.P. (1996) Lexikon der Hilfsstoffe für Pharmazie, Kosmetik und angrenzende Gebiete. *Editio Cantor Verlag Aulenhof*, **4. Auflage**.
- Fimmel, S., Saborowski, A., Orfanos, C.E. and Zouboulis, C.C. (2000) Development of efficient transient transfection systems for introducing antisense oligonucleotides into human epithelial skin cells. *Horm Res*, **54**, 306-311.
- Freitas, C. and Müller, R.H. (1999) Correlation between long-term stability of solid lipid nanoparticles (SLN) and crystallinity of the lipid phase. *Eur J Pharm Biopharm*, **47**, 125-132.
- Frith, R.G. and Phillipou, G. (1985) 15-Hydroxycyproterone acetate and cyproterone acetate levels in plasma and urine. *J Chromatogr*, **338**, 179-186.
- Fritsch, M., Orfanos, C.E. and Zouboulis, C.C. (2001) Sebocytes are the key regulators of androgen homeostasis in human skin. *J Invest Dermatol*, **116**, 793-800.
- Gao, W., Bohl, C.E. and Dalton, J.T. (2005) Chemistry and structural biology of androgen receptor. *Chem Rev*, **105**, 3352-3370.
- Gloor, M., Wirth, H. and Swoboda, U. (1981) Is sebosuppression by cimetidine an antiandrogenic effect? *Acta Derm Venereol*, **61**, 262-264.
- Golditch, I.M. and Price, V.H. (1990) Treatment of hirsutism with cimetidine. *Obstet Gynecol*, **75**, 911-913.
- Gollnick, H.P. and Krauthaim, A. (2003) Topical treatment in acne: current status and future aspects. *Dermatology*, **206**, 29-36.
- Gruber, D.M., Sator, M.O., Joura, E.A., Kokoschka, E.M., Heinze, G. and Huber, J.C. (1998) Topical cyproterone acetate treatment in women with acne: a placebo-controlled trial. *Arch Dermatol*, **134**, 459-463.
- Gunther, C., Kecskes, A., Staks, T. and Tauber, U. (1998) Percutaneous absorption of methylprednisolone aceponate following topical application of Advantan lotion on intact, inflamed and stripped skin of male volunteers. *Skin Pharmacol Appl Skin Physiol*, **11**, 35-42.
- Guy, R., Ridden, C. and Kealey, T. (1996) The improved organ maintenance of the human sebaceous gland: modeling in vitro the effects of epidermal growth factor, androgens, estrogens, 13-cis retinoic acid, and phenol red. *J Invest Dermatol*, **106**, 454-460.
- Gysler, A., Kleuser, B., Sippl, W., Lange, K., Korting, H.C. and Holtje, H.D. (1999) Skin penetration and metabolism of topical glucocorticoids in reconstructed epidermis and in excised human skin. *Pharm Res*, **16**, 1386-1391.
- Gysler, A., Lange, K., Korting, H.C. and Schafer-Korting, M. (1997) Prednicarbate biotransformation in human foreskin keratinocytes and fibroblasts. *Pharm Res*, **14**, 793-797.

- Hamilton, J.B. (1958). Effects of castration in adolescent and young adult males upon further changes in the proportions of bare and hairy scalp. *J Clin Endocrinol Matab* **20**, 1308.
- Hatziantoniou, S., Rallis, M., Demetzos, C. and Papaioannou, G.T. (2000) Pharmacological activity of natural lipids on a skin barrier disruption model. *Pharmacol Res*, **42**, 55-59.
- Henzen, C. (2001) Hirsutimus. *Schweiz Med Forum*, **44**, 1104-1108.
- Hibberts, N.A., Howell, A.E. and Randall, V.A. (1998) Balding hair follicle dermal papilla cells contain higher levels of androgen receptors than those from non-balding scalp. *J Endocrinol*, **156**, 59-65.
- Hibino, T. and Nishiyama, T. (2004) Role of TGF-beta2 in the human hair cycle. *J Dermatol Sci*, **35**, 9-18.
- Hillmer, A.M., Hanneken, S., Ritzmann, S., Becker, T., Freudenberg, J., Brockschmidt, F.F., Flaquer, A., Freudenberg-Hua, Y., Jamra, R.A., Metzen, C., Heyn, U., Schweiger, N., Betz, R.C., Blaumeiser, B., Hampe, J., Schreiber, S., Schulze, T.G., Hennies, H.C., Schumacher, J., Propping, P., Ruzicka, T., Cichon, S., Wienker, T.F., Kruse, R. and Nothen, M.M. (2005) Genetic variation in the human androgen receptor gene is the major determinant of common early-onset androgenetic alopecia. *Am J Hum Genet*, **77**, 140-148.
- Hoffmann, R. and Happle, R. (1999) [Pathogenetic significance of 5-alpha reductase isoenzymes for androgenetic alopecia]. *Hautarzt*, **50**, 165-173.
- Hu, F.Q., Yuan, H., Zhang, H.H. and Fang, M. (2002) Preparation of solid lipid nanoparticles with clobetasol propionate by a novel solvent diffusion method in aqueous system and physicochemical characterization. *Int J Pharm*, **239**, 121-128.
- Hueber, F., Schaefer, H. and Wepierre, J. (1994) Role of transepidermal and transfollicular routes in percutaneous absorption of steroids: in vitro studies on human skin. *Skin Pharmacol*, **7**, 237-244.
- Inui, S., Itami, S., Pan, H.J. and Chang, C. (2000) Lack of androgen receptor transcriptional activity in human keratinocytes. *J Dermatol Sci*, **23**, 87-92.
- Itami, S., Kurata, S. and Takayasu, S. (1995) Androgen induction of follicular epithelial cell growth is mediated via insulin-like growth factor-I from dermal papilla cells. *Biochem Biophys Res Commun*, **212**, 988-994.
- Jacobi, U., Weigmann, H.J., Ulrich, J., Sterry, W. and Lademann, J. (2005) Estimation of the relative stratum corneum amount removed by tape stripping. *Skin Res Technol*, **11**, 91-96.
- Jat, P.S., Noble, M.D., Ataliotis, P., Tanaka, Y., Yannoutsos, N., Larsen, L. and Kioussis, D. (1991) Direct derivation of conditionally immortal cell lines from an H-2Kb-tsA58 transgenic mouse. *Proc Natl Acad Sci U S A*, **88**, 5096-5100.
- Jenning, V. (1999) Feste Lipid-Nanopartikel (SLN<sup>TM</sup>) als Trägersysteme für die dermale Applikation von Retinol: Wirkstoffinkorporation, -freisetzung und Struktur. *Dissertation Freie Universität Berlin*.
- Jenning, V., Mader, K. and Gohla, S.H. (2000a) Solid lipid nanoparticles (SLN) based on binary mixtures of liquid and solid lipids: a (1)H-NMR study. *Int J Pharm*, **205**, 15-21.
- Jenning, V., Schafer-Korting, M. and Gohla, S. (2000b) Vitamin A-loaded solid lipid nanoparticles for topical use: drug release properties. *J Control Release*, **66**, 115-126.

- Jores, K., Mehnert, W. and Mäder, K. (2003) Physicochemical investigations on solid lipid nanoparticles and on oil-loaded solid lipid nanoparticles: a nuclear magnetic resonance and electron spin resonance study. *Pharm Res*, **20**, 1274-1283.
- Kahlig, H., Valenta, C., Dampfhart, U. and Auner, B.G. (2005) Rheology and NMR self-diffusion experiments as well as skin permeation of diclofenac-sodium and cyproterone acetate of new gel preparations. *J Pharm Sci*, **94**, 288-296.
- Kalariya, M., Padhi, B.K., Chougule, M. and Misra, A. (2005) Clobetasol propionate solid lipid nanoparticles cream for effective treatment of eczema: formulation and clinical implications. *Indian J Exp Biol*, **43**, 233-240.
- Kealey, T., Lee, C.M., Thody, A.J. and Coaker, T. (1986) The isolation of human sebaceous glands and apocrine sweat glands by shearing. *Br J Dermatol*, **114**, 181-188.
- König, K., Ehlers, A., Stracke, F., Riemann, I. (2006) In vivo Drug Screening in Human Skin Using Femtosecond Laser Multiphoton Tomography. *Skin Pharmacol Physiol*, **19**, 78-88.
- Kremer, F. and Schoenhals, A. (2002) Broadband dielectric spectroscopy. *Springer, Berlin*, 35-56.
- Labrie, F., Luu-The, V., Labrie, C., Pelletier, G. and El-Alfy, M. (2000) Intracrinology and the skin. *Horm Res*, **54**, 218-229.
- Lademann, J., Otberg, N., Richter, H., Weigmann, H.J., Lindemann, U., Schaefer, H. and Sterry, W. (2001) Investigation of follicular penetration of topically applied substances. *Skin Pharmacol Appl Skin Physiol*, **14 Suppl 1**, 17-22.
- Lademann, J., Schaefer, H., Otberg, N., Teichmann, A., Blume-Peytavi, U. and Sterry, W. (2004) [Penetration of microparticles into human skin.]. *Hautarzt*, **55**, 1117-1119.
- Lemay, A. and Poulin, Y. (2002) Oral contraceptives as anti-androgenic treatment of acne. *J Obstet Gynaecol Can*, **24**, 559-567.
- Liang, T., Rasmusson, G.H. and Brooks, J.R. (1983) 12. Androgens: Pharmacodynamics and antagonists. Biochemical and biological studies with 4-aza-steroidal 5 alpha-reductase inhibitors. *J Steroid Biochem*, **19**, 385-390.
- Lieb, L.M., Liimatta, A.P., Bryan, R.N., Brown, B.D. and Krueger, G.G. (1997) Description of the intrafollicular delivery of large molecular weight molecules to follicles of human scalp skin in vitro. *J Pharm Sci*, **86**, 1022-1029.
- Lissak, A., Sorokin, Y., Calderon, I., Dirnfeld, M., Lioz, H. and Abramovici, H. (1989) Treatment of hirsutism with cimetidine: a prospective randomized controlled trial. *Fertil Steril*, **51**, 247-250.
- List, H.J., Lozano, C., Lu, J., Danielsen, M., Wellstein, A. and Riegel, A.T. (1999) Comparison of chromatin remodeling and transcriptional activation of the mouse mammary tumor virus promoter by the androgen and glucocorticoid receptor. *Exp Cell Res*, **250**, 414-422.
- Loffler, H., Dreher, F. and Maibach, H.I. (2004) Stratum corneum adhesive tape stripping: influence of anatomical site, application pressure, duration and removal. *Br J Dermatol*, **151**, 746-752.
- Lombardi Borgia, S., Regehly, M., Sivaramakrishnan, R., Mehnert, W., Korting, H.C., Danker, K., Roder, B., Kramer, K.D. and Schafer-Korting, M. (2005) Lipid nanoparticles for skin penetration enhancement-correlation to drug localization within the particle matrix as determined by fluorescence and parelectric spectroscopy. *J Control Release*, **110**, 151-163.

- Luderschmidt, C., Jawny, J. and Eiermann, W. (1987) Relative binding affinity at metribolone androgenic binding sites of various antiandrogenic agents. *Arzneimittelforschung*, **37**, 1262-1265.
- Maia, C.S., Mehnert, W. and Schafer-Korting, M. (2000) Solid lipid nanoparticles as drug carriers for topical glucocorticoids. *Int J Pharm*, **196**, 165-167.
- Matias, J.R. and Gaillard, M. (1995) Local inhibition of sebaceous gland growth by topically applied RU 58841. *Ann N Y Acad Sci*, **761**, 56-65.
- Mehnert, W. and Mäder, K. (2001) Solid lipid nanoparticles: production, characterization and applications. *Adv Drug Deliv Rev*, **47**, 165-196.
- Menard, R.H., Martin, H.F., Stripp, B., Gillette, J.R. and Bartter, F.C. (1974) Spironolactone and cytochrome P-450: impairment of steroid hydroxylation in the adrenal cortex. *Life Sci*, **15**, 1639-1648.
- Mordon, S., Sumian, C. and Devoisselle, J.M. (2003) Site-specific methylene blue delivery to pilosebaceous structures using highly porous nylon microspheres: an experimental evaluation. *Lasers Surg Med*, **33**, 119-125.
- Morgan, C.J., Renwick, A.G. and Friedmann, P.S. (2003) The role of stratum corneum and dermal microvascular perfusion in penetration and tissue levels of water-soluble drugs investigated by microdialysis. *Br J Dermatol*, **148**, 434-443.
- Müller, R.H., Radtke, M. and Wissing, S.A. (2002) Solid lipid nanoparticles (SLN) and nanostructured lipid carriers (NLC) in cosmetic and dermatological preparations. *Adv Drug Deliv Rev*, **54 Suppl 1**, S131-155.
- Münster, U., Nakamura, C., Haberland, A., Jores, K., Mehnert, W., Rummel, S., Schaller, M., Korting, H.C., Zouboulis Ch, C., Blume-Peytavi, U. and Schafer-Korting, M. (2005) RU 58841-myristate--prodrug development for topical treatment of acne and androgenetic alopecia. *Pharmazie*, **60**, 8-12.
- Nakamura, C. (2004) Steuerung der kutanen Glucocorticoid-Resorption: Einfluss von festen Lipid-Nanopartikeln. *Dissertation Freie Universität Berlin*.
- Nicolau, G., Baughman, R.A., Tonelli, A., McWilliams, W., Schiltz, J. and Yacobi, A. (1987) Deposition of viprostol (a synthetic PGE2 vasodilator) in the skin following topical administration to laboratory animals. *Xenobiotica*, **17**, 1113-1120.
- Nikkari, T. (1974) Comparative chemistry of sebum. *J Invest Dermatol*, **62**, 257-267.
- Obana, N., Chang, C. and Uno, H. (1997) Inhibition of hair growth by testosterone in the presence of dermal papilla cells from the frontal bald scalp of the postpubertal stump-tailed macaque. *Endocrinology*, **138**, 356-361.
- Otberg, N., Richter, H., Schaefer, H., Blume-Peytavi, U., Sterry, W. and Lademann, J. (2003) Visualisation of Topically Applied Fluorescent Dyes in Hair Follicles by Laser Scanning Microscopy. *Laser Physics*, **13**, 761-764.
- Paus, R. and Cotsarelis, G. (1999) The biology of hair follicles. *N Engl J Med*, **341**, 491-497.
- Philpott, M.P., Sanders, D.A. and Kealey, T. (1994) Effects of insulin and insulin-like growth factors on cultured human hair follicles: IGF-I at physiologic concentrations is an important regulator of hair follicle growth in vitro. *J Invest Dermatol*, **102**, 857-861.
- Plettenberg, A., Meigel, W. and I, M. (1999) Dermatologie an der Schwelle zum neuen Jahrtausend. *berichte von der 40. Tagung der Deutschen dermatologischen Gesellschaft*, 427.
- Price, V.H., Roberts, J.L., Hordinsky, M., Olsen, E.A., Savin, R., Bergfeld, W., Fiedler, V., Lucky, A., Whiting, D.A., Pappas, F., Culbertson, J., Kotey, P., Meehan, A.

- and Waldstreicher, J. (2000) Lack of efficacy of finasteride in postmenopausal woman with androgenetic alopecia. *J Am Acad Dermatol*, **43**, 768-776.
- Radtke, M. (2002) Nanostructured lipid carriers (NLC): Untersuchungen zur Struktur, Wirkstoffinkorporation und Stabilität. *Dissertation Freie Universität Berlin*.
- Randall, V.A., Thornton, M.J. and Messenger, A.G. (1992) Cultured dermal papilla cells from androgen-dependent human hair follicles (e.g. beard) contain more androgen receptors than those from non-balding areas of scalp. *J Endocrinol*, **133**, 141-147.
- Rosenfield, R.L., Deplewski, D., Kentsis, A. and Ciletti, N. (1998) Mechanisms of androgen induction of sebocyte differentiation. *Dermatology*, **196**, 43-46.
- Rudman, S.M., Philpott, M.P., Thomas, G.A. and Kealey, T. (1997) The role of IGF-I in human skin and its appendages: morphogen as well as mitogen? *J Invest Dermatol*, **109**, 770-777.
- Rushton, D.H., Norris, J. and Ramsay, I.D. (1996) Topical 0,05% finasteride significantly reduced serum DHT concentrations, but had no influence in preventing the expression of genetic hair loss in men. *Hair Research for the Next Millenium*, 359-362.
- Sanna, V., Kirschvink, N., Gustin, P., Gavini, E., Roland, I., Delattre, L. and Evrard, B. (2004) Preparation and in vivo toxicity study of solid lipid microparticles as carrier for pulmonary administration. *AAPS PharmSciTech*, **5**, e27.
- Santos Maia, C., Mehnert, W., Schaller, M., Korting, H.C., Gysler, A., Haberland, A. and Schafer-Korting, M. (2002) Drug targeting by solid lipid nanoparticles for dermal use. *J Drug Target*, **10**, 489-495.
- Savin, R.C. (1987) Use of topical minoxidil in the treatment of male pattern baldness. *J Am Acad Dermatol*, **16**, 696-704.
- Sawaya, M.E. (1997) Clinical updates in hair. *Dermatol Clin*, **15**, 37-43.
- Sawaya, M.E. and Penneys, N.S. (1992) Immunohistochemical distribution of aromatase and 3B-hydroxysteroid dehydrogenase in human hair follicle and sebaceous gland. *J Cutan Pathol*, **19**, 309-314.
- Schaefer, H., Watts and Brod. (1990) Prediction of percutaneous Penetration: Methods, Measurements and Modelling.
- Schäfer-Korting, M., Kleuser, B., Ahmed, M., Holtje, H.D. and Korting, H.C. (2005) Glucocorticoids for human skin: new aspects of the mechanism of action. *Skin Pharmacol Physiol*, **18**, 103-114.
- Schmalfuß, U., Neubert, R. and Wohlrab, W. (1996) Modification of drug penetration into human skin using microemulsions. *J Control Release*, **46**, 279-285.
- Schneider, I.M., Wohlrab, W. and Neubert, R. (1997) [Fatty acids and the epidermis]. *Hautarzt*, **48**, 303-310.
- Schreiber, S., Mahmoud, A., Vuia, A., Rubbelke, M.K., Schmidt, E., Schaller, M., Kandarova, H., Haberland, A., Schafer, U.F., Bock, U., Korting, H.C., Liebsch, M. and Schafer-Korting, M. (2005) Reconstructed epidermis versus human and animal skin in skin absorption studies. *Toxicol In Vitro*, **19**, 813-822.
- Serafini, P.C., Catalino, J. and Lobo, R.A. (1985) The effect of spironolactone on genital skin 5 alpha-reductase activity. *J Steroid Biochem*, **23**, 191-194.
- Sivaramakrishnan, R., Nakamura, C., Mehnert, W., Korting, H.C., Kramer, K.D. and Schafer-Korting, M. (2004) Glucocorticoid entrapment into lipid carriers--characterisation by paelectric spectroscopy and influence on dermal uptake. *J Control Release*, **97**, 493-502.
- Slayden, S.M., Moran, C., Sams, W.M., Jr., Boots, L.R. and Azziz, R. (2001) Hyperandrogenemia in patients presenting with acne. *Fertil Steril*, **75**, 889-892.

- Stenn, K.S. and Paus, R. (2001) Controls of hair follicle cycling. *Physiol Rev*, **81**, 449-494.
- Strauss, J.S., Stranieri, A.M., Farrell, L.N. and Downing, D.T. (1980) The effect of marked inhibition of sebum production with 13cis-retinoic acid on skin surface lipid composition. *J Invest Dermatol*, **74**, 66-67.
- Thiboutot, D., Martin, P., Volikos, L. and Gilliland, K. (1998) Oxidative activity of the type 2 isozyme of 17beta-hydroxysteroid dehydrogenase (17beta-HSD) predominates in human sebaceous glands. *J Invest Dermatol*, **111**, 390-395.
- Toll, R., Jacobi, U., Richter, H., Lademann, J., Schaefer, H. and Blume-Peytavi, U. (2004) Penetration profile of microspheres in follicular targeting of terminal hair follicles. *J Invest Dermatol*, **123**, 168-176.
- Trotta, M., Cavalli, R., Carlotti, M.E., Battaglia, L. and Debernardi, F. (2005) Solid lipid micro-particles carrying insulin formed by solvent-in-water emulsion-diffusion technique. *Int J Pharm*, **288**, 281-288.
- Trueb, R.M. and Meyer, J.C. (2000) Male-pattern baldness in men with X-linked recessive ichthyosis. *Dermatology*, **200**, 247-249.
- Tur, E., Maibach, H.I. and Guy, R.H. (1991) Percutaneous penetration of methyl nicotinate at three anatomic sites: evidence for an appendageal contribution to transport? *Skin Pharmacol*, **4**, 230-234.
- Valenta, C. and Janisch, M. (2003) Permeation of cyproterone acetate through pig skin from different vehicles with phospholipids. *Int J Pharm*, **258**, 133-139.
- Van Neste, D., Fuh, V., Sanchez-Pedreno, P., Lopez-Bran, E., Wolff, H., Whiting, D., Roberts, J., Kopera, D., Stene, J.J., Calvieri, S., Tosti, A., Prens, E., Guarrera, M., Kanojia, P., He, W. and Kaufman, K.D. (2000) Finasteride increases anagen hair in men with androgenetic alopecia. *Br J Dermatol*, **143**, 804-810.
- Weigmann, H., Lademann, J., Meffert, H., Schaefer, H. and Sterry, W. (1999) Determination of the horny layer profile by tape stripping in combination with optical spectroscopy in the visible range as a prerequisite to quantify percutaneous absorption. *Skin Pharmacol Appl Skin Physiol*, **12**, 34-45.
- Weindl, G., Schafer-Korting, M., Schaller, M. and Korting, H.C. (2005) Peroxisome proliferator-activated receptors and their ligands: entry into the post-glucocorticoid era of skin treatment? *Drugs*, **65**, 1919-1934.
- Werner, S., Smola, H., Liao, X., Longaker, M.T., Krieg, T., Hofschneider, P.H. and Williams, L.T. (1994) The function of KGF in morphogenesis of epithelium and reepithelialization of wounds. *Science*, **266**, 819-822.
- Westesen, K. and Siekmann, B. (1997) Investigation of the gel formation of phospholipid-stabilised solid lipid nanoparticles. *Int J Pharm*, **151**, 35-45.
- Xia, L.Q., Zouboulis, C., Detmar, M., Mayer-da-Silva, A., Stadler, R. and Orfanos, C.E. (1989) Isolation of human sebaceous glands and cultivation of sebaceous gland-derived cells as an in vitro model. *J Invest Dermatol*, **93**, 315-321.
- Yodo, K., Saisho, S., Shimozawa, K. and Yata, J. (1988) A reversed-phase high-performance liquid chromatographic method for the simultaneous determination of serum concentrations of cyproterone acetate and 15 beta-hydroxycyproterone acetate. *Endocrinol Jpn*, **35**, 143-148.
- Zimmermann, E. and Müller, R.H. (2001) Electrolyte- and pH-stabilities of aqueous solid lipid nanoparticle (SLN) dispersions in artificial gastrointestinal media. *Eur J Pharm Biopharm*, **52**, 203-210.
- Zouboulis, C.C. (2001) Is acne vulgaris a genuine inflammatory disease? *Dermatology*, **203**, 277-279.



- Zouboulis, C.C. (2004) Acne and sebaceous gland function. *Clin Dermatol*, **22**, 360-366.
- Zouboulis, C.C., Korge, B., Akamatsu, H., Xia, L.Q., Schiller, S., Gollnick, H. and Orfanos, C.E. (1991) Effects of 13-cis-retinoic acid, all-trans-retinoic acid, and acitretin on the proliferation, lipid synthesis and keratin expression of cultured human sebocytes in vitro. *J Invest Dermatol*, **96**, 792-797.
- Zouboulis, C.C. and Piquero-Martin, J. (2003) Update and future of systemic acne treatment. *Dermatology*, **206**, 37-53.
- Zouboulis, C.C., Seltmann, H., Hiroi, N., Chen, W., Young, M., Oeff, M., Scherbaum, W.A., Orfanos, C.E., McCann, S.M. and Bornstein, S.R. (2002) Corticotropin-releasing hormone: an autocrine hormone that promotes lipogenesis in human sebocytes. *Proc Natl Acad Sci U S A*, **99**, 7148-7153.
- Zouboulis, C.C., Seltmann, H., Neitzel, H. and Orfanos, C.E. (1999) Establishment and characterization of an immortalized human sebaceous gland cell line (SZ95). *J Invest Dermatol*, **113**, 1011-1020.
- Zouboulis, C.C., Xia, L., Akamatsu, H., Seltmann, H., Fritsch, M., Hornemann, S., Ruhl, R., Chen, W., Nau, H. and Orfanos, C.E. (1998) The human sebocyte culture model provides new insights into development and management of seborrhoea and acne. *Dermatology*, **196**, 21-31.
- Zouboulis Ch, C., Saborowski, A. and Boschnakow, A. (2005) Zileuton, an oral 5-lipoxygenase inhibitor, directly reduces sebum production. *Dermatology*, **210**, 36-38.
- zur Muhlen, A., Schwarz, C. and Mehnert, W. (1998) Solid lipid nanoparticles (SLN) for controlled drug delivery-drug release and release mechanism. *Eur J Pharm Biopharm*, **45**, 149-155.