

6 References

- (1993). "Identification and characterization of the tuberous sclerosis gene on chromosome 16. The European Chromosome 16 Tuberous Sclerosis Consortium." *Cell* **75**(7): 1305-15.
- Abraham, R. T. and G. J. Wiederrecht (1996). "Immunopharmacology of rapamycin." *Annu Rev Immunol* **14**: 483-510.
- Alessi, D. R., M. T. Kozlowski, et al. (1998). "3-Phosphoinositide-dependent protein kinase 1 (PDK1) phosphorylates and activates the p70 S6 kinase in vivo and in vitro." *Curr Biol* **8**(2): 69-81.
- Andrade, M. A. and P. Bork (1995). "HEAT repeats in the Huntington's disease protein." *Nat Genet* **11**(2): 115-6.
- Barbet, N. C., U. Schneider, et al. (1996). "TOR controls translation initiation and early G1 progression in yeast." *Mol Biol Cell* **7**(1): 25-42.
- Beugnet, A., A. R. Tee, et al. (2003). "Regulation of targets of mTOR (mammalian target of rapamycin) signalling by intracellular amino acid availability." *Biochem J* **372**(Pt 2): 555-66.
- Biondi, R. M., A. Kieloch, et al. (2001). "The PIF-binding pocket in PDK1 is essential for activation of S6K and SGK, but not PKB." *Embo J* **20**(16): 4380-90.
- Bosotti, R., A. Isacchi, et al. (2000). "FAT: a novel domain in PIK-related kinases." *Trends Biochem Sci* **25**(5): 225-7.
- Brown, E. J., M. W. Albers, et al. (1994). "A mammalian protein targeted by G1-arresting rapamycin-receptor complex." *Nature* **369**(6483): 756-8.
- Brunn, G. J., P. Fadden, et al. (1997). "The mammalian target of rapamycin phosphorylates sites having a (Ser/Thr)-Pro motif and is activated by antibodies to a region near its COOH terminus." *J Biol Chem* **272**(51): 32547-50.
- Brunn, G. J., C. C. Hudson, et al. (1997). "Phosphorylation of the translational repressor PHAS-I by the mammalian target of rapamycin." *Science* **277**(5322): 99-101.
- Burnett, P. E., R. K. Barrow, et al. (1998). "RAFT1 phosphorylation of the translational regulators p70 S6 kinase and 4E-BP1." *Proc Natl Acad Sci U S A* **95**(4): 1432-7.
- Cafferkey, R., P. R. Young, et al. (1993). "Dominant missense mutations in a novel yeast protein related to mammalian phosphatidylinositol 3-kinase and VPS34 abrogate rapamycin cytotoxicity." *Mol Cell Biol* **13**(10): 6012-23.
- Cantley, L. C. and B. G. Neel (1999). "New insights into tumor suppression: PTEN suppresses tumor formation by restraining the phosphoinositide 3-kinase/AKT pathway." *Proc Natl Acad Sci U S A* **96**(8): 4240-5.
- Cheatham, L., M. Monfar, et al. (1995). "Structural and functional analysis of pp70S6k." *Proc Natl Acad Sci U S A* **92**(25): 11696-700.
- Chen, J., X. F. Zheng, et al. (1995). "Identification of an 11-kDa FKBP12-rapamycin-binding domain within the 289-kDa FKBP12-rapamycin-

- associated protein and characterization of a critical serine residue." Proc Natl Acad Sci U S A **92**(11): 4947-51.
- Chen, R. H., C. Sarnecki, et al. (1992). "Nuclear localization and regulation of erk- and rsk-encoded protein kinases." Mol Cell Biol **12**(3): 915-27.
- Chiu, M. I., H. Katz, et al. (1994). "RAPT1, a mammalian homolog of yeast Tor, interacts with the FKBP12/rapamycin complex." Proc Natl Acad Sci U S A **91**(26): 12574-8.
- Choi, K. M., L. P. McMahon, et al. (2003). "Two Motifs in the Translational Repressor PHAS-I Required for Efficient Phosphorylation by Mammalian Target of Rapamycin and for Recognition by Raptor." J Biol Chem **278**(22): 19667-73.
- Chou, M. M. and J. Blenis (1996). "The 70 kDa S6 kinase complexes with and is activated by the Rho family G proteins Cdc42 and Rac1." Cell **85**(4): 573-83.
- Chung, J., C. J. Kuo, et al. (1992). "Rapamycin-FKBP specifically blocks growth-dependent activation of and signaling by the 70 kd S6 protein kinases." Cell **69**(7): 1227-36.
- Coffer, P. J. and J. R. Woodgett (1994). "Differential subcellular localisation of two isoforms of p70 S6 protein kinase." Biochem Biophys Res Commun **198**(2): 780-6.
- Conlon, I. and M. Raff (1999). "Size control in animal development." Cell **96**(2): 235-44.
- De Benedetti, A., S. Joshi-Barve, et al. (1991). "Expression of antisense RNA against initiation factor eIF-4E mRNA in HeLa cells results in lengthened cell division times, diminished translation rates, and reduced levels of both eIF-4E and the p220 component of eIF-4F." Mol Cell Biol **11**(11): 5435-45.
- Dennis, P. B., A. Jaeschke, et al. (2001). "Mammalian TOR: a homeostatic ATP sensor." Science **294**(5544): 1102-5.
- Dennis, P. B., N. Pullen, et al. (1996). "The principal rapamycin-sensitive p70(s6k) phosphorylation sites, T-229 and T-389, are differentially regulated by rapamycin-insensitive kinase kinases." Mol Cell Biol **16**(11): 6242-51.
- Di Como, C. J. and K. T. Arndt (1996). "Nutrients, via the Tor proteins, stimulate the association of Tap42 with type 2A phosphatases." Genes Dev **10**(15): 1904-16.
- Diggle, T. A., C. Schmitz-Peiffer, et al. (1991). "Evidence that insulin activates casein kinase 2 in rat epididymal fat- cells and that this may result in the increased phosphorylation of an acid-soluble 22 kDa protein." Biochem J **279**(Pt 2): 545-51.
- Dolinski, K., S. Muir, et al. (1997). "All cyclophilins and FK506 binding proteins are, individually and collectively, dispensable for viability in *Saccharomyces cerevisiae*." Proc Natl Acad Sci U S A **94**(24): 13093-8.

- Downward, J. (1998). "Lipid-regulated kinases: some common themes at last." *Science* **279**(5351): 673-4.
- Dufner, A., M. Andjelkovic, et al. (1999). "Protein kinase B localization and activation differentially affect S6 kinase 1 activity and eukaryotic translation initiation factor 4E-binding protein 1 phosphorylation." *Mol Cell Biol* **19**(6): 4525-34.
- Dufner, A. and G. Thomas (1999). "Ribosomal S6 kinase signaling and the control of translation." *Exp Cell Res* **253**(1): 100-9.
- Duncan, R., S. C. Milburn, et al. (1987). "Regulated phosphorylation and low abundance of HeLa cell initiation factor eIF-4F suggest a role in translational control. Heat shock effects on eIF-4F." *J Biol Chem* **262**(1): 380-8.
- Elit, L. (2002). "CCI-779 Wyeth." *Curr Opin Investig Drugs* **3**(8): 1249-53.
- Fadden, P., T. A. Haystead, et al. (1997). "Identification of phosphorylation sites in the translational regulator, PHAS-I, that are controlled by insulin and rapamycin in rat adipocytes." *J Biol Chem* **272**(15): 10240-7.
- Fang, Y., M. Vilella-Bach, et al. (2001). "Phosphatidic acid-mediated mitogenic activation of mTOR signaling." *Science* **294**(5548): 1942-5.
- Fingar, D. C., S. Salama, et al. (2002). "Mammalian cell size is controlled by mTOR and its downstream targets S6K1 and 4EBP1/eIF4E." *Genes Dev* **16**(12): 1472-87.
- Gao, X., Y. Zhang, et al. (2002). "Tsc tumour suppressor proteins antagonize amino-acid-TOR signalling." *Nat Cell Biol* **4**(9): 699-704.
- Gingras, A. C., S. P. Gygi, et al. (1999). "Regulation of 4E-BP1 phosphorylation: a novel two-step mechanism." *Genes Dev* **13**(11): 1422-37.
- Gingras, A. C., S. G. Kennedy, et al. (1998). "4E-BP1, a repressor of mRNA translation, is phosphorylated and inactivated by the Akt(PKB) signaling pathway." *Genes Dev* **12**(4): 502-13.
- Gingras, A. C., B. Raught, et al. (2001). "Hierarchical phosphorylation of the translation inhibitor 4E-BP1." *Genes Dev* **15**(21): 2852-64.
- Gingras, A. C., B. Raught, et al. (1999). "eIF4 initiation factors: effectors of mRNA recruitment to ribosomes and regulators of translation." *Annu Rev Biochem* **68**: 913-63.
- Gingras, A. C., B. Raught, et al. (2001). "Control of translation by the target of rapamycin proteins." *Prog Mol Subcell Biol* **27**: 143-74.
- Gingras, A. C., B. Raught, et al. (2001). "Regulation of translation initiation by FRAP/mTOR." *Genes Dev* **15**(7): 807-26.
- Goncharova, E. A., D. A. Goncharov, et al. (2002). "Tuberin regulates p70 S6 kinase activation and ribosomal protein S6 phosphorylation. A role for the TSC2 tumor suppressor gene in pulmonary lymphangioleiomyomatosis (LAM)." *J Biol Chem* **277**(34): 30958-67.
- Gothel, S. F. and M. A. Marahiel (1999). "Peptidyl-prolyl cis-trans isomerases, a superfamily of ubiquitous folding catalysts." *Cell Mol Life Sci* **55**(3): 423-36.

- Gout, I., T. Minami, et al. (1998). "Molecular cloning and characterization of a novel p70 S6 kinase, p70 S6 kinase beta containing a proline-rich region." *J Biol Chem* **273**(46): 30061-4.
- Grammer, T. C., L. Cheatham, et al. (1996). "The p70S6K signalling pathway: a novel signalling system involved in growth regulation." *Cancer Surv* **27**: 271-92.
- Groot, R. P. d., L. M. Ballou, et al. (1994). "Positive regulation of the cAMP-responsive activator CREM by the p70 S6 Kinase: an alternative route to mitogen-induced gene expression." *Cell* **79**: 81-91.
- Groves, M. R. and D. Barford (1999). "Topological characteristics of helical repeat proteins." *Curr Opin Struct Biol* **9**(3): 383-9.
- Han, J. W., R. B. Pearson, et al. (1995). "Rapamycin, wortmannin, and the methylxanthine SQ20006 inactivate p70s6k by inducing dephosphorylation of the same subset of sites." *J Biol Chem* **270**(36): 21396-403.
- Hara, K., Y. Maruki, et al. (2002). "Raptor, a binding partner of target of rapamycin (TOR), mediates TOR action." *Cell* **110**(2): 177-89.
- Hara, K., K. Yonezawa, et al. (1998). "Amino acid sufficiency and mTOR regulate p70 S6 kinase and eIF-4E BP1 through a common effector mechanism." *J Biol Chem* **273**(23): 14484-94.
- Harada, H., J. S. Andersen, et al. (2001). "p70S6 kinase signals cell survival as well as growth, inactivating the pro-apoptotic molecule BAD." *Proc Natl Acad Sci U S A* **98**(17): 9666-70.
- Hardie, D. G. and S. A. Hawley (2001). "AMP-activated protein kinase: the energy charge hypothesis revisited." *Bioessays* **23**(12): 1112-9.
- Hartwell, L. H. (1967). "Macromolecule synthesis in temperature-sensitive mutants of yeast." *J Bacteriol* **93**(5): 1662-70.
- Hashemolhosseini, S., Y. Nagamine, et al. (1998). "Rapamycin inhibition of the G1 to S transition is mediated by effects on cyclin D1 mRNA and protein stability." *J Biol Chem* **273**(23): 14424-9.
- Haystead, T. A., C. M. Haystead, et al. (1994). "Phosphorylation of PHAS-I by mitogen-activated protein (MAP) kinase. Identification of a site phosphorylated by MAP kinase in vitro and in response to insulin in rat adipocytes." *J Biol Chem* **269**(37): 23185-91.
- Heesom, K. J., M. B. Avison, et al. (1998). "Insulin-stimulated kinase from rat fat cells that phosphorylates initiation factor 4E-binding protein 1 on the rapamycin-insensitive site (serine-111)." *Biochem J* **336**(Pt 1): 39-48.
- Heesom, K. J. and R. M. Denton (1999). "Dissociation of the eukaryotic initiation factor-4E/4E-BP1 complex involves phosphorylation of 4E-BP1 by an mTOR-associated kinase." *FEBS Lett* **457**(3): 489-93.
- Heitman, J., N. R. Movva, et al. (1991). "Targets for cell cycle arrest by the immunosuppressant rapamycin in yeast." *Science* **253**(5022): 905-9.
- Helliwell, S. B., P. Wagner, et al. (1994). "TOR1 and TOR2 are structurally and functionally similar but not identical phosphatidylinositol kinase homologues in yeast." *Mol Biol Cell* **5**(1): 105-18.

- Herbert, T. P., A. R. Tee, et al. (2002). "The extracellular signal-regulated kinase pathway regulates the phosphorylation of 4E-BP1 at multiple sites." *J Biol Chem* **277**(13): 11591-6.
- Hidalgo, M. and E. K. Rowinsky (2000). "The rapamycin-sensitive signal transduction pathway as a target for cancer therapy." *Oncogene* **19**(56): 6680-6.
- Hornstein, E., A. Git, et al. (1999). "The expression of poly(A)-binding protein gene is translationally regulated in a growth-dependent fashion through a 5'-terminal oligopyrimidine tract motif." *J Biol Chem* **274**(3): 1708-14.
- Huang, S. and P. J. Houghton (2002). "Inhibitors of mammalian target of rapamycin as novel antitumor agents: from bench to clinic." *Curr Opin Investig Drugs* **3**(2): 295-304.
- Iiboshi, Y., P. J. Papst, et al. (1999). "Amino acid-dependent control of p70(s6k). Involvement of tRNA aminoacylation in the regulation." *J Biol Chem* **274**(2): 1092-9.
- Inoki, K., Y. Li, et al. (2002). "TSC2 is phosphorylated and inhibited by Akt and suppresses mTOR signalling." *Nat Cell Biol* **4**(9): 648-57.
- Isotani, S., K. Hara, et al. (1999). "Immunopurified mammalian target of rapamycin phosphorylates and activates p70 S6 kinase alpha in vitro." *J Biol Chem* **274**(48): 34493-8.
- Jacinto, E., B. Guo, et al. (2001). "TIP41 interacts with TAP42 and negatively regulates the TOR signaling pathway." *Mol Cell* **8**(5): 1017-26.
- Jefferies, H. B., S. Fumagalli, et al. (1997). "Rapamycin suppresses 5'TOP mRNA translation through inhibition of p70s6k." *Embo J* **16**(12): 3693-704.
- Jeno, P., L. M. Ballou, et al. (1988). "Identification and characterization of a mitogen-activated S6 kinase." *Proc Natl Acad Sci U S A* **85**(2): 406-10.
- Jiang, Y. and J. R. Broach (1999). "Tor proteins and protein phosphatase 2A reciprocally regulate Tap42 in controlling cell growth in yeast." *Embo J* **18**(10): 2782-92.
- Kawamata, S., H. Sakaida, et al. (1998). "The upregulation of p27Kip1 by rapamycin results in G1 arrest in exponentially growing T-cell lines." *Blood* **91**(2): 561-9.
- Keith, C. T. and S. L. Schreiber (1995). "PIK-related kinases: DNA repair, recombination, and cell cycle checkpoints." *Science* **270**(5233): 50-1.
- Khan, A., A. M. Pepio, et al. (2001). "Serotonin activates S6 kinase in a rapamycin-sensitive manner in Aplysia synaptosomes." *J Neurosci* **21**(2): 382-91.
- Kim, D. H., D. D. Sarbassov, et al. (2002). "mTOR interacts with raptor to form a nutrient-sensitive complex that signals to the cell growth machinery." *Cell* **110**(2): 163-75.
- Kim, D. H., D. Sarbassov dos, et al. (2003). "GbetaL, a positive regulator of the rapamycin-sensitive pathway required for the nutrient-sensitive interaction between raptor and mTOR." *Mol Cell* **11**(4): 895-904.

- Kimura, N., C. Tokunaga, et al. (2003). "A possible linkage between AMP-activated protein kinase (AMPK) and mammalian target of rapamycin (mTOR) signalling pathway." *Genes Cells* **8**(1): 65-79.
- Koh, H., K. Jee, et al. (1999). "Cloning and characterization of a nuclear S6 kinase, S6 kinase-related kinase (SRK); a novel nuclear target of Akt." *Oncogene* **18**(36): 5115-9.
- Kohn, A. D., A. Barthel, et al. (1998). "Construction and characterization of a conditionally active version of the serine/threonine kinase Akt." *J Biol Chem* **273**(19): 11937-43.
- Koltin, Y., L. Faucette, et al. (1991). "Rapamycin sensitivity in *Saccharomyces cerevisiae* is mediated by a peptidyl-prolyl cis-trans isomerase related to human FK506-binding protein." *Mol Cell Biol* **11**(3): 1718-23.
- Kozak, M. (1991). "Structural features in eukaryotic mRNAs that modulate the initiation of translation." *J Biol Chem* **266**(30): 19867-70.
- Kunz, J., R. Henriquez, et al. (1993). "Target of rapamycin in yeast, TOR2, is an essential phosphatidylinositol kinase homolog required for G1 progression." *Cell* **73**(3): 585-96.
- Kwiatkowski, D. J., H. Zhang, et al. (2002). "A mouse model of TSC1 reveals sex-dependent lethality from liver hemangiomas, and up-regulation of p70S6 kinase activity in Tsc1 null cells." *Hum Mol Genet* **11**(5): 525-34.
- Laser, M., V. S. Kasi, et al. (1998). "Differential activation of p70 and p85 S6 kinase isoforms during cardiac hypertrophy in the adult mammal." *J Biol Chem* **273**(38): 24610-9.
- Lazaris-Karatzas, A., K. S. Montine, et al. (1990). "Malignant transformation by a eukaryotic initiation factor subunit that binds to mRNA 5' cap." *Nature* **345**(6275): 544-7.
- Lee-Fruman, K. K., C. J. Kuo, et al. (1999). "Characterization of S6K2, a novel kinase homologous to S6K1." *Oncogene* **18**(36): 5108-14.
- Lin, T. A., X. Kong, et al. (1994). "PHAS-I as a link between mitogen-activated protein kinase and translation initiation." *Science* **266**(5185): 653-6.
- Lin, T. A., X. Kong, et al. (1995). "Control of PHAS-I by insulin in 3T3-L1 adipocytes. Synthesis, degradation, and phosphorylation by a rapamycin-sensitive and mitogen- activated protein kinase-independent pathway." *J Biol Chem* **270**(31): 18531-8.
- Lin, T. A. and J. C. Lawrence, Jr. (1997). "Control of PHAS-I phosphorylation in 3T3-L1 adipocytes: effects of inhibiting protein phosphatases and the p70S6K signalling pathway." *Diabetologia* **40 Suppl 2**: S18-24.
- Liu, Z., T. Sekito, et al. (2001). "RTG-dependent mitochondria to nucleus signaling is negatively regulated by the seven WD-repeat protein Lst8p." *Embo J* **20**(24): 7209-19.
- Loewith, R., E. Jacinto, et al. (2002). "Two TOR Complexes, Only One of which Is Rapamycin Sensitive, Have Distinct Roles in Cell Growth Control." *Mol Cell* **10**(3): 457-68.

- Mader, S., H. Lee, et al. (1995). "The translation initiation factor eIF-4E binds to a common motif shared by the translation factor eIF-4 gamma and the translational repressors 4E-binding proteins." *Mol Cell Biol* **15**(9): 4990-7.
- Manning, B. D., A. R. Tee, et al. (2002). "Identification of the tuberous sclerosis complex-2 tumor suppressor gene product tuberin as a target of the phosphoinositide 3-kinase/akt pathway." *Mol Cell* **10**(1): 151-62.
- Marcotrigiano, J., A. C. Gingras, et al. (1999). "Cap-dependent translation initiation in eukaryotes is regulated by a molecular mimic of eIF4G." *Mol Cell* **3**(6): 707-16.
- Martin, K. A., S. S. Schalm, et al. (2001). "Regulation of ribosomal S6 kinase 2 by effectors of the phosphoinositide 3-kinase pathway." *J Biol Chem* **276**(11): 7884-91.
- Martin, K. A., S. S. Schalm, et al. (2001). "Ribosomal S6 kinase 2 inhibition by a potent C-terminal repressor domain is relieved by mitogen-activated protein-extracellular signal-regulated kinase kinase-regulated phosphorylation." *J Biol Chem* **276**(11): 7892-8.
- Marx, S. O., J. Gaburjakova, et al. (2001). "Coupled gating between cardiac calcium release channels (ryanodine receptors)." *Circ Res* **88**(11): 1151-8.
- Miron, M. and N. Sonenberg (2001). "Regulation of translation via TOR signaling: insights from *Drosophila melanogaster*." *J Nutr* **131**(11): 2988S-93S.
- Montagne, J., T. Radimerski, et al. (2001). "Insulin signaling: lessons from the *Drosophila* tuberous sclerosis complex, a tumor suppressor." *Sci STKE* **2001**(105): PE36.
- Montagne, J., M. J. Stewart, et al. (1999). "Drosophila S6 kinase: a regulator of cell size." *Science* **285**(5436): 2126-9.
- Moser, B. A., P. B. Dennis, et al. (1997). "Dual requirement for a newly identified phosphorylation site in p70s6k." *Mol Cell Biol* **17**(9): 5648-55.
- Mothe-Satney, I., G. J. Brunn, et al. (2000). "Mammalian target of rapamycin-independent phosphorylation of PHAS-I in four (S/T)P sites detected by phospho-specific antibodies." *J Biol Chem* **275**(43): 33836-43.
- Mothe-Satney, I., D. Yang, et al. (2000). "Multiple mechanisms control phosphorylation of PHAS-I in five (S/T)P sites that govern translational repression." *Mol Cell Biol* **20**(10): 3558-67.
- Murata, K., J. Wu, et al. (1997). "B cell receptor-associated protein alpha4 displays rapamycin-sensitive binding directly to the catalytic subunit of protein phosphatase 2A." *Proc Natl Acad Sci U S A* **94**(20): 10624-9.
- Nanahoshi, M., T. Nishiuma, et al. (1998). "Regulation of protein phosphatase 2A catalytic activity by alpha4 protein and its yeast homolog Tap42." *Biochem Biophys Res Commun* **251**(2): 520-6.
- Nave, B. T., M. Ouwendijk, et al. (1999). "Mammalian target of rapamycin is a direct target for protein kinase B: identification of a convergence point for opposing effects of insulin and amino-acid deficiency on protein translation." *Biochem J* **344 Pt 2**: 427-31.

- Nojima, H., C. Tokunaga, et al. (2003). "The mammalian target of rapamycin (mTOR) partner, raptor, binds the mTOR substrates p70 S6 kinase and 4E-BP1 through their TOR signaling (TOS) motif." *J Biol Chem* **278**(18): 15461-4.
- Oldham, S. and E. Hafen (2003). "Insulin/IGF and target of rapamycin signaling: a TOR de force in growth control." *Trends Cell Biol* **13**(2): 79-85.
- Oldham, S., J. Montagne, et al. (2000). "Genetic and biochemical characterization of dTOR, the Drosophila homolog of the target of rapamycin." *Genes Dev* **14**(21): 2689-94.
- Pause, A., N. Methot, et al. (1994). "Dominant negative mutants of mammalian translation initiation factor eIF-4A define a critical role for eIF-4F in cap-dependent and cap-independent initiation of translation." *Embo J* **13**(5): 1205-15.
- Pearson, R. B., P. B. Dennis, et al. (1995). "The principal target of rapamycin-induced p70s6k inactivation is a novel phosphorylation site within a conserved hydrophobic domain." *Embo J* **14**(21): 5279-87.
- Pearson, R. B. and G. Thomas (1995). "Regulation of p70s6k/p85s6k and its role in the cell cycle." *Prog Cell Cycle Res* **1**: 21-32.
- Pende, M., S. C. Kozma, et al. (2000). "Hypoinsulinaemia, glucose intolerance and diminished beta-cell size in S6K1-deficient mice." *Nature* **408**(6815): 994-7.
- Peterson, R. T., P. A. Beal, et al. (2000). "FKBP12-rapamycin-associated protein (FRAP) autophosphorylates at serine 2481 under translationally repressive conditions." *J Biol Chem* **275**(10): 7416-23.
- Peterson, R. T., B. N. Desai, et al. (1999). "Protein phosphatase 2A interacts with the 70-kDa S6 kinase and is activated by inhibition of FKBP12-rapamycin-associated protein." *Proc Natl Acad Sci U S A* **96**(8): 4438-42.
- Pham, P. T., S. J. Heydrick, et al. (2000). "Assessment of cell-signaling pathways in the regulation of mammalian target of rapamycin (mTOR) by amino acids in rat adipocytes." *J Cell Biochem* **79**(3): 427-41.
- Potter, C. J., H. Huang, et al. (2001). "Drosophila Tsc1 functions with Tsc2 to antagonize insulin signaling in regulating cell growth, cell proliferation, and organ size." *Cell* **105**(3): 357-68.
- Poulin, F., A. C. Gingras, et al. (1998). "4E-BP3, a new member of the eukaryotic initiation factor 4E-binding protein family." *J Biol Chem* **273**(22): 14002-7.
- Pullen, N., P. B. Dennis, et al. (1998). "Phosphorylation and activation of p70s6k by PDK1." *Science* **279**(5351): 707-10.
- Pullen, N. and G. Thomas (1997). "The modular phosphorylation and activation of p70s6k." *FEBS Lett* **410**(1): 78-82.
- Rameh, L. E. and L. C. Cantley (1999). "The role of phosphoinositide 3-kinase lipid products in cell function." *J Biol Chem* **274**(13): 8347-50.
- Raught, B. and A. C. Gingras (1999). "eIF4E activity is regulated at multiple levels." *Int J Biochem Cell Biol* **31**(1): 43-57.

- Reinhard, C., A. Fernandez, et al. (1994). "Nuclear localization of p85s6k: functional requirement for entry into S phase." *Embo J* **13**(7): 1557-65.
- Roberg, K. J., S. Bickel, et al. (1997). "Control of amino acid permease sorting in the late secretory pathway of *Saccharomyces cerevisiae* by SEC13, LST4, LST7 and LST8." *Genetics* **147**(4): 1569-84.
- Rohde, J., J. Heitman, et al. (2001). "The TOR kinases link nutrient sensing to cell growth." *J Biol Chem* **276**(13): 9583-6.
- Romanelli, A., V. C. Dreisbach, et al. (2002). "Characterization of phosphatidylinositol 3-kinase-dependent phosphorylation of the hydrophobic motif site Thr(389) in p70 S6 kinase 1." *J Biol Chem* **277**(43): 40281-9.
- Romanelli, A., K. A. Martin, et al. (1999). "p70 S6 kinase is regulated by protein kinase Czeta and participates in a phosphoinositide 3-kinase-regulated signalling complex." *Mol Cell Biol* **19**(4): 2921-8.
- Rousseau, D., A. C. Gingras, et al. (1996). "The eIF4E-binding proteins 1 and 2 are negative regulators of cell growth." *Oncogene* **13**(11): 2415-20.
- Sabatini, D. M., R. K. Barrow, et al. (1999). "Interaction of RAFT1 with gephyrin required for rapamycin-sensitive signaling." *Science* **284**(5417): 1161-4.
- Sabatini, D. M., H. Erdjument-Bromage, et al. (1994). "RAFT1: a mammalian protein that binds to FKBP12 in a rapamycin- dependent fashion and is homologous to yeast TORs." *Cell* **78**(1): 35-43.
- Sabers, C. J., M. M. Martin, et al. (1995). "Isolation of a protein target of the FKBP12-rapamycin complex in mammalian cells." *JBC* **270**: 815-822.
- Sachs, A. B., P. Sarnow, et al. (1997). "Starting at the beginning, middle, and end: translation initiation in eukaryotes." *Cell* **89**(6): 831-8.
- Saucedo, L. J., X. Gao, et al. (2003). "Rheb promotes cell growth as a component of the insulin/TOR signalling network." *Nat Cell Biol* **5**(6): 566-71.
- Schalm, S. S. and J. Blenis (2002). "Identification of a conserved motif required for mTOR signaling." *Curr Biol* **12**(8): 632-9.
- Schmelzle, T. and M. N. Hall (2000). "TOR, a central controller of cell growth." *Cell* **103**(2): 253-62.
- Schmidt, E. V. (1999). "The role of c-myc in cellular growth control." *Oncogene* **18**(19): 2988-96.
- Schreiber, S. L. (1991). "Chemistry and biology of the immunophilins and their immunosuppressive ligands." *Science* **251**(4991): 283-7.
- Schwab, M. S., S. H. Kim, et al. (1999). "p70(S6K) controls selective mRNA translation during oocyte maturation and early embryogenesis in *Xenopus laevis*." *Mol Cell Biol* **19**(4): 2485-94.
- Scott, P. H., G. J. Brunn, et al. (1998). "Evidence of insulin-stimulated phosphorylation and activation of the mammalian target of rapamycin mediated by a protein kinase B signaling pathway." *Proc Natl Acad Sci U S A* **95**(13): 7772-7.
- Sehgal, S. N. (1995). "Rapamune (Sirolimus, rapamycin): an overview and mechanism of action." *Ther Drug Monit* **17**(6): 660-5.

- Sekulic, A., C. C. Hudson, et al. (2000). "A direct linkage between the phosphoinositide 3-kinase-AKT signaling pathway and the mammalian target of rapamycin in mitogen-stimulated and transformed cells." *Cancer Res* **60**(13): 3504-13.
- Serruys, P. W., E. Regar, et al. (2002). "Rapamycin eluting stent: the onset of a new era in interventional cardiology." *Heart* **87**(4): 305-7.
- Shah, O. J., J. C. Anthony, et al. (2000). "4E-BP1 and S6K1: translational integration sites for nutritional and hormonal information in muscle." *Am J Physiol Endocrinol Metab* **279**(4): E715-29.
- Shima, H., M. Pende, et al. (1998). "Disruption of the p70(s6k)/p85(s6k) gene reveals a small mouse phenotype and a new functional S6 kinase." *Embo J* **17**(22): 6649-59.
- Stan, R., M. M. McLaughlin, et al. (1994). "Interaction between FKBP12-rapamycin and TOR involves a conserved serine residue." *J Biol Chem* **269**(51): 32027-30.
- Stewart, M. J., C. O. Berry, et al. (1996). "The Drosophila p70s6k homolog exhibits conserved regulatory elements and rapamycin sensitivity." *Proc Natl Acad Sci U S A* **93**(20): 10791-6.
- Stocker, H., T. Radimerski, et al. (2003). "Rheb is an essential regulator of S6K in controlling cell growth in Drosophila." *Nat Cell Biol* **5**(6): 559-66.
- Takata, M., W. Ogawa, et al. (1999). "Requirement for Akt (protein kinase B) in insulin-induced activation of glycogen synthase and phosphorylation of 4E-BP1 (PHAS-1)." *J Biol Chem* **274**(29): 20611-8.
- Tang, H., E. Hornstein, et al. (2001). "Amino acid-induced translation of TOP mRNAs is fully dependent on phosphatidylinositol 3-kinase-mediated signaling, is partially inhibited by rapamycin, and is independent of S6K1 and rpS6 phosphorylation." *Mol Cell Biol* **21**(24): 8671-83.
- Tee, A. R., D. C. Fingar, et al. (2002). "Tuberous sclerosis complex-1 and -2 gene products function together to inhibit mammalian target of rapamycin (mTOR)-mediated downstream signaling." *Proc Natl Acad Sci U S A* **99**(21): 13571-6.
- Tee, A. R., Manning B. D., Roux P. R., Cantley L. C., and Blenis J. (2003). *Curr Biol*.
- Tee, A. R. and C. G. Proud (2002). "Caspase cleavage of initiation factor 4E-binding protein 1 yields a dominant inhibitor of cap-dependent translation and reveals a novel regulatory motif." *Mol Cell Biol* **22**(6): 1674-83.
- Tsukiyama-Kohara, K., S. M. Vidal, et al. (1996). "Tissue distribution, genomic structure, and chromosome mapping of mouse and human eukaryotic initiation factor 4E-binding proteins 1 and 2." *Genomics* **38**(3): 353-63.
- von Manteuffel, S. R., P. B. Dennis, et al. (1997). "The insulin-induced signalling pathway leading to S6 and initiation factor 4E binding protein 1 phosphorylation bifurcates at a rapamycin-sensitive point immediately upstream of p70s6k." *Mol Cell Biol* **17**(9): 5426-36.

- von Manteuffel, S. R., A. C. Gingras, et al. (1996). "4E-BP1 phosphorylation is mediated by the FRAP-p70s6k pathway and is independent of mitogen-activated protein kinase." *Proc Natl Acad Sci U S A* **93**(9): 4076-80.
- Wang, X., L. E. Campbell, et al. (1998). "Amino acid availability regulates p70 S6 kinase and multiple translation factors." *Biochem J* **334**(Pt 1): 261-7.
- Weng, Q. P., K. Andrabi, et al. (1995). "Phosphatidylinositol 3-kinase signals activation of p70 S6 kinase in situ through site-specific p70 phosphorylation." *Proc Natl Acad Sci U S A* **92**(12): 5744-8.
- Weng, Q. P., M. Kozlowski, et al. (1998). "Regulation of the p70 S6 kinase by phosphorylation in vivo. Analysis using site-specific anti-phosphopeptide antibodies." *J Biol Chem* **273**(26): 16621-9.
- Williams, M. R., J. S. Arthur, et al. (2000). "The role of 3-phosphoinositide-dependent protein kinase 1 in activating AGC kinases defined in embryonic stem cells." *Curr Biol* **10**(8): 439-48.
- Wilson, K. F., W. J. Wu, et al. (2000). "Cdc42 stimulates RNA splicing via the S6 kinase and a novel S6 kinase target, the nuclear cap-binding complex." *J Biol Chem* **275**(48): 37307-10.
- Wu, X., K. Senechal, et al. (1998). "The PTEN/MMAC1 tumor suppressor phosphatase functions as a negative regulator of the phosphoinositide 3-kinase/Akt pathway." *Proc Natl Acad Sci U S A* **95**(26): 15587-91.
- Yang, D. Q. and M. B. Kastan (2000). "Participation of ATM in insulin signalling through phosphorylation of eIF-4E-binding protein 1." *Nat Cell Biol* **2**(12): 893-8.
- Yokogami, K., S. Wakisaka, et al. (2000). "Serine phosphorylation and maximal activation of STAT3 during CNTF signaling is mediated by the rapamycin target mTOR." *Curr Biol* **10**(1): 47-50.
- Zhang, H., J. P. Stallok, et al. (2000). "Regulation of cellular growth by the Drosophila target of rapamycin dTOR." *Genes Dev* **14**(21): 2712-24.
- Zhang, Y., X. Gao, et al. (2003). "Rheb is a direct target of the tuberous sclerosis tumour suppressor proteins." *Nat Cell Biol* **5**(6): 578-81.