

Literaturverzeichnis

- Anfinsen, C. B. (1973). Principles that govern the folding of protein chains. *Science*, *181*, 223–230.
- Anfinsen, C. B., Haber, E., Sela, M., & White, F. H. (1961). The kinetics of formation of native ribonuclease during oxidation of the reduced polypeptide chain. *Proc. Natl. Acad. Sci. USA*, *47*, 1309–1314.
- Avbelj, F. (1992). Use of a potential of mean force to analyze free energy contributions in protein folding. *Biochemistry*, *31*(27), 6290–6297.
- Bastolla, U., Farwer, J., Knapp, E. W., & Vendruscolo, M. (2001). How to guarantee optimal stability for most representative structures in the protein data bank. *Proteins*, *44*(2), 79–96.
- Berman, H. M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T. N., Weissig, H., Shindyalov, I. N., & Bourne, P. E. (2000). The protein data bank. *Nucleic Acids Research*, *28*, 235–242.
- Bowie, J. U., Luthy, R., & Eisenberg, D. (1991). A method to identify protein sequences that fold into a known 3-dimensional structure. *Science*, *253*, 164–170.
- Brooks, B. R., Bruccoleri, R. E., Olafson, B. D., States, D. J., Swaminathan, S., & Karplus, M. (1983). Charmm: A program for macromolecular energy, minimization, and dynamics calculations. *J. Comp. Chem.*, *4*, 187–217.
- Bryant, S. H., & Lawrence, C. E. (1993). An empirical energy function for threading protein sequence through the folding motif. *Proteins*, *16*(1), 92–112.
- Brünger, A., Clore, G., Gronenborn, A., & Karplus, M. (1986). Three-dimensional structure of proteins determined by molecular dynamics with interproton distance restraints: application to crambin. *Proc. Natl. Acad. Sci. USA*, *83*, 3801–3805.

- Burkert, U., & Allinger, N. L. (1982). *Molecular Mechanics*. American Chemical Society, Washington, DC.
- Chan, H., & Dill, K. (1990). Origins of structure in globular proteins. *Proc. Natl. Acad. Sci. USA*, *87*, 6388–6392.
- Covell, D. G., & Jernigan, R. L. (1990). Conformations of folded proteins in restricted spaces. *Biochemistry*, *29*(13), 3287–3294.
- Crippen, G., & Havel, T. (1988). *Distance geometry and molecular conformation*. Wiley, New York.
- de Groot, B. L., & Grubmüller, H. (2001). Water permeation across biological membranes: Mechanism and dynamics of aquaporin-1 and glpf. *Science*, *294*, 2353–2357.
- Dima, R., Settanni, G., Micheletti, C., Banavar, J., & Maritan, A. (2000). Extraction of interaction potentials between amino acids from native protein structures. *J. Chem. Phys.*, *112*, 9151–9166.
- Eisenberg, D., & McLachlan, A. D. (1986). Solvation energy in protein folding and binding. *Nature*, *319*, 199–203.
- Fisher, D., Rice, D., Bowie, J. U., & Eisenberg, D. (1996). Assigning amino acid sequences to 3-dimensional protein folds. *FASEB J.*, *10*, 126–136.
- Foresman, J. B., & Frisch, A. (1996). *Exploring Chemistry With Electronic Structure Methods*. Gaussian Inc, Pittsburgh, PA.
- Godzik, A., Kolinski, A., & Skolnick, J. (1995). Are proteins ideal mixtures of amino acids? analysis of energy parameter sets. *Protein Sci.*, *4*(10), 2107–2117.
- Godzik, A., Skolnik, J., & Kolinski, A. (1993). Regularities in interaction patterns of globular proteins. *Protein Eng.*, *6*, 801–810.
- Goldstein, R., Luthey-Schulten, Z., & Wolynes, P. (1992a). Optimal protein-folding codes from spin-glass theory. *Proc. Natl. Acad. Sci. USA*, *89*, 4918–4922.
- Goldstein, R., Luthey-Schulten, Z., & Wolynes, P. (1992b). Protein tertiary structure recognition using optimized hamiltonians with local interactions. *Proc. Natl. Acad. Sci. USA*, *89*, 9029–9033.

- Hao, M., & Scheraga, H. (1996). How optimization of potential function affects protein folding. *Proc. Natl. Acad. Sci. USA*, *93*, 4984–4989.
- Hendlich, M., Lackner, P., Weitckus, S., Floekner, H., Froschauer, R., Gottsbacher, K., Casari, G., & Sippl, M. (1990). Identification of native protein folds amongst a large number of incorrect models. The calculation of low energy conformations from potentials of mean force. *J. Mol. Biol.*, *216*, 167–180.
- Hinds, D., & Levitt, M. (1994). Exploring conformational space with a simple lattice model for protein structure. *J. Mol. Biol.*, *243*, 668–682.
- Holm, L., & Sander, C. (1992). Evaluation of protein models by atomic solvation preference. *J. Mol. Biol.*, *225*(1), 93–105.
- Holm, L., & Sander, C. (1993). Protein structure comparison by alignment of distance matrices. *J. Mol. Biol.*, *233*, 123–138.
- Jelsch, C., Teeter, M. M., Lamzin, V., Pichon-Pesme, V., Blessing, R. H., & Lecomte, C. (2000). Accurate protein crystallography at ultra-high resolution: valence electron distribution in crambin. *Proc. Natl. Acad. Sci. USA*, *97*(7), 3171–3176.
- Jernigan, R. L., & Bahar, I. (1996). Structure-derived potentials and protein simulations. *Curr. Opin. Struct. Biol.*, *6*(2), 195–209.
- Jones, D. T., & Thornton, W. R. T. J. M. (1992). A new approach to protein fold recognition. *Nature*, *358*, 86–89.
- Kabsch, W. (1976). A solution for the best rotation to relate two sets of vectors. *Acta Crystallographica, A* *32*, 922–923.
- Karlin, S., Zuker, M., & Brocchieri, L. (1994). Measuring residue associations in protein structures. Possible implications for protein folding. *J. Mol. Biol.*, *239*(2), 227–248.
- Kocher, J. P., Rooman, M. J., & Wodak, S. J. (1994). Factors influencing the ability of knowledge-based potentials to identify native sequence-structure matches. *J. Mol. Biol.*, *235*(5), 1598–1613.
- Koretke, K., Luthey-Shulten, Z., & Wolynes, P. (1998). Self-consistently optimized energy functions for protein structure prediction by molecular dynamics. *Proc. Natl. Acad. Sci. USA*, *89*, 2932–2937.

- Levitt, M. (1976). A simplified representation of protein conformations for rapid simulation of protein folding. *J. Mol. Biol.*, 104(1), 59–107.
- Lifson, S., & Sander, C. (1979). Antiparallel and parallel beta-strands differ in amino acid residue preferences. *Nature*, 282, 109–111.
- M. Vendruscolo, E. D. (1998). Pairwise contact potentials are unsuitable for protein folding. *J. Chem. Phys.*, 109, 11101–11108.
- Madej, T., & Mossing, M. C. (1993). Hamiltonians for protein tertiary structure prediction based on three-dimensional environment principles. *J. Mol. Biol.*, 233(3), 480–487.
- Maiorov, V. N., & Crippen, G. M. (1992a). Contact potential that recognizes the correct folding of globular proteins. *J. Mol. Biol.*, 227(3), 876–888.
- Maiorov, V. N., & Crippen, G. M. (1992b). Contact potential that recognizes the correct folding of globular proteins. *J. Mol. Biol.*, 227(3), 876–88.
- Mirny, L., & Domany, E. (1996). Protein fold recognition and dynamics in the space of contact maps. *Proteins*, 26, 391–410.
- Mirny, L., & Shakhnovich, E. (1996). How to derive a protein folding potential? a new approach to an old problem. *J. Mol. Biol.*, 264, 1164–1179.
- Miyazawa, S., & Jernigan, R. (1985). Estimation of effective interresidue contact energies from protein crystal structures: quasi chemical approximation. *Macromolecules*, 18, 534–552.
- Ortiz, A. R., Kolinski, A., & Skolnick, J. (1998). Nativelike topology assembly of small proteins using predicted restraints in monte carlo simulations. *Proc. Natl. Acad. Sci. USA*, 95, 1020–1025.
- Park, B., & Levitt, M. (1996). Energy functions that discriminate x-ray and near native folds from well-constructed decoys. *J. Mol. Biol.*, 258(2), 367–392.
- Park, B., & Levitt, M. (1995). The complexity and accuracy of discrete state models of protein structure. *J. Mol. Biol.*, 249, 493–507.
- Salamov, A., & Solovyev, V. (1995). Prediction of protein secondary structure by combining nearest-neighbor algorithms and multiple sequence alignments. *J. Mol. Biol.*, 247, 11–15.

- Sippl, M. J. (1990). Calculation of conformational ensembles from potentials of mean force. An approach to the knowledge-based prediction of local structures in globular proteins. *J. Mol. Biol.*, 213(4), 859–83.
- Sippl, M. J. (1993a). Boltzmann's principle, knowledge-based mean fields and protein folding. An approach to the computational determination of protein structures. *J. Comput. Aided. Mol. Des.*, 7(4), 473–501.
- Sippl, M. J. (1993b). Recognition of errors in three-dimensional structures of proteins. *Proteins*, 17(4), 355–62.
- Sippl, M. J. (1995). Knowledge-based potentials for proteins. *Curr. Opin. Struct. Biol.*, 5(2), 229–235.
- Sippl, M. J., & Weitckus, S. (1992). Detection of native-like models for amino acid sequences of unknown three-dimensional structure in a data base of known protein conformations. *Proteins*, 13(3), 258–71.
- Skolnick, J., Jaroszweski, L., Kolinski, A., & Godzik, A. (1997). Derivation and testing of pair potentials for protein folding. when is the quasi-chemical approximation correct? *Prot. Sci.*, 6, 676–688.
- Thomas, P., & Dill, K. (1996). An iterative method for extracting energy-like quantities from protein structures. *Proc. Natl. Acad. Sci. USA*, 93, 11628–11633.
- V. Maiorov, G. C. (1992). Contact potential that recognizes the correct folding of globular proteins *J. Mol. Biol.*, 227, 876–888.
- Vagedes, P., Rabenstein, B., Aqvist, J., Marelius, J., & Knapp, E. W. (2000). The deacylation step of acetylcholinesterase: Computer simulation studies. *J. Am. Chem. Soc.*, 122, 12254–12262.
- van Gunsteren, W. F., & Berendsen, H. J. C. (1977). Algorithms for macromolecular dynamics and constraint dynamics. *Molecular Physics*, 34, 1311–1327.
- van Mourik, J., Clementi, C., Maritan, A., Seno, F., & Banavar, J. (1999). Determination of interaction potentials of amino acids from native protein structures: Tests on simple lattice models. *J. Chem. Phys.*, 110, 10123–10133.
- Vendruscolo, M., Kussel, E., & Domany, E. (1997a). Recovery of protein structure from contact maps. *Fold. Des.*, 2, 295–306.

- Vendruscolo, M., Kussel, E., & Domany, E. (1997b). Recovery of protein structure from contact maps. *Fold. Des.*, 2, 295–306.
- Vendruscolo, M., Najmanovich, R., & Domany, E. (2000). Can a pairwise contact potential stabilize native protein folds against decoys obtained by threading? *Proteins*, 38, 134–148.
- V.N. Maiorov, G. C. (1994). Learning about protein folding via potential functions. *Proteins*, 20, 167–173.
- Voet, D., & Voet, J. G. (1995). *Biochemistry* (2 edition). Wiley, New York.
- Wallqvist, A., & Ullner, M. (1994). A simplified amino acid potential for use in structure predictions of proteins. *Proteins*, 18(3), 267–280.
- Warshel, A., & Levitt, M. (1976). Theoretical studies of enzymic reactions: dielectric, electrostatic and steric stabilization of the carbonium ion in the reaction of lysozyme. *J. Mol. Biol.*, 103, 227–249.
- Zhang, K. Y., & Eisenberg, D. (1994). The three-dimensional profile method using residue preference as a continuous function of residue environment. *Protein Sci.*, 3(4), 687–695.