

## Referenzen:

1. Abbruzzese, M., Favale, E., Leandri, M. & Ratto, S. New subcortical components of the cerebral somatosensory evoked potential in man. *Acta Neurol Scand* **58**, 325-32 (1978).
2. Adrian, E. D. & Matthews, B. H. C. The interpretation of potential waves in the cortex. *J Physiol* **81**, 440-471 (1934).
3. Allison, T., McCarthy, G., Wood, C. C. & Jones, S. J. Potentials evoked in human and monkey cerebral cortex by stimulation of the median nerve. A review of scalp and intracranial recordings. *Brain* **114 ( Pt 6)**, 2465-503 (1991).
4. Allison, W. Recovery functions of somatosensory evoked responses in man. *Electroencephalogr Clin Neurophysiol* **14**, 331-343 (1962).
5. Baker, S. N., Curio, G. & Lemon, R. N. EEG oscillations at 600 Hz are macroscopic markers for cortical spike bursts. *J Physiol* **550**, 529-34 (2003).
6. Berger, H. Über das Elektrencephalogramm des Menschen. *Archiv für Psychiatrie und Nervenkrankheiten* **87**, 527-570 (1929).
7. Borck, C. in *Instrument-Experiment. Historische Studien* (ed. Meinel, C.) 118-127 (Diepholz, Berlin, 2000).
8. Buchner, H., Ludwig, I., Veldkamp, R., Willmes, K. & Ferbert, A. [Topography of early cortical median nerve somatosensory evoked potentials: results for routine use of the method]. *EEG EMG Z Elektroenzephalogr Elektromyogr Verwandte Geb* **23**, 203-9 (1992).

9. Buchner, H., Ludwig, I., Waberski, T., Wilmes, K. & Ferbert, A. Hemispheric asymmetries of early cortical somatosensory evoked potentials revealed by topographic analysis. *Electromyogr Clin Neurophysiol* **35**, 207-15 (1995).
10. Calvin, W. H. & Loeser, J. D. Doublet and burst firing patterns within the dorsal column nuclei of cat and man. *Exp Neurol* **48**, 406-26 (1975).
11. Caton, R. The electrical currents of the brain. *British Medical Journal* **28**, 278 (1875).
12. Cracco, R. Q. & Cracco, J. B. Somatosensory evoked potential in man: far field potentials. *Electroencephalogr Clin Neurophysiol* **41**, 460-6 (1976).
13. Curio, G., Mackert, B. M., Abraham-Fuchs, K. & W, H. in *Oscillatory event related brain dynamics* (ed. Pantev, C.) 205 - 217 (Plenum Press, New York, 1994a).
14. Curio, G., Mackert, B. M., Burghoff, M., Koetitz, R., Abraham-Fuchs, K. & Harer, W. Localization of evoked neuromagnetic 600 Hz activity in the cerebral somatosensory system. *Electroencephalogr Clin Neurophysiol* **91**, 483-7 (1994b).
15. Curio, G., Mackert, B. M., Burghoff, M., Neumann, J., Nolte, G., Scherg, M. & Marx, P. Somatotopic source arrangement of 600 Hz oscillatory magnetic fields at the human primary somatosensory hand cortex. *Neurosci Lett* **234**, 131-4 (1997).
16. Curio, G. High frequency (600 Hz) bursts of spike-like activities generated in the human cerebral somatosensory system. *Electroencephalogr Clin Neurophysiol Suppl* **49**, 56-61 (1999).
17. Curio, G. Linking 600-Hz "spikelike" EEG/MEG wavelets ("sigma-bursts") to cellular substrates: concepts and caveats. *J Clin Neurophysiol*. **17**, 377-396 (2000).
18. Dawson, G. A summation technique for the detection of small evoked potentials. *Electroencephalogr Clin Neurophysiol* **6**, 65-84 (1954).

19. Dingledine, R. & McBain, C. J. in *Basic Neurochemistry* (ed. Siegel, G. J.) 315- 334 (Lippincott, Williams & Wilkins, Philadelphia, Baltimore, New York, London, Buenos Aires, Hong Kong, Sydney, Tokyo, 1999).
20. Dudel, J. in *Neuro- und Sinnesphysiologie* (ed. Schmidt, R. F.) 31- 58 (Springer Verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo, 1993).
21. Eisen, A., Roberts, K., Low, M., Hoirch, M. & Lawrence, P. Questions regarding the sequential neural generator theory of the somatosensory evoked potential raised by digital filtering. *Electroencephalogr Clin Neurophysiol* **59**, 388-95 (1984).
22. Elbert, T., Pantev, C., Wienbruch, C., Rockstroh, B. & Taub, E. Increased cortical representation of the fingers of the left hand in string players. *Science* **270**, 305-7 (1995).
23. Emerson, R. G., Sgro, J. A., Pedley, T. A. & Hauser, W. A. State-dependent changes in the N20 component of the median nerve somatosensory evoked potential. *Neurology* **38**, 64-8 (1988).
24. Emori, T., Yamada, T., Seki, Y., Yasuhara, A., Ando, K., Honda, Y., Leis, A. A. & Vachatimanont, P. Recovery functions of fast frequency potentials in the initial negative wave of median SEP. *Electroencephalogr Clin Neurophysiol* **78**, 116-23 (1991).
25. Engel, A. K., Roelfsema, P. R., Fries, P., Brecht, M. & Singer, W. Role of the temporal domain for response selection and perceptual binding. *Cereb Cortex* **7**, 571-82 (1997).
26. Engel, A. K. & Singer, W. Temporal binding and the neural correlates of sensory awareness. *Trends Cogn Sci* **5**, 16-25 (2001).

27. Flor, H., Elbert, T., Knecht, S., Wienbruch, C., Pantev, C., Birbaumer, N., Larbig, W. & Taub, E. Phantom-limb pain as a perceptual correlate of cortical reorganization following arm amputation. *Nature* **375**, 482-4 (1995).
28. Galarreta, M. & Hestrin, S. A network of fast-spiking cells in the neocortex connected by electrical synapses. *Nature* **402**, 72-5 (1999).
29. Glenn, L. L. & Steriade, M. Discharge rate and excitability of cortically projecting intralaminar thalamic neurons during waking and sleep states. *J Neurosci* **2**, 1387-404 (1982).
30. Gobbele, R., Buchner, H. & Curio, G. High-frequency (600 Hz) SEP activities originating in the subcortical and cortical human somatosensory system. *Electroencephalogr Clin Neurophysiol* **108**, 182-9 (1998).
31. Gobbele, R., Buchner, H., Scherg, M. & Curio, G. Stability of high-frequency (600 Hz) components in human somatosensory evoked potentials under variation of stimulus rate--evidence for a thalamic origin. *Clin Neurophysiol* **110**, 1659-63 (1999).
32. Gobbele, R., Waberski, T. D., Kuelkens, S., Sturm, W., Curio, G. & Buchner, H. Thalamic and cortical high-frequency (600 Hz) somatosensory-evoked potential (SEP) components are modulated by slight arousal changes in awake subjects. *Exp Brain Res* **133**, 506-513 (2000).
33. Gobbele, R., Waberski, T. D., Thyerlei, D., Thissen, M., Darvas, F., Klostermann, F., Curio, G. & Buchner, H. Functional dissociation of a subcortical and cortical component of high-frequency oscillations in human somatosensory evoked potentials by motor interference. *Neurosci Lett* **350**, 97-100 (2003).
34. Gray, C. M. & McCormick, D. A. Chattering cells: superficial pyramidal neurons contributing to the generation of synchronous oscillations in the visual cortex. *Science* **274**, 109-13 (1996).

35. Green, J. B., Nelson, A. V. & Michael, D. Digital zero-phase-shift filtering of short-latency somatosensory evoked potentials. *Electroencephalogr Clin Neurophysiol* **63**, 384-8 (1986).
36. Greenwood, P. M. & Goff, W. R. Modification of median nerve somatic evoked potentials by prior median nerve, peroneal nerve, and auditory stimulation. *Electroencephalogr Clin Neurophysiol* **68**, 295-302 (1987).
37. Guido, W. & Weyand, T. Burst responses in thalamic relay cells of the awake behaving cat. *J Neurophysiol* **74**, 1782-6 (1995).
38. Halboni, P., Kaminski, R., Gobbele, R., Zuchner, S., Waberski, T. D., Herrmann, C. S., Topper, R. & Buchner, H. Sleep stage dependant changes of the high-frequency part of the somatosensory evoked potentials at the thalamus and cortex. *Clin Neurophysiol* **111**, 2277-84 (2000).
39. Hashimoto, I., Mashiko, T. & Imada, T. Somatic evoked high-frequency magnetic oscillations reflect activity of inhibitory interneurons in the human somatosensory cortex. *Electroencephalogr Clin Neurophysiol* **100**, 189-203 (1996).
40. Hashimoto, I., Kimura, T., Fukushima, T., Iguchi, Y., Saito, Y., Terasaki, O. & Sakuma, K. Reciprocal modulation of somatosensory evoked N20m primary response and high-frequency oscillations by interference stimulation. *Clin Neurophysiol* **110**, 1445-51 (1999).
41. Hashimoto, I. High-frequency oscillations of somatosensory evoked potentials and fields. *J Clin Neurophysiol* **17**, 309-20 (2000).
42. Haueisen, J., Heuer, T., Nowak, H., Liepert, J., Weiller, C., Okada, Y. & Curio, G. The influence of lorazepam on somatosensory-evoked fast frequency (600 Hz) activity in MEG. *Brain Res* **874**, 10-4 (2000).

43. Hendry, S. H., Hsiao, S. S. & Brown, M. C. in *Fundamental neuroscience* (ed. Zigmond, M. J.) 577- 590 (Academic Press, San Diego, London, 2003).
44. Huerta, P. T. & Lisman, J. E. Bidirectional synaptic plasticity induced by a single burst during cholinergic theta oscillation in CA1 in vitro. *Neuron* **15**, 1053-63 (1995).
45. Ikeda, H., Leyba, L., Bartolo, A., Wang, Y. & Okada, Y. C. Synchronized spikes of thalamocortical axonal terminals and cortical neurons are detectable outside the pig brain with MEG. *J Neurophysiol* **87**, 626-30 (2002).
46. Inoue, K., Hashimoto, I. & Nakamura, S. High-frequency oscillations in human posterior tibial somatosensory evoked potentials are enhanced in patients with Parkinson's disease and multiple system atrophy. *Neurosci Lett* **297**, 89-92 (2001).
47. Jones, M. S. & Barth, D. S. Sensory-evoked high-frequency (gamma-band) oscillating potentials in somatosensory cortex of the unanesthetized rat. *Brain Res* **768**, 167-76 (1997).
48. Jones, M. S. & Barth, D. S. Spatiotemporal organization of fast (>200 Hz) electrical oscillations in rat Vibrissa/Barrel cortex. *J Neurophysiol* **82**, 1599-609 (1999).
49. Jones, M. S., MacDonald, K. D., Choi, B., Dudek, F. E. & Barth, D. S. Intracellular correlates of fast (>200 Hz) electrical oscillations in rat somatosensory cortex. *J Neurophysiol* **84**, 1505-18 (2000).
50. Jones, M. S. & Barth, D. S. Effects of bicuculline methiodide on fast (>200 Hz) electrical oscillations in rat somatosensory cortex. *J Neurophysiol* **88**, 1016-25 (2002).
51. Julien, R. M. in *Drogen und Psychopharmaka* (ed. Julien, R. M.) 77- 100 (Spektrum Akademischer Verlag, Heidelberg, Berlin, Oxford, 1997).
52. Jung, P., Baumgartner, U., Bauermann, T., Magerl, W., Gawehn, J., Stoeter, P. & Treede, R. D. Asymmetry in the human primary somatosensory cortex and handedness. *Neuroimage* **19**, 913-23 (2003).

53. Katayama, Y. & Tsubokawa, T. Somatosensory evoked potentials from the thalamic sensory relay nucleus (VPL) in humans: correlations with short latency somatosensory evoked potentials recorded at the scalp. *Electroencephalogr Clin Neurophysiol* **68**, 187-201 (1987).
54. Klostermann, F., Nolte, G., Losch, F. & Curio, G. Differential recruitment of high frequency wavelets (600 Hz) and primary cortical response (N20) in human median nerve somatosensory evoked potentials. *Neurosci Lett* **256**, 101-4 (1998).
55. Klostermann, F., Funk, T., Vesper, J. & Curio, G. Spatiotemporal characteristics of human intrathalamic high-frequency (>400Hz) SEP components. *Neuroreport* **10**, 3627-31 (1999a).
56. Klostermann, F., Nolte, G. & Curio, G. Multiple generators of 600 Hz wavelets in human SEP unmasked by varying stimulus rates. *Neuroreport* **10**, 1625-9 (1999b).
57. Klostermann, F., Funk, T., Vesper, J., Siedenberg, R. & Curio, G. Propofol narcosis dissociates human intrathalamic and cortical high-frequency (> 400 hz) SEP components. *Neuroreport* **11**, 2607-10 (2000a).
58. Klostermann, F., Funk, T., Vesper, J., Siedenberg, R. & Curio, G. Double-pulse stimulation dissociates intrathalamic and cortical high-frequency (>400Hz) SEP components in man. *Neuroreport* **11**, 1295-9 (2000b).
59. Klostermann, F., Gobbele, R., Buchner, H., Siedenberg, R. & Curio, G. Differential gating of slow postsynaptic and high-frequency spike-like components in human somatosensory evoked potentials under isometric motor interference. *Brain Res* **922**, 95-103 (2001a).
60. Klostermann, F., Nolte, G. & Curio, G. Independent short-term variability of spike-like (600 Hz) and postsynaptic (N20) cerebral SEP components. *Neuroreport* **12**, 349-352 (2001b).

61. Klostermann, F., Gobbele, R., Buchner, H. & Curio, G. Intrathalamic non-propagating generators of high-frequency (1000 Hz) somatosensory evoked potential (SEP) bursts recorded subcortically in man. *Clin Neurophysiol* **113**, 1001-5 (2002).
62. Kojima, Y., Uozumi, T., Akamatsu, N., Matsunaga, K., Urasaki, E. & Tsuji, S. Somatosensory evoked high frequency oscillations recorded from subdural electrodes. *Clin Neurophysiol* **112**, 2261-4 (2001).
63. Lisman, J. E. Bursts as a unit of neural information: making unreliable synapses reliable. *Trends Neurosci* **20**, 38-43 (1997).
64. Loewi, O. Über humorale Übertragbarkeit der Herzmuskelwirkung. *Pflügers Archiv* **189**, 239- 242 (1921).
65. Losch, F. *Entwicklung nicht-invasiver, topographisch hochauflösender MEG/EEG-Messverfahren im thalamokortikalen somatosensiblen System des Menschen* (Dissertation Freie Universität Berlin, 2001).
66. Maccabee, P. J., Pinkhasov, E. I. & Cracco, R. Q. Short latency somatosensory evoked potentials to median nerve stimulation: effect of low frequency filter. *Electroencephalogr Clin Neurophysiol* **55**, 34-44 (1983).
67. Mackert, B. M., Weisenbach, S., Nolte, G. & Curio, G. Rapid recovery (20 ms) of human 600 Hz electroencephalographic wavelets after double stimulation of sensory nerves. *Neurosci Lett* **286**, 83-6 (2000).
68. Maegaki, Y., Najm, I., Terada, K., Morris, H. H., Bingaman, W. E., Kohaya, N., Takenobu, A., Kadonaga, Y. & Luders, H. O. Somatosensory evoked high-frequency oscillations recorded directly from the human cerebral cortex. *Clin Neurophysiol* **111**, 1916-26 (2000).
69. Mainen, Z. F. & Sejnowski, T. J. Reliability of spike timing in neocortical neurons. *Science* **268**, 1503-6 (1995).

70. McCormick, D. A., Connors, B. W., Lighthall, J. W. & Prince, D. A. Comparative electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. *J Neurophysiol* **54**, 782-806 (1985).
71. McCormick, D. A. & Feeser, H. R. Functional implications of burst firing and single spike activity in lateral geniculate relay neurons. *Neuroscience* **39**, 103-13 (1990).
72. Misulis, K. E. in *Clinical Neurophysiology* (ed. Head, T. C.) 193- 230 (Butterworth and Heinemann, Burlington, MA, 2003).
73. Mochizuki, H., Machii, K., Terao, Y., Furubayashi, T., Hanajima, R., Enomoto, H., Uesugi, H., Shiio, Y., Kamakura, K., Kanazawa, I. & Ugawa, Y. Recovery function of and effects of hyperventilation on somatosensory evoked high-frequency oscillation in Parkinson's disease and myoclonus epilepsy. *Neurosci Res* **46**, 485-92 (2003).
74. Mountcastle, V. B. The columnar organization of the neocortex. *Brain* **120**, 701- 722 (1997).
75. Nakano, S. & Hashimoto, I. The later part of high-frequency oscillations in human somatosensory evoked potentials is enhanced in aged subjects. *Neurosci Lett* **276**, 83-6 (1999).
76. Nakano, S. & Hashimoto, I. High-frequency oscillations in human somatosensory evoked potentials are enhanced in school children. *Neurosci Lett* **291**, 113-6 (2000).
77. Olsen, R. W. & DeLorey, T. M. in *Basic Neurochemistry* (ed. Siegel, G. J.) 335- 346 (Lippincott, Williams & Wilkins, Philadelphia, Baltimore, New York, London, Buenos Aires, Hong Kong, Sydney, Tokyo, 1999).
78. Ozaki, I., Suzuki, C., Yaegashi, Y., Baba, M., Matsunaga, M. & Hashimoto, I. High frequency oscillations in early cortical somatosensory evoked potentials. *Electroencephalogr Clin Neurophysiol* **108**, 536-42 (1998).

79. Ozaki, I., Yaegashi, Y., Kimura, T., Baba, M., Matsunaga, M. & Hashimoto, I. Dipole orientation differs between high frequency oscillations and N20m current sources in human somatosensory evoked magnetic fields to median nerve stimulation. *Neurosci Lett* **310**, 41-4 (2001).
80. Ozaki, I. & Hashimoto, I. in *Recent Advances in Human Brain Mapping* (ed. Yamazaki, K.) 175-180 (Elsevier, 2002).
81. Perry, E. & Young, A. in *Neurochemistry of consciousness* (ed. Young, A.) 3- 24 (John Benjamins Publishing Company, Amsterdam, Philadelphia, 2002).
82. Prestor, B., Gnidovec, B. & Golob, P. Long sensory tracts (cuneate fascicle) in cervical somatosensory evoked potential after median nerve stimulation. *Electroencephalogr Clin Neurophysiol* **104**, 470-9 (1997).
83. Rao, S. M., Binder, J. R., Bandettini, P. A., Hammeke, T. A., Yetkin, F. Z., Jesmanowicz, A., Lisk, L. M., Morris, G. L., Mueller, W. M. & Estkowski, L. D. Functional magnetic resonance imaging of complex human movements. *Neurology* **43**, 2311-2318 (1993).
84. Rechtschaffen, A. & Kales, A. *A manual standardized terminology, techniques and scoring system for sleep stages of human subjects* (Public health service;, Washington, 1968).
85. Restuccia, D., Valeriani, M., Grassi, E., Mazza, S. & Tonali, P. Dissociated changes of somatosensory evoked low-frequency scalp responses and 600 Hz bursts after single-dose administration of lorazepam. *Brain Res* **946**, 1-11 (2002).
86. Restuccia, D., Della Marca, G., Valeriani, M., Rubino, M., Paciello, N., Vollono, C., Capuano, A. & Tonali, P. Influence of cholinergic circuitries in generation of high-frequency somatosensory evoked potentials. *Clin Neurophysiol* **114**, 1538-48 (2003).

87. Rieke, Warland, Ruyter, d., Steveninck & Bialek. *Spikes, Exploring the neural code* (Cambridge, 1997).
88. Rossini, P. M., Narici, L., Martino, G., Pasquarelli, A., Peresson, M., Pizzella, V., Tecchio, F. & Romani, G. L. Analysis of interhemispheric asymmetries of somatosensory evoked magnetic fields to right and left median nerve stimulation. *Electroencephalogr Clin Neurophysiol* **91**, 476-82 (1994).
89. Sakuma, K. & Hashimoto, I. High-frequency magnetic oscillations evoked by posterior tibial nerve stimulation. *Neuroreport* **10**, 227-30 (1999).
90. Schluter, N. D., Rushworth, M. F., Passingham, R. E. & Mills, K. R. Temporary interference in human lateral premotor cortex suggests dominance for the selection of movements. A study using transcranial magnetic stimulation. *Brain* **121**, 785-799 (1998).
91. Shimazu, H., Kaji, R., Tsujimoto, T., Kohara, N., Ikeda, A., Kimura, J. & Shibasaki, H. High-frequency SEP components generated in the somatosensory cortex of the monkey. *Neuroreport* **11**, 2821-6 (2000).
92. Singer, W. & Gray, C. M. Visual feature integration and the temporal correlation hypothesis. *Annu Rev Neurosci* **18**, 555-86 (1995).
93. Singer, W. in *Der Beobachter im Gehirn* (ed. Singer, W.) 87-111 (Suhrkamp, Frankfurt, 2002).
94. Sonoo, M., Genba-Shimizu, K., Mannen, T. & Shimizu, T. Detailed analysis of the latencies of median nerve somatosensory evoked potential components, 2: Analysis of subcomponents of the P13/14 and N20 potentials. *Electroencephalogr Clin Neurophysiol* **104**, 296-311 (1997).
95. Steriade, M., Curro Dossi, R. & Contreras, D. Electrophysiological properties of intralaminar thalamocortical cells discharging rhythmic (approximately 40 HZ) spike-

- bursts at approximately 1000 HZ during waking and rapid eye movement sleep. *Neuroscience* **56**, 1-9 (1993).
96. Steriade, M. Corticothalamic resonance, states of vigilance and mentation. *Neuroscience* **101**, 243-76 (2000).
97. Stöhr, M., Dichgans, J. & Buettner, U. *Evozierte Potentiale* (Springer, 1996).
98. Suzuki, I. & Mayanagi, Y. Intracranial recording of short latency somatosensory evoked potentials in man: identification of origin of each component. *Electroencephalogr Clin Neurophysiol* **59**, 286-96 (1984).
99. Swadlow, H. A. Influence of VPM afferents on putative inhibitory interneurons in S1 of the awake rabbit: evidence from cross-correlation, microstimulation, and latencies to peripheral sensory stimulation. *J Neurophysiol* **73**, 1584-99 (1995).
100. Swadlow, H. A., Beloozerova, I. N. & Sirota, M. G. Sharp, local synchrony among putative feed-forward inhibitory interneurons of rabbit somatosensory cortex. *J Neurophysiol* **79**, 567-82 (1998).
101. Tamas, G., Buhl, E. H., Lorincz, A. & Somogyi, P. Proximally targeted GABAergic synapses and gap junctions synchronize cortical interneurons. *Nat Neurosci* **3**, 366-71 (2000).
102. Tanosaki, M., Hashimoto, I., Iguchi, Y., Kimura, T., Takino, R., Kurobe, Y., Haruta, Y. & Hoshi, Y. Specific somatosensory processing in somatosensory area 3b for human thumb: a neuromagnetic study. *Clin Neurophysiol* **112**, 1516-22 (2001).
103. Tanosaki, M., Kimura, T., Takino, R., Iguchi, Y., Suzuki, A., Kurobe, Y., Haruta, Y., Hoshi, Y. & Hashimoto, I. Movement interference attenuates somatosensory high-frequency oscillations: contribution of local axon collaterals of 3b pyramidal neurons. *Clin Neurophysiol* **113**, 993-1000 (2002).

104. Tecchio, F., Rossini, P. M., Pizzella, V., Cassetta, E., Pasqualetti, P. & Romani, G. L. A neuromagnetic normative data set for hemispheric sensory hand cortical representations and their interhemispheric differences. *Brain Res Brain Res Protoc* **2**, 306-14 (1998).
105. Urasaki, E., Genmoto, T., Akamatsu, N., Wada, S. & Yokota, A. The effects of stimulus rates on high frequency oscillations of median nerve somatosensory-evoked potentials - direct recording study from the human cerebral cortex. *Clin Neurophysiol* **113**, 1794-7 (2002).
106. Wikstrom, H., Roine, R. O., Salonen, O., Aronen, H. J., Virtanen, J., Ilmoniemi, R. J. & Huttunen, J. Somatosensory evoked magnetic fields to median nerve stimulation: interhemispheric differences in a normal population. *Electroencephalogr Clin Neurophysiol* **104**, 480-7 (1997).
107. Yamada, T., Kameyama, S., Fuchigami, Y., Nakazumi, Y., Dickins, Q. S. & Kimura, J. Changes of short latency somatosensory evoked potential in sleep. *Electroencephalogr Clin Neurophysiol* **70**, 126-36 (1988).
108. Zschocke, S. in *Klinische Elektroenzephalographie* (ed. Zschocke, S.) 3- 15 (Springer Verlag, Berlin, Heidelberg, New York, Barcelona, Hongkong, London, Mailand, Paris, Tokio, 2002a).
109. Zschocke, S. *Klinische Elektroenzephalographie, 2. Auflage* (Springer, Berlin, Heidelberg, New Yorck, 2002b).