

7. Literaturverzeichnis

Arnqvist HJ, Bornfeldt KE, Chen Y, Lindstrom T. The insulin-like growth factor system in vascular smooth muscle: interaction with insulin and growth factors. *Metabolism* 1995; 44(10 Suppl 4):58-66.

Berman DM, Gilman AG. Mammalian RGS proteins: barbarians at the gate. *J Biol Chem* 1998; 273(3):1269-1272.

Bianchi BR, Lynch KJ, Touma E et al. Pharmacological characterization of recombinant human and rat P2X receptor subtypes. *Eur J Pharmacol* 1999; 376(1-2):127-138.

Buhl AM, Johnson NL, Dhanasekaran N, Johnson GL. G alpha 12 and G alpha 13 stimulate Rho-dependent stress fiber formation and focal adhesion assembly. *J Biol Chem* 1995; 270(42):24631-24634.

Burnstock G. Purinergic signaling and vascular cell proliferation and death. *Arterioscler Thromb Vasc Biol* 2002; 22(3):364-373.

Campbell GR, Campbell JH. Smooth muscle phenotypic changes in arterial wall homeostasis: implications for the pathogenesis of atherosclerosis. *Exp Mol Pathol* 1985; 42(2):139-162.

Camps M, Carozzi A, Schnabel P, Scheer A, Parker PJ, Gierschik P. Isozyme-selective stimulation of phospholipase C-beta 2 by G protein beta gamma-subunits. *Nature* 1992; 360(6405):684-686.

Carafoli E, Brini M. Calcium pumps: structural basis for and mechanism of calcium transmembrane transport. *Curr Opin Chem Biol* 2000; 4(2):152-161.

Commu D, Robaye B, Boeynaems JM. Pharmacological characterization of the human P2Y11 receptor. *Br J Pharmacol* 1999; 128(6):1199-1206.

Daub H, Wallasch C, Lankenau A, Herrlich A, Ullrich A. Signal characteristics of G protein-transactivated EGF receptor. *EMBO J* 1997; 16(23):7032-7044.

Di VF. Dr. Jekyll/Mr. Hyde: the dual role of extracellular ATP. *J Auton Nerv Syst* 2000; 81(1-3):59-63.

Dohlman HG, Thorner J. RGS proteins and signaling by heterotrimeric G proteins. *J Biol Chem* 1997; 272(7):3871-3874.

Edgecombe M, Craddock HS, Smith DC, McLennan AG, Fisher MJ. Diadenosine polyphosphate-stimulated gluconeogenesis in isolated rat proximal tubules. *Biochem J* 1997; 323 (Pt 2):451-456.

Erlinge D, Edvinsson L, Brunkwall J, Yee F, Wahlestedt C. Human neuropeptide Y Y1 receptor antisense oligodeoxynucleotide specifically inhibits neuropeptide Y-evoked vasoconstriction. *Eur J Pharmacol* 1993; 240(1):77-80.

Erlinge D, Hou M, Webb TE, Barnard EA, Moller S. Phenotype changes of the vascular smooth muscle cell regulate P2 receptor expression as measured by quantitative RT-PCR. *Biochem Biophys Res Commun* 1998; 248(3):864-870.

Exton JH. Regulation of phosphoinositide phospholipases by hormones, neurotransmitters, and other agonists linked to G proteins. *Annu Rev Pharmacol Toxicol* 1996; 36:481-509.

Filippov AK, Webb TE, Barnard EA, Brown DA. Dual coupling of heterologously-expressed rat P2Y6 nucleotide receptors to N-type Ca²⁺ and M-type K⁺ currents in rat sympathetic neurones. *Br J Pharmacol*. 1999 Feb; 126(4):1009-17.

Flodgaard H, Klenow H. Abundant amounts of diadenosine 5',5"-P₁,P₄-tetraphosphate are present and releasable, but metabolically inactive, in human platelets. *Biochem J* 1982; 208(3):737-742.

Gohla A, Harhammer R, Schultz G. The G-protein G13 but not G12 mediates signaling from lysophosphatidic acid receptor via epidermal growth factor receptor to Rho. *J Biol Chem* 1998; 273(8):4653-4659.

Groschel-Stewart U, Bardini M, Robson T, Burnstock G. Localisation of P2X5 and P2X7 receptors by immunohistochemistry in rat stratified squamous epithelia. *Cell Tissue Res* 1999; 296(1):599-605.

Grynkiewicz G, Poenie M, Tsien RY. A new generation of Ca²⁺ indicators with greatly improved fluorescence properties. *J Biol Chem* 1985; 260(6):3440-3450.

Harada H, Chan CM, Loesch A, Unwin R, Burnstock G. Induction of proliferation and apoptotic cell death via P2Y and P2X receptors, respectively, in rat glomerular mesangial cells. *Kidney Int* 2000; 57(3):949-958.

Heidenreich S, Tepel M, Schluter H, Harrach B, Zidek W. Regulation of rat mesangial cell growth by diadenosine phosphates. *J Clin Invest* 1995; 95(6):2862-2867.

Huwiler A, van RG, Wartmann M, Pfeilschifter J. Stimulation by extracellular ATP and UTP of the stress-activated protein kinase cascade in rat renal mesangial cells. *Br J Pharmacol* 1997; 120(5):807-812.

Huwiler A, Wartmann M, van den BH, Pfeilschifter J. Extracellular nucleotides activate the p38-stress-activated protein kinase cascade in glomerular mesangial cells. *Br J Pharmacol* 2000; 129(3):612-618.

Inesi G, Sagara Y. Thapsigargin, a high affinity and global inhibitor of intracellular Ca²⁺ transport ATPases. *Arch Biochem Biophys* 1992; 298(2):313-317.

Ip JH, Fuster V, Badimon L, Badimon J, Taubman MB, Chesebro JH. Syndromes of accelerated atherosclerosis: role of vascular injury and smooth muscle cell proliferation. *J Am Coll Cardiol* 1990; 15(7):1667-1687.

Ives HE, Schultz GS, Galardy RE, Jamieson JD. Preparation of functional smooth muscle cells from the rabbit aorta. *J Exp Med* 1978; 148(5):1400-1413.

Iwata K, Haruki S, Kimura T. High-performance liquid chromatographic determination of diadenosine 5',5"-p1,p4-tetraphosphate with precolumn fluorescence derivatization and its application to metabolism study in whole blood. *J Chromatogr B Biomed Appl* 1995; 667(2):339-343.

Jackson CL, Raines EW, Ross R, Reidy MA. Role of endogenous platelet-derived growth factor in arterial smooth muscle cell migration after balloon catheter injury. *Arterioscler Thromb* 1993; 13(8):1218-1226.

Jankowski J, Hagemann J, Yoon MS et al. Increased vascular growth in hemodialysis patients induced by platelet-derived diadenosine polyphosphates. *Kidney Int* 2001; 59(3):1134-1141.

Janssens R, Paindavoine P, Parmentier M, Boeynaems JM. Human P2Y2 receptor polymorphism: identification and pharmacological characterization of two allelic variants. *Br J Pharmacol* 1999; 127(3):709-716.

Jawien A, Bowen-Pope DF, Lindner V, Schwartz SM, Clowes AW. Platelet-derived growth factor promotes smooth muscle migration and intimal thickening in a rat model of balloon angioplasty. *J Clin Invest* 1992; 89(2):507-511.

Jiang H, Wu D, Simon MI. Activation of phospholipase C beta 4 by heterotrimeric GTP-binding proteins. *J Biol Chem* 1994; 269(10):7593-7596.

Jiang Q, Guo D, Lee BX et al. A mutational analysis of residues essential for ligand recognition at the human P2Y1 receptor. *Mol Pharmacol* 1997; 52(3):499-507.

Katz A, Wu D, Simon MI. Subunits beta gamma of heterotrimeric G protein activate beta 2 isoform of phospholipase C. *Nature* 1992; 360(6405):686-689.

Kennedy C, Qi AD, Herold CL, Harden TK, Nicholas RA. ATP, an agonist at the rat P2Y(4) receptor, is an antagonist at the human P2Y(4) receptor. *Mol Pharmacol* 2000; 57(5):926-931.

Laugwitz KL, Allgeier A, Offermanns S et al. The human thyrotropin receptor: a heptahelical receptor capable of stimulating members of all four G protein families. *Proc Natl Acad Sci U S A* 1996; 93(1):116-120.

Lee CH, Park D, Wu D, Rhee SG, Simon MI. Members of the Gq alpha subunit gene family activate phospholipase C beta isozymes. *J Biol Chem* 1992; 267(23):16044-16047.

Luthje J, Ogilvie A. The presence of diadenosine 5',5'''-P1,P3-triphosphate (Ap3A) in human platelets. *Biochem Biophys Res Commun* 1983; 115(1):253-260.

Luttrell LM, la Rocca GJ, van BT, Luttrell DK, Lefkowitz RJ. Gbetagamma subunits mediate Src-dependent phosphorylation of the epidermal growth factor receptor. A scaffold for G protein-coupled receptor-mediated Ras activation. *J Biol Chem* 1997; 272(7):4637-4644.

Lynch KJ, Touma E, Niforatos W et al. Molecular and functional characterization of human P2X(2) receptors. *Mol Pharmacol* 1999; 56(6):1171-1181.

Mallat Z, Tedgui A. Apoptosis in the vasculature: mechanisms and functional importance. *Br J Pharmacol* 2000; 130(5):947-962.

Marshall CJ. Specificity of receptor tyrosine kinase signaling: transient versus sustained extracellular signal-regulated kinase activation. *Cell* 1995; 80(2):179-185.

Martin F, Pintor J, Rovira JM, Ripoll C, Miras-Portugal MT, Soria B. Intracellular diadenosine polyphosphates: a novel second messenger in stimulus-secretion coupling. *FASEB J* 1998; 12(14):1499-1506.

Mateo J, Miras-Portugal MT, Rotllan P. Ecto-enzymatic hydrolysis of diadenosine polyphosphates by cultured adrenomedullary vascular endothelial cells. *Am J Physiol* 1997; 273(3 Pt 1):C918-C927.

Moghal N, Sternberg PW. Multiple positive and negative regulators of signaling by the EGF-receptor. *Curr Opin Cell Biol* 1999; 11(2):190-196.

Nicholas RA. Identification of the P2Y(12) receptor: a novel member of the P2Y family of receptors activated by extracellular nucleotides. *Mol Pharmacol* 2001; 60(3):416-420.

Offermanns S, Laugwitz KL, Spicher K, Schultz G. G proteins of the G12 family are activated via thromboxane A2 and thrombin receptors in human platelets. *Proc Natl Acad Sci U S A* 1994; 91(2):504-508.

Pintor J, az-Hernandez M, Bustamante C, Gualix J, de Terreros FJ, Miras-Portugal MT. Presence of dinucleotide and ATP receptors in human cerebrocortical synaptic terminals. *Eur J Pharmacol* 1999; 366(2-3):159-165.

Pintor J, Peral A, Hoyle CH et al. Effects of diadenosine polyphosphates on tear secretion in New Zealand white rabbits. *J Pharmacol Exp Ther* 2002; 300(1):291-297.

Ralevic V, Hoyle CH, Burnstock G. Pivotal role of phosphate chain length in vasoconstrictor versus vasodilator actions of adenine dinucleotides in rat mesenteric arteries. *J Physiol* 1995; 483 (Pt 3):703-713.

Reusch HP, Schaefer M, Plum C, Schultz G, Paul M. Gbeta gamma mediate differentiation of vascular smooth muscle cells. *J Biol Chem* 2001; 276(22):19540-19547.

Reusch HP, Zimmermann S, Schaefer M, Paul M, Moelling K. Regulation of Raf by Akt controls growth and differentiation in vascular smooth muscle cells. *J Biol Chem* 2001; 276(36):33630-33637.

Ridley AJ, Hall A. The small GTP-binding protein rho regulates the assembly of focal adhesions and actin stress fibers in response to growth factors. *Cell* 1992; 70(3):389-399.

Ross R, Glomset JA. The pathogenesis of atherosclerosis (first of two parts). *N Engl J Med* 1976; 295(7):369-377.

Ross R, Glomset JA. The pathogenesis of atherosclerosis (second of two parts). *N Engl J Med* 1976; 295(8):420-425.

Safrany ST, Ingram SW, Cartwright JL et al. The diadenosine hexaphosphate hydrolases from *Schizosaccharomyces pombe* and *Saccharomyces cerevisiae* are homologues of the human diphosphoinositol polyphosphate phosphohydrolase. Overlapping substrate specificities in a MutT-type protein. *J Biol Chem* 1999; 274(31):21735-21740.

Satterwhite CM, Farrelly AM, Bradley ME. Chemotactic, mitogenic, and angiogenic actions of UTP on vascular endothelial cells. *Am J Physiol* 1999; 276(3 Pt 2):H1091-H1097.

Schachter JB, Li Q, Boyer JL, Nicholas RA, Harden TK. Second messenger cascade specificity and pharmacological selectivity of the human P2Y1-purinoceptor. *Br J Pharmacol* 1996; 118(1):167-173.

Schluter H, Offers E, Bruggemann G et al. Diadenosine phosphates and the physiological control of blood pressure. *Nature* 1994; 367(6459):186-188.

Schwartz SM, Ross R. Cellular proliferation in atherosclerosis and hypertension. *Prog Cardiovasc Dis* 1984; 26(5):355-372.

Steinmetz M, Schlatter E, Boudier HA, Rahn KH, De Mey JG. Diadenosine polyphosphates cause contraction and relaxation in isolated rat resistance arteries. *J Pharmacol Exp Ther* 2000; 294(3):1175-1181.

Sternweis PC, Smrcka AV. Regulation of phospholipase C by G proteins. *Trends Biochem Sci* 1992; 17(12):502-506.

Taylor SJ, Chae HZ, Rhee SG, Exton JH. Activation of the beta 1 isozyme of phospholipase C by alpha subunits of the Gq class of G proteins. *Nature* 1991; 350(6318):516-518.

Tepel M, Jankowski J, Schluter H et al. Diadenosine polyphosphates' action on calcium and vessel contraction. *Am J Hypertens* 1997; 10(12 Pt 1):1404-1410.

Thomas WA, Reiner JM, Florentin FA, Lee KT, Lee WM. Population dynamics of arterial smooth muscle cells. V. Cell proliferation and cell death during initial 3 months in atherosclerotic lesions induced in swine by hypercholesterolemic diet and intimal trauma. *Exp Mol Pathol* 1976; 24(3):360-374.

Tsukamoto A, Kaneko Y. Thapsigargin, a Ca(2+)-ATPase inhibitor, depletes the intracellular Ca2+ pool and induces apoptosis in human hepatoma cells. *Cell Biol Int* 1993; 17(10):969-970.

van Corven EJ, Hordijk PL, Medema RH, Bos JL, Moolenaar WH. Pertussis toxin-sensitive activation of p21ras by G protein-coupled receptor agonists in fibroblasts. *Proc Natl Acad Sci U S A* 1993; 90(4):1257-1261.

van der GM, Jankowski J, Schluter H, Zidek W, Tepel M. Mediation of the vasoactive properties of diadenosine tetraphosphate via various purinoceptors. *J Hypertens* 1998; 16(12 Pt 2):1939-1943.

van der GM, Khattab M, Borgel J, Schluter H, Zidek W. Differential effects of diadenosine phosphates on purinoceptors in the rat isolated perfused kidney. Br J Pharmacol 1997; 120(8):1453-1460.

von DA, Ogilvie A. Ecto-diadenosine 5',5"-P1,P4-tetraphosphate (Ap4A)-hydrolase is expressed as an ectoenzyme in a variety of mammalian and human cells and adds new aspects to the turnover of Ap4A. Biofactors 2000; 11(3):179-187.

Vouret-Craviari V, Van Obberghen-Schilling E, Scimeca JC, Van OE, Pouyssegur J. Differential activation of p44mapk (ERK1) by alpha-thrombin and thrombin-receptor peptide agonist. Biochem J 1993; 289 (Pt 1):209-214.

Wu D, Katz A, Simon MI. Activation of phospholipase C beta 2 by the alpha and beta gamma subunits of trimeric GTP-binding protein. Proc Natl Acad Sci U S A 1993; 90(11):5297-5301.