

## Literaturverzeichnis

- Alleva EF, Santucci D. Psychosocial vs. "physical" stress situations in rodents and humans: role of neurotrophins. *Physiol Behav* 2001, 73: 313-20
- Aloe L, Bracci Laudiero L, Alleva E, Lambiase A, Micera A, Tirassa P. Emotional stress induced by parachute jumping enhances blood nerve growth factor levels and the distribution of nerve growth factor receptors in lymphocytes. *Proc Natl Acad Sci USA* 1994, 91: 10440-10444
- Aloe L, Bracci-Laudiero L, Manni L. The expanding role of nerve growth factor: from neurotrophic activity to immunologic diseases. *Allergy* 1997, 52: 883-894
- Aloe L, Tuveri MA, Guerra P, Pinna L, Tirassa P, Micera A, Alleva E. Changes in human plasma nerve growth factor level after chronic alcohol consumption and withdrawal. *Alcohol Clin Exp Res* 1996, 3: 462-466
- Altar CA. Neurotrophins and depression. *Trends Pharmacol Sci* 1999, 20: 59-61
- Andreasson S, Allebeck P, Engstrom A, Rydberg U (1987) Cannabis and schizophrenia. A longitudinal study of Swedish conscripts. *Lancet* 2: 1483-1486
- Ang ET, Wong PT, Moochhala S, Ng YK. Neuroprotection associated with running: is it a result of increased endogenous neurotrophic factors? *Neuroscience* 2003, 118: 334-345
- Araujo DM, Lapchak PA, Robitaille Y, Gauthier S, Quirion R. Differential alteration of various cholinergic markers in cortical and subcortical regions of human brain in Alzheimer's disease. *J Neurochem* 1988, 50: 1914-1923
- Bersani G, Iannitelli A, Maselli P, Pancheri P, Aloe L, Angelucci F, Alleva E. Low nerve growth factor plasma levels in schizophrenic patients: a preliminary study. *Schizophrenia Res* 1999, 37: 201-203
- Biber K, Walden J, Gebicke-Harter P, Berger M, van Calker D. Adenosine A1 receptor-mediated activation of phospholipase C in cultured astrocytes depends on the level of receptor expression. *J Neurosci* 1996, 17: 4956-4964
- Bonini S, Lambiase A, Bonini S, Angelucci F, Magrini L, Manni L and Aloe L. Circulating nerve growth factor levels are increased in humans with allergic diseases and asthma. *Proc Natl Acad Sci USA* 1995, 93: 10955-10960
- Chen B, Dowlatshahi D, MacQueen GM, Wang JF, Young LT. Increased hippocampal BDNF immunoreactivity in subjects treated with antidepressant medication. *Biol Psychiatry* 2001, 50: 260-5
- Chrousos GP, Gold PW. The concept of stress and stress systems disorders. *JAMA* 1992, 267: 1244-1252
- Ciccarelli R, di Iorio P, Bruno V, Battaglia G, Dálimonte I, D'Onofrio M, Nicoletti F, Caciagli F. Activation of A (1) adenosine of mGlu3 metabotropic glutamate receptors enhances the release of nerve growth factor and S-100beta protein from cultured astrocytes. *Glia* 1999, 27: 275-281
- Cirulli F. Role of environmental factors on brain development and nerve growth factor expression. *Physiol Behav* 2001, 73: 321-330
- Cotman CW, Berchtold NC. Exercise: a behavioural intervention to enhance brain health and plasticity. *Trends Neurosci* 2003, 25: 295-301
- De Santis S, Pace A, Bove L, Cognetti F, Properzi F, Fiore M, Triaca V, Savarese A, Simone MD, Jandolo B, Manzione L, Aloe L. Patients treated with antitumor drugs displaying neurological deficits are characterized by a low circulating level of nerve growth factor. *Clin Cancer Res* 2000, 6: 90-95
- Duggan C, Sham P, Lee A, Minne C, Murray R. Neuroticism: a vulnerability marker for depression evidence from a family study. *J Affect Disord* 1995, 35: 139-43
- Duman RS, Heninger GR, Nestler EJ. A molecular and cellular theory of depression. *Arch Gen Psychiatry* 1997, 54: 597-606
- Duman RS, Malberg J, Nakagawa S and D'Sa C. Neuronal plasticity and survival in mood disorders. *Biol Psychiatry* 2000, 48: 732-739

- Fantini F, Magnini C, Bracci Laudiero L and Pincelli CT: Nerve growth factor is increased in psoriatic skin. *J Invest Dermatol* 1995, 105: 854-855
- Finkbeiner S. CREB couples neurotrophin signals to survival messages. *Neuron* 2000, 25:11-14
- Friedman WJ, Greene LA. Neurotrophin signaling via Trks and p75. *Exp Cell Res* 1999, 253: 131-42
- Garrido R, King-Pospisil K, Son KW, Hennig B, Toborek M. Nicotine upregulates nerve growth factor expression and prevents apoptosis of cultured spinal cord neurons. *Neurosci Res* 2003 47: 349-55
- Hadjiconstantinou M, McGuire L, Duchemin AM, Laskowski B, Kiecolt-Glaser J, Glaser R. Changes in plasma nerve growth factor levels in older adults associated with chronic stress. *J Neuroimmunol* 2001, 116: 102-106
- Hamada A, Watanabe N, Ohtomo H, Matsuda H. Nerve growth factor enhances survival and cytotoxic activity of human eosinophils. *Br J Haematol* 1996, 93: 299-302
- Hefti F, Weiner WJ. Nerve growth factor and Alzheimer's disease. *Ann Neurol* 1986, 20: 275-281
- Heinz A, Hermann D, Smolka MN, Rieks M, Graf KJ, Pohlau D, Kuhn W, Bauer M. Effects of acute psychological stress on adhesion molecules, interleukins and sex hormones: implications for coronary heart disease. *Psychopharmacology (Berl)* 2003, 165: 111-7
- Hellweg R, Gericke CA, Jendroska K, Hartung HD, Cervós-Navarro, J. NGF content in the cerebral cortex of non-demented patients with amyloid-plaques and in symptomatic Alzheimer's disease. *Int J Devl Neuroscience* 1998b, 16: 787-794
- Hellweg R, Hock C, Hartung HD. An improved rapid and highly sensitive enzyme immunoassay for nerve growth factor. *Technique, J Meth Cell Mol Biol* 1989, 1: 43-49
- Hellweg R, Jockers-Scherübl M. Neurotrophic factors in memory disorders. *Life Sci* 1994, 55: 2165-2169
- Hellweg R, von Arnim CAF, Büchner M, Huber R, Riepe MW. Neuroprotection and neuronal dysfunction upon repetitive inhibition of oxidative phosphorylation. *Exp Neurol* 2003, 183: 346-354
- Hellweg R, von Richthofen S, Anders D, Baethge C, Röpke S, Hartung HD, Gericke CA. The time course of nerve growth factor content in different neuropsychiatric diseases - a unifying hypothesis. *J Neural Transm* 1998a, 105: 871-903
- Hellweg R. Der nerve growth factor bei neuropsychiatrischen Erkrankungen - ein pleiotroper Modulator mit peripherer und zentralnervöser Wirkung (Habilitationsschrift am Fachbereich Humanmedizin der Freien Universität Berlin im Fachbereich Psychiatrie) 1997
- Herman JP, Cullinan WE. Neurocircuitry of Stress: central control of the hypothalamic-pituitary-adrenocortical axis. *Trends Neurosci* 1997, 20: 78-84
- Hohlfeld R, Kerschensteiner M, Stadelmann C, Lassmann H, Wekerle H. The neuroprotective effect of inflammation: implications for the therapy of multiple sclerosis. *J Neuroimmunol* 2000, 107: 161-166
- Hörtnagl H, Hellweg R. Pathophysiological aspects of human neurodegenerative diseases. *Wien Klin Wochenschr* 1997, 109: 623-635
- Jensen J, Mikkelsen J, Mørk A. Increased adenylyl cyclase type 1 mRNA, but not adenylyl cyclase type 2 in the rat hippocampus following antidepressant treatment. *Eur Neuropsychopharmacol* 2000, 10: 105-111
- Jockers-Scherübl MC, Matthies U, Danker-Hopfe H, Lang UE, Hellweg R. Chronic cannabis abuse raises nerve growth factor concentrations in drug-naïve schizophrenic patients. *Psychopharmacology (im Druck)*
- Jockers-Scherübl MC, Zouboulis CC, Boegner F, Hellweg R. Is nerve growth factor a serum marker for neurological and psychiatric complications in Behcet's disease? *Lancet* 1996, 347: 982
- Jonnala RR, Terry AV, Buccafusco JJ. Nicotine increases the expression of high affinity nerve growth factor receptors in both in vitro and in vivo. *Life Sci* 2002, 70: 1543-54
- Karege F, Perret G, Bondolfi G, Schwald M, Bertschy G, Aubry JM. Decreased serum brain-derived neurotrophic factor levels in major depressed patients. *Psychiatry Res* 2002, 109: 143-8

- Keegan BM, Noseworthy JH. Multiple sclerosis. *Ann Rev Med* 2002, 53: 285-302
- Kerschensteiner M, Gallmeier E, Behrens L, Klinkert WEF, Kolbeck R, Hoppe E, Oropenza-Wekerle RL, Bartke I, Stadelmann C, Lassmann H, Wekerle H, Hohlfeld H. Activated human T cells, B cells and monocytes produce brain-derived neurotrophic factor (BDNF) in vitro and in brain lesions: a neuroprotective role of inflammation? *J Exp Med* 1999, 189: 865-870
- Korschning S, Thoenen H. Two-site enzyme immunoassay for nerve growth factor. In: Barnes D, Sirbasku Da (Hrsg). *Methods in Enzymology: Peptide growth factors*. Academic Press Orlando 1987, 147: 167-85
- Lambiase A, Bonini S, Bonini S, Micera A, Magrini L, Bracci Laudiero L, Aloe L. Increased plasma levels of nerve growth factor in vernal keratoconjunctivitis and relationship to conjunctival mast cells. *Invest Ophthalmol Vis Sci* 1995, 36: 2127-2132
- Leung DYM. Atopic dermatitis: New insights and opportunities for therapeutic intervention. *J Allergy Clin Immunol* 2000, 105: 860-876
- Levi-Montalcini R, Skaper S, Dal Toso R, Petrelli L, Leon A. Nerve growth factor: from neurotrophin to neurokine. *Trends Neurosci* 1996, 19: 514-520
- Linker RA, Mäurer M, Gaupp S, Martini R, Holtmann B, Giess R, Rieckmann P, Lassmann H, Toyka KV, Sendtner M, Gold R. CNTF is a major protective factor in demyelinating CNS disease: A neurotrophic cytokine as modulator of neuroinflammation. *Nature Med* 2002, 6: 620-624
- Luppi P, Levi-Montalcini R, Bracci-Laudiero L, Bertolini A, Arletti R, Tavernari D, Vigneti E, Aloe L. NGF is released into plasma during human pregnancy: an oxytocin-mediated response? *Neuroreport* 1993, 4: 1063-1065
- Ma QP, Woolf CJ. The progressive tactile hyperalgesia induced by peripheral inflammation is nerve growth factor dependent. *Neuroreport* 1997, 8: 804-810
- MacQueen GM, Ramakrishnan K, Croll SD, Siuciak JA, Yu G, Young LT, Fahnestock M. Performance of heterozygous brain-derived neurotrophic factor knockout mice on behavioral analogues of anxiety, nociception, and depression. *Behav Neurosci* 2001, 115: 1145-53
- Mahadik SP, Parikh V, Khan MM, Buckley PF. Differential effects of antipsychotics on nerve growth factor in rat brain. *NCDEU Meeting Abstracts* 2002 Session II-8
- Malcangio M, Lessmann V. A common thread for pain and memory synapses? Brain-derived neurotrophic factor and trkB receptors. *Trends Pharmacol Sci* 2003, 24: 116-21
- Manji HK, Quiroz JA, Sporn J, Payne JL, Denicoff K, Gray N, Zarate CA, Charney DS. Enhancing neuronal plasticity and cellular resilience to develop novel, improved therapeutics for difficult-to-treat depression. *Biol Psychiatry* 2003, 53: 707-742
- McEwen BS. The neurobiology of stress: from serendipity to clinical relevance. *Brain Res* 2000, 886: 172-189
- Mostert S, Kesselring J. Effects of a short-term exercise training program on aerobic fitness, fatigue, health perception, and activity level of subjects with multiple sclerosis. *Mult Scler* 2002, 8: 161-168
- Mufson EJ, Bothwell M, Kordower JH. Loss of nerve growth factor receptor-containing neurons in Alzheimer's disease: a quantitative analysis across subregions of the basal forebrain. *Exp Neurol* 1990, 105: 221-232
- Mumford C, Compston A. Problems with rating scales for multiple sclerosis: a novel approach - the CAMBS score. *J Neurol* 1983, 240: 209-215
- Nemeroff CB. The corticotropin-releasing factor (CRF) hypothesis of depression: new findings and new directions. *Mol Psychiatry* 1996, 1: 336-42
- Neves-Pereira M, Mundo E, Muglia P, King N, Macciardi F, Kennedy JL. The brain-derived neurotrophic factor gene confers susceptibility to bipolar disorder: evidence from a family-based association study. *Am J Hum Genet* 2002, 71: 651-5
- Otten U, Scully JL, Ehrhard PB, Gadiant RA. Neurotrophins : signals between the nervous and immune system, *Prog Brain Res* 1994, 103: 293-305

- Parikh V, Evans DR, Khan MM, Mahadik SP. Nerve growth factor in never-medicated first-episode psychotic and medicated chronic schizophrenic patients: possible implications for treatment outcome. *Schizophr Res* 2003, 60: 117-23
- Petajan JH, White AT. Recommendations for physical activity in patients with multiple sclerosis. *Sports Med* 1999, 27: 179-191
- Rajkowska G. Postmortem studies in mood disorders indicate altered numbers of neurons and glial cells. *Biol Psychiatry* 2000, 48: 766-77
- Russo-Neustadt A, Beard RC, Cotman CW. Exercise, antidepressant medications, and enhanced brain derived neurotrophic factor expression. *Neuropharmacology* 1999, 21: 679-682
- Russo-Neustadt A, Ha T, Ramirez R, Kesslak JP. Physical activity-antidepressant treatment combination: impact on brain-derived neurotrophic factor and behavior in an animal model. *Behav Brain Res* 2001, 120: 87-95
- Russo-Neustadt A. Brain-derived neurotrophic factor, behavior, and new directions for the treatment of mental disorders. *Semin Clin Neuropsychiatry* 2003, 8: 109-18
- Russo-Neustadt AA, Beard RC, Huang YM, Cotman CW. Physical activity and antidepressant treatment potentiate the expression of specific brain-derived neurotrophic factor transcripts in the rat hippocampus. *Neuroscience* 2002, 101: 305-312
- Saarelainen T, Hendolin P, Lucas G, Koponen E, Sairanen M, MacDonald E, Agerman K, Haapasalo A, Nawa H, Aloyz R, Ernfors P, Castren E. Activation of the TrkB neurotrophin receptor is induced by antidepressant drugs and is required for antidepressant-induced behavioral effects. *J Neurosci* 2003, 23: 349-57
- Schildkraut JJ. The catecholamine hypothesis of affective disorders: a review of supporting evidence. *Am J Psychiatry* 1965, 122: 509-22
- Schinder AF, Poo M. The neurotrophin hypothesis for synaptic plasticity. *Trends Neurosci* 2000, 23: 639-645
- Schwartz M, Moalem G, Leibowitz-Amit R, Cohen IR. Innate and adaptive immune responses can be beneficial for CNS repair. *Trends Neurosci* 1999, 22: 295-299
- Scott SA, Crutcher KA. Nerve growth factor and Alzheimer's disease. *Rev Neurosci* 1994, 5: 179-211
- Sen S, Nesse RM, Stoltenberg SF, Li S, Gleiberman L, Chakravarti A. A BDNF coding variant is associated with the NEO personality inventory domain neuroticism, a risk factor for depression. *Neuropsychopharmacol* 2003, 28: 397-401
- Serrano Sanchez T, Robinson R, Agramonte M, Lorigados M, Pedre L, Diaz Armesto I, Gonzalez Fraguera ME, Dorta-Contreras AJ. Endogenous nerve growth factor in patients with Alzheimer's disease. *Rev Neurol* 2001, 32: 825-828
- Serrano T, Lorigados LC, Armenteros S. Nerve growth factor levels in normal human sera. *Neuroreport* 1996, 8: 179-181
- Sheline YI. 3D MRI studies of neuroanatomic changes in unipolar major depression: the role of stress and medical comorbidity. *Biol Psychiatry* 2000, 48: 791-800
- Sheridan JF, Stark JL, Avitsur R, Padgett DA. Social disruption, immunity, and susceptibility to viral infection. Role of glucocorticoid insensitivity and NGF. *Ann N Y Acad Sci* 2000, 917: 894-905
- Shimizu E, Hashimoto K, Okamura N, Koike K, Komatsu N, Kumakiri C. Alterations of serum levels of brain-derived neurotrophic factor (BDNF) in depressed patients with or without antidepressants. *Biol Psychiatry* 2003, 54: 70-5
- Shirayama Y, Chen AC, Nakagawa S, Russell DS, Duman RS. Brain-derived neurotrophic factor produces antidepressant effects in behavioral models of depression. *J Neurosci* 2002, 22: 3251-3261
- Siegel GJ, Chauhan NB. Neurotrophic factors in Alzheimer's and Parkinson's disease brain. *Brain Res Rev* 2000, 33: 199-227

- Siuciak JA, Lewis DR, Wiegand SJ, Lindsay RM. Antidepressant-like effect of brain-derived neurotrophic factor (BDNF). *Pharmacol Biochem Behav* 1997, 56: 131–137
- Skaper SD, Pollock M, Facci L. Mast cells differentially express and release active high molecular weight neurotrophins. *Brain Res Mol Brain Res* 2001, 97: 177-185
- Solomon GF. Psychoneuroimmunology: Interactions between central nervous system and immune system. *J Neurosci Res* 1987, 18: 1-9
- Stadelmann C, Kerschensteiner M, Misgeld T, Brück W, Hohlfeld R, Lassman H. BDNF and gp145trkB in multiple sclerosis brain lesions. Neuroprotective interactions between immune cells and neuronal cells? *Brain* 2002, 125: 75-85
- Tafet GE, Bernardini R. Psychoneuroendocrinological links between depression and chronic stress. *Prog Neuropsychopharmacol Biol Psychiatry* 2003, 27: 893-903
- Terry AV, Hill WD, Parikh V, Waller JL, Evans DR, Mahadik SP. Differential effects of haloperidol, risperidone, and clozapine exposure on cholinergic markers and spatial learning performance in rats. *Neuropsychopharmacology* 2003, 28: 300-9
- Thoenen H, Barde YA, Davies AM, Johnson JE. Neurotrophic factors and neuronal death. *Ciba Found Symp* 1987a, 126: 82–95
- Thoenen H. Neurotrophins and neuronal plasticity. *Science* 1995, 270: 593-8
- Tyler WJ, Alonso M, Bramham CR, Pozzo-Miller LD. From acquisition to consolidation: on the role of brain-derived neurotrophic factor signaling in hippocampal-dependent learning. *Learn Mem* 2002, 9: 224-37
- Valdo P, Stegagno C, Mazzucco S, Zuliani E, Zanusso GL, Moretto G, Raine CS, Bonetti B. Enhanced expression of NGF receptors in multiple sclerosis lesions. *J Neuropathol Exp Neurol* 2002, 61: 91-98
- Villoslada P, Hauser SL, Bartke I, Unger J, Heald N, Rosenberg D, Cheung SW, Mobley WC, Fisher S, Genain CP. Human nerve growth factor protects common marmosets against autoimmune encephalomyelitis by switching the balance of T helper cell type 1 and 2 cytokines within the nervous system. *J Exp Med* 2000, 191: 1799-1806
- Whitehouse PJ, Price DL, Struble RG, Clark AW, Coyle JT, Delon MR. Alzheimer's disease and senile dementia: loss of neurons in the basal forebrain. *Science* 1982, 215: 1237-1239