

10 Literaturverzeichnis

- Aalto-Setala, K., Fisher, E.A., Chen, X., Chajek-Shaul, T., H., T., Z., R., Walsh, A., Ramakrishnan, R., Ginsberg, H.N., Breslow, J.L., Mechanism of hypertriglyceridemia in human apolipoprotein (apo) CIII transgenic mice. Diminished very low density lipoprotein fractional catabolic rate associated with increased apo CIII and reduced apo E on the particles, *J. Clin. Invest.* 90, 1889-1900 (1992).
- Abbott, N.J., Romero, I.A., Transporting therapeutics across the blood-brain barrier, *Mol. Med. Today* 2, 106-113 (1996).
- Absalom, D.R., Opsonins and dysopsonins: an overview, *Methods Enzymol.* 132, 281-318 (1986).
- Adams, P.D., Lopez, P., Sellers, W.R., Kaelin, W.G., Jr., Fluorescence-activated cell sorting of transfected cells, *Methods Enzymol.* 283, 59-72 (1997).
- Aggerbeck, L.P., Wetterau, J.R., Weisgraber, K.H., Wu, C.S., Lindgren, F.T., Human apolipoprotein E3 in aqueous solution. II. Properties of the amino- and carboxyl-terminal domains, *J. Biol. Chem.* 263, 6249-6258 (1988).
- Allen, T.M., Hansen, C., Martin, F., Redemann, C., Yau-Young, A., Liposomes containing synthetic lipid derivates of poly(ethylene glycol) show prolonged circulating half-lives in vivo, *Biochim. Biophys. Acta* 1066, 29-36 (1991).
- Allen, T.M., Mehra, T., Hansen, C., Chin, Y.C., Stealth liposomes: an improved sustained release system for 1-beta-D-arabinofuranosylcytosine, *Cancer Res.* 52, 2431-2439 (1992).
- Altieri, D.C., Mannuccio, P.M., Capitanio, A.M., Binding of fibrinogen to human monocytes, *J. Clin. Invest.* 78(1986).
- Alyautdin, R.N., Gothier, D., Petrov, V., Kharkevich, D., Kreuter, J., Analgesic activity of the hexapeptide dalargin adsorbed on the surface of polysorbate 80-coated poly(butyl cyanoacrylate) nanoparticles, *Eur. J. Pharm. Biopharm.* 41, 44-48 (1995).
- Alyautdin, R.N., Petrov, V.E., Langer, K., Berthold, A., Kharkevich, D.A., Kreuter, J., Delivery of loperamide across the blood-brain barrier with polysorbate 80-coated polybutylcyanoacrylate nanoparticles, *Pharm. Res.* 14, 325-328 (1997).
- Alyautdin, R.N., Petrov, V.E., Langer, K., Berthold, A., Kreuter, J., Kharkevich, D.A., The delivery of loperamide to the brain by using polybutyl cyanoacrylate nanoparticles, *Eksp. Klin. Farmakol.* 61, 17-20 (1998a).

- Alyautdin, R.N., Tezikov, E.B., Ramge, P., Kharkevich, D.A., Begley, D.J., Kreuter, J., Significant entry of tubocurarine into the brain of rats by adsorption to polysorbate 80-coated polybutylcyanoacrylate nanoparticles: an in situ brain perfusion study, *J. Microencapsul.* 15, 67-74 (1998b).
- Amersham, Gel filtration: Principles and Methods, (1993).
- Anderson, N.L., Anderson, N.G., A two-dimensional gel database of human plasma proteins, *Electrophoresis* 12, 883-906 (1991).
- Appel, R.D., Hochstrasser, D.F., Funk, M., Vargas, J.R., Pellegrini, C., Müller, A.F., Scherrer, J.R., The MELANIE project: from a biopsy to automatic protein map interpretation by computer, *Electrophoresis* 14, 1223-1231 (1991).
- Appel, R.D., Palagi, P.M., Walther, D., Vargas, J.R., Sanchez, J.C., Ravier, F., Pasquali, C., Hochstrasser, D.F., Melanie II--a third-generation software package for analysis of two-dimensional electrophoresis images: I. Features and user interface, *Electrophoresis* 18, 2724-2734 (1997a).
- Appel, R.D., Vargas, J.R., Palagi, P.M., Walther, D., Hochstrasser, D.F., Melanie II--a third-generation software package for analysis of two-dimensional electrophoresis images: II. Algorithms, *Electrophoresis* 18, 2735-2748 (1997b).
- Arai, T., Norde, W., The behaviour of some model proteins at solid-liquid interfaces 2, Sequential and competitive adsorption, *Colloids Surfaces* 51, 186-216 (1990).
- Archambault, J.G., Brash, J.L., Protein resistant polyurethane surfaces by chemical grafting of PEO: amino-terminated PEO as grafting reagent, *Colloids Surf B Biointerfaces* 39, 9-16 (2004).
- Audus, K.L., Borchardt, R.T., Characteristics of the large neutral amino acid transport system of bovine brain microvessel endothelial cell monolayers, *J. Neurochem.* 47, 484-488 (1986).
- Batrakova, E.V., Li, S., Miller, D.W., Kabanov, A.V., Pluronic P85 increases permeability of a broad spectrum of drugs in polarized BBMEC and Caco-2 cell monolayers, *Pharm. Res.* 16, 1366-1372 (1999).
- Bauer, B., In vitro Zellkulturmodelle der Blut-Hirn Schranke zur Untersuchung der permeation und P-Glycoprotein-Interaktion von Arzneistoffen, Dissertation, Ruprecht-Karls-Universität Heidelberg (2002).
- Bauer, K.H., Frömmig, K.-H., Führer, C., Lehrbuch der Pharmazeutischen Technologie, Wissenschaftliche Verlagsgesellschaft mbH Stuttgart, (2002).
- Begley, D.J., The blood-brain barrier: principles for targeting peptides and drugs to the central nervous system, *J. Pharm. Pharmacol.* 48, 136-46 (1996).
- Behnke, A.R., Yarborough, O.D., Physiologic studies of helium, *US nav. med. Bull.* 36, 542-548 (1938).

- Bellinzona, M., Roser, F., Matthies, C., Samii, M., Saini, M., Biopolymer-mediated suramin chemotherapy in the treatment of experimental brain tumours, *Acta Oncol.* 43, 259-263 (2004).
- Benita, S., Levy, M.Y., Submicron emulsions as colloidal drug carriers for intravenous administration: comprehensive physicochemical characterization, *J. Pharm. Sci.* 82, 1069-1079 (1993).
- Bergström, K., Holmberg, K., Safranj, A., Hoffman, A.S., Edgell, M.J., Kozlowski, A., Hovanes, B.A., Harris, J.M., Reduction of fibrinogen adsorption on PEG-coated polystyrene surfaces, *J. Biomed. Mater. Res.* 26, 779-790 (1992).
- Bickel, U., Yoshikawa, T., Landaw, E.M., Faull, K.F., Pardridge, W.M., Pharmacologic effects in vivo in brain by vector-mediated peptide drug delivery, *Proc. Natl. Acad. Sci. U S A* 90, 2618-2622 (1993).
- Bisgaier, C.L., Siebenkas, M.V., Williams, K.J., Effects of apolipoproteins A-IV and A-I on the uptake of phospholipid liposomes by hepatocytes, *J. Biol. Chem.* 264, 862-866 (1989).
- Bittner, B., Guenzi, A., Fullhardt, P., Zuercher, G., Gonzalez, R.C., Mountfield, R.J., Improvement of the bioavailability of colchicine in rats by co-administration of D-alpha-tocopherol polyethylene glycol 1000 succinate and a polyethoxylated derivative of 12-hydroxy-stearic acid, *Arzneimittelforschung* 52, 684-688 (2002).
- Bjellqvist, B., Ek, K., Righetti, P.G., Gianazza, E., Gorg, A., Westermeier, R., Postel, W., Isoelectric focusing in immobilized pH gradients: principle, methodology and some applications, *J. Biochem. Biophys. Methods* 6, 317-339 (1982).
- Bjellqvist, B., Pasquali, C., Ravier, F., Sanchez, J.C., Hochstrasser, D.F., A nonlinear wide-range immobilized pH gradient for two-dimensional electrophoresis and its definition in a relevant pH scale, *Electrophoresis* 14, 1357-1365 (1993).
- Blunk, T., Hochstrasser, D.F., Sanchez, J.-C., Müller, B.W., Müller, R.H., Colloidal carriers for intravenous drug targeting: plasma protein adsorption patterns on surface-modified latex particles evaluated by two-dimensional polyacrylamide gel electrophoresis, *Electrophoresis* 14, 1382-1387 (1993).
- Blunk, T., Plasmaproteinadsorption auf kolloidalen Arzneistoffträgern, Dissertation, Christian-Albrechts-Universität zu Kiel (1994).
- Blunk, T., Lück, M., Diederichs, J.E., Müller, B.W., H., M.R., Plasma Protein Adsorption on Polystyrene Particles of Different Size Modified with Poloxamer 407. *Intern. Symp. Control. Rel. Bioact. Mater.* Vol. 22 47-48 (1995).
- Blunk, T., Lück, M., Calvör, A., Hochstrasser, D.F., Sanchez, J.-C., Müller, B.W., H., M.R., Kinetics of plasma protein adsorption on model particles for controlled drug delivery and drug targeting, *Eur. J. Pharm. Biopharm.* 42, 262-268 (1996).

- Bock, T.K., Lucks, J.S., Kleinebudde, P., Müller, R.H., Müller, B.W., High Pressure Homogenisation of Parenteral Fat Emulsions - Influence of Process Parameters on Emulsion Quality, *Eur. J. Pharm. Biopharm.* 40 (3), 157-160 (1994).
- Bogman, K., Erne-Brand, F., Alsenz, J., Drewe, J., The role of surfactants in the reversal of active transport mediated by multidrug resistance proteins, *J. Pharm. Sci.* 92, 1250-1261 (2003).
- Bogman, K., Zysset, Y., Degen, L., Hopfgartner, G., Gutmann, H., Alsenz, J., Drewe, J., P-glycoprotein and surfactants: effect on intestinal talinolol absorption, *Clin. Pharmacol. Ther.* 77, 24-32 (2005).
- Borchard, G., Audus, K.L., Shi, F., Kreuter, J., Uptake of surfactant coated poly(methyl)methacrylate nanoparticles by bovine brain microvessel endothelial cell monolayers, *Int. J. Pharm.* 110, 29-35 (1994).
- Borchard, G., Arzneistofftransport an der Bluthirnschranke, in: Müller, R. H., Hildebrand, G. E. (Hrsg.), *Pharmazeutische Technologie: Moderne Arzneiformen, Wissenschaftliche Verlagsgesellschaft, Stuttgart*, (1998)
- Bradford, M.M., A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding, *Anal Biochem* 72, 248-54 (1976).
- Brash, J.L., Ten Hove, P., Effect of plasma dilution on adsorption of fibrinogen to solid surfaces, *Thromb. Haemost.* 51, 326-330 (1984).
- Brash, J.L., Protein adsorption at the solid-solution interface in relation to blood-material interactions, in: Brash, J. L., Horbett, T. A. (Hrsg.), *Proteins at Interfaces: Physicochemical and Biochemical studies*, Am. Chem. Soc., Washington D. C., USA, 490-506 (1987)
- Brash, J.L., Ten Hove, P., Protein adsorption studies on 'standard' polymeric materials, *J. Biomater. Sci. Polym. Ed.* 4, 591-599 (1993).
- Brash, J.L., Exploiting the current paradigm of blood-material interactions for the rational design of blood-compatible materials, *J. Biomater. Sci. Polym. Ed.* 11, 1135-46 (2000).
- Brat, D.J., Windebank, A.J., Brimijoin, S., Emulsifier for intravenous cyclosporin inhibits neurite outgrowth, causes deficits in rapid axonal transport and leads to structural abnormalities in differentiating N1E.115 neuroblastoma, *J. Pharmacol. Exp. Ther.* 261, 803-810 (1992).
- Breemhaar, W., Brinkman, E., Ellens, D.J., Beugeling, T., Bantjes, A., Preferential adsorption of high density lipoprotein from blood plasma onto biomaterial surfaces, *Biomaterials* 5, 269-274 (1984).

- Brewster, M.E., Anderson, W.R., Helton, D.O., Bodor, N., Pop, E., Dose-dependent brain delivery of zidovudine through the use of a zidovudine chemical delivery system, *Pharm. Res.* 12, 796-8 (1995).
- Broadwell, R.D., Baker-Cairns, B.J., Friden, P.M., Oliver, C., Villegas, J.C., Transcytosis of protein through the mammalian cerebral epithelium and endothelium. III. Receptor-mediated transcytosis through the blood-brain barrier of blood-borne transferrin and antibody against the transferrin receptor, *Exp. Neurol.* 142, 47-65 (1996).
- Brown, M.S., Goldstein, J.L., A receptor-mediated pathway for cholesterol homeostasis, *Science* 232, 34-47 (1986).
- Brunner, U., Edelgas intravenös verbessert die Anästhesie, *Pharmazeutische Zeitung* 32(1999).
- Bucke, W.E., Leitzke, S., Diederichs, J.E., Borner, K., Hahn, H., Ehlers, S., Müller, R.H., Surface-modified amikacin-liposomes: organ distribution and interaction with plasma proteins, *J. Drug. Target.* 5, 99-108 (1997).
- Burnham, W.R., Hansrani, P.K., Knott, C.E., Cook, J.A., Davis, S.S., Stability of fat emulsions based intravenous feeding mixture, *Int. J. Pharm.* 13, 9-22 (1983).
- Buttle, I., O/W-Emulsionen für die intravenöse Applikation von Arzneistoffen, Dissertation, Freie Universität Berlin (2004).
- Camner, P., Lundborg, M., Lastbom, L., Gerde, P., Gross, N., Jarstrand, C., Experimental and calculated parameters on particle phagocytosis by alveolar macrophages, *J. Appl. Physiol.* 92, 2608-2616 (2002).
- Cardin, A.D., Hirose, N., Blankenship, D.T., Jackson, R.L., Harmony, J.A., Sparrow, D.A., Sparrow, J.T., Binding of a high reactive heparin to human apolipoprotein E: identification of two heparin-binding domains, *Biochem. Biophys. Res. Commun.* 134, 783-789 (1986).
- Carignano, M.A., Szleifer, I.I., Prevention of protein adsorption by flexible and rigid chain molecules, *Colloids Surf. B. Biointerfaces* 18, 169-182 (2000).
- Cattel, L., Ceruti, M., Dosio, F., From conventional to stealth liposomes: a new frontier in cancer chemotherapy, *Tumori* 89, 237-249 (2003).
- Chesnay, S., Huang, L., Structure and function of lipid-DNA complexes for gene delivery, *Annu. Rev. Biophys. Biomol. Struct.* 29, 27-47 (2000).
- Chonn, A., Semple, S.C., Cullis, P.R., Association of blood proteins with large unilamellar liposomes in vivo. Relation to circulation lifetimes, *J. Biol. Chem.* 267, 18759-18765 (1992).
- Coester, C., Kreuter, J., von Briesen, H., Langer, K., Preparation of avidin-labelled gelatin nanoparticles as carriers for biotinylated peptide nucleic acid (PNA), *Int. J. Pharm.* 196, 147-149 (2000a).

- Coester, C., Development of a new carrier system for oligonucleotides and plasmids based on gelatin nanoparticles, *NewDrugs* 1, 14-17 (2003a).
- Coester, C., Nanopartikel aus Gelatine und Humanem Serum Albumin, *Bioforum* 11, 714-716 (2003b).
- Coester, C.J., Langer, K., van Briesen, H., Kreuter, J., Gelatin nanoparticles by two step desolvation--a new preparation method, surface modifications and cell uptake, *J. Microencapsul.* 17, 187-193 (2000b).
- Cook, B.C., Retzinger, G.S., Elution of fibrinogen and other plasma protein from unmodified and from lecithin-coated polystyrene-divinylbenzene beads, *J. Colloid Interf. Sci.* 153, 1-12 (1992).
- Cordon-Cardo, C., O'Brien, J.P., Casals, D., Rittman-Grauer, L., Biedler, J.L., Melamed, M.R., Bertino, J.R., Multidrug-resistance gene (P-glycoprotein) is expressed by endothelial cells at blood-brain barrier sites, *Proc. Natl. Acad. Sci. U S A* 86, 695-698 (1989).
- Cornaire, G., Woodley, J., Hermann, P., Cloarec, A., Arellano, C., Houin, G., Impact of excipients on the absorption of P-glycoprotein substrates in vitro and in vivo, *Int. J. Pharm.* 278, 119-131 (2004).
- Cornelius, R.M., Brash, J.L., Adsorption from plasma and buffer of single- and two-chain high molecular weight kininogen to glass and sulfonated polyurethane surfaces, *Biomaterials* 20, 341-50 (1999).
- Cornelius, R.M., Archambault, J., Brash, J.L., Identification of apolipoprotein A-I as a major adsorbate on biomaterial surfaces after blood or plasma contact, *Biomaterials* 23, 3583-3587 (2002a).
- Cornelius, R.M., Archambault, J.G., Berry, L., Chan, A.K., Brash, J.L., Adsorption of proteins from infant and adult plasma to biomaterial surfaces, *J. Biomed. Mater. Res.* 60, 622-632 (2002b).
- Cornelius, R.M., Sanchez, J., Olsson, P., Brash, J.L., Interactions of antithrombin and proteins in the plasma contact activation system with immobilized functional heparin, *J. Biomed. Mater. Res. A* 67, 475-483 (2003).
- Cullen, S.C., Gross, E.G., The anesthetic properties of xenon in animals and human beings, with additional observations on krypton, *Science* 113, 580-582 (1951).
- Davis, S.S., Pharmaceutical aspects of i. v. fat emulsions, *J. Hosp. Pharm.* 32, 149-170 (1974).
- Davis, S.S., Colloids as drug-delivery systems, *Pharmaceutical Technology* 5, 71-88 (1981).

- Davis, S.S., Illum, L., Moghimi, S.M., Davies, M.C., Porter, C.J.Ü., Muir, I.S., Brindley, A.M., Christy, N.M., Norman, M.E., Williams, P., Dunn, S.E., Microspheres for targeting drugs to specific body sites, *J. Contr. Rel.* 24, 157-163 (1993).
- Degussa, Firmenschrift: Tego® Care 450, (2003).
- Dehouck, B., Dehouck, M.P., Fruchart, J.C., Cecchelli, R., Upregulation of the low density lipoprotein receptor at the blood-brain barrier: intercommunications between brain capillary endothelial cells and astrocytes, *J. Cell Biol.* 126, 465-473 (1994).
- Dehouck, B., Fenart, L., Dehouck, M.P., Pierce, A., Torpier, G., Cecchelli, R., A new function for the LDL receptor: transcytosis of LDL across the blood-brain barrier, *J. Cell. Biol.* 138, 877-889 (1997).
- Diederichs, J.E., Plasma protein adsorption patterns on liposomes: establishment of analytical procedure, *Electrophoresis* 17, 607-611 (1996).
- Dingler, A., Feste Lipid-Nanopartikel als kolloidale Wirkstoffträgersysteme zur dermalen Applikation, Dissertation, Freie Universität Berlin (1998).
- Dintaman, J.M., Silverman, J.A., Inhibition of P-glycoprotein by D-alpha-tocopheryl polyethylene glycol 1000 succinate (TPGS), *Pharm. Res.* 16, 1550-1556 (1999).
- Doenicke, A.W., Roizen, M.F., Hoernecke, R., Lorenz, W., Ostwald, P., Solvent for etomidate may cause pain and adverse effects, *Br J Anaesth* 83, 464-6 (1999).
- Doolittle, N.D., Abrey, L.E., Ferrari, N., Hall, W.A., Laws, E.R., McLendon, R.E., Muldoon, L.L., Peereboom, D., Peterson, D.R., Reynolds, C.P., Senter, P., Neuwelt, E.A., Targeted delivery in primary and metastatic brain tumors: summary report of the seventh annual meeting of the Blood-Brain Barrier Disruption Consortium, *Clin. Cancer Res.* 8, 1702-1709 (2002).
- Drewes, L.R., What is the blood-brain barrier? A molecular perspective. Cerebral vascular biology, *Adv. Exp. Med. Biol.* 474, 111-122 (1999).
- Drin, G., Rousselle, C., Scherrmann, J.M., Rees, A.R., Temsamani, J., Peptide delivery to the brain via adsorptive-mediated endocytosis: advances with SynB vectors, *AAPS PharmSci.* 4, E26 (2002).
- Drion, N., Lemaire, M., Lefauconnier, J.M., Scherrmann, J.M., Role of P-glycoprotein in the blood-brain transport of colchicine and vinblastine, *J. Neurochem.* 67, 1688-93 (1996).
- Driscoll, D.F., Giampietro, K., Wichelhaus, D.P., Peterss, H., Nehne, L., Niemann, W., Bistrian, B.R., Physicochemical stability assessments of lipid emulsions of varying oil composition, *Clin. Nutr.* 20, 151-157 (2001).

- Dunay, I.R., Heimesaat, M.M., Bushrab, F.N., Müller, R.H., Stocker, H., Arasteh, K., Kurowski, M., Fitzner, R., Borner, K., Liesenfeld, O., Atovaquone maintenance therapy prevents reactivation of toxoplasmic encephalitis in a murine model of reactivated toxoplasmosis, *Antimicrob. Agents Chemother.* 48, 4848-4854 (2004).
- Dutta, S., Ebling, W.F., Emulsion formulation reduces propofol's dose requirements and enhances safety, *Anesthesiology* 87, 1394-1405 (1997).
- Ehrlich, P., *Collected Studies on Immunity*, John Wiley & Sons, New York, (1906).
- FDA, Food additive database, Internet: <http://vm.cfsan.fda.gov/~dms/eafus.html> (2005).
- Fenart, L., Buee-Scherrer, V., Descamps, L., Duhem, C., Poullain, M.G., Cecchelli, R., Dehouck, M.P., Inhibition of P-glycoprotein: rapid assessment of its implication in blood-brain barrier integrity and drug transport to the brain by an in vitro model of the blood-brain barrier, *Pharm. Res.* 15, 993-1000 (1998).
- Fiedler, H.P., *Lexikon der Hilfsstoffe für Pharmazie, Kosmetik und angrenzender Gebiete*, 4. Auflage, Edition Cantor Verlag, Aulendorf, (1996).
- Fishman, J.B., Rubin, J.B., Handrahan, J.V., Connor, J.R., Fine, R.E., Receptor-mediated transcytosis of transferrin across the blood-brain barrier, *J. Neurosci. Res.* 18, 299-304 (1987).
- Forth, W., Henschler, D., Rummel, W., *Allgemeine und spezielle Pharmakologie und Toxikologie*, Urban & Fischer Verlag, München, (2001).
- Franks, N.P., Dickinson, R., de Sousa, S.L., Hall, A.C., Lieb, W.R., How does xenon produce anaesthesia?, *Nature* 396, 324 (1998).
- Freitas, C., Müller, R.H., Stability determination of solid lipid nanoparticles (SLN) in aqueous dispersion after addition of electrolyte, *J. Microencapsul.* 16, 59-71 (1999).
- Friden, P.M., Walus, L.R., Musso, G.F., Taylor, M.A., Malfroy, B., Starzyk, R.M., Anti-transferrin receptor antibody and antibody-drug conjugates cross the blood-brain barrier, *Proc. Natl. Acad. Sci. U S A* 88, 4771-4775 (1991).
- Friese, A., Seiler, E., Quack, G., Lorenz, B., Kreuter, J., Increase of the duration of the anticonvulsive activity of a novel NMDA receptor antagonist using poly(butylcyanoacrylate) nanoparticles as a parenteral controlled release system, *Eur. J. Pharm. Biopharm.* 49, 103-109 (2000).
- Fukuta, M., Okada, H., Iinuma, S., Yanai, S., Toguchi, H., Insulin fragments as a carrier for peptide delivery across the blood-brain barrier, *Pharm. Res.* 11, 1681-1688 (1994).

- Fundaro, A., Cavalli, R., Bargoni, A., Vighetto, D., Zara, G.P., Gasco, M.R., Non-stealth and stealth solid lipid nanoparticles (SLN) carrying doxorubicin: pharmacokinetics and tissue distribution after i.v. administration to rats, *Pharmacol Res* 42, 337-43 (2000).
- Gäbler, R., Schmidt, J., Xenon - Eine neue Alternative?, www.tu-dresden.de/medkai/989913.pdf, Klinik und Poliklinik für Anästhesiologie und Intensivtherapie, TU Dresden (1999).
- Galazka, V., Khan, A.M., Davies, M.C., Müller, R.H., Effect of gamma-irradiation and plasma etching on the surface of polymers, *Archiv der Pharmazie* 320, 980 (1987).
- Gasco, M.R., Method for producing solid lipid microspheres having a narrow size distribution, U. S. Patent 5,250,236 (1993)
- Georgieff, M., Mückter, H., Fröba, G., Bäder, S., Liebl, B., Marx, T., Xenon statt Lachgas?, *Deutsches Ärzteblatt* 94, A2202-2205, B1792-1795, C1658-1661 (1997).
- Georgieff, M., Froeba, G., Müller, R.H., Buttle, I., Wachter, U., Intravenous xenon anaesthesia. *6th EuroSIVA Meeting on Intravenous Anaesthesia* (Glasgow, 2003).
- Gessner, A., Waicz, R., Lieske, A., Paulke, B.R., Mäder, K., Müller, R.H., Nanoparticles with decreasing surface hydrophobicities: influence on plasma protein adsorption, *Int. J. Pharm.* 196, 245-249 (2000).
- Gessner, A., Untersuchungen zur Proteinadsorption auf kolloidalen Modellpartikeln und Arzneistoffträgern zur parenteralen Anwendung, Dissertation, Freie Universität Berlin (2001).
- Gessner, A., Olbrich, C., Schröder, W., Kayser, O., Müller, R.H., The role of plasma proteins in brain targeting: species dependent protein adsorption patterns on brain-specific lipid drug conjugate (LDC) nanoparticles, *Int. J. Pharm.* 214, 87-91 (2001).
- Gessner, A., Lieske, A., Paulke, B.R., Müller, R.H., Influence of surface charge density on protein adsorption on polymeric nanoparticles: analysis by two-dimensional electrophoresis, *Eur. J. Pharm. Biopharm.* 54, 165-170 (2002).
- Gessner, A., Lieske, A., Paulke, B.R., Müller, R.H., Functional groups on polystyrene model nanoparticles: influence on protein adsorption, *J. Biomed. Mater. Res.* 65A, 319-26 (2003).
- Gessner, A., Paulke, B.-R., Göppert, T.M., Müller, R.H., Protein rejecting properties of PEG-grafted nanoparticles: Influence of PEG-chain length and surface density evaluated by two-dimensional electrophoresis and bicinchoninic acid (BCA)-protein assay, *Die Pharmazie* (im Druck 2005).

- Geyer, R.P., Watkin, D.M., Matthews, L.W., Stare, F.J., Parenteral nutrition. XI. Studies with stable and unstable fat emulsions administered intravenously, Proc. Soc. Exp. Biol. Med. 77, 872-876 (1951).
- Goldstein, G.W., Betz, A.L., Recent advances in understanding brain capillary function, Ann. Neurol. 14, 389-395 (1983).
- Göppert, T.M., Müller, R.H., Alternative sample preparation prior to two-dimensional electrophoresis protein analysis on solid lipid nanoparticles, Electrophoresis 25, 134-140 (2004a).
- Göppert, T.M., Müller, R.H., Influence of different lipids on plasma protein patterns of solid lipid nanoparticles. *AAPS Annual Meeting, Baltimore* (2004b).
- Göppert, T.M., Müller, R.H., Polysorbate-stabilized solid lipid nanoparticles as colloidal carriers for intravenous targeting of drugs to the brain: Comparison of plasma protein adsorption patterns, J. Drug Target. 13 (3), 179-187 (2005a).
- Göppert, T.M., Müller, R.H., Adsorption kinetics of plasma proteins on solid lipid nanoparticles for drug targeting, Int. J. Pharm. accepted(2005b).
- Göppert, T.M., Müller, R.H., Protein adsorption patterns on poloxamer- and poloxamine-stabilized solid lipid nanoparticles (SLN), Eur. J. Pharm. Biopharm. 60, 361-372 (2005c).
- Göppert, T.M., Souto, E., Müller, R.H., Comparison of plasma protein adsorption patterns on solid lipid nanoparticles (SLN) for intravenous drug targeting dependent on their age. *CRS-Meeting, Miami* (2005).
- Görg, A., Postel, W., Gunther, S., The current state of two-dimensional electrophoresis with immobilized pH gradients, Electrophoresis 9, 531-546 (1988a).
- Görg, A., Postel, W., Gunther, S., Horizontal two-dimensional electrophoresis with immobilized pH gradients using PhastSystem, Electrophoresis 9, 57-59 (1988b).
- Görg, A., Boguth, G., Obermaier, C., Scheibe, B., Weiss, W., Booklet of the "Hands-on Course. ICES '95 Meeting (Paris, 1995).
- Görg, A., Boguth, G., Obermaier, C., Weiss, W., Two-dimensional electrophoresis of proteins in an immobilized pH 4-12 gradient, Electrophoresis 19, 1516-1519 (1998).
- Goto, T., Nakata, Y., Morita, S., Will xenon be a stranger or a friend?: the cost, benefit, and future of xenon anesthesia, Anesthesiology 98, 1-2 (2003).
- Graham, D.J., Phillips, M.C., Proteins at liquid interfaces, I, Kinetics of adsorption and surface denaturation, J. Coll. Interf. Sci. 70, 403 (1979).

- Gref, R., Minamitake, Y., Peracchia, M.T., Trubetskoy, V., Torchilin, V., Langer, R., Biodegradable long-circulating polymeric nanospheres, *Science* 263, 1600-1603 (1994).
- Gref, R., Lück, M., Quellec, P., Marchand, M., Dellacherie, E., Harnisch, S., Blunk, T., Müller, R.H., 'Stealth' corona-core nanoparticles surface modified by polyethylene glycol (PEG): influences of the corona (PEG chain length and surface density) and of the core composition on phagocytic uptake and plasma protein adsorption, *Colloids Surf. B Biointerfaces* 18, 301-313 (2000).
- Gulyaev, A.E., Gelperina, S.E., Skidan, I.N., Antropov, A.S., Kivman, G.Y., Kreuter, J., Significant transport of doxorubicin into the brain with polysorbate 80-coated nanoparticles, *Pharm. Res.* 16, 1564-1569 (1999).
- Habgood, M.D., Begley, D.J., Abbott, N.J., Determinants of passive drug entry into the central nervous system, *Cell Mol. Neurobiol.* 20, 231-253 (2000).
- Haeberli, A., Human Protein Data, VCH Verlagsgesellschaft, Weinheim, (1995).
- Hailer, S., Wolfram, G., Influence of artificial fat emulsions on the composition of serum lipoproteins in humans, *Am. J. Clin. Nutr.* 43, 225-233 (1986).
- Hallberg, D., Holm, I., Obel, A.L., Schuberth, O., Wretlind, A., Fat emulsion for complete intravenous nutrition, *Postgrad. Med.* 42, A149-52 (1967).
- Hanash, S.M., Strahler, J.R., Somerlot, L., Postel, W., Görg, A., Two-dimensional electrophoresis with immobilized pH gradients in the first dimension: protein focussing as a function of time, *Electrophoresis* 8, 229-234 (1987).
- Harnisch, S., Vergleichende Untersuchungen zur Plasmaproteinadsorption auf O/W-Emulsionen und Polymer-Partikeln zur parenteralen Anwendung, Dissertation, Freie Universität Berlin (1998).
- Harnisch, S., Müller, R.H., Plasma protein adsorption patterns on emulsions for parenteral administration: establishment of a protocol for two-dimensional polyacrylamide electrophoresis, *Electrophoresis* 19, 349-354 (1998).
- Harnisch, S., Müller, R.H., Adsorption kinetics of plasma proteins on oil-in-water emulsions for parenteral nutrition, *Eur. J. Pharm. Biopharm.* 49, 41-46 (2000).
- Haynes, P., Miller, I., Aebersold, R., Gemeiner, M., Eberini, I., Lovati, M.R., Manzoni, C., Vignati, M., Gianazza, E., Proteins of rat serum: I. Establishing a reference two-dimensional electrophoresis map by immunodetection and microbore high performance liquid chromatography-electrospray mass spectrometry, *Electrophoresis* 19, 1484-1492 (1998).
- Hedemann, H., Lück, M., Blunk, T., Frokjaer, S., Müller, R.H., Fat emulsions based on structured lipids (1,3-specific triglycerides) : An investigation of the in vitro interaction with plasma proteins, *Clin. Nutr.* 15, 175-178 (1996).

- Hildebrand, G.E., Tack, J., Harnisch, S., Method for producing morphologically uniform micro and nanoparticles using micromixers, (2000)
- Hochstrasser, D.F., Harrington, M.G., Hochstrasser, A.C., Miller, M.J., Merril, C.R., Methods for increasing the resolution of two-dimensional protein electrophoresis, *Anal. Biochem.* 173, 424-435 (1988).
- Hochstrasser, D.F., Merril, C.R., 'Catalysts' for polyacrylamide gel polymerization and detection of proteins by silver staining, *Appl. Theor. Electrophor.* 1, 35-40 (1988).
- Hochstrasser, D.F., Tissot, J.D., in: Chrambach, A., Dunn, M. J., Radola, B. J. (Hrsg.), *Advances in Electrophoresis* Vol. 4, VCH Verlagsgesellschaft, Weinheim, 270-361 (1991)
- Horbett, T.A., Brash, J.L., Proteins at interfaces: Current issues and future prospects, in: Brash, J. L., Horbett, T. A. (Hrsg.), *Proteins at Interfaces*, Am. Chem. Soc., Washington D. C., USA, 1-33 (1987)
- Hsu, M.J., Juliano, R.L., Interactions of liposomes with the reticuloendothelial system. II: Nonspecific and receptor-mediated uptake of liposomes by mouse peritoneal macrophages, *Biochim Biophys Acta* 720, 411-419 (1982).
- Huai-Yun, H., Secret, D.T., Mark, K.S., Carney, D., Brandquist, C., Elmquist, W.F., Miller, D.W., Expression of multidrug resistance-associated protein (MRP) in brain microvessel endothelial cells, *Biochem. Biophys. Res. Commun.* 243, 816-820 (1998).
- Hunt, J.E., Simpson, R.J., Krilis, S.A., Identification of a region of beta 2-glycoprotein I critical for lipid binding and anti-cardiolipin antibody cofactor activity, *Proc Natl Acad Sci U S A* 90, 2141-5 (1993).
- Illum, L., Davis, S.S., Targeting of colloidal particles to the bone marrow, *Life Sci.* 40, 1553-1560 (1987).
- Illum, L., Davis, S.S., Müller, R.H., Mak, E., West, P., The organ distribution and circulation time of intravenously injected colloidal carriers sterically stabilized with a block copolymer--poloxamine 908, *Life Sci.* 40, 367-374 (1987).
- Innerarity, T.L., Pitas, R.E., Mahley, R.W., Binding of arginine-rich (E) apoprotein after recombination with phospholipid vesicles to the low density lipoprotein receptors of fibroblasts, *J. Biol. Chem.* 254, 4186-4190 (1979).
- Innerarity, T.L., Friedlander, E.J., Rall, S.C., Jr., Weisgraber, K.H., Mahley, R.W., The receptor-binding domain of human apolipoprotein E. Binding of apolipoprotein E fragments, *J. Biol. Chem.* 258, 12341-12347 (1983).
- Jahangir, R., McCloskey, C.B., Mc Clung, W.G., Labow, R.S., Brash, J.L., Santerre, J.P., The influence of protein adsorption and surface modifying macromolecules on the hydrolytic degradation of a poly(ether-urethane) by cholesterol esterase, *Biomaterials* 24, 121-130 (2003).

- Janzer, R.C., Raff, M.C., Astrocytes induce blood-brain barrier properties in endothelial cells, *Nature* 325, 253-257 (1987).
- Jenning, V., Gysler, A., Schäfer-Korting, M., Gohla, S.H., Vitamin A loaded solid lipid nanoparticles for topical use: occlusive properties and drug targeting to the upper skin, *Eur. J. Pharm. Biopharm.* 49, 211-218 (2000).
- Jenning, V., Gohla, S.H., Encapsulation of retinoids in solid lipid nanoparticles (SLN), *J. Microencapsul.* 18, 149-158 (2001).
- Jeon, S.L., Andrade, J.D., Protein-surface interactions in the presence of polyethylene oxide, II. Effect of protein size, *J. Coll. Interf. Sci.* 142, 159-166 (1991).
- Jeon, S.L., Lee, J.H., Andrade, J.D., De Gennes, P.G., Protein-surface interactions in the presence of polyethylene oxide, I. Simplified theory, *J. Coll. Interf. Sci.* 142, 149-158 (1991).
- Ji, Z.S., Fazio, S., Lee, Y.L., Mahley, R.W., Secretion-capture role for apolipoprotein E in remnant lipoprotein metabolism involving cell surface heparan sulfate proteoglycans, *J. Biol. Chem.* 269, 2764-2772 (1994).
- Juliano, R.L., Factors affecting the clearance kinetics and tissue distribution of liposomes, microspheres and emulsions, *Adv. Drug Deliv. Rev.* 2, 31-54 (1988).
- Jungblut, P., Wittmann-Liebold, B., Protein analysis on a genomic scale, *J. Biotechnol.* 41, 111-20 (1995).
- Kamboh, M.I., Mehdi, H., Genetics of apolipoprotein H (beta2-glycoprotein I) and anionic phospholipid binding, *Lupus* 7 (2), 10-13 (1998).
- Kartner, N., Evernden-Porelle, D., Bradley, G., Ling, V., Detection of P-glycoprotein in multidrug-resistant cell lines by monoclonal antibodies, *Nature* 316, 820-823 (1985).
- Kazatchkine, M.D., Carreno, M.P., Activation of the complement system at the interface between blood and artificial surfaces, *Biomaterials* 9, 30-35 (1988).
- Klang, S.H., Parnas, M., Benita, S., Emulsions as drug carriers - possibilities, limitations and future perspectives, in: Müller, R. H., Benita, S., Böhm, B. (Hrsg.), *Emulsions and Nanosuspensions for the Formulation of Poorly Soluble Drugs*, medpharm Scientific Publishers Stuttgart, (1998)
- Klimke, R., Silbernagl, S., *Lehrbuch der Physiologie*, Georg Thieme Verlag, Stuttgart, (2003).
- Klose, J., Proteinmapping by combined isoelectric focussing and electrophoresis of mouse tissues - A novel approach to testing for induced point mutation in mammals, *Hummangenetik* 26, 211-234 (1975).

- Klose, J., Kobalz, U., Two-dimensional electrophoresis of proteins: An update protocol and implications for a functional analysis of the genome, *Electrophoresis* 16, 1034-1059 (1995).
- Kostner, K., Beyond LDL-Cholesterol: New Treatments Raising HDL-Cholesterol or Enhancing Reverse Cholesterol Transport, *Journal für Kardiologie* 9 (7-8), 328-331 (2002).
- Krause, K.P., Müller, R.H., Production and characterisation of highly concentrated nanosuspensions by high pressure homogenisation, *Int. J. Pharm.* 214, 21-24 (2001).
- Kreuter, J., Evaluation of nanoparticles as drug-delivery systems. III: materials, stability, toxicity, possibilities of targeting, and use, *Pharm. Acta Helv.* 58, 242-250 (1983).
- Kreuter, J., Wilson, C.G., Fry, J.R., Paterson, P., Ratcliffe, J.H., Toxicity and association of polycyanoacrylate nanoparticles with hepatocytes, *J. Microencapsul.* 1, 253-257 (1984).
- Kreuter, J., Nanoparticles, in: Swarbrick, J., Boylan, J. C. (Hrsg.), *Encyclopedia of Pharmaceutical Technology*, Vol. 10, Marcel Dekker, New York, S. 165-190 (1994)
- Kreuter, J., Alyautdin, R.N., Kharkevich, D.A., Ivanov, A.A., Passage of peptides through the blood-brain barrier with colloidal polymer particles (nanoparticles), *Brain Res.* 674, 171-174 (1995).
- Kreuter, J., Petrov, V.E., Kharkevich, D.A., Alyautdin, R.N., Influence of the type of surfactant on the analgesic effects induced by the peptide dalargin after its delivery across the blood-brain barrier using surfactant-coated nanoparticles, *J. Contr. Rel.* 49, 81-87 (1997).
- Kreuter, J., Nanoparticulate systems for brain delivery of drugs, *Adv. Drug Deliv. Rev.* 47, 65-81 (2001).
- Kreuter, J., Shamenkov, D., Petrov, V., Ramge, P., Cychutek, K., Koch-Brandt, C., Alyautdin, R.N., Apolipoprotein-mediated transport of nanoparticle-bound drugs across the blood-brain barrier, *J. Drug Target.* 10, 317-325 (2002).
- Kreuter, J., Ramge, P., Petrov, V., Hamm, S., Gelperina, S.E., Engelhardt, B., Alyautdin, R.N., von Briesen, H., Begley, D.J., Direct evidence that polysorbate-80-coated poly(butylcyanoacrylate) nanoparticles deliver drugs to the CNS via specific mechanisms requiring prior binding of drug to the nanoparticles, *Pharm. Res.* 20, 409-416 (2003).
- Kumagai, A.K., Eisenberg, J.B., Pardridge, W.M., Absorptive-mediated endocytosis of cationized albumin and a beta-endorphin-cationized albumin chimeric peptide by isolated brain capillaries. Model system of blood-brain barrier transport, *J. Biol. Chem.* 262, 15214-15219 (1987).

- Kurrat, R., Wälivaara, B., Marti, A., Textor, M., Tengvall, P., Ramsden, J.J., Spencer, N.D., Plasma protein adsorption on titanium: Comparative in situ studies using optical waveguide lightmode spectroscopy and ellipsometrie, *Colloids Surf. B Biointerfaces* 11, 187-201 (1998).
- Kusuhara, H., Sugiyama, Y., Role of transporters in the tissue-selective distribution and elimination of drugs: transporters in the liver, small intestine, brain and kidney, *J. Control. Release* 78, 43-54 (2002).
- Lachmann, B., Armbruster, S., Schairer, W., Landstra, M., Trouwborst, A., Van Daal, G.J., Kusuma, A., Erdmann, W., Safety and efficacy of xenon in routine use as an inhalational anaesthetic, *Lancet* 335, 1413-1415 (1990).
- Lee, J.H., Andrade, J.D., Surface properties of aqueous PEO/PPO block copolymer surfactants, in: Andrade, J. D. (Hrsg.), *Polymer Surface Dynamics*, Plenum Press, New York, 119-136 (1988)
- Lee, J.H., Kopecek, J., Andrade, J.D., Protein-resistant surfaces prepared by PEO-containing block copolymer surfactants, *J. Biomed. Mater. Res.* 23, 351-368 (1989).
- Lenaerts, V., Couvreur, P., Christiaens-Leyh, D., Joiris, E., Roland, M., Rollman, B., Speiser, P., Degradation of poly (isobutyl cyanoacrylate) nanoparticles, *Biomaterials* 5, 65-68 (1984).
- Leroux, J.C., De Jaeghere, F., Anner, B., Doelker, E., Gurny, R., An investigation on the role of plasma and serum opsonins on the internalization of biodegradable poly(D,L-lactic acid) nanoparticles by human monocytes, *Life Sci* 57, 695-703 (1995).
- Leyke, S., Göppert, T.M., Köhler-Sokolowska, W., Paulke, B.-R., Müller, R.H., Presber, W., Improved effect of Nanoparticles on cells infected by Toxoplasma gondii after coating with serumproteins, (2005).
- Leyke, S., Dissertation in Vorbereitung, Freie Universität Berlin (2006).
- Lherm, C., Müller, R.H., Puisieux, F., Couvreur, P., Alkylcyanoacrylate drug carriers: II. Cytotoxicity of cyanoacrylate nanoparticles with different alkyl chain length, *Int. J. Pharm.* 84, 13-22 (1992).
- Li, J.T., Caldwell, K.D., Plasma protein interaction with PluronicTM-treated colloids, *Colloids Surf. B Biointerfaces* 7, 19-22 (1996).
- Lind, K., Kresse, M., Müller, R.H., Comparison of protein adsorption patterns onto differently charged hydrophilic superparamagnetic iron oxide particles obtained in vitro and ex vivo, *Electrophoresis* 22, 3514-3521 (2001a).
- Lind, K., Kresse, M., Müller, R.H., Evaluation of desorption of proteins adsorbed to hydrophilic surfaces by two-dimensional electrophoresis, *Proteomics* 1, 1059-1066 (2001b).

- Lind, K., MRT-Kontrastmittel: Proteinadsorption und In-Vivo-Studien von stärkederivatstabilisierten superparamagnetischen Eisenoxid-Partikeln (SPIO), Dissertation, Freie Universität Berlin (2002).
- Lind, K., Kresse, M., Debus, N.P., Müller, R.H., A novel formulation for superparamagnetic iron oxide (SPIO) particles enhancing MR lymphography: comparison of physicochemical properties and the in vivo behaviour, *J. Drug Target.* 10, 221-230 (2002).
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., Randall, R.J., Protein measurement with the Folin phenol reagent, *J. Biol. Chem.* 193, 265-275 (1951).
- Lucarelli, M., Gennarelli, M., Cardelli, P., Novelli, G., Scarpa, S., Dallapiccola, B., Strom, R., Expression of receptors for native and chemically modified low-density lipoproteins in brain microvessels, *FEBS Lett.* 401, 53-58 (1997).
- Lück, M., Plasmaproteinadsorption als ein möglicher Schlüsselfaktor für eine kontrollierte Arzneistoffapplikation mit partikulären Trägern, Dissertation, Freie Universität Berlin (1997).
- Lück, M., Schröder, W., Harnisch, S., Thode, K., Blunk, T., Paulke, B.R., Kresse, M., Müller, R.H., Identification of plasma proteins facilitated by enrichment on particulate surfaces: analysis by two-dimensional electrophoresis and N-terminal microsequencing, *Electrophoresis* 18, 2961-2967 (1997).
- Lück, M., Paulke, B.R., Schröder, W., Blunk, T., Müller, R.H., Analysis of plasma protein adsorption on polymeric nanoparticles with different surface characteristics, *J. Biomed. Mater. Res.* 39, 478-485 (1998a).
- Lück, M., Pistel, K.F., Li, Y.X., Blunk, T., Müller, R.H., Kissel, T., Plasma protein adsorption on biodegradable microspheres consisting of poly(D,L-lactide-co-glycolide), poly(L-lactide) or ABA triblock copolymers containing poly(oxyethylene). Influence of production method and polymer composition, *J. Control. Release* 55, 107-120 (1998b).
- Lück, M., Schröder, W., Paulke, B.R., Blunk, T., Müller, R.H., Complement activation by model drug carriers for intravenous application: determination by two-dimensional electrophoresis, *Biomaterials* 20, 2063-2068 (1999).
- Lynch, C., 3rd, Baum, J., Tenbrinck, R., Xenon anesthesia, *Anesthesiology* 92, 865-868 (2000).
- Mahley, R.W., Apolipoprotein E: cholesterol transport protein with expanding role in cell biology, *Science* 240, 622-630 (1988).
- Marty, J.J., Oppenheim, R.C., Speiser, P., Nanoparticles--a new colloidal drug delivery system, *Pharm. Acta Helv.* 53, 17-23 (1978).

- Mehnert, W., zur Mühlen, A., Dingler, A., Weyhers, H., Müller, R.H., Solid lipid nanoparticles (SLN) - ein neuartiger Wirkstoff-Carrier für Kosmetika und Pharmazeutika, II. Wirkstoffinkorporation, Freisetzung und Sterilisierbarkeit, *Pharm. Ind.* 59, 511-514 (1997).
- Mehnert, W., Mäder, K., Solid lipid nanoparticles: production, characterization and applications, *Adv. Drug Del. Rev.* 47, 165-196 (2001).
- Meresse, S., Delbart, C., Fruchart, J.C., Cecchelli, R., Low-density lipoprotein receptor on endothelium of brain capillaries, *J. Neurochem.* 53, 340-345 (1989).
- Merril, C.R., Golman, D., van Keuren, M.L., Simplified silver protein detection and image enhancement methods in polyacrylamide gels, *Electrophoresis* 3, 17-23 (1982).
- Messer, Stoffdatenblatt Xenon, Messer Griesheim GmbH, Krefeld (2000).
- Miller, D.W., Batrakova, E.V., Waltner, T.O., Alakhov, V., Kabanov, A.V., Interactions of pluronic block copolymers with brain microvessel endothelial cells: evidence of two potential pathways for drug absorption, *Bioconjug. Chem.* 8, 649-657 (1997).
- Miller, G., Drug targeting. Breaking down barriers, *Science* 297, 1116-1118 (2002).
- Miller, M.J., Computer-assisted analysis of two-dimensional gel electrophoretograms, in: Chrambach, A., Dunn, M. J., Radola, B. J. (Hrsg.), *Advances in Electrophoresis*, Vol. 3, VCH Verlagsgesellschaft, Weinheim/New York, (1989)
- Moghimi, S.M., Muir, I.S., Illum, L., Davis, S.S., Kolb-Bachofen, V., Coating particles with a block co-polymer (poloxamine-908) suppresses opsonization but permits the activity of dysopsonins in the serum, *Biochim. Biophys. Acta* 1179, 157-165 (1993).
- Moghimi, S.M., Mechanisms regulating body distribution of nanospheres conditioned with pluronic and tetronic block co-polymers, *Adv. Drug Deliv. Rev.* 16, 183-193 (1995a).
- Moghimi, S.M., Exploiting bone marrow microvascular structure for drug delivery and future therapies, *Adv. Drug Del. Rev.* 17, 61-73 (1995b).
- Moghimi, S.M., Re-establishing the long circulatory behaviour of poloxamine-coated particles after repeated intravenous administration: applications in cancer drug delivery and imaging, *Biochim. Biophys. Acta* 1472, 399-403 (1999).
- Möschwitzer, J., Achleitner, G., Pomper, H., Müller, R.H., Development of an intravenously injectable chemically stable aqueous omeprazole formulation using nanosuspension technology, *Eur. J. Pharm. Biopharm.* 58, 615-9 (2004).

- Muckerheide, A., Apple, R.J., Pesce, A.J., Michael, J.G., Cationization of protein antigens. I. Alteration of immunogenic properties, *J. Immunol.* 138, 833-837 (1987).
- Müller, B.W., Müller, R.H., Particle Size Analysis of Latex Suspensions and Microemulsions by Photon Correlation Spectroscopy, *J. Pharm. Sci.* 73, 1915-1918 (1984).
- Müller, R.H., Heinemann, S., Surface Modelling of Microparticles as Parenteral Systems with High Tissue Affinity, in: Gurny, R., Junginger, H. E. (Hrsg.), *Bioadhesion - Possibilities and Future Trends*, Wissenschaftliche Verlagsgesellschaft, Stuttgart, Germany, 202-214 (1989)
- Müller, R.H., Colloidal Carriers for Controlled Drug Delivery and Targeting - Modification, Characterization and in vivo Distribution, Wissenschaftliche Verlagsgesellschaft Stuttgart, CRC Press Boca Raton, (1991).
- Müller, R.H., Mehnert, W., Lucks, J.S., Schwarz, C., zur Mühlen, A., Weyhers, H., Freitas, C., Rühl, D., Solid Lipid Nanoparticles (SLN) - An Alternative Colloidal Carrier System for Controlled Drug Delivery, *Eur. J. Pharm. Biopharm.* 41, 62-69 (1995).
- Müller, R.H., Zetapotential und Partikelladung in der Laborpraxis, APV Paperback, Wissenschaftliche Verlagsgesellschaft Stuttgart, (1996).
- Müller, R.H., Lucks, J.S., Arzneistoffträger aus festen Lipidteilchen, Feste Lipidnanosphären (SLN), European Patent EP 0605497 (1996)
- Müller, R.H., Maassen, S., Weyhers, H., Mehnert, W., Phagocytic uptake and cytotoxicity of solid lipid nanoparticles (SLN) sterically stabilized with poloxamine 908 and poloxamer 407, *J. Drug Target.* 4, 161-170 (1996a).
- Müller, R.H., Rühl, D., Runge, S.A., Biodegradation of solid lipid nanoparticles as a function of lipase incubation time, *Int. J. Pharm.* 144, 115-121 (1996b).
- Müller, R.H., Schuhmann, R., Teilchengrößenmessung in der Laborpraxis, APV Paperback, Wissenschaftliche Verlagsgesellschaft Stuttgart, (1996).
- Müller, R.H., Surface Hydrophobicity - Determination by Rose Bengal (RB) Adsorption Methods, in: Müller, R. H., Mehnert, W. (Hrsg.), *Particle and Surface Characterisation Methods*, Wissenschaftliche Verlagsgesellschaft Stuttgart, 215-228 (1997)
- Müller, R.H., Rühl, D., Runge, S., Schulze-Forster, K., Mehnert, W., Cytotoxicity of solid lipid nanoparticles as a function of the lipid matrix and the surfactant, *Pharm. Res.* 14, 458-462 (1997a).
- Müller, R.H., Weyers, H., zur Mühlen, A., Dingler, A., Mehnert, W., Solid lipid nanoparticles - ein neuartiger Wirkstoff-Carrier für Kosmetika und Pharmazeutika, *Pharm. Ind.* 59, 423-427 (1997b).

- Müller, R.H., Feste Lipidnanopartikel (SLN), in: Müller, R. H., Hildebrand, G. E. (Hrsg.), Pharmazeutische Technologie: Moderne Arzneiformen, Wissenschaftliche Verlagsgesellschaft, Stuttgart, (1998)
- Müller, R.H., Olbrich, C., Solid Lipid Nanoparticles: Phagocytic Uptake, in vitro Cytotoxicity and in vitro Biodegradation, 1st Communication, Pharm. Ind. 61, 462-467 (1999).
- Müller, R.H., Mäder, K., Gohla, S., Solid lipid nanoparticles (SLN) for controlled drug delivery - a review of the state of the art, Eur. J. Pharm. Biopharm. 50, 161-77 (2000).
- Müller, R.H., Lück, M., Kreuter, J., Medicament excipient particles for tissue specific application of a medicament, PCT-application PCT/EP98/064299 (P53601) (2001)
- Müller, R.H., Schmidt, S., PathFinder technology for the delivery of drugs to the brain, NewDrugs 2, 38-42 (2002).
- Müller, R.H., Wissing, S.A., Lipopearls for topical delivery of active compounds and controlled release, in: Rathbone, M. J., Hadgraft, J., Roberts, M. S. (Hrsg.), Modified Release Drug Delivery Systems, Marcel Dekker, 571-587 (2003)
- Müller, R.H., Schmidt, S., Buttle, I., Akkar, A., Schmitt, J., Bromer, S., SolEmuls-novel technology for the formulation of i.v. emulsions with poorly soluble drugs, Int. J. Pharm. 269, 293-302 (2004).
- Müller, R.H., Souto, E., Göppert, T.M., Gohla, S., Production of biofunctionalized solid lipid nanoparticles (SLN[®]) for site-specific drug delivery, in: Kumar, C. (Hrsg.), Nanofabrication towards biomedical applications - Biofunctionalization of nanomaterials, (2005)
- Musser, J.B., Fontana, J.L., Mongan, P.D., The anesthetic and physiologic effects of an intravenous administration of a halothane lipid emulsion (5% vol/vol), Anesth. Analg. 88, 671-675 (1999).
- Mutschler, E., Arzneimittelwirkungen, Wissenschaftliche Verlagsgesellschaft, Stuttgart, (2001).
- Nielsen, B.L., Brown, L.R., The basis for colored silver-protein complex formation in stained polyacrylamide gels, Anal. Biochem. 141, 311-315 (1984).
- Norman, M.E., Williams, P., Illum, L., In vivo evaluation of protein adsorption to sterically stabilised colloidal carriers, J. Biomed. Mater. Res. 27, 861-866 (1993).
- Oakley, B.R., Kirsch, D.R., Morris, N.R., A simplified ultrasensitive silver stain for detecting proteins in polyacrylamide gels, Anal. Biochem. 105, 361-363 (1980).

- O'Farrel, P.H., High resolution two-dimensional electrophoresis of proteins, *J. Biol. Chem.* 250, 4007-4021 (1975).
- Ogawara, K., Furumoto, K., Nagayama, S., Minato, K., Higaki, K., Kai, T., Kimura, T., Pre-coating with serum albumin reduces receptor-mediated hepatic disposition of polystyrene nanosphere: implications for rational design of nanoparticles, *J Control Release* 100, 451-5 (2004).
- Olbrich, C., Bakowsky, U., Lehr, C.M., Müller, R.H., Kneuer, C., Cationic solid-lipid nanoparticles can efficiently bind and transfect plasmid DNA, *J. Control. Release* 77, 345-355 (2001).
- Oldendorf, W.H., Lipid solubility and drug penetration of the blood brain barrier, *Proc. Soc. Exp. Biol. Med.* 147, 813-815 (1974).
- Oleschuk, R.D., McComb, M.E., Chow, A., Ens, W., Standing, K.G., Perreault, H., Marois, Y., King, M., Characterization of plasma proteins adsorbed onto biomaterials. By MALDI-TOFMS, *Biomaterials* 21, 1701-1710 (2000).
- Olivier, J.C., Fenart, L., Chauvet, R., Pariat, C., Cecchelli, R., Couet, W., Indirect evidence that drug brain targeting using polysorbate 80-coated polybutylcyanoacrylate nanoparticles is related to toxicity, *Pharm. Res.* 16, 1836-42 (1999).
- Olson, A.D., Miller, M.J., Elsie 4: quantitative computer analysis of sets of two-dimensional gel electrophoretograms, *Anal. Biochem.* 169, 49-70 (1988).
- O'Mullane, J.E., Artursson, P., Tomlinson, E., Biopharmaceutics of microparticulate drug carriers, *Ann. NY Acad. Sci.* 507, 120-140 (1987).
- Pan, G., Liss, P.S., Metastable-Equilibrium Adsorption Theory I. Theoretical, *J. Coll. Interf. Sci.* 201, 71-76 (1998a).
- Pan, G., Liss, P.S., Metastable-Equilibrium Adsorption Theory II. Experimental, *J. Coll. Interf. Sci.* 201, 77-85 (1998b).
- Pardridge, W.M., Eisenberg, J., Yang, J., Human blood-brain barrier insulin receptor, *J. Neurochem.* 44, 1771-8 (1985).
- Pardridge, W.M., Triguero, D., Buciak, J., Yang, J., Evaluation of cationized rat albumin as a potential blood-brain barrier drug transport vector, *J. Pharmacol. Exp. Ther.* 255, 893-899 (1990).
- Pardridge, W.M., Kang, Y.S., Buciak, J.L., Yang, J., Human insulin receptor monoclonal antibody undergoes high affinity binding to human brain capillaries in vitro and rapid transcytosis through the blood-brain barrier in vivo in the primate, *Pharm. Res.* 12, 807-816 (1995).
- Pardridge, W.M., Physiologic-based strategies for protein drug delivery to the brain, *J. Contr. Rel.* 39, 281-286 (1996).

- Pardridge, W.M., Drug and gene targeting to the brain with molecular Trojan horses, Nat. Rev. Drug Discov. 1, 131-139 (2002).
- Pardridge, W.M., Blood-brain barrier drug targeting: the future of brain drug development, Mol. Interv. 3, 90-105, 51 (2003).
- Park, Y.J., Nah, S.H., Lee, J.Y., Jeong, J.M., Chung, J.K., Lee, M.C., Yang, V.C., Lee, S.J., Surface-modified poly(lactide-co-glycolide) nanospheres for targeted bone imaging with enhanced labeling and delivery of radioisotope, J Biomed Mater Res A 67, 751-60 (2003).
- Patel, H.M., Serum opsonins and liposomes: their interaction and opsonophagocytosis, Crit. Rev. Ther. Drug Carrier Syst. 9, 39-90 (1992).
- Patravale, V.B., Ambarkhane, A.V., Study of solid lipid nanoparticles with respect to particle size distribution and drug loading, Pharmazie 58, 392-395 (2003).
- Paulke, B.-R., Härtig, W., Brückner, G., Synthesis of nanoparticles for brain cell labelling in vivo, Acta Polymerica 43, 288-291 (1992).
- Peracchia, M.T., Harnisch, S., Pinto-Alphandary, H., Gulik, A., Dedieu, J.C., Desmaele, D., d'Angelo, J., Müller, R.H., Couvreur, P., Visualization of in vitro protein-rejecting properties of PEGylated stealth polycyanoacrylate nanoparticles, Biomaterials 20, 1269-1275 (1999).
- Petri, B., Dissertation in Vorbereitung, Freie Universität Berlin, Berlin (2005).
- Pitas, R.E., Boyles, J.K., Lee, S.H., Hui, D., Weisgraber, K.H., Lipoproteins and their receptors in the central nervous system. Characterization of the lipoproteins in cerebrospinal fluid and identification of apolipoprotein B,E(LDL) receptors in the brain, J. Biol. Chem. 262, 14352-14360 (1987).
- Poehling, H.-M., Neuhof, V., Visualization of proteins with a silver "stain": a critical analysis, Electrophoresis 2, 141-147 (1981).
- Pommier, C.G., O'Shea, J., Chused, T., Takahashi, T., Ochoa, M., Nutman, T.B., Bianco, C., Brown, E.J., Differentiation stimuli induce receptors for plasma fibronectin on the human myelomonocytic cell line HL-60, Blood 64, 858-866 (1984).
- Porter, C.J., Moghimi, S.M., Illum, L., Davis, S.S., The polyoxyethylene/polyoxypropylene block co-polymer poloxamer-407 selectively redirects intravenously injected microspheres to sinusoidal endothelial cells of rabbit bone marrow, FEBS Lett 305, 62-66 (1992).
- Price, M.E., Cornelius, R.M., Brash, J.L., Protein adsorption to polyethylene glycol modified liposomes from fibrinogen solution and from plasma, Biochim. Biophys. Acta 1512, 191-205 (2001).
- Prokai, L., Prokai-Tatrai, K., Bodor, N., Targeting drugs to the brain by redox chemical delivery systems, Med. Res. Rev. 20, 367-416 (2000).

- Rabilloud, T., Mechanisms of protein silver staining in polyacrylamide gels: a 10-year synthesis, *Electrophoresis* 11, 785-794 (1990).
- Rabilloud, T., A comparison between low background silver diammine and silver nitrate protein stains, *Electrophoresis* 13, 429-439 (1992).
- Radtke, M., Müller, R.H., NLC® – Nanostructured lipid carriers: the new generation of lipid drug carriers, *NewDrugs* 2, 48-52 (2001).
- Rall, S.C., Jr., Weisgraber, K.H., Innerarity, T.L., Mahley, R.W., Structural basis for receptor binding heterogeneity of apolipoprotein E from type III hyperlipoproteinemic subjects, *Proc. Natl. Acad. Sci. U S A* 79, 4696-4700 (1982).
- Raman, B., Cheung, A., Marten, M.R., Quantitative comparison and evaluation of two commercially available, two-dimensional electrophoresis image analysis software packages, Z3 and Melanie, *Electrophoresis* 23, 2194-2202 (2002).
- Rapoport, S.I., Osmotic opening of the blood-brain barrier, *Ann. Neurol.* 24, 677-684 (1988).
- Rege, B.D., Kao, J.P., Polli, J.E., Effects of nonionic surfactants on membrane transporters in Caco-2 cell monolayers, *Eur. J. Pharm. Sci.* 16, 237-246 (2002).
- Reinhard, E., Pharmazeutische Biologie, Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart, (1990).
- Reyle-Hahn, M., Rossaint, R., Xenon--a new anesthetic, *Anaesthesist* 49, 869-874 (2000).
- Righetti, P.G., Drysdale, J.W., Small-scale fractionation of proteins and nucleic acids by isoelectric focusing in polyacrylamide gels, *Ann. N Y Acad. Sci.* 209, 163-187 (1973).
- Rimpler, H., Lipide, in: Rimpler, H. (Hrsg.), *Biogene Arzneistoffe*, Deutscher Apotheker Verlag, Stuttgart, 137-195 (1999)
- Rossaint, R., Reyle-Hahn, M., Schulte Am Esch, J., Scholz, J., Scherpereel, P., Vallet, B., Giunta, F., Del Turco, M., Erdmann, W., Tenbrinck, R., Hammerle, A.F., Nagele, P., Multicenter randomized comparison of the efficacy and safety of xenon and isoflurane in patients undergoing elective surgery, *Anesthesiology* 98, 6-13 (2003).
- Rosseneu, M., Structure and Function of Apolipoproteins, CRC Press, Boca Raton, (1992).
- RoteListe®, Arzneimittelverzeichnis für Deutschland, Rote Liste Service GmbH, Frankfurt/Main, Editio Cantor Verlag, Aulendorf, (2005).

- Rousselle, C., Clair, P., Lefauconnier, J.M., Kaczorek, M., Scherrmann, J.M., Temsamani, J., New advances in the transport of doxorubicin through the blood-brain barrier by a peptide vector-mediated strategy, *Mol. Pharmacol.* 57, 679-686 (2000).
- Rousselle, C., Smirnova, M., Clair, P., Lefauconnier, J.M., Chavanieu, A., Calas, B., Scherrmann, J.M., Temsamani, J., Enhanced delivery of doxorubicin into the brain via a peptide-vector-mediated strategy: saturation kinetics and specificity, *J. Pharmacol. Exp. Ther.* 296, 124-131 (2001).
- Rowinsky, E.K., Onetto, N., Canetta, R.M., Arbuck, S.G., Taxol: the first of the taxanes, an important new class of antitumor agents, *Semin. Oncol.* 19, 646-662 (1992).
- Saito, H., Okuhira, K., Tsuchimoto, N., Vertut-Doi, A., Matsumoto, C., Tanimoto, T., Okada, S., Handa, T., Modulation of apolipoprotein E-mediated plasma clearance and cell uptake of emulsion particles by cholestryl ester, *Lipids* 36, 27-33 (2001).
- Salminen, E., Kankuri, M., Nuutila, J., Lilius, E.M., Pellimiemi, T.T., Modulation of IgG and complement receptor expression of phagocytes in kidney cancer patients during treatment with interferon-alpha, *Anticancer Res.* 21, 2049-2055 (2001).
- Sanovich, E., Bartus, R.T., Friden, P.M., Dean, R.L., Le, H.Q., Brightman, M.W., Pathway across blood-brain barrier opened by the bradykinin agonist, RMP-7, *Brain Res.* 705, 125-35 (1995).
- Sasol, Produktinformation DYNASANE. (2003).
- Sauer, I., Dunay, I., Weisgraber, K.H., Bienert, M., Dathe, M., An Apolipoprotein E-derived peptide mediates uptake of sterically stabilized liposomes into brain capillary endothelial cells, submitted (2004).
- Scanu, A.M., Plasma apolipoproteins: Gene structure, function, and variants, in: Putman, F. W. (Hrsg.), *The Plasma Proteins. Structure, Function, and Genetic Control*, Vol. V. Academic Press, Inc., New York, 141-190 (1987)
- Schechter, G., CNS diseases congress: advances in therapeutics, tools and trials. 28-29 June 2004, Strategic Research Institute, Philadelphia, PA, USA, *Expert Rev. Neurother.* 4, 747-9 (2004).
- Scherer, D., Einfluß von Polybutylcyanoacrylat-Nanopartikeln auf die orale Adsorption von Arzneistoffen, Dissertation, Johann Wolfgang Goethe-Universität, Frankfurt am Main (1992).
- Schmidt, S., Gessner, A., Müller, R.H., Comparison of plasma protein patterns on emulsions for parenteral nutrition dependent on their age. *Intern. Symp. Control. Rel. Bioact. Mater.* 28 480-481 (2001).
- Schmidt, S., Parenterale O/W-Emulsionen: Plasmaproteininteraktion und Inkorporation von Arzneistoffen, Dissertation, Freie Universität Berlin (2002).

- Schöler, N., Zimmermann, E., Katzfey, U., Hahn, H., Müller, R.H., Liesenfeld, O., Preserved solid lipid nanoparticles (SLN) at low concentrations do cause neither direct nor indirect cytotoxic effects in peritoneal macrophages, *Int. J. Pharm.* 196, 235-239 (2000).
- Schöler, N., Krause, K., Kayser, O., Müller, R.H., Borner, K., Hahn, H., Liesenfeld, O., Atovaquone nanosuspensions show excellent therapeutic effect in a new murine model of reactivated toxoplasmosis, *Antimicrob. Agents Chemother.* 45, 1771-1779 (2001).
- Schröder, U., Sabel, B.A., Nanoparticles, a drug carrier system to pass the blood-brain barrier, permit central analgesic effects of i.v. dalargin injections, *Brain Res.* 710, 121-124 (1996).
- Schuhmann, R., Physikalische Stabilität parenteraler FettEmulsionen, Dissertation, Freie Universität Berlin (1995).
- Schwarz, C., Mehnert, W., Lucks, J.S., Müller, R.H., Solid lipid nanoparticles (SLN) for controlled drug delivery. I. Production, characterization and sterilization, *J. Contr. Rel.* 30, 83-96 (1994).
- Schwarz, C., Feste Lipid Nanopartikel: Herstellung, Charakterisierung, Arzneistoffinkorporation und -freisetzung, Dissertation, Freie Universität Berlin (1995).
- Schwarze, S.R., Ho, A., Vocero-Akbani, A., Dowdy, S.F., In vivo protein transduction: delivery of a biologically active protein into the mouse, *Science* 285, 1569-1572 (1999).
- Schwick, H.G., Heide, K., Immunochemistry and immunology of collagen and gelatin, *Bibl. Haematol.* 33, 111-125 (1969).
- Seelig, A., A general pattern for substrate recognition by P-glycoprotein, *Eur. J. Biochem.* 251, 252-261 (1998).
- Segrest, J.P., Garber, D.W., Brouillette, C.G., Harvey, S.C., Anantharamaiah, G.M., The amphipathic alpha helix: a multifunctional structural motif in plasma apolipoproteins, *Adv. Protein Chem.* 45, 303-369 (1994).
- Shapiro, A.L., Vinuela, E., Maizel, J.V., Jr., Molecular weight estimation of polypeptide chains by electrophoresis in SDS-polyacrylamide gels, *Biochem. Biophys. Res. Commun.* 28, 815-820 (1967).
- Shen, B.W., Lipid-protein interaction at solid-water interface. Adsorption of human apo-high density lipoprotein to amphiphilic interfaces, *J. Biol. Chem.* 260, 1032-1039 (1985).

- Shrivastava, P., Hanibuchi, M., Yano, S., Parajuli, P., Tsuruo, T., Sone, S., Circumvention of multidrug resistance by a quinoline derivative, MS-209, in multidrug-resistant human small-cell lung cancer cells and its synergistic interaction with cyclosporin A or verapamil, *Cancer Chemother. Pharmacol.* 42, 483-490 (1998).
- Siekmann, B., Westesen, K., Investigations on solid lipid nanoparticles prepared by precipitation in o/w emulsions, *Eur. J. Pharm. Biopharm.* 43, 104-109 (1996).
- Skehel, J.M., Schneider, K., Murphy, N., Graham, A., Benson, G.M., Cutler, P., Camilleri, P., Phenotyping apolipoprotein E*3-leiden transgenic mice by two-dimensional polyacrylamide gel electrophoresis and mass spectrometric identification, *Electrophoresis* 21, 2540-2545 (2000).
- Smith, A., Hunneyball, I.M., Evaluation of poly (lactic acid) as a biodegradable drug delivery system for parenteral administration, *Int. J. Pharm.* 30, 215-220 (1986).
- Smith, P.K., Krohn, R.I., Hermanson, G.T., Mallia, A.K., Gartner, F.H., Provenzano, M.D., Fujimoto, E.K., Goeke, N.M., Olson, B.J., Klenk, D.C., Measurement of protein using bicinchoninic acid, *Anal. Biochem.* 150, 76-85 (1985).
- Sokol, R.J., Butler-Simon, N.A., Bettis, D., Smith, D.J., Silverman, A., Tocopheryl polyethylene glycol 1000 succinate therapy for vitamin E deficiency during chronic childhood cholestasis: neurologic outcome, *J. Pediatr.* 111, 830-836 (1987).
- Sokol, R.J., Johnson, K.E., Karrer, F.M., Narkewicz, M.R., Smith, D., Kam, I., Improvement of cyclosporin absorption in children after liver transplantation by means of water-soluble vitamin E, *Lancet* 338, 212-214 (1991).
- Speiser, P.P., Nanoparticles and liposomes: a state of the art, *Methods Find Exp Clin Pharmacol* 13, 337-42 (1991).
- Speiser, P.P., Nanopartikel, in: Müller, R. H., Hildebrand, G. E. (Hrsg.), *Pharmazeutische Technologie: Moderne Arzneiformen, Wissenschaftliche Verlagsgesellschaft, Stuttgart*, (1998)
- Steer, C.J., Klausner, R.D., Clathrin-coated pits and coated vesicles: functional and structural studies, *Hepatology* 3, 437-454 (1983).
- Steiniger, S.C., Kreuter, J., Khalansky, A.S., Skidan, I.N., Bobruskin, A.I., Smirnova, Z.S., Severin, S.E., Uhl, R., Kock, M., Geiger, K.D., Gelperina, S.E., Chemotherapy of glioblastoma in rats using doxorubicin-loaded nanoparticles, *Int. J. Cancer* 109, 759-767 (2004).
- Stich, T.M., Determination of protein covalently bound to agarose supports using bicinchoninic acid, *Anal. Biochem.* 191, 343-346 (1990).
- Stolnik, S., Illum, L., Davis, S.S., Long circulating microparticle drug carriers, *Adv. Drug Del. Rev.* 16, 195-214 (1995).

- Stroschneider, E., Margreiter, J., Pfeiffer, K.P., Eggenreich, U., Balogh, D., Comparative study of propofol emulsions for anesthesia in varicose vein operations, *Anesthesiol. Intensivmed. Notfallmed. Schmerzther.* 33, 781-785 (1998).
- Tabatt, K., Kneuer, C., Sameti, M., Olbrich, C., Muller, R.H., Lehr, C.M., Bakowsky, U., Transfection with different colloidal systems: comparison of solid lipid nanoparticles and liposomes, *J. Control. Release* 97, 321-332 (2004).
- Tadros, T.F., Vincent, B., Influence of temperature and electrolytes on the adsorption behaviour of poly(ethylene oxide)-poly(propylene oxide) block copolymer on polystyrene latex and the stability of the polymer coated particles, *J. Phys. Chem.* 84, 1575 (1980).
- Tan, J.S., Butterfield, D.E., Voycheck, C.L., Caldwell, K.D., Li, J.T., Surface modification of nanoparticles by PEO/PPO block copolymers to minimize interactions with blood components and prolong blood circulation in rats, *Biomaterials* 14, 823-833 (1993).
- Thiebaut, F., Tsuruo, T., Hamada, H., Gottesman, M.M., Pastan, I., Willingham, M.C., Cellular localization of the multidrug-resistance gene product P-glycoprotein in normal human tissues, *Proc. Natl. Acad. Sci. U S A* 84, 7735-7738 (1987).
- Thiele, L., Diederichs, J.E., Reszka, R., Merkle, H.P., Walter, E., Competitive adsorption of serum proteins at microparticles affects phagocytosis by dendritic cells, *Biomaterials* 24, 1409-1418 (2003).
- Thode, K., Spezifische Kontrastmittel für die Magnetresonanz-Tomographie: Physikochemische Charakterisierung und Studien zur Plasmaproteinadsorption, Dissertation, Freie Universität Berlin (1996).
- Thode, K., Müller, R.H., Theory and Set Up of Hydropobic Interaction Chromatography (HIC), in: Müller, R. H., Mehnert, W. (Hrsg.), *Particle and Surface Characterisation Methods*, Wissenschaftliche Verlagsgesellschaft Stuttgart, 229-234 (1997)
- Tice, T.R., Gilley, R.M., Preparation of injectable controlled-release microcapsules by a solvent-evaporation process, *J. Contr. Rel.* 2, 343 (1985).
- Tracy, R.P., Currie, R.M., Young, D.S., Two-dimensional gel electrophoresis of serum specimens from a normal population, *Clin. Chem.* 28, 890-899 (1982).
- Tröster, S.D., Kreuter, J., Contact angle of surfactants with a potential to alter the body distribution of colloidal drug carriers on poly (methyl methacrylate) surfaces, *Int. J. Pharm.* 45, 91-100 (1988).
- Tröster, S.D., Müller, U., Kreuter, J., Modification of the body distribution of poly(methyl methacrylate) nanoparticles in rats by coating with surfactants, *Int. J. Pharm.* 61, 85-100 (1990).

- Truong-Le, V.L., August, J.T., Leong, K.W., Controlled gene delivery by DNA-gelatin nanospheres, *Hum. Gene Ther.* 9, 1709-1717 (1998).
- van Gelder, W., Cleton-Soeteman, M.I., Huijskes-Heins, M.I., van Run, P.R., van Eijk, H.G., Transcytosis of 6.6-nm gold-labeled transferrin: an ultrastructural study in cultured porcine blood-brain barrier endothelial cells, *Brain Res.* 746, 105-116 (1997).
- Van Leuven, F., Marynen, P., Sottrup-Jensen, L., Cassiman, J.J., Van den Berghe, H., The receptor-binding domain of human alpha 2-macroglobulin. Isolation after limited proteolysis with a bacterial proteinase, *J. Biol. Chem.* 261, 11369-11373 (1986).
- van Raamsdonk, J.M., Cornelius, R.M., Brash, J.L., Chang, P.L., Deterioration of polyamino acid-coated alginate microcapsules in vivo, *J. Biomater. Sci. Polym. Ed.* 13, 863-884 (2002).
- van Zuylen, L., Verweij, J., Sparreboom, A., Role of formulation vehicles in taxane pharmacology, *Invest. New Drugs* 19, 125-141 (2001).
- Vandorpe, J., Schacht, E., Dunn, S., Hawley, A., Stolnik, S., Davis, S.S., Garnett, M.C., Davies, M.C., Illum, L., Long circulating biodegradable poly(phosphazene) nanoparticles surface modified with poly(phosphazene)-poly(ethylene oxide) copolymer, *Biomaterials* 18, 1147-1152 (1997).
- Vasko, K.A., Riley, A.M., DeWall, R.A., Poloxalkol R (pluronic F-68): a priming solution for cardiopulmonary bypass, *Trans. Am. Soc. Artif. Intern. Organs* 18, 526-531, 545 (1972).
- Verrecchia, T., Spenlehauer, G., Bazile, D.V., Murry-Brelier, A., Archimbaud, Y., Veillard, M., Non-stealth (poly(lactic acid/albumin)) and stealth (poly(lactic acid-polyethylene glycol)) nanoparticles as injectable drug carriers, *J. Contr. Rel.* 36, 49-61 (1995).
- Vogel, W., Bomford, A., Young, S., Williams, R., Heterogeneous distribution of transferrin receptors on parenchymal and nonparenchymal liver cells: biochemical and morphological evidence, *Blood* 69, 264-270 (1987).
- Vroman, L., Adams, A.L., Fischer, G.C., Munoz, P.C., Interaction of high molecular weight kininogen, factor XII, and fibrinogen in plasma at interfaces, *Blood* 55, 156-159 (1980).
- Vroman, L., Adams, A.L., Adsorption of proteins out of plasma and solutions in narrow spaces, *J. Coll. Interf. Sci.* 111, 391-402 (1986).
- Wang, J.X., Sun, X., Zhang, Z.R., Enhanced brain targeting by synthesis of 3',5'-dioctanoyl-5-fluoro-2'-deoxyuridine and incorporation into solid lipid nanoparticles, *Eur. J. Pharm. Biopharm.* 54, 285-290 (2002).
- Weber, C., Coester, C., Kreuter, J., Langer, K., Desolvation process and surface characterisation of protein nanoparticles, *Int. J. Pharm.* 194, 91-102 (2000).

- Weigt, H., Georgieff, M., Beyer, C., Georgieff, E.M., Kuhse, J., Fohr, K.J., Lipid emulsions reduce NMDA-evoked currents, *Neuropharmacology* 47, 373-380 (2004).
- Weigt, H.U., Georgieff, M., Beyer, C., Wachter, U., Fohr, K.J., Xenon incorporated in a lipid emulsion inhibits NMDA receptor channels, *Acta Anaesthesiol Scand* 47, 1119-1124 (2003).
- Weisgraber, K.H., Innerarity, T.L., Harder, K.J., Mahley, R.W., Milne, R.W., Marcel, Y.L., Sparrow, J.T., The receptor-binding domain of human apolipoprotein E. Monoclonal antibody inhibition of binding, *J. Biol. Chem.* 258, 12348-12354 (1983).
- Weisgraber, K.H., Mahley, R.W., Kowal, R.C., Herz, J., Goldstein, J.L., Brown, M.S., Apolipoprotein C-I modulates the interaction of apolipoprotein E with beta-migrating very low density lipoproteins (beta-VLDL) and inhibits binding of beta-VLDL to low density lipoprotein receptor-related protein, *J. Biol. Chem.* 265, 22453-22459 (1990).
- Weisgraber, K.H., Apolipoprotein E: structure-function relationships, *Adv. Protein Chem.* 45, 249-302 (1994).
- Wesemeyer, H., Müller, B.W., Müller, R.H., Adsorption of ethoxylated surfactants on nanoparticles. II. Determination of adsorption enthalpy by microcalorimetry, *Int. J. Pharm.* 89, 33-40 (1993).
- Westermeier, R., Elektrophorese-Praktikum, VCH Verlagsgesellschaft, Weinheim, (1990).
- Westesen, K., Bunjes, H., Do nanoparticles prepared from lipids solid at room temperature always possess a solid matrix?, *Int. J. Pharm.* 115, 129-131 (1995).
- Wetterau, J.R., Aggerbeck, L.P., Rall, S.C., Jr., Weisgraber, K.H., Human apolipoprotein E3 in aqueous solution. I. Evidence for two structural domains, *J. Biol. Chem.* 263, 6240-6248 (1988).
- Weyhers, H., Feste Lipid Nanopartikel (SLN) für die gewebsspezifische Arzneistoffapplikation, Dissertation, Freie Universität Berlin (1995).
- Weyhers, H., Ehlers, S., Mehnert, W., Hahn, H., Müller, R.H., Solid Lipid Nanoparticles - Determination of in vivo toxicity. *Proc. 1st World Meeting APGI/APV* 489-490 (Budapest, 1995).
- Wilkens, D.J., Meyers, P.A., Studies on the relationship between the electrophoretic properties of colloids and their blood clearance and organ distribution in the rat, *Brit. J. Exp. Path.* 47, 568-576 (1966).
- Wissing, S.A., Kayser, O., Müller, R.H., Solid lipid nanoparticles for parenteral drug delivery, *Adv. Drug Deliv. Rev.* 56, 1257-1272 (2004).

- Wojciechowski, P., Brash, J.L., The Vroman effect in tube geometry: the influence of flow on protein adsorption measurements, *J. Biomater. Sci. Polym. Ed.* 2, 203-16 (1991).
- Wretlind, A., Development of fat emulsions, *JPEN J. Parenter. Enteral. Nutr.* 5, 230-235 (1981).
- Wurm, H., beta 2-Glycoprotein-I (apolipoprotein H) interactions with phospholipid vesicles, *Int. J. Biochem.* 16, 511-515 (1984).
- Yamazaki, A., Winnik, F.M., Cornelius, R.M., Brash, J.L., Modification of liposomes with N-substituted polyacrylamides: identification of proteins adsorbed from plasma, *Biochim. Biophys. Acta* 1421, 103-115 (1999).
- Yang, S.C., Lu, L.F., Cai, Y., Zhu, J.B., Liang, B.W., Yang, C.Z., Body distribution in mice of intravenously injected camptothecin solid lipid nanoparticles and targeting effect on brain, *J. Control. Release* 59, 299-307 (1999).
- Zalipsky, S., Hansen, C.B., Lopes de Menezes, D.E., Allen, T.M., Long-circulating, polyethylene-glycol-grafted immunoliposomes, *J. Contr. Rel.* 39, 153-161 (1996).
- Zara, G.P., Cavalli, R., Bargoni, A., Fundaro, A., Vighetto, D., Gasco, M.R., Intravenous administration to rabbits of non-stealth and stealth doxorubicin-loaded solid lipid nanoparticles at increasing concentrations of stealth agent: pharmacokinetics and distribution of doxorubicin in brain and other tissues, *J. Drug Target.* 10, 327-335 (2002).
- Zhang, F., Kang, E.T., Neoh, K.G., Wang, P., Tan, K.L., Surface modification of stainless steel by grafting of poly(ethylene glycol) for reduction in protein adsorption, *Biomaterials* 22, 1541-8 (2001).
- Zillies, J.C., Ludwig-Maximilians-Universität, München, Persönliche Mitteilung, Juli. (2005).
- Zillies, J.C., Dissertation in Vorbereitung, Ludwig-Maximilians-Universität, München (2006).
- Zimmermann, E., Feste Lipidnanopartikel (SLN) als kolloidaler Arzneistoffträger für die orale und intravenöse Applikation eines chemisch instabilen Wirkstoffs, Dissertation, Freie Universität Berlin (2001).
- Zimmermann, I., Possibilities and limitations of laser light scattering techniques for particle size analysis. *Colloidal Drug Carriers - 1st Expert Meeting* (1995).
- zur Mühlen, A., Feste Lipidnanopartikel mit prolongierter Wirkstoffliberation: Herstellung, Langzeitstabilität, Charakterisierung, Freisetzungsvorhalten und -mechanismen, Dissertation, Freie Universität Berlin (1996).

zur Mühlen, A., Schwarz, C., Mehnert, W., Solid lipid nanoparticles (SLN) for controlled drug delivery--drug release and release mechanism, Eur. J. Pharm. Biopharm. 45, 149-155 (1998).

Zwiorek, K., Kloeckner, J., Wagner, E., Coester, C., Gelatin nanoparticles as a new and simple gene delivery system, J. Pharm. Pharm. Sci. 7, 22-28 (2005).